

digoxin into guinea pigs increases the K content of the ventricles and serum. Toxic doses produce a further rise in serum K whereas the amount of K in the ventricles does not differ significantly from that of untreated animals.

Intravenous infusion into guinea pigs of KCl solution (8.4–134 mM) simultaneously with digoxin (56 µg/ml) fails to influence the lethal dose of digoxin, but with 268 mM solution of the salt the lethal dose of digoxin is decreased ten-fold.

The time of onset of the positive inotropic response of the isolated right ventricle of the guinea pig, stimulated electrically, to a toxic dose of digoxin is not influenced when the KCl concentration of the bathing fluid is raised from 5.6 to 13.5 mM.

The rate of increase of the amplitude of contraction is markedly reduced but the maximum attained is unaltered and the life of the ventricle is greatly prolonged.

The duration of the therapeutic action of digoxin in the failing heart-lung preparation of the guinea pig is increased if the K content of the circulating fluid is increased from 5.6 to 6.1 mM, but considerably decreased when it is raised to 12.1 mM.

These data suggest that: (1) the extracardiac tissues are more sensitive than the myocardium to the action of digoxin on K flux, and (2) that K, depending on its concentration in the heart, can increase or decrease the cardiotoxicity of digoxin.

45 Pharmacology of Chlordiazepoxide (Librium) and Analogues. L. O. RANDALL and B. KAPPELL (U.S.A.).

Chlordiazepoxide (Librium) is the first member of a new chemical class of compounds which has shown unique taming effects in animals⁽¹⁾ and powerful anti-anxiety effects in human subjects.⁽²⁾ Other pharmacological effects first observed in animals and later confirmed in human subjects include anticonvulsant activity and muscle relaxant effects at low doses, with sedative effects and appetite-stimulating effects at high doses. Certain pharmacological effects which were minimal in animal tests were also minimal in human subjects including effects on blood pressure, heart rate and autonomic nervous system.

Analogues of chlordiazepoxide (7-chloro-2-methylamino-5-phenyl-3H-1:4-benzodiazepine 4-oxide hydrochloride) showed a similar type of activity in muscle relaxant and anticonvulsant tests. Screening tests include the inclined screen test in mice, the fighting mouse test,⁽³⁾ the anti-strychnine test and the gross behaviour test in cats. Benzodiazepines having activity of interest include 7-chloro-1-methyl-5-phenyl-3H-1:4-benzodiazepin-2(1H)-one; 7-chloro-2-oxo-5-phenyl-1:2-dihydro-3H-1:4-benzodiazepine 4-oxide; 5-phenyl-7-trifluoro-methyl-3H-1:4-benzodiazepin-2(1H)-one; 7-chloro-2-oxo-5-phenyl-1:2-dihydro-3H-1:4-benzodiazepine; 7-chloro-5-(2-chlorophenyl)-3H-1:4-benzodiazepin-2(1H)-one and 7-nitro-5-phenyl-3H-1:4-benzodiazepin-2(1H)-one.

1. RANDALL, L. O., SCHALLEK, W., HEISE, G. A., KEITH, E. F. and BAGDON, R. E. (1960), *J. Pharmacol.*, **129**, 163.
2. TOBIN, J. M. and LEWIS, N. D. C. (1960), *J. Amer. med. Ass.*, **174**, 96.
3. TEDESCHI, R. C., TEDESCHI, D. H., MUCHA, A., COOK, L., MATTIS, P. A. and FELLOWS, C. J. (1958), *J. Pharmacol.*, **125**, 28.

46 The Effects of Ethanol and Chlordiazepoxide in Altering Autonomic Responses Evoked by Isocortical and Paleocortical Stimulation. M. N. CARROLL, JR., E. C. HOFF, J. F. KELL, JR. and C. G. SUTER (U.S.A.).

Unipolar stimulation of the anterior sigmoid gyrus (I), limbic lobe (II), hypothalamus (III), N. Von Bechterew (IV), amygdala (V), and central gray (VI) loci in flaxedilized cats evoked pressor responses, cardiac arrhythmias (nodal and ventricular extrasystoles and ventricular tachycardia with indications of coronary insufficiency), mydriasis and striking inhibition of mobility of the pyloric sphincter and ileum. Salivation was elicited from subcortical loci II, III, IV and VI, and retraction of the nictitating membrane was noted from I, III and VI. Lacrimation was not observed from any of the areas studied.

The intravenous administration of chlordiazepoxide (10–20 mg/kg) elicited a transient hypotension and slight bradycardia whereas 10 per cent ethanol (3 ml/kg, i.v.) produced increased pyloric mobility and slight to marked alterations in the electrocardiogram. Chlordiazepoxide greatly attenuated pressor responses from all brain areas studied whereas ethanol exerted weak anti-pressor effects in areas I, II and strong blocking action against central stimulation of III. Centrally-induced cardiac arrhythmias were completely blocked by chlordiazepoxide in loci II and V whereas ethanol partially attenuated the response only from II. Neither compound showed any activity against centrally evoked salivation, mydriasis or gastrointestinal inhibition.

Inasmuch as chlordiazepoxide lacks the undesirable side effects of ethanol, its prophylactic potential against centrally-induced arrhythmias and pressor responses merits intensive investigation.

47 Effects of Chlordiazepoxide (Librium) and Other Psychopharmacological Agents on "Fixed" Behaviour in Rats. W. T. LIBERSON, A. KAFKA and E. SCHWARTZ (U.S.A.).

Effects of various therapies were investigated on fixation rats⁽¹⁾ with regard to: (1) motor coordination; (2) sensory and symbolic discrimination and speed of their acquisition; (3) reaction times to stimuli during approach, avoidance or escape behaviour; (4) variability of behaviour; (5) compulsive rigidity; and (6) variety of patterns

of learning. While forced guidance may, none of the tested drugs or electric shock therapy has "cured" rats once fixated behaviour became established. At high doses (1 mg/kg) LSD₂₅ blocks goal-directed behaviour. Chlorpromazine, reserpine and meprobamate suppress avoidance and reduce escape. In higher dosages, they also slow down learning and depress approach behaviour. Before establishment of fixation, i.e. during the training period, high doses of tranquilizers prevent fixation in 20–35 per cent of cases. Barbiturates in doses sufficient to affect avoidance interfere with motor behaviour. MOA inhibitors and related agents (iproniazid, amphetamine, Tofranil and Catron) quicken the approach reactions and, in high doses, increase the rate of avoidance behaviour, but do not significantly affect fixation. Chlordiazepoxide (Librium) differs from previous drugs. Although it induces docility, it shortens reaction time of both avoidance and approach to a greater degree than MAO inhibitors. The behaviour becomes very variable, and Librium given during training period prevents fixation in 50 per cent of the rats. Moreover, Librium fosters learning and the effects of guidance and reduces latency times, while most of these were inhibited by chlorpromazine. Thus, although Librium affects fixation prophylactically in greater percentage of cases than other drugs employed, it does not cure established fixation.

1. LIBERSON, FELDMAN and ALLEN (1959), *Neuropsychopharmac.*, **1**, 351; (1959), *J. Neuropsych.*, **1**, 17.

48 On Mechanism of the Antidepressant Action of Imipramine. F. SULSER and B. B. BRODIE (U.S.A.).

The potent antidepressant effect of the iminodibenzyl derivative imipramine is clinically established. Pharmacologically, the drug exerts weak chlorpromazine-like actions in normal animals and men. None of the screening procedures currently used would suggest this drug to be an antidepressant drug. The antidepressant action of imipramine becomes dramatically evident in studies which show that it prevents in rats and dogs the depression (chronic pretreatment) and increased central parasympathetic activity (single dose) evoked by reserpine or a synthetic benzoquinolizine (RO-4 1284). However, imipramine does not prevent the action of chlorpromazine. Studies of the physiological disposition of imipramine indicate that the drug acts through a metabolic product, monomethyl norimipramine. This compound is much more active in blocking reserpine or RO-4 1284; furthermore, it shows, in contrast to imipramine, no delay in its action. It not only prevents the action of reserpine or RO-4 1284 but actually reverses it, the animals displaying considerable hyperactivity. Neither imipramine nor its metabolite block monoamine oxidase; they do not

increase levels of brain serotonin and norepinephrine or prevent reserpine and RO-4 1284 from releasing brain amines. The possibility that imipramine or its metabolite reveal their antidepressant action by blocking free serotonin will be discussed.

49 Unusual Central Depressant Properties of a New Piperidine Compound. T. B. O'DELL, M. D. NAPOLI and J. H. MIRSKY (U.S.A.).

One approach to the study of the complexities of the central nervous system and its functions is through the use of new pharmacodynamic tools. Such tools might be developed by the investigation of properties of new compounds having novel or unusual chemical structures. Analysis of the responses obtained with novel compounds, coupled with known actions of standard drugs, could well provide new leads toward a better understanding of the functioning of the central nervous system.

A series of new piperidine derivatives has been investigated using a battery of standard pharmacological testing procedures. An unusual combination of effects on the central nervous system has been revealed from these studies. One compound, in particular, 4-(3-indolyloethyl)-1-phenethylpiperidine hydrochloride demonstrated strong analgesic activity comparable to that found with morphine, and could also produce a tranquilizing effect which resembled that produced by reserpine. These effects could be separated dose-wise with analgesic being apparent at doses below those required to induce the tranquilization responses. True sedative effects (i.e. sleep) could not be induced regardless of the magnitude of the dose administered. The duration of these responses, especially the tranquilizing action, was very prolonged and was a function of the dose employed. Thus, compared to the actions on the central nervous system of known drugs, this new compound appears to possess a unique combination of properties.

50 Influence of a New Group of Tranquillizers, Derivatives of Dibenzosuberene, on the Central Nervous System. Z. VOTAVA, J. METYŠOVÁ and M. SOUŠKOVÁ (Czechoslovakia).

In the group of 16 new dibenzosuberene and dibenzosuberene analogues of chlorprothixene, interesting correlations between the chemical structure and pharmacological effect on the central nervous system were found. Tests on prolongation of thiopental anaesthesia, rotating rod, depression of the body temperatures, anti-pentazol action and tentative conditioned reflexes revealed that the double bond in the middle ring of dibenzosuberene increased the tranquillizing activity. Further rise of the activity was produced by the introduction of chlorine in the position two of the cycle. The most promising drugs of this type were chloroheptadiene and chloroheptatriene.