

available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/biochempharm

Subject index Volume 72

ABC

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

Acacetin

Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293

accumulation

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

Acetaldehyde

Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616

Acetylcholinesterase

Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344

Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358

Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597

Actin cytoskeleton

STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32

Actin polymerization

8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells, 541

Activators

TPA-induced up-regulation of activator protein-1 can be inhibited or enhanced by analogs of the natural product curcumin, 928

Acute lung injury

Effective attenuation of acute lung injury in vivo and the formyl peptide-induced neutrophil activation in vitro by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

Addiction

Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919

Adenosine

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Adhesion molecules GPIIb

Identification and characterization of triosephosphate isomerase that specifically interacts with the integrin α IIb cytoplasmic domain, 551

Adipogenesis

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

Adiponectin

Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616

Adiposity

Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity in vivo, 42

ADP-induced platelet aggregation

Protein S-glutathionylation and platelet anti-aggregating activity of disulfiram, 608

Adrenoreceptors

Tyramine in the assessment of regional adrenergic function, 1724

AG490

Suppression of IL-1 β expression by the Jak 2 inhibitor AG490 in cerulein-stimulated pancreatic acinar cells, 1555

Agranulocytosis

Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential in vitro system for the study of agranulocytosis, 783

AH receptor

Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267

Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention, 795

Akt

Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423

Alkylating agents

Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11

Alkylating drugs

Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432

Allodynia

Anti-allodynic property of flavonoid myricitrin in models of persistent inflammatory and neuropathic pain in mice, 1707

Allosteric

Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor₁ (CRF₁) receptor, 244

Allyl isothiocyanate

Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104

Alternative exon splicing

Tissue specific glucocorticoid receptor expression, a role for alternative first exon usage?, 1529

Alternative pathway

The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161

AMD3100

Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588

AMP-activated protein kinase

Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91

Analogues

Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulinotropic polypeptide, 719

Androgen receptor

Epigenetic regulation of nuclear steroid receptors, 1589

Andrographolide

Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132

Angiotensin

Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176

Animal models

Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197

Antagonist

Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588

Anti-angiogenesis

The angiosuppressive effects of 20(R)- ginsenoside Rg₃, 437

Anticancer

Synthesis of microtubule-interfering halogenated nescapine analogs that perturb mitosis in cancer cells followed by cell death, 415

Anti-cholinergic

Synergistic effects of the anti-cholinergic R,R-glycopyrrolate with anti-inflammatory drugs, 1690

Antidepressant

Pharmacological characterization of recombinant N-type calcium channel (Ca_v2.2) mediated calcium mobilization using FLIPR, 770

Anti-inflammatory and anti-tumor activity

Inhibition of NF- κ B activation through targeting I κ B kinase by celastrol, a quinone methide triterpenoid, 1311

Antimalarial agents

Dual-acting diamine antiparasmodial and chloroquine resistance modulating agents, 156

Antimycotic

Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463

Antioxidant

TPA-induced up-regulation of activator protein-1 can be inhibited or enhanced by analogs of the natural product curcumin, 928

Anti-allodynic property of flavonoid myricitrin in models of persistent inflammatory and neuropathic pain in mice, 1707

Antiproliferative activity

Mitochondrial permeability transition induced by novel pyridothiopyranopyrimidine derivatives: Potential new antimitochondrial antitumour agents, 1657

Antiproliferative effect

Cellular pharmacokinetics and pharmacodynamics of dipyridamole in vascular smooth muscle cells, 956

Anti-tumor activity

Distinct functions of IRF-3 and IRF-7 in IFN- α gene regulation and control of anti-tumor activity in primary macrophages, 1469

Anti-tumor drugs

NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366

AP-1

TPA-induced up-regulation of activator protein-1 can be inhibited or enhanced by analogs of the natural product curcumin, 928

2',4',6'-Tris(methoxymethoxy) chalcone attenuates hepatic stellate cell proliferation by a heme oxygenase-dependent pathway, 1322

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Apigenin-induced-apoptosis

Apigenin-induced-apoptosis is mediated by the activation of PKC δ and caspases in leukemia cells, 681

Apoptosis

Dipeptidyl peptidase II and leukocyte cell death, 70

Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132

P-glycoprotein enhances TRAIL-triggered apoptosis in multidrug resistant cancer cells by interacting with the death receptor DR5, 293

In vitro and *in vivo* studies of a novel potential anticancer agent of isochaihulactone on human lung cancer A549 cells, 308

Apoptotic cell death induction and angiogenesis inhibition in large established medullary thyroid carcinoma xenografts by Ret inhibitor RPI-1, 405

Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486

Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530

Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616

Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671

Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761

The methionine salvage pathway compound 4-methylthio-2-oxobutanate causes apoptosis independent of down-regulation of ornithine decarboxylase, 806

Role of 5-lipoxygenase pathway in the regulation of RAW 264.7 macrophage proliferation, 1022

Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054

- Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090
- Regulation of death and survival in astrocytes by ADP activating P2Y₁ and P2Y₁₂ receptors, 1031
- NF- κ B in solid tumors, 1142
- Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197
- Synergistic induction of apoptosis by sulindac and arsenic trioxide in human lung cancer A549 cells via reactive oxygen species-dependent down-regulation of survivin, 1228
- Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382
- Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343
- Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423
- Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668
- R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697
- Arsenic trioxide**
- Synergistic induction of apoptosis by sulindac and arsenic trioxide in human lung cancer A549 cells via reactive oxygen species-dependent down-regulation of survivin, 1228
- Arsenite**
- Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761
- Aryl hydrocarbon receptor**
- Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267
- Ah receptor: Dioxin-mediated toxic responses as hints to deregulated physiologic functions, 393
- AS101**
- The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423
- ASBT**
- Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polyps, 729
- Ascorbate**
- Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671
- Asthma**
- Synergistic effects of the anti-cholinergic R,R-glycopyrrolate with anti-inflammatory drugs, 1690
- ATM**
- NF- κ B activation by double-strand breaks, 1132
- ATP synthase**
- Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911
- Autoimmunity**
- Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453
- Autonomic nervous system**
- Tyramine in the assessment of regional adrenergic function, 1724
- Autophagy**
- Dipeptidyl peptidase II and leukocyte cell death, 70
- 5-Aza-2'-deoxycytidine**
- Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577
- Azoxymethane**
- Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
- AZT**
- 3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239
- Baboon placenta**
- Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730
- Bacterial toxins**
- Listeria monocytogenes* induced Rac1-dependent signal transduction in endothelial cells, 1367
- Bax**
- Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132
- Bcl-2 family**
- Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132
- BCL-3**
- Deregulated NF- κ B activity in haematological malignancies, 1069
- Benzo[a]pyrene**
- Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343
- Benzyl isothiocyanate**
- Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217
- Bergerat fold**
- The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207
- Bid**
- Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132
- Binding**
- Recognition forces in ligand-protein complexes: Blending information from different sources, 1633
- Bioactivation**
- Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential *in vitro* system for the study of agranulocytosis, 783
- Bioavailability**
- Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042
- Bioreactor**
- Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358
- Blastogenesis**
- Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486

- Bone morphogenetic protein**
Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332
- Bone regeneration**
Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268
- Bone resorption**
Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184
- BORIS**
Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577
- Bovine aortic endothelial cells**
Crocetin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558
- Brain endothelial cells**
The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869
- Breast cancer**
Paradoxical effects of the phage display-derived peptide antagonist IGF-F1-1 on insulin-like growth factor-1 receptor signaling, 53
Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332
TGF- β in cancer and as a therapeutic target, 523
Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530
Human equilibrative nucleoside transporter-1 (hENT1) is required for the transcriptomic response of the nucleoside-derived drug 5'-DFUR in breast cancer MCF7 cells, 1646
- Breast cancer cells**
Novel amidine analogue of melphalan as a specific multi-functional inhibitor of growth and metabolism of human breast cancer cells, 320
- BRET**
Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355
- Bucillamine**
Bucillamine induces glutathione biosynthesis via activation of the transcription factor Nrf2, 455
- Budding**
Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- Bupleurum scorzonrifolium***
In vitro and *in vivo* studies of a novel potential anticancer agent of isochaihulactone on human lung cancer A549 cells, 308
- C-1305**
Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668
- Ca²⁺ mobilisation**
Diverse signalling by different chemokines through the chemokine receptor CCR5, 739
- Caco-2 cells**
Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974
- Calcium mobilization**
Pharmacological characterization of recombinant N-type calcium channel (Ca_v2.2) mediated calcium mobilization using FLIPR, 770
- Cancer**
Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671
Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054
NF- κ B in solid tumors, 1142
Epigenetic silencing of MHC2TA transcription in cancer, 1570
- Cancer therapy**
Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237
- Cancer-testis antigen**
Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577
- Cannabinoids**
The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869
R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenyl-methanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697
- Capsaicin**
Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104
- Carbamate prophylaxis**
Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344
- Carcinogenesis**
Ah receptor: Dioxin-mediated toxic responses as hints to deregulated physiologic functions, 393
S100A8 and S100A9 in inflammation and cancer, 1622
- Carcinogenic metals**
Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91
- Cardiovascular disease prevention**
Insulin sensitizers may prevent metabolic inflammation, 125
- Caspase**
Apigenin-induced-apoptosis is mediated by the activation of PKC δ and caspases in leukemia cells, 681
- Caspase-3**
Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761
- Caspases**
P-glycoprotein enhances TRAIL-triggered apoptosis in multidrug resistant cancer cells by interacting with the death receptor DR5, 293
The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423
Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668
- CCL2**
Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463
- CCL27**
Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463

CCL5

Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463

Cdc25

Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217

cDNA microarray

Human equilibrative nucleoside transporter-1 (hENT1) is required for the transcriptomic response of the nucleoside-derived drug 5'-DFUR in breast cancer MCF7 cells, 1646

Celastrol

Inhibition of NF- κ B activation through targeting I κ B kinase by celastrol, a quinone methide triterpenoid, 1311

A549 cells

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

Cell adhesion

The influence of various structural parameters of semi-synthetic sulfated polysaccharides on the P-selectin inhibitory capacity, 474

Cell cycle

Synthesis of microtubule-interfering halogenated nospapine analogs that perturb mitosis in cancer cells followed by cell death, 415

Role of 5-lipoxygenase pathway in the regulation of RAW 264.7 macrophage proliferation, 1022

Inhibition of phosphatidylinositol 3-kinase-mediated glucose metabolism coincides with resveratrol-induced cell cycle arrest in human diffuse large B-cell lymphomas, 1246

Cell cycle arrest

In vitro and *in vivo* studies of a novel potential anticancer agent of isochaihulactone on human lung cancer A549 cells, 308

Cell cycle inhibition

Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217

Cell death

Enrichment of non-synchronized cells in the G1, S and G2 phases of the cell cycle for the study of apoptosis, 1396

Cell growth

Role of 5-lipoxygenase pathway in the regulation of RAW 264.7 macrophage proliferation, 1022

Cell migration

New approaches to blockade of α 4-integrins, proven therapeutic targets in chronic inflammation, 1460

Cell sorting

Enrichment of non-synchronized cells in the G1, S and G2 phases of the cell cycle for the study of apoptosis, 1396

Cell survival

Radioccontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334

Cell-cycle

Differential processing of antitumour-active and antitumour-inactive *trans* platinum compounds by SKOV-3 ovarian cancer cells, 280

 β -Cells

The diabetogenic antibiotic streptozotocin modifies the tryptic digest pattern for peptides of the enzyme O-GlcNAc-selective N-acetyl- β -D-glucosaminidase that contain amino acid residues essential for enzymatic activity, 710

Cellular signalling

NF- κ B activation by double-strand breaks, 1132

NF- κ B activation by reactive oxygen species: Fifteen years later, 1493

Central and peripheral nervous system

NF- κ B functions in the nervous system: From development to disease, 1180

CEP-1347

Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197

Cerulein

Suppression of IL-1 β expression by the Jak 2 inhibitor AG490 in cerulein-stimulated pancreatic acinar cells, 1555

A431 cervix squamous cell carcinoma

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

Chemokine receptor CCR5

Diverse signalling by different chemokines through the chemokine receptor CCR5, 739

Chemokines

Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588

Diverse signalling by different chemokines through the chemokine receptor CCR5, 739

Inflammation and cancer: How hot is the link?, 1605

Chemoprevention

Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention, 795

Chemoresistance

Enrichment of non-synchronized cells in the G1, S and G2 phases of the cell cycle for the study of apoptosis, 1396

Chemosensitizers

Dual-acting diamine antiplasmodial and chloroquine resistance modulating agents, 156

Chemotaxis

Effective attenuation of acute lung injury *in vivo* and the formyl peptide-induced neutrophil activation *in vitro* by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

Chemotherapy

Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polyps, 729

Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054

8-Chloro-adenosine

8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells, 541

Chloroquine

Dual-acting diamine antiplasmodial and chloroquine resistance modulating agents, 156

CHO-IR cells

PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279

Cholesterol

Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749

Cholestyramine

Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749

Chondrocyte

Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382

- Chromatin**
Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114
- Chromatin structure**
Chromatin structure and epigenetics, 1563
- Chronic inflammation**
New approaches to blockade of α 4-integrins, proven therapeutic targets in chronic inflammation, 1460
- Chronopharmacotherapy**
Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237
- Cinnamaldehyde**
Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104
- Circadian rhythm**
Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237
- Cisplatin**
XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166
- Citrus pectin**
Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
- Class II transactivator**
Epigenetic silencing of *MHC2TA* transcription in cancer, 1570
- Cleavage**
Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- Clinical trial failure**
Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197
- Clozapine**
Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential in vitro system for the study of agranulocytosis, 783
- Cofactor**
Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114
- Collagen**
The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227
- Collagen biosynthesis**
Novel amidine analogue of melphalan as a specific multifunctional inhibitor of growth and metabolism of human breast cancer cells, 320
- Colon cancer**
Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
- Competitive inhibition**
3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239
- Computer simulation**
Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344
- Connective tissue growth factor**
STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32
- Conotoxins**
Toxin insights into nicotinic acetylcholine receptors, 661
- Contacts**
Recognition forces in ligand–protein complexes: Blending information from different sources, 1633
- COPD**
Synergistic effects of the anti-cholinergic R,R-glycopyrrolate with anti-inflammatory drugs, 1690
- Corticotropin releasing factor**
Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor₁ (CRF₁) receptor, 244
- COX**
Inflammation and cancer: How hot is the link?, 1605
- COX-2**
Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701
The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869
- Crocetin**
Crocetin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558
- Crosslinks**
Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11
- Cryptotanshinone**
Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680
- Crystallography**
Recognition forces in ligand–protein complexes: Blending information from different sources, 1633
- Curcumin**
Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62
TPA-induced up-regulation of activator protein-1 can be inhibited or enhanced by analogs of the natural product curcumin, 928
Regulation of inflammation and redox signaling by dietary polyphenols, 1439
Curcumin regulates signal transducer and activator of transcription (STAT) expression in K562 cells, 1547
- CXCR4**
Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588
- Cyclooxygenase-2**
Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin *in vivo*: NF- κ B and AP-1 as prime targets, 1506
- Cyclooxygenase-2 (COX-2)**
Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293
- Cyclosporin A**
Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042
- CYL-26z**
Effective attenuation of acute lung injury *in vivo* and the formyl peptide-induced neutrophil activation *in vitro* by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

CYP2B1

Cytochrome P450 reductase dependent inhibition of cytochrome P450 2B1 activity: Implications for gene directed enzyme prodrug therapy, 893

CYP2C8

Cloning of a cDNA encoding a novel marmoset CYP2C enzyme, expression in yeast cells and characterization of its enzymatic functions, 1738

CYP3A

Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042

CYP3A23

Selective induction of intestinal CYP3A23 by $1\alpha,25$ -dihydroxyvitamin D₃ in rats, 385

Cytochrome P450

Selective induction of intestinal CYP3A23 by $1\alpha,25$ -dihydroxyvitamin D₃ in rats, 385

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

Cytochrome P450 reductase dependent inhibition of cytochrome P450 2B1 activity: Implications for gene directed enzyme prodrug therapy, 893

Cytochrome P450 reductase

Cytochrome P450 reductase dependent inhibition of cytochrome P450 2B1 activity: Implications for gene directed enzyme prodrug therapy, 893

Cytokine receptor

Jaks and cytokine receptors—An intimate relationship, 1538

Cytokines

Extending the nuclear roles of I κ B kinase subunits, 1081

NF- κ B in solid tumors, 1142

Listeria monocytogenes induced Rac1-dependent signal transduction in endothelial cells, 1367

NF- κ B activation by reactive oxygen species: Fifteen years later, 1493

Cytotoxicity

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366

Cellular pharmacokinetics and pharmacodynamics of dipyrindamole in vascular smooth muscle cells, 956

Daidzein

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

DARPP-32

Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919

De novo purine synthesis

Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816

15-Deoxy- $\Delta^{12,14}$ -prostaglandin

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Depression

Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor₁ (CRF₁) receptor, 244

Development

NF- κ B functions in the nervous system: From development to disease, 1180

Dexamethasone

Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042

5'-DFUR

Human equilibrative nucleoside transporter-1 (hENT1) is required for the transcriptomic response of the nucleoside-derived drug 5'-DFUR in breast cancer MCF7 cells, 1646

DHFR

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

Diabetes

Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176

The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965

Differentiation

Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530

p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417

Diffuse large B cell lymphoma

Inhibition of phosphatidylinositol 3-kinase-mediated glucose metabolism coincides with resveratrol-induced cell cycle arrest in human diffuse large B-cell lymphomas, 1246

1 $\alpha,25$ -Dihydroxyvitamin D₃

Selective induction of intestinal CYP3A23 by $1\alpha,25$ -dihydroxyvitamin D₃ in rats, 385

7,12-Dimethylbenz[a]anthracene (DMBA)

Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293

Dioxin

Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267

Dioxin receptor

Ah receptor: Dioxin-mediated toxic responses as hints to deregulated physiologic functions, 393

Dipyridamole

Cellular pharmacokinetics and pharmacodynamics of dipyridamole in vascular smooth muscle cells, 956

Dissociation rate constant

Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure-activity relationships and insurmountable antagonism, 838

Disulfiram

Protein S-glutathionylation and platelet anti-aggregating activity of disulfiram, 608

DNA damage

Extending the nuclear roles of I κ B kinase subunits, 1081

DNA hypomethylation

Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577

DNA methylation

Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114

Chromatin structure and epigenetics, 1563

Epigenetic silencing of MHC2TA transcription in cancer, 1570

DNA repair

Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11

- XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166
- DNA topoisomerase inhibitors
- NF- κ B activation by double-strand breaks, 1132
- Docking simulations
- Toxin insights into nicotinic acetylcholine receptors, 661
- Dopamine
- Tyramine in the assessment of regional adrenergic function, 1724
- Double-strand breaks
- NF- κ B activation by double-strand breaks, 1132
- Down-regulation
- Proteasome-independent down-regulation of estrogen receptor- α (ER α) in breast cancer cells treated with 4,4'-dihydroxy-*trans*-stilbene, 573
- DPP
- Dipeptidyl peptidase II and leukocyte cell death, 70
- Drug resistance
- Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11
- Dual-acting diamine antiplasmodial and chloroquine resistance modulating agents, 156
- Receptor tyrosine kinase (RTK) inhibition is effective in chemosensitising EGFR-expressing drug resistant human ovarian cancer cell lines when used in combination with cytotoxic agents, 941
- Dual perfusion
- Effect of human serum albumin on transplacental transfer of glyburide, 632
- EGCG
- Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850
- EGF receptor
- Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485
- EGFR expressing
- Receptor tyrosine kinase (RTK) inhibition is effective in chemosensitising EGFR-expressing drug resistant human ovarian cancer cell lines when used in combination with cytotoxic agents, 941
- Electrophile-responsive element (EpRE)
- Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217
- Endocrine pancreas
- The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965
- Endocytic recycling compartment
- PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279
- Endogenous Glucose production
- Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198
- Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757
- Endoplasmic reticulum
- Green tea flavonols inhibit glucosidase II, 640
- Endothelial cells
- Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176
- BE-I-PLA2, a novel acidic phospholipase A₂ from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inducer of prostaglandin I₂ release by endothelial cells, 377
- Listeria monocytogenes* induced Rac1-dependent signal transduction in endothelial cells, 1367
- Endothelial nitric oxide synthase
- Crocetin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558
- Enhanced sensitivity
- Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816
- Enzyme product inhibition
- Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256
- Epidermis
- p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417
- Epigenetics
- Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114
- Chromatin structure and epigenetics, 1563
- Epigenetic regulation of nuclear steroid receptors, 1589
- Epstein Barr Virus
- Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- Equilibrative nucleoside transporter-1
- Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816
- ERCC1
- XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166
- ERK
- Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268
- 2',4',6'-Tris(methoxymethoxy) chalcone attenuates hepatic stellate cell proliferation by a heme oxygenase-dependent pathway, 1322
- Erk1/2
- Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382
- Estrogen
- Down-regulation of estrogen receptor- α in MCF-7 human breast cancer cells after proteasome inhibition, 566
- Estrogen receptor
- Epigenetic regulation of nuclear steroid receptors, 1589
- Estrogen receptor- α
- Proteasome-independent down-regulation of estrogen receptor- α (ER α) in breast cancer cells treated with 4,4'-dihydroxy-*trans*-stilbene, 573
- Estrogen receptor- α down-regulation
- Down-regulation of estrogen receptor- α in MCF-7 human breast cancer cells after proteasome inhibition, 566
- Etoposide
- The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207
- Experimental autoimmune encephalomyelitis
- R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697

Experimental colitis

Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432

Expression

Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267

FDH

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

FERM domain

Jaks and cytokine receptors—An intimate relationship, 1538

Fetus

Expression of glutathione S-transferases in fetal lung and liver tissue from parental strains and F1 crosses between C57BL/6 and BALB/c F1 mice following in utero exposure to 3-methylcholanthrene, 115

Fibrogenesis

The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227

Flavonoids

Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184

Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217

Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850

Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850

Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship, 1010

Fluorescence imaging plate reader

Pharmacological characterization of recombinant N-type calcium channel (Ca_v2.2) mediated calcium mobilization using FLIPR, 770

Flupentixol

Biochemical and pharmacological properties of an allosteric modulator site of the human P-glycoprotein (ABCB1), 145

Folate

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

Formyl peptide receptor-like 1

Differential production of leukotriene B₄ or prostaglandin E₂ by WKYMVm or serum amyloid A via formyl peptide receptor-like 1, 860

G protein-coupled receptors

Regulation of death and survival in astrocytes by ADP activating P2Y₁ and P2Y₁₂ receptors, 1031

G/F-actin ratio

8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells, 541

G2/M arrest

8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells, 541

Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668

Gemini Vitamin D₃

Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332

Gene expression

The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227

Differential processing of antitumour-active and antitumour-inactive *trans* platinum compounds by SKOV-3 ovarian cancer cells, 280

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

Regulation of the hypertonic stress response and other cellular functions by the Rel-like transcription factor NFAT5, 1597

Gene regulation

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

Epigenetic silencing of MHC2TA transcription in cancer, 1570

Genistein

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

Gestational diabetes

Effect of human serum albumin on transplacental transfer of glyburide, 632

Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730

GHKL family

The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207

Ginsenosides

The angiosuppressive effects of 20(R)- ginsenoside Rg₃, 437

Glioblastoma

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Glucocorticoid receptor

Tissue specific glucocorticoid receptor expression, a role for alternative first exon usage?, 1529

Glucocorticoids

Regulation of inflammation and redox signaling by dietary polyphenols, 1439

Glucose homeostasis

Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulinotropic polypeptide, 719

Glucose metabolism

Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198

Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757

Glucose tracers

Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198

Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757

Glucose-6 phosphatase

Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198

Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757

Glucose-dependent insulinotropic polypeptide

Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulinotropic polypeptide, 719

Glucose-lowering

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

Glucosidase II

Green tea flavonols inhibit glucosidase II, 640

Glutamate–cysteine ligase (GCL)

Bucillamine induces glutathione biosynthesis via activation of the transcription factor Nrf2, 455

Glutathione (GSH)

Bucillamine induces glutathione biosynthesis via activation of the transcription factor Nrf2, 455

Glutathione S-transferases

Expression of glutathione S-transferases in fetal lung and liver tissue from parental strains and F1 crosses between C57BL/6 and BALB/c F1 mice following in utero exposure to 3-methylcholanthrene, 115

Glyburide

Effect of human serum albumin on transplacental transfer of glyburide, 632

Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730

Glycerol

Tyramine in the assessment of regional adrenergic function, 1724

Glycolysis

Inhibition of phosphatidylinositol 3-kinase-mediated glucose metabolism coincides with resveratrol-induced cell cycle arrest in human diffuse large B-cell lymphomas, 1246

GM1

Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343

Gomisin A

Gomisin A alters substrate interaction and reverses P-glycoprotein-mediated multidrug resistance in HepG2-DR cells, 824

Gonadotropin-releasing hormone

Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure–activity relationships and insurmountable antagonism, 838

G-protein coupled receptor

Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588

Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure–activity relationships and insurmountable antagonism, 838

Grb14

Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355

Green tea

The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227

Green tea flavonols inhibit glucosidase II, 640

Growth suppressor

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

GSH

2',4',6'-Tris(methoxymethoxy) chalcone attenuates hepatic stellate cell proliferation by a heme oxygenase-dependent pathway, 1322

[³⁵S]GTP γ S binding

Diverse signalling by different chemokines through the chemokine receptor CCR5, 739

Gut

Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polyps, 729

GW9662

Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity *in vivo*, 42

HDAC inhibitors

Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701

Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577

Heart protection

Uridine-5'-triphosphate (UTP) reduces infarct size and improves rat heart function after myocardial infarct, 949

Heat shock proteins

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

Heme oxygenase

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

Hemodialysis arteriovenous graft stenosis

Cellular pharmacokinetics and pharmacodynamics of dipyridamole in vascular smooth muscle cells, 956

hENT1

Human equilibrative nucleoside transporter-1 (hENT1) is required for the transcriptomic response of the nucleoside-derived drug 5'-DFUR in breast cancer MCF7 cells, 1646

Hepa-1c1c7 cells

Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217

Heparin

The influence of various structural parameters of semi-synthetic sulfated polysaccharides on the P-selectin inhibitory capacity, 474

Hepatic stellate cell

2',4',6'-Tris(methoxymethoxy) chalcone attenuates hepatic stellate cell proliferation by a heme oxygenase-dependent pathway, 1322

Hepatitis C

Recruitment of an interferon molecular signaling complex to the mitochondrial membrane: Disruption by hepatitis C virus NS3-4A protease, 1477

Hepatocyte nuclear factor 1 alpha

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

HI-6

Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597

HIF-1 α

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992

High content screening

PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279

High fat diet-induced model

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

High-density culture

Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382

High-fat diet

Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity *in vivo*, 42

Histamine

Multiple role of histamine H₁-receptor–PKC–MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375

Histamine H₁-receptor

Multiple role of histamine H₁-receptor–PKC–MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375

Histatin

Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974

Histone

Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114

Epigenetic regulation of nuclear steroid receptors, 1589

Histone deacetylase

Regulation of inflammation and redox signaling by dietary polyphenols, 1439

Histone modification

Epigenetic silencing of MHC2TA transcription in cancer, 1570

Histones

Listeria monocytogenes induced Rac1-dependent signal transduction in endothelial cells, 1367

Hodgkin disease

Deregulated NF- κ B activity in haematological malignancies, 1069

Homology models

Toxin insights into nicotinic acetylcholine receptors, 661

HSP90

The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207

5-HT_{1A} receptor

Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498

5-HT synthesis

Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498

Human adipose stromal cells

Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268

Human aortic smooth muscle cells

Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680

Human lung cancer cell

8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells, 541

Human parvovirus B19

Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453

Human placenta

Effect of human serum albumin on transplacental transfer of glyburide, 632

Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730

Human platelets

Protein S-glutathionylation and platelet anti-aggregating activity of disulfiram, 608

Human serum albumin

Effect of human serum albumin on transplacental transfer of glyburide, 632

Huperzine A

Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344

HUVEC

The angiosuppressive effects of 20(R)- ginsenoside Rg₃, 437

Hydrogen peroxide

Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761

Hydrophobic effect

Recognition forces in ligand–protein complexes: Blending information from different sources, 1633

hydrophobicity

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

Hypertonic stress

Regulation of the hypertonic stress response and other cellular functions by the Rel-like transcription factor NFAT5, 1597

Hypoxia

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992

Hypoxia-inducible factor-1

Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91

IFN

Curcumin regulates signal transducer and activator of transcription (STAT) expression in K562 cells, 1547

Ifosfamide

Cytochrome P450 reductase dependent inhibition of cytochrome P450 2B1 activity: Implications for gene directed enzyme prodrug therapy, 893

Igf-I

Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382

IGF-I receptor

Novel amidine analogue of melphalan as a specific multi-functional inhibitor of growth and metabolism of human breast cancer cells, 320

- I κ B**
Deregulated NF- κ B activity in haematological malignancies, 1069
- I κ B kinase**
Inhibition of NF- κ B activation through targeting I κ B kinase by celastrol, a quinone methide triterpenoid, 1311
Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin *in vivo*: NF- κ B and AP-1 as prime targets, 1506
- IKK**
Extending the nuclear roles of I κ B kinase subunits, 1081
Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054
Deregulated NF- κ B activity in haematological malignancies, 1069
The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161
Enrichment of non-synchronized cells in the G1, S and G2 phases of the cell cycle for the study of apoptosis, 1396
- IL-1 β**
Suppression of IL-1 β expression by the Jak 2 inhibitor AG490 in cerulein-stimulated pancreatic acinar cells, 1555
- IL-2 receptor**
Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432
- Imatinib**
Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237
- Iminoether, SKOV-3 tumour cells**
Differential processing of antitumour-active and antitumour-inactive *trans* platinum compounds by SKOV-3 ovarian cancer cells, 280
- Immune subversion**
Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453
- Immunoprecipitation**
Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382
- In situ glutathione measurement**
Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- In situ loop**
Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042
- Incontinentia pigmenti**
NF- κ B and inflammation in genetic disease, 1153
- Indole-3-carbinol**
Indole-3-carbinol mediated cell cycle arrest of LNCaP human prostate cancer cells requires the induced production of activated p53 tumor suppressor protein, 1714
- Inducible nitric oxide synthase (iNOS)**
Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293
- Induction**
Selective induction of intestinal CYP3A23 by 1 α ,25-dihydroxyvitamin D₃ in rats, 385
- Inflammation**
The influence of various structural parameters of semi-synthetic sulfated polysaccharides on the P-selectin inhibitory capacity, 474
- Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship, 1010
Extending the nuclear roles of I κ B kinase subunits, 1081
Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090
NF- κ B in solid tumors, 1142
NF- κ B and inflammation in genetic disease, 1153
The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161
Therapeutic and preventive properties of quercetin in experimental arthritis correlate with decreased macrophage inflammatory mediators, 1304
S100A8 and S100A9 in inflammation and cancer, 1622
Synergistic effects of the anti-cholinergic R,R-glycopyrrolate with anti-inflammatory drugs, 1690
R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenyl-methanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697
Anti-allodynic property of flavonoid myricitrin in models of persistent inflammatory and neuropathic pain in mice, 1707
- Inhibition**
Green tea flavonols inhibit glucosidase II, 640
Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902
- Inhibitors**
Down-regulation of estrogen receptor- α in MCF-7 human breast cancer cells after proteasome inhibition, 566
TPA-induced up-regulation of activator protein-1 can be inhibited or enhanced by analogs of the natural product curcumin, 928
- Innate immunity**
Toll-like receptors: From the discovery of NF κ B to new insights into transcriptional regulations in innate immunity, 1102
- iNOS**
Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992
Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship, 1010
- Insulin receptor**
Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355
- Insulin resistance**
Insulin sensitizers may prevent metabolic inflammation, 125
- Insulin secretion**
The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965
- Insulin signalling**
PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279
- Insulin-like growth factor antagonist**
Paradoxical effects of the phage display-derived peptide antagonist IGF-F1-1 on insulin-like growth factor-1 receptor signaling, 53

Insulin-like growth factor binding proteins

Paradoxical effects of the phage display-derived peptide antagonist IGF-F1-1 on insulin-like growth factor-1 receptor signaling, 53

Insulin-like growth factor-1

Paradoxical effects of the phage display-derived peptide antagonist IGF-F1-1 on insulin-like growth factor-1 receptor signaling, 53

 β_1 -Integrin

Novel amidine analogue of melphalan as a specific multifunctional inhibitor of growth and metabolism of human breast cancer cells, 320

Integrins

Identification and characterization of triosephosphate isomerase that specifically interacts with the integrin α IIb cytoplasmic domain, 551

New approaches to blockade of $\alpha 4$ -integrins, proven therapeutic targets in chronic inflammation, 1460

Interferon

Recruitment of an interferon molecular signaling complex to the mitochondrial membrane: Disruption by hepatitis C virus NS3-4A protease, 1477

Interferon gamma

Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184

Interleukins

Multiple role of histamine H_1 -receptor-PKC-MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375

Inflammation and cancer: How hot is the link?, 1605

Intermolecular forces

Recognition forces in ligand-protein complexes: Blending information from different sources, 1633

Intestinal absorption

Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974

Intestine

Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polyps, 729

Intracellular pH

Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343

[125 I]iodoarylazidoprazosin (or [125 I]IAAP)

Biochemical and pharmacological properties of an allosteric modulator site of the human P-glycoprotein (ABCB1), 145

Ionizing radiation

NF- κ B activation by double-strand breaks, 1132

IRF

Recruitment of an interferon molecular signaling complex to the mitochondrial membrane: Disruption by hepatitis C virus NS3-4A protease, 1477

IRF-3

Distinct functions of IRF-3 and IRF-7 in IFN- α gene regulation and control of anti-tumor activity in primary macrophages, 1469

IRF-7

Distinct functions of IRF-3 and IRF-7 in IFN- α gene regulation and control of anti-tumor activity in primary macrophages, 1469

Iron

Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343

IRS-1

Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355

Ischemia

Uridine-5'-triphosphate (UTP) reduces infarct size and improves rat heart function after myocardial infarct, 949

Islet beta-cell

The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965

Islets

The diabetogenic antibiotic streptozotocin modifies the tryptic digest pattern for peptides of the enzyme O-GlcNAc-selective N-acetyl- β -D-glucosaminidase that contain amino acid residues essential for enzymatic activity, 710

Isochaihulactone

In vitro and *in vivo* studies of a novel potential anticancer agent of isochaihulactone on human lung cancer A549 cells, 308

Isoflavone

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

Jak

STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32

Jak/Stat

Suppression of IL-1 β expression by the Jak 2 inhibitor AG490 in cerulein-stimulated pancreatic acinar cells, 1555

Janus kinase

Jaks and cytokine receptors—An intimate relationship, 1538

JNK

Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090

K562

Curcumin regulates signal transducer and activator of transcription (STAT) expression in K562 cells, 1547

Keratinocytes

Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463

p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417

6-Ketocholestanol

Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911

Keto-acid

The methionine salvage pathway compound 4-methylthio-2-oxobutanate causes apoptosis independent of down-regulation of ornithine decarboxylase, 806

Kidney

Radiocontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334

Kinase

Radiocontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334

Kinetic analysis

Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344

Kinetics

Green tea flavonols inhibit glucosidase II, 640

Konjac mannan

Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749

- KR-62980
Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530
- K-ras activity
Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749
- K-vitamin derivative
Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217
- Leucovorin
Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256
- Leukemia
Apigenin-induced-apoptosis is mediated by the activation of PKC δ and caspases in leukemia cells, 681
Deregulated NF- κ B activity in haematological malignancies, 1069
Chromatin structure and epigenetics, 1563
- Leukotriene
Role of 5-lipoxygenase pathway in the regulation of RAW 264.7 macrophage proliferation, 1022
- Leukotriene receptor
Role of 5-lipoxygenase pathway in the regulation of RAW 264.7 macrophage proliferation, 1022
- Ligand binding
Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor₁ (CRF₁) receptor, 244
- Lipid peroxidation
Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343
- Lipolysis
Tyramine in the assessment of regional adrenergic function, 1724
- Lipopolysaccharide (LPS)
Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293
- Listeria
Listeria monocytogenes induced Rac1-dependent signal transduction in endothelial cells, 1367
- Liver
Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090
- Liver cancer
Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217
- LNCaP
Indole-3-carbinol mediated cell cycle arrest of LNCaP human prostate cancer cells requires the induced production of activated p53 tumor suppressor protein, 1714
- Lovastatin
Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749
- LOX
Inflammation and cancer: How hot is the link?, 1605
- LPS
Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62
- NF- κ B activation by reactive oxygen species: Fifteen years later, 1493
- LTB₄
Differential production of leukotriene B₄ or prostaglandin E₂ by WKYMVm or serum amyloid A via formyl peptide receptor-like 1, 860
- Lung
Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749
- Lung cancer
Synergistic induction of apoptosis by sulindac and arsenic trioxide in human lung cancer A549 cells via reactive oxygen species-dependent down-regulation of survivin, 1228
- Lung tumors
Expression of glutathione S-transferases in fetal lung and liver tissue from parental strains and F1 crosses between C57BL/6 and BALB/c F1 mice following in utero exposure to 3-methylcholanthrene, 115
- Lupus
Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453
- Lymphoma
Deregulated NF- κ B activity in haematological malignancies, 1069
- Lysophosphatidic acid
STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32
- Macrophages
Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship, 1010
Therapeutic and preventive properties of quercetin in experimental arthritis correlate with decreased macrophage inflammatory mediators, 1304
Distinct functions of IRF-3 and IRF-7 in IFN- α gene regulation and control of anti-tumor activity in primary macrophages, 1469
- Major histocompatibility complex class II
Epigenetic silencing of MHC2TA transcription in cancer, 1570
- Malaria
Dual-acting diamine antiplasmodial and chloroquine resistance modulating agents, 156
- MAP kinases
Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355
Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485
- MAPK
Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217
Multiple role of histamine H₁-receptor-PKC-MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375
- Marmoset
Cloning of a cDNA encoding a novel marmoset CYP2C enzyme, expression in yeast cells and characterization of its enzymatic functions, 1738
- Mass spectrometry
Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730
- Mathematical modeling
Target-mediated drug disposition and dynamics, 1

Matrix metalloproteinase-9

Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680

MCF10

Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332

MCF-7 cells

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

Mdm2

Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981

MDR1 gene

P-glycoprotein enhances TRAIL-triggered apoptosis in multidrug resistant cancer cells by interacting with the death receptor DR5, 293

mdr1a/1b knockout mice

Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042

Medium spiny neuron

Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919

Medulla oblongata

Sodium tanshinone IIA sulfonate derived from *Salvia miltiorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582

Melphalan

Novel amidine analogue of melphalan as a specific multifunctional inhibitor of growth and metabolism of human breast cancer cells, 320

Menadione

Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671

6-Mercaptopurine

Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816

Metabolic activation

NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366

Metabolic inflammation

Insulin sensitizers may prevent metabolic inflammation, 125

Metabolic syndrome

Insulin sensitizers may prevent metabolic inflammation, 125

Metabolism

Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730

Metformin

Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176

Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198

Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757

Methionine

The methionine salvage pathway compound 4-methylthio-2-oxobutanate causes apoptosis independent of down-regulation of ornithine decarboxylase, 806

Methyl mercaptopurineriboside

Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816

3-Methylcholanthrene

Expression of glutathione S-transferases in fetal lung and liver tissue from parental strains and F1 crosses between C57BL/6 and BALB/c F1 mice following in utero exposure to 3-methylcholanthrene, 115

Methylthioadenosine phosphorylase

The methionine salvage pathway compound 4-methylthio-2-oxobutanate causes apoptosis independent of down-regulation of ornithine decarboxylase, 806

Methylxanthines

Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902

Mice

Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749

Microarray

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

Microglia

Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992

Microtubule destabilizing

In vitro and *in vivo* studies of a novel potential anticancer agent of isochailulactone on human lung cancer A549 cells, 308

Migration

Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680

Mitochondria

Critical role of pro-apoptotic Bcl-2 family members in andrographolide-induced apoptosis in human cancer cells, 132

Salicylate modulates oxidative stress in the rat colon: A proteomic approach, 204

Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343

Mitochondrial permeability transition induced by novel pyridothiopyranopyrimidine derivatives: Potential new antimitochondrial antitumor agents, 1657

Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668

Mitochondrial dysfunction

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

Mitochondrial membrane potential

Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911

Mitochondrial toxicity

3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239

- Mitogen-activated protein kinases (MAPKs)**
 Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
 Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin *in vivo*: NF- κ B and AP-1 as prime targets, 1506
- Mitogen-activated protein (MAPK)**
 Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293
- Mitomycin C**
 XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166
 NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366
- mitoNEET**
 Insulin sensitizers may prevent metabolic inflammation, 125
- Mitotic arrest**
 Synthesis of microtubule-interfering halogenated noscapine analogs that perturb mitosis in cancer cells followed by cell death, 415
- Mixed Ah receptor/Nrf2 activators**
 Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention, 795
- Molecular dynamics**
 Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498
- Mouse model**
 Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
- Mouse skin**
 Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin *in vivo*: NF- κ B and AP-1 as prime targets, 1506
- mTOR**
 Hypoxia-induced iNOS expression in microglia is regulated by the PI3-kinase/Akt/mTOR signaling pathway and activation of hypoxia inducible factor-1 α , 992
- Multidrug resistance**
 Biochemical and pharmacological properties of an allosteric modulator site of the human P-glycoprotein (ABCB1), 145
 Gomisin A alters substrate interaction and reverses P-glycoprotein-mediated multidrug resistance in HepG2-DR cells, 824
- Multidrug-resistance-associated protein 2(Mrp2/MRP2)**
 Bucillamine induces glutathione biosynthesis via activation of the transcription factor Nrf2, 455
- Multiple Myeloma**
 The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423
- Multiple sclerosis**
 R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenyl-methanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697
- Murine thymocytes**
 Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486
- Mutagenesis**
 Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor₁ (CRF₁) receptor, 244
- MyD88**
 Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62
 Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (-)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850
- Myricitrin**
 Anti-allodynic property of flavonoid myricitrin in models of persistent inflammatory and neuropathic pain in mice, 1707
- NAD⁺**
 Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902
- Necrosis**
 Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902
 Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197
- Necrosis**
 Dipeptidyl peptidase II and leukocyte cell death, 70
 Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671
- Nephrotoxicity**
 Radiocontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334
- Neurodegenerative diseases**
 NF- κ B functions in the nervous system: From development to disease, 1180
- Neurofibromin**
 Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485
- Neurons and glia**
 NF- κ B functions in the nervous system: From development to disease, 1180
- Neuropathy**
 Anti-allodynic property of flavonoid myricitrin in models of persistent inflammatory and neuropathic pain in mice, 1707
- Neutrophils**
 Effective attenuation of acute lung injury *in vivo* and the formyl peptide-induced neutrophil activation *in vitro* by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749
 Differential production of leukotriene B₄ or prostaglandin E₂ by WKYMVm or serum amyloid A via formyl peptide receptor-like 1, 860
- NFAT5**
 Regulation of the hypertonic stress response and other cellular functions by the Rel-like transcription factor NFAT5, 1597
- NF κ B**
 Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293
 Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432
- NF- κ B**
 Extending the nuclear roles of I κ B kinase subunits, 1081

- Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054
- Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090
- Deregulated NF- κ B activity in haematological malignancies, 1069
- Toll-like receptors: From the discovery of NF κ B to new insights into transcriptional regulations in innate immunity, 1102
- Keeping up NF- κ B appearances: Epigenetic control of immunity or inflammation-triggered epigenetics, 1114
- NF- κ B activation by double-strand breaks, 1132
- NF- κ B in solid tumors, 1142
- NF- κ B and inflammation in genetic disease, 1153
- The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161
- NF- κ B functions in the nervous system: From development to disease, 1180
- Inhibition of NF- κ B activation through targeting I κ B kinase by celastrol, a quinone methide triterpenoid, 1311
- Regulation of inflammation and redox signaling by dietary polyphenols, 1439
- NF- κ B activation by reactive oxygen species: Fifteen years later, 1493
- 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J_2 as a potential endogenous regulator of redox-sensitive transcription factors, 1516
- Inflammation and cancer: How hot is the link?, 1605
- S100A8 and S100A9 in inflammation and cancer, 1622
- NGF**
- Multiple role of histamine H_1 -receptor-PKC-MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375
- Niacin**
- Elevated K-ras activity with cholestyramine and lovastatin, but not konjac mannan or niacin in lung—Importance of mouse strain, 1749
- Nicotinic acetylcholine receptor (nAChR)**
- Toxin insights into nicotinic acetylcholine receptors, 661
- NIK**
- The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161
- Nitric oxide**
- BE-I-PLA2, a novel acidic phospholipase A_2 from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inducer of prostaglandin I_2 release by endothelial cells, 377
- Crocetin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558
- Nonlinear pharmacokinetics**
- Target-mediated drug disposition and dynamics, 1
- Nonpeptide antagonist**
- Single amino acid residue determinants of non-peptide antagonist binding to the corticotropin-releasing factor $_1$ (CRF $_1$) receptor, 244
- Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure-activity relationships and insurmountable antagonism, 838
- Norepinephrine**
- Tyramine in the assessment of regional adrenergic function, 1724
- NR3C1**
- Tissue specific glucocorticoid receptor expression, a role for alternative first exon usage?, 1529
- Nrf2**
- Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention, 795
- Regulation of inflammation and redox signaling by dietary polyphenols, 1439
- 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J_2 as a potential endogenous regulator of redox-sensitive transcription factors, 1516
- NRH:quinone oxidoreductase 2**
- NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366
- NSAIDs**
- Synergistic induction of apoptosis by sulindac and arsenic trioxide in human lung cancer A549 cells via reactive oxygen species-dependent down-regulation of survivin, 1228
- Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257
- N-type calcium channel**
- Pharmacological characterization of recombinant N-type calcium channel (Ca $_v$ 2.2) mediated calcium mobilization using FLIPR, 770
- Nuclear factor-erythroid 2 p45-related factor (Nrf2)**
- Bucillamine induces glutathione biosynthesis via activation of the transcription factor Nrf2, 455
- Nuclear factor- κ B (NF- κ B)**
- Suppression of endotoxin-induced proinflammatory responses by citrus pectin through blocking LPS signaling pathways, 1001
- Nuclear morphologies**
- Different fates of intracellular glutathione determine different modalities of apoptotic nuclear vesiculation, 1405
- Nuclear receptors**
- Epigenetic regulation of nuclear steroid receptors, 1589
- Nucleoside reverse transcriptase inhibitors (NRTIs)**
- 3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239
- Nucleoside transporter**
- Human equilibrative nucleoside transporter-1 (hENT1) is required for the transcriptomic response of the nucleoside-derived drug 5'-DFUR in breast cancer MCF7 cells, 1646
- Nucleotides**
- Regulation of death and survival in astrocytes by ADP activating P2Y $_1$ and P2Y $_{12}$ receptors, 1031
- Nucleus**
- Extending the nuclear roles of I κ B kinase subunits, 1081
- Nucleus accumbens**
- Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919
- Nutlin-3**
- Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
- ob/ob mice**
- Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulintropic polypeptide, 719
- Obesity**
- Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity in vivo, 42
- The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965

- Obidoxime**
Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358
Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597
- OCT4**
Epigenetic control of CTCFL/BORIS and OCT4 expression in urogenital malignancies, 1577
- O-GlcNAcase**
The diabetogenic antibiotic streptozotocin modifies the tryptic digest pattern for peptides of the enzyme O-GlcNAc-selective N-acetyl- β -D-glucosaminidase that contain amino acid residues essential for enzymatic activity, 710
- OGT**
The diabetogenic antibiotic streptozotocin modifies the tryptic digest pattern for peptides of the enzyme O-GlcNAc-selective N-acetyl- β -D-glucosaminidase that contain amino acid residues essential for enzymatic activity, 710
- Organophosphorus compounds**
Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344
- Ortho-7**
Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597
- Osteoclasts**
Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761
- Osteogenic differentiation**
Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268
- Osteoporosis**
Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184
- Ovarian cancer**
Receptor tyrosine kinase (RTK) inhibition is effective in chemosensitising EGFR-expressing drug resistant human ovarian cancer cell lines when used in combination with cytotoxic agents, 941
- Ovariectomy**
Sodium tanshinone IIA sulfonate derived from *Slavia militorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582
- Oxidants**
Regulation of inflammation and redox signaling by dietary polyphenols, 1439
- Oxidative stress**
Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486
Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice, 671
Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902
- Oxidized low density lipoprotein**
Crocin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558
- p100**
Deregulated NF- κ B activity in haematological malignancies, 1069
- p100/p52**
The alternative NF- κ B pathway from biochemistry to biology: Pitfalls and promises for future drug development, 1161
- p19/ARF**
Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
- p21^{waf1/cip1}**
Indole-3-carbinol mediated cell cycle arrest of LNCaP human prostate cancer cells requires the induced production of activated p53 tumor suppressor protein, 1714
- P2Y receptors**
Uridine-5'-triphosphate (UTP) reduces infarct size and improves rat heart function after myocardial infarct, 949
- P450 GDEPT**
Cytochrome P450 reductase dependent inhibition of cytochrome P450 2B1 activity: Implications for gene directed enzyme prodrug therapy, 893
- p53**
Circumvention and reactivation of the p53 oncogene checkpoint in mouse colon tumors, 981
Indole-3-carbinol mediated cell cycle arrest of LNCaP human prostate cancer cells requires the induced production of activated p53 tumor suppressor protein, 1714
- p53 family members**
p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417
- Pancreatic acinar cells**
Suppression of IL-1 β expression by the Jak 2 inhibitor AG490 in cerulein-stimulated pancreatic acinar cells, 1555
- Paraoxon**
Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358
- Parkinson's disease**
Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616
Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197
- Partial least squares**
Salicylate modulates oxidative stress in the rat colon: A proteomic approach, 204
- PDGF receptor**
Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237
- Pentagalloyl glucose**
Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974
- Perfusion**
Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358
- Permeability transition**
Mitochondrial permeability transition induced by novel pyridothiopyranopyrimidine derivatives: Potential new antimitochondrial antitumour agents, 1657
- Permeability transition pore**
Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911
- Peroxisome proliferator activated receptor**
Epigenetic regulation of nuclear steroid receptors, 1589
- Peroxisome proliferator-activated receptor γ**
Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity in vivo, 42

15-PGDH

Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701

PGE₂

Differential production of leukotriene B₄ or prostaglandin E₂ by WKYMVm or serum amyloid A via formyl peptide receptor-like 1, 860

PGE₂

The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869

P-glycoprotein

P-glycoprotein enhances TRAIL-triggered apoptosis in multidrug resistant cancer cells by interacting with the death receptor DR5, 293

Gomisin A alters substrate interaction and reverses P-glycoprotein-mediated multidrug resistance in HepG2-DR cells, 824

Site-dependent contributions of P-glycoprotein and CYP3A to cyclosporin A absorption, and effect of dexamethasone in small intestine of mice, 1042

Pharmacodynamics

Target-mediated drug disposition and dynamics, 1

Pharmacokinetics

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

Pharmacology

Pharmacological characterization of recombinant N-type calcium channel (Ca_v2.2) mediated calcium mobilization using FLIPR, 770

Phenolic compound

Salicylate modulates oxidative stress in the rat colon: A proteomic approach, 204

Phenothiazine

Biochemical and pharmacological properties of an allosteric modulator site of the human P-glycoprotein (ABCB1), 145

Phenotyping

Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486

Phorbol ester

Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin in vivo: NF- κ B and AP-1 as prime targets, 1506

Phosphatidylinositol 3-kinase (PI3K/Akt)

Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293

Phosphatidylinositol 3-kinase

Inhibition of phosphatidylinositol 3-kinase-mediated glucose metabolism coincides with resveratrol-induced cell cycle arrest in human diffuse large B-cell lymphomas, 1246

Phospholipase A₂

BE-I-PLA2, a novel acidic phospholipase A₂ from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inducer of prostaglandin I₂ release by endothelial cells, 377

Phospholipase C

Regulation of death and survival in astrocytes by ADP activating P2Y₁ and P2Y₁₂ receptors, 1031

Physiologic responses

Ah receptor: Dioxin-mediated toxic responses as hints to deregulated physiologic functions, 393

Physostigmine

Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344

Phytochemicals

Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62

Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention, 795

Phytoestrogen

Metabolism of the soyabean isoflavone daidzein by CYP1A2 and the extra-hepatic CYPs 1A1 and 1B1 affects biological activity, 624

PI3K γ

Effective attenuation of acute lung injury in vivo and the formyl peptide-induced neutrophil activation in vitro by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

Piperine

Modulation of cadmium induced alterations in murine thymocytes by piperine: Oxidative stress, apoptosis, phenotyping and blastogenesis, 486

PKC

Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176

Apigenin-induced-apoptosis is mediated by the activation of PKC δ and caspases in leukemia cells, 681

Multiple role of histamine H₁-receptor-PKC-MAPK signalling pathway in histamine-stimulated nerve growth factor synthesis and secretion, 1375

PKC δ

p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417

Platelet

BE-I-PLA2, a novel acidic phospholipase A₂ from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inducer of prostaglandin I₂ release by endothelial cells, 377

Identification and characterization of triosephosphate isomerase that specifically interacts with the integrin α IIb cytoplasmic domain, 551

Platelet receptors

BE-I-PLA2, a novel acidic phospholipase A₂ from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inducer of prostaglandin I₂ release by endothelial cells, 377

Platinum anticancer drugs

Differential processing of antitumour-active and antitumour-inactive *trans* platinum compounds by SKOV-3 ovarian cancer cells, 280

Platinum drugs

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

Ploidy

Enrichment of non-synchronized cells in the G1, S and G2 phases of the cell cycle for the study of apoptosis, 1396

PML-RAR α

Chromatin structure and epigenetics, 1563

Poly(ADP-ribose)polymerase-1

Caffeine metabolites are inhibitors of the nuclear enzyme poly(ADP-ribose)polymerase-1 at physiological concentrations, 902

Polyamines

The methionine salvage pathway compound 4-methylthio-2-oxobutanate causes apoptosis independent of down-regulation of ornithine decarboxylase, 806

Polysaccharides

The influence of various structural parameters of semi-synthetic sulfated polysaccharides on the P-selectin inhibitory capacity, 474

PPAR γ

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J_2 as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Pregnancy

Identification of glyburide metabolites formed by hepatic and placental microsomes of humans and baboons, 1730

Prenylation

Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485

Privileged structures

Dual-acting diamine antiplasmodial and chloroquine resistance modulating agents, 156

Prolactin releasing peptide (PrRP)

Sodium tanshinone IIA sulfonate derived from *Slavia militorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582

Proliferation

Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268

Proline-rich protein

Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974

Prostaglandin E_2

Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104

Suppressive effects of antimycotics on tumor necrosis factor- α -induced CCL27, CCL2, and CCL5 production in human keratinocytes, 463

Prostaglandin I_2

BE-1-PLA2, a novel acidic phospholipase A_2 from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inductor of prostaglandin I_2 release by endothelial cells, 377

Prostate

Ras inhibition results in growth arrest and death of androgen-dependent and androgen-independent prostate cancer cells, 427

Prostate cancer

TGF- β in cancer and as a therapeutic target, 523

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

Indole-3-carbinol mediated cell cycle arrest of LNCaP human prostate cancer cells requires the induced production of activated p53 tumor suppressor protein, 1714

Prostate carcinoma

Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91

Protease

Proteasome-independent down-regulation of estrogen receptor- α (ER α) in breast cancer cells treated with 4,4'-dihydroxy-*trans*-stilbene, 573

Proteasome

Down-regulation of estrogen receptor- α in MCF-7 human breast cancer cells after proteasome inhibition, 566

Proteasome-independent down-regulation of estrogen receptor- α (ER α) in breast cancer cells treated with 4,4'-dihydroxy-*trans*-stilbene, 573

Can NF- κ B be a target for novel and efficient anti-cancer agents?, 1054

Protein kinase C

Regulation of death and survival in astrocytes by ADP activating P2Y $_1$ and P2Y $_{12}$ receptors, 1031

Protein phosphatase

Fluorinated Cpd 5, a pure arylating K-vitamin derivative, inhibits human hepatoma cell growth by inhibiting Cdc25 and activating MAPK, 1217

Protein thiols

Protein S-glutathionylation and platelet anti-aggregating activity of disulfiram, 608

Protein trafficking

Jaks and cytokine receptors—An intimate relationship, 1538

Protein tyrosine phosphatase-1B

PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279

PSCA

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

P-selectin

The influence of various structural parameters of semi-synthetic sulfated polysaccharides on the P-selectin inhibitory capacity, 474

Pt-DNA binding

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

PTEN

Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530

PTP1B

Interaction between the insulin receptor and Grb14: A dynamic study in living cells using BRET, 1355

Pyridostigmine

Kinetic analysis of the protection afforded by reversible inhibitors against irreversible inhibition of acetylcholinesterase by highly toxic organophosphorus compounds, 344

Pyridothioipyranopyrimidine derivatives

Mitochondrial permeability transition induced by novel pyridothioipyranopyrimidine derivatives: Potential new antimitochondrial antitumour agents, 1657

Quercetin

Effect of salivary proteins on the transport of tannin and quercetin across intestinal epithelial cells in culture, 974

Quercetin, a flavonoid, inhibits proliferation and increases osteogenic differentiation in human adipose stromal cells, 1268

Therapeutic and preventive properties of quercetin in experimental arthritis correlate with decreased macrophage inflammatory mediators, 1304

Cloning of a cDNA encoding a novel marmoset CYP2C enzyme, expression in yeast cells and characterization of its enzymatic functions, 1738

Quiescent cell proline dipeptidase

Dipeptidyl peptidase II and leukocyte cell death, 70

Quinone methide triterpenoid

Inhibition of NF- κ B activation through targeting I κ B kinase by celastrol, a quinone methide triterpenoid, 1311

Quinones

NRH:quinone oxidoreductase 2 (NQO2) catalyzes metabolic activation of quinones and anti-tumor drugs, 366

R,R-Glycopyrrolate

Synergistic effects of the anti-cholinergic R,R-glycopyrrolate with anti-inflammatory drugs, 1690

R-8-OH-DPAT

Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498

Rac1

Listeria monocytogenes induced Rac1-dependent signal transduction in endothelial cells, 1367

Radicicol

The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207

RAGE

S100A8 and S100A9 in inflammation and cancer, 1622

Ras

Ras inhibition results in growth arrest and death of androgen-dependent and androgen-independent prostate cancer cells, 427

Effective attenuation of acute lung injury in vivo and the formyl peptide-induced neutrophil activation in vitro by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485

Rat

Sodium tanshinone IIA sulfonate derived from *Slavia miliorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582

Rat liver

Green tea flavonols inhibit glucosidase II, 640

RAW 264.7 monocyte/macrophages

Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293

Reactive oxygen species

Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91

Protective effect of monosialoganglioside GM1 against chemically induced apoptosis through targeting of mitochondrial function and iron transport, 1343

NF- κ B activation by reactive oxygen species: Fifteen years later, 1493

Mitochondrial permeability transition induced by novel pyridothiopyranopyrimidine derivatives: Potential new antimitochondrial antitumour agents, 1657

Real-time determination

Development of a dynamic model for real-time determination of membrane-bound acetylcholinesterase activity upon perfusion with inhibitors and reactivators, 358

A₃ receptors

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Receptor binding

Antagonism of peroxisome proliferator-activated receptor γ prevents high-fat diet-induced obesity in vivo, 42

Receptor internalisation

Diverse signalling by different chemokines through the chemokine receptor CCR5, 739

Receptor internalization

PTP1B-dependent insulin receptor phosphorylation/residency in the endocytic recycling compartment of CHO-IR cells, 1279

Receptor kinetics

Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure–activity relationships and insurmountable antagonism, 838

Receptor tyrosine kinase inhibition

Receptor tyrosine kinase (RTK) inhibition is effective in chemosensitising EGFR-expressing drug resistant human ovarian cancer cell lines when used in combination with cytotoxic agents, 941

Receptors

The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227

Redox regulation

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Redox status

Salicylate modulates oxidative stress in the rat colon: A proteomic approach, 204

Regulatory T cell

Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432

Rel transcription factors

Regulation of the hypertonic stress response and other cellular functions by the Rel-like transcription factor NFAT5, 1597

resistance

Different accumulation of cisplatin, oxaliplatin and JM216 in sensitive and cisplatin-resistant human cervical tumour cells, 693

Resistance

Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polypos, 729

Resistance mechanism

Leucovorin-induced resistance against FDH growth suppressor effects occurs through DHFR up-regulation, 256

Respiratory burst

Effective attenuation of acute lung injury in vivo and the formyl peptide-induced neutrophil activation in vitro by CYL-26z through the phosphoinositide 3-kinase γ pathway, 749

Resveratrol

Inhibition of phosphatidylinositol 3-kinase-mediated glucose metabolism coincides with resveratrol-induced cell cycle arrest in human diffuse large B-cell lymphomas, 1246

Regulation of inflammation and redox signaling by dietary polyphenols, 1439

Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin *in vivo*: NF- κ B and AP-1 as prime targets, 1506

Ret

Apoptotic cell death induction and angiogenesis inhibition in large established medullary thyroid carcinoma xenografts by Ret inhibitor RPI-1, 405

R-Flurbiprofen

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

Rg₃

The angiosuppressive effects of 20(R)- ginsenoside Rg₃, 437

Rheumatic arthritis

Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184

Rheumatoid arthritis

Therapeutic and preventive properties of quercetin in experimental arthritis correlate with decreased macrophage inflammatory mediators, 1304

Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453

RIG-I

Recruitment of an interferon molecular signaling complex to the mitochondrial membrane: Disruption by hepatitis C virus NS3-4A protease, 1477

ROS

Metformin reduces angiotensin-mediated intracellular production of reactive oxygen species in endothelial cells through the inhibition of protein kinase C, 176

Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616

Rosiglitazone

Differential anti-proliferative actions of peroxisome proliferator-activated receptor- γ agonists in MCF-7 breast cancer cells, 530

RPI-1

Apoptotic cell death induction and angiogenesis inhibition in large established medullary thyroid carcinoma xenografts by Ret inhibitor RPI-1, 405

S100

S100A8 and S100A9 in inflammation and cancer, 1622

S-15176

Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911

S-8-OH-DPAT

Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498

Salvia miltiorrhizae (*Salvia miltiorrhiza* Bunge)

Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680

Schizophrenia

Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential in vitro system for the study of agranulocytosis, 783

Serum amyloid A

Differential production of leukotriene B₄ or prostaglandin E₂ by WKYMVm or serum amyloid A via formyl peptide receptor-like 1, 860

S-Glutathionylated proteins

Protein S-glutathionylation and platelet anti-aggregating activity of disulfiram, 608

SH2 domain

Jaks and cytokine receptors—An intimate relationship, 1538

SH-SY5Y

Adiponectin protects human neuroblastoma SH-SY5Y cells against acetaldehyde-induced cytotoxicity, 616

Signal transduction

The phyto-chemical (–)-epigallocatechin gallate suppresses gene expression of epidermal growth factor receptor in rat hepatic stellate cells *in vitro* by reducing the activity of Egr-1, 227

Ras inhibition results in growth arrest and death of androgen-dependent and androgen-independent prostate cancer cells, 427

Signaling

Radiocontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334

Unscrambling the role of human parvovirus B19 signaling in systemic autoimmunity, 1453

Signaling pathways

The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869

SiRNA

XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166

Skin

p63 and p73, members of the p53 gene family, transactivate PKC δ , 1417

Slavia miltiorrhiza

Sodium tanshinone IIA sulfonate derived from *Slavia miltiorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582

SLC

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

Smad3

Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332

Small intestine

Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin, 198

Selective induction of intestinal CYP3A23 by 1 α ,25-dihydroxyvitamin D₃ in rats, 385

Erratum to: "Glucose utilization is suppressed in the gut of insulin-resistant high fat-fed rats and is restored by metformin" [Biochem. Pharmacol. 72 (2) (2006) 198–203], 1757

Smooth muscle cell

Cellular pharmacokinetics and pharmacodynamics of dipyridamole in vascular smooth muscle cells, 956

Snake venoms

BE-I-PLA2, a novel acidic phospholipase A₂ from *Bothrops erythromelas* venom: Isolation, cloning and characterization as potent anti-platelet and inductor of prostaglandin I₂ release by endothelial cells, 377

Sodium channel inhibitor

Pharmacological characterization of recombinant N-type calcium channel (Ca_v2.2) mediated calcium mobilization using FLIPR, 770

Sodium tanshinone IIA sulfonate

Sodium tanshinone IIA sulfonate derived from *Slavia miltiorrhiza* Bunge up-regulate the expression of prolactin releasing peptide (PrRP) in the medulla oblongata in ovariectomized rats, 582

Solvated systems

Recognition forces in ligand-protein complexes: Blending information from different sources, 1633

Sox9

Igf-I extends the chondrogenic potential of human articular chondrocytes *in vitro*: Molecular association between Sox9 and Erk1/2, 1382

Src kinase

STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32

Stable luciferase reporter cell lines

Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217

STAT

Jaks and cytokine receptors—An intimate relationship, 1538

Curcumin regulates signal transducer and activator of transcription (STAT) expression in K562 cells, 1547

STAT3

STAT3-independent inhibition of lysophosphatidic acid-mediated upregulation of connective tissue growth factor (CTGF) by cucurbitacin I, 32

Statin

Molecular targets for emerging anti-tumor therapies for neurofibromatosis type 1, 1485

Stereoselectivity

Stereoselectivity of 8-OH-DPAT toward the serotonin 5-HT_{1A} receptor: Biochemical and molecular modeling study, 498

Streptozotocin

The diabetogenic antibiotic streptozotocin modifies the tryptic digest pattern for peptides of the enzyme O-GlcNAc-selective N-acetyl- β -D-glucosaminidase that contain amino acid residues essential for enzymatic activity, 710

Stromal cell-derived factor

Characterization of the molecular pharmacology of AMD3100: A specific antagonist of the G-protein coupled chemokine receptor, CXCR4, 588

Stromal cells

Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential in vitro system for the study of agranulocytosis, 783

Structural change

Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597

Structure-activity analysis

Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure-activity relationship, 1010

Structure-activity relationships (SAR)

Toxin insights into nicotinic acetylcholine receptors, 661
Kinetics of nonpeptide antagonist binding to the human gonadotropin-releasing hormone receptor: Implications for structure-activity relationships and insurmountable antagonism, 838

Structure-function

Paradoxical effects of the phage display-derived peptide antagonist IGF-F1-1 on insulin-like growth factor-1 receptor signaling, 53

Substrate interaction

Gomisin A alters substrate interaction and reverses P-glycoprotein-mediated multidrug resistance in HepG2-DR cells, 824

Survivin

Synergistic induction of apoptosis by sulindac and arsenic trioxide in human lung cancer A549 cells via reactive oxygen species-dependent down-regulation of survivin, 1228
The immunomodulator AS101 induces growth arrest and apoptosis in Multiple Myeloma: Association with the Akt/Survivin pathway, 1423

Sustained delivery

Cellular pharmacokinetics and pharmacodynamics of dipyridamole in vascular smooth muscle cells, 956

Synaptic plasticity

Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919

T cells

R-(+)-[2,3-Dihydro-5-methyl-3-(4-morpholinylmethyl)-pyrrolo-[1,2,3-de]-1,4-benzoxazin-6-yl]-1-naphthalenyl-methanone (WIN-2) ameliorates experimental autoimmune encephalomyelitis and induces encephalitogenic T cell apoptosis: Partial involvement of the CB₂ receptor, 1697

Tachykinins

Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104

Target-mediated drug disposition

Target-mediated drug disposition and dynamics, 1

Tartrate-resistant acid phosphatase (TRAP)

Involvement of hydrogen peroxide in the differentiation and apoptosis of preosteoclastic cells exposed to arsenite, 761

TBK1

Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850

TCDD

Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267

TCH346

Recent clinical failures in Parkinson's disease with apoptosis inhibitors underline the need for a paradigm shift in drug discovery for neurodegenerative diseases, 1197

12-O-Tetradecanoylphorbol 13-acetate (TPA)

Acacetin suppressed LPS-induced up-expression of iNOS and COX-2 in murine macrophages and TPA-induced tumor promotion in mice, 1293

2,3,7,8-Tetrachlorodibenzo-p-dioxin

Regulating the regulator: Factors that control levels and activity of the aryl hydrocarbon receptor, 267

Tetrafluoroethylcysteine

Cytosolic heat shock proteins and heme oxygenase-1 are preferentially induced in response to specific and localized intramitochondrial damage by tetrafluoroethylcysteine, 80

TGF- β

Gene expression profiling changes induced by a novel Gemini Vitamin D derivative during the progression of breast cancer, 332

TGF- β in cancer and as a therapeutic target, 523

Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701

Therapeutic targets

NF- κ B in solid tumors, 1142

New approaches to blockade of α 4-integrins, proven therapeutic targets in chronic inflammation, 1460

Thiazolidinediones

Insulin sensitizers may prevent metabolic inflammation, 125

6-Thioguanine

Molecular mechanisms underlying the enhanced sensitivity of thiopurine-resistant T-lymphoblastic cell lines to methyl mercaptopurineriboside, 816

Thiol modification

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Thioxanthene

Biochemical and pharmacological properties of an allosteric modulator site of the human P-glycoprotein (ABCB1), 145

Thymidine

3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239

Thymidine kinase 2

3'-Azido-3'-deoxythymidine (AZT) is a competitive inhibitor of thymidine phosphorylation in isolated rat heart and liver mitochondria, 239

Thyroid carcinoma

Apoptotic cell death induction and angiogenesis inhibition in large established medullary thyroid carcinoma xenografts by Ret inhibitor RPI-1, 405

TMEV

The synthetic cannabinoid WIN 55,212-2 increases COX-2 expression and PGE₂ release in murine brain-derived endothelial cells following Theiler's virus infection, 869

TNF

Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship, 1010

Mechanisms of crosstalk between TNF-induced NF- κ B and JNK activation in hepatocytes, 1090

Inflammation and cancer: How hot is the link?, 1605

TNF receptor

Alkylating drugs applied in non-cytotoxic doses as a novel compounds targeting inflammatory signal pathway, 1432

TNF- α

Cryptotanshinone from *Salvia miltiorrhiza* BUNGE has an inhibitory effect on TNF- α -induced matrix metalloproteinase-9 production and HASMC migration via down-regulated NF- κ B and AP-1, 1680

Tolbutamide

Cloning of a cDNA encoding a novel marmoset CYP2C enzyme, expression in yeast cells and characterization of its enzymatic functions, 1738

Toll-like receptor 4

Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62

Toll-like receptors

Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850

Toll-like receptors: From the discovery of NF κ B to new insights into transcriptional regulations in innate immunity, 1102

Topiramate

The antiepileptic drug topiramate preserves metabolism-secretion coupling in insulin secreting cells chronically exposed to the fatty acid oleate, 965

Topoisomerase II alpha

Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11

Topoisomerase II beta

Topoisomerase II beta levels are a determinant of melphalan-induced DNA crosslinks and sensitivity to cell death, 11

Topoisomerases

Novel amidine analogue of melphalan as a specific multifunctional inhibitor of growth and metabolism of human breast cancer cells, 320

Toxic responses

Ah receptor: Dioxin-mediated toxic responses as hints to deregulated physiologic functions, 393

Toxicity

Clozapine bioactivation induces dose-dependent, drug-specific toxicity of human bone marrow stromal cells: A potential in vitro system for the study of agranulocytosis, 783

TRAIL

P-glycoprotein enhances TRAIL-triggered apoptosis in multidrug resistant cancer cells by interacting with the death receptor DR5, 293

TRAMP

Gene expression profiling in R-flurbiprofen-treated prostate cancer: R-Flurbiprofen regulates prostate stem cell antigen through activation of AKT kinase, 1257

Trans geometry

Differential processing of antitumour-active and antitumour-inactive *trans* platinum compounds by SKOV-3 ovarian cancer cells, 280

Transcription factors

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

NF- κ B functions in the nervous system: From development to disease, 1180

Resveratrol modulates phorbol ester-induced pro-inflammatory signal transduction pathways in mouse skin in vivo: NF- κ B and AP-1 as prime targets, 1506

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J₂ as a potential endogenous regulator of redox-sensitive transcription factors, 1516

Transformation

Ras inhibition results in growth arrest and death of androgen-dependent and androgen-independent prostate cancer cells, 427

Transforming growth factor-beta

TGF- β in cancer and as a therapeutic target, 523

Transient transfection

Newly constructed stable reporter cell lines for mechanistic studies on electrophile-responsive element-mediated gene expression reveal a role for flavonoid planarity, 217

Transplacental carcinogenesis

Expression of glutathione S-transferases in fetal lung and liver tissue from parental strains and F1 crosses between C57BL/6 and BALB/c F1 mice following in utero exposure to 3-methylcholanthrene, 115

Transplacental transfer

Effect of human serum albumin on transplacental transfer of glyburide, 632

Transporter

Alterations in transporter expression in liver, kidney, and duodenum after targeted disruption of the transcription factor HNF1 α , 512

trans-Stilbene

Proteasome-independent down-regulation of estrogen receptor- α (ER α) in breast cancer cells treated with 4,4'-dihydroxy-*trans*-stilbene, 573

Triazoloacridinones

Induction of G2/M phase arrest and apoptosis of human leukemia cells by potent antitumor triazoloacridinone C-1305, 1668

TRIF

Inhibition of homodimerization of Toll-like receptor 4 by curcumin, 62

Suppression of MyD88- and TRIF-dependent signaling pathways of toll-like receptor by (–)-epigallocatechin-3-gallate, a polyphenol component of green tea, 850

Triosephosphate isomerase

Identification and characterization of triosephosphate isomerase that specifically interacts with the integrin α IIb cytoplasmic domain, 551

2',4',6'-Tris(methoxymethoxy) chalcone

2',4',6'-Tris(methoxymethoxy) chalcone attenuates hepatic stellate cell proliferation by a heme oxygenase-dependent pathway, 1322

TRPA1

Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104

TRPV1

Contractile mechanisms coupled to TRPA1 receptor activation in rat urinary bladder, 104

Tubular cells

Radioccontrast media cause dephosphorylation of Akt and downstream signaling targets in human renal proximal tubular cells, 1334

Tubulin-binding

Synthesis of microtubule-interfering halogenated noscaine analogs that perturb mitosis in cancer cells followed by cell death, 415

Tumor

Expression of transporters potentially involved in the targeting of cytostatic bile acid derivatives to colon cancer and polyps, 729

Tumor necrosis factor

Attenuation of osteoclastogenesis and osteoclast function by apigenin, 184

Regulation of death and survival in astrocytes by ADP activating P2Y₁ and P2Y₁₂ receptors, 1031

Tumor suppressor

Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701

Type 2 diabetes

Evaluation of the antidiabetic activity of DPP IV resistant N-terminally modified versus mid-chain acylated analogues of glucose-dependent insulinotropic polypeptide, 719

Type I IFN

Distinct functions of IRF-3 and IRF-7 in IFN- α gene regulation and control of anti-tumor activity in primary macrophages, 1469

Type II DNA topoisomerase

The HSP90 and DNA topoisomerase VI inhibitor radicicol also inhibits human type II DNA topoisomerase, 1207

Tyrosine kinase

Basis for dosing time-dependent change in the anti-tumor effect of imatinib in mice, 1237

Tyrosine-kinase inhibitor

Apoptotic cell death induction and angiogenesis inhibition in large established medullary thyroid carcinoma xenografts by Ret inhibitor RPI-1, 405

Uncoupler

Effect of the mitochondrial transition pore inhibitor, S-15176, on rat liver mitochondria: ATP synthase modulation and mitochondrial uncoupling induction, 911

5' untranslated region

Tissue specific glucocorticoid receptor expression, a role for alternative first exon usage?, 1529

Val-boro-Pro

Dipeptidyl peptidase II and leukocyte cell death, 70

Vascular endothelial growth factor

Critical roles of AMP-activated protein kinase in the carcinogenic metal-induced expression of VEGF and HIF-1 proteins in DU145 prostate carcinoma, 91

Vascular relaxation function

Crocetin improves endothelium-dependent relaxation of thoracic aorta in hypercholesterolemic rabbit by increasing eNOS activity, 558

VDR

Selective induction of intestinal CYP3A23 by 1 α ,25-dihydroxyvitamin D₃ in rats, 385

VEGF

Adenosine modulates vascular endothelial growth factor expression via hypoxia-inducible factor-1 in human glioblastoma cells, 19

Ventral tegmental area

Coincident signaling in mesolimbic structures underlying alcohol reinforcement, 919

Viral evasion

Recruitment of an interferon molecular signaling complex to the mitochondrial membrane: Disruption by hepatitis C virus NS3-4A protease, 1477

Weight gain

KR-62980: A novel peroxisome proliferator-activated receptor γ agonist with weak adipogenic effects, 446

WKY/Mvm

Differential production of leukotriene B₄ or prostaglandin E₂ by WKY/Mvm or serum amyloid A via formyl peptide receptor-like 1, 860

Wnt

Histone deacetylase inhibitors and transforming growth factor- β induce 15-hydroxyprostaglandin dehydrogenase expression in human lung adenocarcinoma cells, 701

XPA

XPA versus ERCC1 as chemosensitising agents to cisplatin and mitomycin C in prostate cancer cells: Role of ERCC1 in homologous recombination repair, 166

X-ray crystallography

Crystal structures of acetylcholinesterase in complex with HI-6, Ortho-7 and obidoxime: Structural basis for differences in the ability to reactivate tabun conjugates, 597

X-ray protein structure

Toxin insights into nicotinic acetylcholine receptors, 661

Yeast cell

Cloning of a cDNA encoding a novel marmoset CYP2C enzyme, expression in yeast cells and characterization of its enzymatic functions, 1738