



Accumulation of thymidine-derived sugars in thymidine phosphorylase overexpressing cells

I.V. Bijnsdorp^a, K. Azijli^a, E.E. Jansen^b, M.M. Wamelink^b, C. Jakobs^b, E.A. Struys^b,
M. Fukushima^{c,1}, F.A.E. Kruijdt^d, G.J. Peters^{a,2,*}

^a VU University Medical Center, Department Medical Oncology, De Boelelaan 1117 CCA 1.42, 1081 HV Amsterdam, The Netherlands

^b VU University Medical Center, Department of Clinical Chemistry, Metabolic Unit, Amsterdam, The Netherlands

^c Tokushima Research Center, Taiho Pharmaceuticals Co., Ltd., Tokushima, Japan

^d University Medical Center Groningen, Department of Medical Oncology, Groningen, The Netherlands

ARTICLE INFO

Article history:

Received 9 February 2010

Accepted 11 May 2010

Keywords:

Thymidine phosphorylase

Angiogenesis

Deoxyribose

Thymidine phosphorylase inhibitor

ABSTRACT

Thymidine phosphorylase (TP) is often overexpressed in cancer and potentially plays a role in the stimulation of angiogenesis. The exact mechanism of angiogenesis induction is unclear, but is postulated to be related to thymidine-derived sugars. TP catalyzes the conversion of thymidine (TdR) to thymine and deoxyribose-1-phosphate (dR-1-P), which can be converted to dR-5-P, glyceraldehyde-3-phosphate (G3P) or deoxyribose (dR). However, it is unclear which sugar accumulates in this reaction. Therefore, in the TP overexpressing Colo320 TP1 and RT112/TP cells we determined by LC-MS/MS which sugars accumulated, their subcellular localization (using ³H-TdR) and whether dR was secreted from the cells. In both TP-overexpressing cell lines, dR-1-P and dR-5-P accumulated intracellularly at high levels and dR was secreted extensively by the cells. A specific inhibitor of TP completely blocked TdR conversion, and thus no sugars were formed. To examine whether these sugars may be used for the production of angiogenic factors or other products, we determined with ³H-TdR in which subcellular location these sugars accumulated. TdR-derived sugars accumulated in the cytoskeleton and to some extent in the cell membrane, while incorporation into the DNA was responsible for trapping in the nucleus. In conclusion, various metabolic routes were entered, of which the TdR-derived sugars accumulated in the cytoskeleton and membrane. Future studies should focus on which exact metabolic pathway is involved in the induction of angiogenesis.

© 2010 Elsevier Inc. All rights reserved.

1. Introduction

The platelet derived endothelial cell growth factor (PD-ECGF) or thymidine phosphorylase (TP) is often overexpressed in human cancers, which has been related to a higher microvessel density, a higher tumor stage and the induction of metastasis [1–4]. In various *in vitro* studies, TP had a chemotactic effect on endothelial cells [5–8]. *In vivo*, TP induced angiogenesis using various cancer cell lines that were transfected with TP [9,10]. The role of TP in angiogenesis seems evident, but its exact mechanism is still

unknown and it is postulated to be related to the sugars that are formed from thymidine (TdR) degradation by TP.

Deoxy- α -D-ribose-1-phosphate (dR-1-P) and thymine are the first products that are formed by the phosphorylysis of TdR by TP (Fig. 1) [11]. After its formation, dR-1-P rapidly disappears, possibly by conversion to 2-deoxy-D-ribose (dR) or 2-deoxyribose-5-phosphate (dR-5-P) [12]. dR-5-P can be converted to glyceraldehyde-3-phosphate (G3P), which can enter the glycolytic or pentose phosphate pathway. dR can form advanced glycation end products (AGE) or reactive oxygen species (ROS) by Schiff base reactions [13]. The Schiff base is formed from either dR or dR-5-P, which can be coupled to an intracellular protein by nonenzymatic condensation between the sugar's aldehyde group and a lysine residue. This Schiff base can subsequently rearrange to an Amadori product, such as an α -hydroxyketone, which may also form an enediol intermediate [14]. These unstable intermediates react via non-enzymatic reactions to form AGEs. During these reactions, specifically in the transition metal-catalyzed auto-oxidation, free radicals are produced [15–17]. The formation of AGE from these reactions could be responsible for the angiogenic activity of TP [13].

Abbreviations: TP, thymidine phosphorylase; TdR, thymidine; G3P, glyceraldehyde-3-phosphate; dR, deoxyribose; dR-1-P, deoxyribose-1-phosphate; dR-5-P, deoxyribose-5-phosphate; TPI, thymidine phosphorylase inhibitor; AGE, advanced glycation endproducts; VEGF, vascular endothelial growth factor; IL-8, interleukin-8.

* Corresponding author. Tel.: +31 20 4442630; fax: +31 20 4443844.

E-mail address: gj.peters@vumc.nl (G.J. Peters).

¹ Employee of Taiho Pharmaceuticals Co., Ltd.

² Received financial support from Taiho Pharmaceuticals Co., Ltd.




Fig. 1. Schematic overview of the conversion of TdR to TdR-related sugars and possible pathways which can stimulate angiogenesis. (A) TdR enters the cell, after which it is converted to thymine and deoxyribose-1-phosphate (dR-1-P). dR-1-P can be further converted to dR or dR-5-P. dR-5-P can enter the pentose phosphate pathway or glycolysis, stimulating the metabolism of the cells, possibly activating angiogenesis. dR is secreted by the cells, after which it can attract endothelial cells and/or be taken up by the endothelial cells, stimulating angiogenesis. Both dR and dR-5-P can undergo Schiff base reactions, first by opening the ring structure of the sugar by anomerization, after which it can undergo a Schiff base reaction where an Amadori product is formed. Subsequently, reactive oxygen species (ROS) are formed, or advanced glycation end products (AGE). Moieties at which [5'-³H]-TdR was labeled and the TdR-derived sugars are indicated with asterisk. TP: Thymidine phosphorylase; 1. thymidine kinase (TK); 2. DNA polymerase; 3. a phosphatase; 4. phospho-pentomutase; 6. deoxyribosephosphatase. (B) Structural formulae of the open ring structure of dR (2-deoxy-D-ribose), a Schiff base and an Amadori product. The open chain aldehyde glycates protein by a non-enzymatic condensation of the sugar to form a Schiff base, which rearranges to form an α -hydroxyketone Amadori product. R = protein (see also ref. [13]).

The enzymatic activity of TP is indispensable for the angiogenic effect, since a competitive inhibitor of TP could block the angiogenic effect [18]. Therefore, substrate and metabolites from the reaction are often used for studying the angiogenic effect of TP [19]. The metabolite dR has previously shown angiogenic activity [6,10] and could elongate the sprouting in the aortic ring assay [19]. In addition, dR induced endothelial cell migration in various studies [5,8]. The association between TP activity and angiogenesis is based on the potential accumulation of TdR-derived sugars. However, little is known about the cellular metabolism of dR-1-P. Although dR is considered the main factor causing the angiogenic switch, the real identity and extent of formation of TdR-derived sugars remains unclear. dR-1-P rapidly disappears after its formation and its conversion could be reduced by addition of dR-5-P [11]. The aim of the present study was to characterize the specific sugars that can be formed by the TP reaction, their subcellular localization and whether they are secreted by the cancer cells, enabling endothelial cells to migrate and invade towards tumor sites.

2. Materials and methods

2.1. Cell culture and chemicals

The human colon carcinoma cell line Colo320 TP1 was a variant of Colo320 (obtained from the ATCC), transfected with TP, as described previously [20]. RT112/TP was a kind gift of Dr. Bicknell

(Oxford, UK). Cells were cultured as monolayers in DMEM, supplemented with 10% heat inactivated FCS and 20 mM Hepes in 25 cm² culture flasks (Greiner Bio-One, Frickenhausen, Germany). Cells were maintained in a humidified 5% CO₂ atmosphere at 37 °C. TPI was provided by Taiho Pharmaceutical, Co., Ltd. (Tokushima, Japan). TdR, thymine, dR, dR-1-P, dR, dR-5-P and G3P were obtained from Sigma Aldrich Chemicals (Zwijndrecht, The Netherlands). These chemicals were dissolved in PBS in stock solutions of 1–20 mM and stored at –20 °C. ¹³C₆-glucose-6-phosphate (prepared as previously described [21]) and ¹³C₂-xylulose were obtained from Omicron biochemicals (South Bend, IN, USA). Ethoxyamine was obtained from Acros (Geel, Belgium) and pyridine from Merck (Darmstadt, Germany). 5'-³H-TdR (specific activity of 18.8 mCi/mmol, conc. 12.9 µg/ml) was purchased from Moravak Biochemicals Inc. (Brea, CA, USA). The exact moiety where the sugar of the TdR is labelled is indicated in Fig. 1. The thymidine phosphorylase antibody was purchased from R&D Systems Inc. (Minneapolis, MN, USA) and the thymidine kinase antibody was purchased from QED Bioscience (San Diego, CA, USA). The secondary antibody, goat- α -mouse-IRDye (800CW; #926-32210 and 680; #926-32220) was obtained from Westburg (Leusden, The Netherlands).

2.2. Treatment of the cells

For measurement of TdR and its sugars, Colo320TP1 and RT112/TP cells were seeded at 2×10^6 cells/flask. After 48 h, cells were

exposed to 100 μM TdR for 0 min, 15 min, 30 min and 60 min. TPI (10 μM) was added 24 h prior to TdR exposure, after which samples were incubated with TdR for 60 min. After treatment, cells were trypsinized, washed in PBS and centrifuged for 5 min at $323 \times g$ at room temperature (RT). The medium and the cell pellets were stored separately at -80°C until sample preparation.

2.3. Measurement of TP enzymatic activity and of TdR and thymine

HPLC measurement of TP enzymatic activity was performed as described previously [22] and is based on the detection of both TdR and thymine [23]. After incubation with thymidine, 25 μl 80% trichloroacetic acid (TCA) was added to the samples, which were left on ice for 20 min. Subsequently, samples were centrifuged at $14,000 \times g$ at 4°C for 10 min. The supernatant was transferred to a new vial, and the pH was neutralized. Samples were mixed and centrifuged for 1 min at $14,000 \times g$. The upper aqueous layer was used for analysis of TdR and thymine by HPLC analysis for nucleosides with UV detection as described previously [23].

2.4. Measurement of sugar-phosphates

For sample preparation, the cell pellet was dissolved in 500 μl MQ and subsequently samples were sonicated $3 \times$ for 3 s on ice. 5 μM $^{13}\text{C}_6$ -glucose-6-phosphate was used as an internal standard. The samples were centrifuged through a filter with 10 kDa cut off (Ultracel YM-10; #42407; Amicon Micron, Millipore, Billerica, MA, USA). Subsequently, the samples were injected into the LC-MS/MS as described previously [24]. The amount of sugars was expressed based on the total protein content of each sample that was measured before centrifugation through the 10 kDa cut off filter. Protein concentration was determined using the Bio-Rad Protein assay (#500-0006; Bio-Rad Laboratories, Veenendaal, The Netherlands) according to manufacture's instruction.

2.5. Measurement of dR

For preparation of the samples for the dR measurement, the cell pellet was dissolved in 150 μl Milli Q water. $^{13}\text{C}_2$ -xylulose was used as internal standard. 17 μl of 20 μM $^{13}\text{C}_2$ -xylulose was added to 50 μl of the sample. To deproteinize the sample, 200 μl methanol was added. After 15 min, the sample was centrifuged for 10 min at $10,000 \times g$ at 4°C . Following this, the supernatant was transferred to a new vial and evaporated to dryness under a slight stream of nitrogen. Ethoxime derivatives of dR were formed by treating the residue with 2 mg ethoxyamine in 100 μl pyridine at 60°C for 30 min. After cooling down to RT, the hydroxygroups were acetylated by adding 100 μl acetic anhydride at 80°C for 60 min. This solution was evaporated to dryness and the residue was redissolved in ethylacetate. 1–2 μl was injected into the GC-MS operating under positive chemical ionization in the single ion monitoring mode.

2.6. Separation of cell compartments

Colo320 TP1 and RT112/TP cells (1.5×10^6 cells) were exposed to 200 μM TdR (hot:cold (1:21) from which the batch of $[5\text{-}^3\text{H}]$ -TdR was mixed with 1 mM unlabeled TdR). After incubation for 1, 6 or 24 h or 1 h plus 24 h $[5\text{-}^3\text{H}]$ -TdR-free medium at 37°C , cell fractions were separated using a ProteoExtract[®] Subcellular Proteome Extraction Kit according to manufacture's instructions (Calbiochem, San Diego, CA). Of every cell fraction, including from the medium above the cells after the designated incubation times, 5 μl was counted. To determine to which fraction (protein or non-protein fraction) secreted $[5\text{-}^3\text{H}]$ -TdR-derived metabolites accumulated, 100 μl of the medium above the cells after the retention

was precipitated with 60 μl 35% TCA for 20 min on ice. Subsequently, samples were centrifuged for 5 min at $300 \times g$ at 4°C and 5 μl of the supernatant (non-protein fraction) was counted. In addition, the pellet (protein fraction) was recovered in 200 μl MQ, of which 5 μl was counted.

2.7. Western blotting

Colo320, Colo320 TP1, RT112 and RT112/TP cells were washed twice with ice-cold PBS and lysed in lysis buffer (Cell Signalling Technology Inc., Denver, USA). Cell lysates were scraped, transferred into a vial and centrifuged at $11,000 \times g$ at 4°C for 10 min. Supernatants were transferred to a new vial and protein amounts were determined by the Bio-Rad assay, according to the manufacturer's instruction (Bio-Rad Laboratories, Veenendaal, The Netherlands). From each condition 30 μg of protein was separated on a 10% SDS-PAGE and electroblotted onto polyvinylidenedifluoride (PVDF) membranes (Millipore ImmobilonTM-FL PVDF, 0.45 μm). Subsequently, the membranes were blocked for 1 h at room temperature (RT) in Odyssey blocking buffer (Odyssey blocking buffer #927-40003, Westburg, Leusden, The Netherlands) and incubated overnight at 4°C with the primary antibodies (dilution 1:1000–10,000 in Odyssey blocking buffer 1:1 diluted with PBS-T (PBS with 0.05% Tween-20)). The membrane was washed 5 times in PBS-T and incubated with the secondary antibodies (1:10,000) for 1 h at RT in the dark. After incubation, the membrane was washed in PBS-T and followed by 5 min washing in PBS without Tween-20 to decrease the background signal. Subsequently, the bands were scanned using an Odyssey Infrared Imager (LI-COR Biosciences, Lincoln, Nebraska USA), 0 mm offset, 84 μm resolution and with high quality [25].

2.8. Statistical analysis

Potential differences between controls and TdR exposed cells were evaluated using the two-tailed Student's *t*-test for paired data. Changes were considered significantly different when $P < 0.05$.

3. Results

3.1. TdR conversion to thymine

In order to determine the extent of TdR conversion, TdR and thymine levels were measured, after incubation with 100 μM TdR for 15, 30 and 60 min. As expected, the parental Colo320 and RT112 cells did not or hardly convert TdR to thymine, since these cells do not or hardly express TP (Fig. 2) [20]. TP was highly expressed in both Colo320 TP1 and RT112/TP cells (Fig. 2), while no band could be detected in Colo320 parental cells and a very faint band was found in RT112 cells. Colo320 and RT112 cells converted thymidine at a rate of 0 ± 0 and 3 ± 0.04 nmol/million cells/h, respectively.




Fig. 2. Western blot showing the expression levels of thymidine phosphorylase in Colo320, Colo320 TP1, RT112 and RT112/TP cells.




Fig. 3. The accumulation of sugars after conversion of TdR by thymidine phosphorylase. (A) Total levels of thymine and total levels of measured sugars. (B) Measurement of the levels of the various TdR-related sugars. (C) Measurement of the levels of dR that was secreted by the cells after degradation of TdR. All values represent means of three independent experiments \pm SEM. Significant differences compared to the control levels are indicated * $P < 0.05$.

Colo320 TP1 and RT112/TP converted thymidine at a rate of 518 ± 40 and 418 ± 34 nmol/million cells/h (Fig. 3A). After 1 h incubation of intact cells with TdR, the extracellular thymine concentrations were 24 and 30 μ M, respectively. The TK protein, which activates TdR for DNA synthesis, was similarly expressed in the TP-transfected cells (data not shown).

3.2. Intracellular accumulation of dR-5-P and dR-

TP degrades TdR into one molecule of thymine and one dR-1-P, which can be metabolized to other sugars (Fig. 1). We measured the intracellular levels of the potential metabolites dR-1-P, dR-5-P, dR and G3P in Colo320, RT112 and the transfected variants, Colo320 TP1 and RT112/TP. In the cells that did not express TP, no sugars accumulated at all (data not shown), which is in agreement with the absence of thymine formation. In general, the sugars accumulated to a much higher extent in RT112/TP cells, compared to Colo320 TP1 cells (Fig. 3A and B). dR-1-P increased rapidly, and was about 10 fold higher in RT112/TP cells compared to Colo320 TP1 cells. dR-1-P was rapidly metabolized to dR-5-P and dR. In both Colo320TP1 and RT112/TP cells, dR-1-P was the major intracellular metabolite (Fig. 3B). In RT112/TP cells, the accumulation of dR-5-P was

lower than the level of dR-1-P. dR accumulated intracellularly to a low extent in Colo320TP1 cells and was only 4 fold increased compared to the control dR level. In RT112/TP cells, dR hardly accumulated. In Colo320 TP1 cells, G3P increased only at very low levels, with a 2 fold increase compared to the control. G3P did not accumulate in RT112/TP cells.

Addition of TPI alone to the cultures did not alter the production of the tested sugars. TPI completely blocked the conversion of TdR in Colo320 TP1 and RT112/TP cells (Fig. 3A), which is in agreement with the formation of the sugars, which were not formed at all (Fig. 3A and B).

3.3. dR is secreted from the cells

dR is an important angiogenic sugar that can be secreted by the cells [7,13]. Therefore, we determined the level of dR in the medium. dR-1-P and dR-5-P can not cross the membrane, because of the negative charge of the phosphate group on these sugars. In the medium above Colo320 TP1 and RT112/TP cells, dR increased in time (Fig. 3C) and was found at higher levels than the other sugars intracellularly (Table 1). Of the total thymine detected, 10 and 13% was measured as extracellular dR after 1 h by Colo320 TP1 and RT112/TP cells, respectively (Table 1). TPI alone did not have

Table 1

Percentage of TdR-related sugars compared to the total converted TdR.

| | Colo320TP1 | | | RT112/TP | | |
|-------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| | 15 | 30 | 60 | 15 | 30 | 60 |
| dR-1-P | 8.9 ± 0.4 [*] | 7.6 ± 2.7 [*] | 4.8 ± 0.6 [*] | 10.7 ± 0.9 [*] | 30.8 ± 2.7 [*] | 32.9 ± 2.6 [*] |
| dR-5-P | 1.9 ± 0.9 | 2.6 ± 1.7 [*] | 4.8 ± 0.6 [*] | 17.3 ± 3.5 [*] | 17.9 ± 2.9 [*] | 3.4 ± 0.9 |
| G3P | 0.5 ± 0.2 | 0.4 ± 0.2 | 0.3 ± 0.2 | 12.2 ± 4.0 | 6.2 ± 1.5 | 3.4 ± 1.4 |
| dR | 0.9 ± 0.4 | 0.6 ± 0.3 | 0.8 ± 0.7 | 2.1 ± 0.7 | 3.4 ± 0.9 | 0.7 ± 0.1 |
| Extracellular dR | 10.8 ± 2.4 [*] | 10.4 ± 2.2 [*] | 10.0 ± 0.3 [*] | 10.3 ± 1.4 [*] | 12.8 ± 0.5 [*] | 13.0 ± 0.1 [*] |
| Total sugars in relation to thymine | 23.2 ± 2.5 [*] | 21.5 ± 4.1 [*] | 20.7 ± 1.7 [*] | 52.5 ± 0.2 [*] | 71.2 ± 4.5 [*] | 53.5 ± 4.1 [*] |
| Total thymine | 100 | 100 | 100 | 100 | 100 | 100 |

Values were calculated relative to the amount of thymine formed and represent means (%) of three independent experiments ± SEM. Significant differences compared to the control levels are indicated.

^{*} $P < 0.05$.

any effect on the intracellular levels of dR, while TPI prevented dR from being formed (Fig. 3C).

3.4. Thymidine-derived sugars and accumulation levels

Since the thymine levels are representative for the total amount of sugars that are formed, we compared each value with the amount of thymine (in which thymine was set at 100% for each time point) (Table 1). In Colo320 TP1 cells, the total amount of measured sugars was much lower than that of RT112/TP cells (Fig. 3A). In Colo320 TP1 cells, about 20% of the total TdR-derived sugars presented intracellularly as dR-1-P, dR-5-P, dR and G3P and extracellular dR. This did not increase after a longer exposure time to TdR, indicating that the sugars are rapidly metabolized to other products. In RT112/TP cells, about 50% of the TdR-derived sugars were measured (Table 1), indicating a slower metabolism of these sugars into other products compared to that in Colo320 TP1 cells.

3.5. Accumulation in the cytoskeleton and nucleus

In order to determine in which cellular compartment the converted products of TdR accumulated, Colo320 TP1 and RT112/TP cells were exposed to [5'-³H]-TdR for different time periods. Subsequently, subcellular cell fractions were separated and the

amount of radioactivity was determined in each compartment (Fig. 4).

In Colo320 TP1 cells, the level of [5'-³H]-products in the cytosolic compartment decreased in time, until it was hardly detectable after 24 h. Products in the membrane fraction increased to some extent after 6 h, while after 24 h hardly any radioactivity was detectable in this fraction. In the nuclear fraction, the radioactivity was increased after 24 h, which was about 55% of the total intracellular radioactivity. This is possibly in part due to metabolic activation of TdR by thymidine kinase, resulting in incorporations into the DNA and nuclear trapping. Interestingly, TdR-derived radioactivity was present in the cytoskeletal protein fraction, which increased in time up to about 45% after 24 h.

In RT112/TP cells, the radioactivity in the cytosolic fraction decreased in time, with about 45% of [5'-³H]-products still detectable after 24 h (Fig. 4). This is in agreement with the lower metabolism of the sugars in these cells, as measured by LC-MS/MS (Table 1). Comparable to Colo320 TP1 cells, the radioactive products in the nuclear fraction increased in RT112/TP cells, although at lower levels; up to 25% after 24 h. Both the membrane and cytoskeleton fraction increased in time, with a maximum of about 25% and 10% after 24 h incubation, respectively.

When the [5'-³H]-TdR was washed away following 1 h incubation, the radioactivity was found to a high extent in the




Fig. 4. Accumulation of TdR-related sugars. (A) Accumulation in subcellular compartments of the cytosol, membrane, nucleus and cytoskeleton. Values represent means of three independent experiments ± SEM. nd = Not detectable. (B) Percentage of secreted TdR-related sugars in the medium above the cells, relative to the total radioactivity intra- and extracellular. Cells were exposed for 1 h to TdR after which the medium was refreshed and incubated for another 24 h. Subsequently the medium was analyzed for TdR-related sugars total, in the protein and in the non-protein fraction. Values are mean percentage of total formed thymine of 3 independent experiments ± SEM.

membrane (9.6 and 13.6%) and cytoskeletal (73.5 and 39.4%) compartments of both Colo320 TP1 and RT112/TP cells, respectively (data not shown). Hardly any of the radioactive label retained in the nuclear fraction, which may be related to a lack of activation of TdR by TK within 1 h exposure. In summary, the TdR-sugars are used in the cytoskeletal and membrane, which may possibly be related to protein-glycation.

3.6. Secreted products are in the non-protein fraction

To determine whether products were secreted after conversion of TdR, Colo320 TP1 and RT112/TP cells were exposed to [5'-³H]-TdR for 1 h, after which the medium was refreshed and cells were washed to remove the excess of [5'-³H]-TdR. The amounts of secreted products were determined by analyzing the amount of radioactivity in the medium fraction. Both Colo320 TP1 and RT112/TP cells secreted radioactive products, with higher levels for RT112/TP (24%) than for Colo320 TP1 (15%) cells.

To analyze whether the secreted products were in the protein or in the non-protein fraction, proteins were precipitated by adding TCA (Fig. 4B). In both Colo320 TP1 and RT112/TP cells, the largest part of the secreted radioactive products was found in the non-protein fraction. This indicates that the sugars may not be used to a high extent for protein-glycation of secreted proteins, but may be more important for processes intracellularly.

4. Discussion

In this study we describe TdR phosphorylation to dR-1-P, which is rapidly followed by its isomerization to dR-5-P or degradation to dR of which the latter was extensively secreted. To our knowledge, the formation of these sugars using human (eukaryotic) cells has not been described earlier. We observed different levels of accumulation of TdR-derived sugars in the two cell types. In addition, the converted TdR was found in the cytoskeleton and to some extent in the cell membrane.

Previously we demonstrated that dR-1-P formed from TdR rapidly disappeared from Colo320 TP1 cells [11]. dR-1-P can be isomerized to dR-5-P mediated by phosphopentomutase (E.C. 5.4.2.7) and is degraded to dR by a phosphatase [10]. dR-5-P can be split into G3P plus acetaldehyde by deoxyriboaldolase (E.C. 4.1.2.4) [12,26]. The formed acetaldehyde may be converted into acetyl-CoA by aldehyde oxidase and acetyl-CoA synthetase [27]. G3P can enter several metabolic pathways, including the glycolytic and the pentose phosphate pathway, and possibly therefore did not accumulate in the cells in our study. Initially, these steps were only described in e.g. *Bacillus cereus* [28]. Enzymatic activity needed for this pathway was also identified in epithelial aneuploid WISH cells [12]. The sugars may also be substrate in the formation of AGE (unpublished data), which may be used for protein glycation, e.g. adding sugar moieties to proteins by non-enzymatic reactions [16,29]. However, whether protein glycation of AGEs are formed in the cells used in our study, remains to be identified. Brown et al. previously reported a possible Schiff base reaction in RT112 cells, and demonstrated the presence of reactive oxygen species by expression levels of the oxidative stress marker heme oxygenase-1 (HO-1) [30].

dR-1-P can also be formed by conversion of purine deoxynucleosides by purine nucleoside phosphorylase (PNP) [31]. Therefore theoretically, PNP may also be an angiogenic enzyme. However, to the best of our knowledge, a role for PNP in angiogenesis has never been described before. Important differences between TP and PNP involve their function: TP overexpression in tumor sites has clearly been related with angiogenesis [2], while PNP is involved in immune function [32,33]. In addition, PNP is hardly overexpressed in cancer [32–34], making it less likely to be an angiogenic enzyme. It might be

possible that other products than the sugars are involved in angiogenesis as well, such as β -amino-iso-butyric acid, which is another downstream metabolite of thymine [19]. However, several enzymes involved in thymine catabolism (e.g. DPD and β -ureidopropionase) are usually low or not detectable in many tumors [35–37]. In our and other panels of colon cancer cell lines, DPD activity was low to not detectable [38,39], in contrast to pancreatic and breast cancer [39–41]. Therefore, it is unlikely that β -amino-iso-butyric acid is involved in the angiogenic activity of TP.

In the present study, we used cancer cells to determine the sugar metabolism in order to stimulate angiogenesis. TP expression in endothelial cells may also be important for their angiogenic properties. However, TP expression is very low in endothelial cells [42,43]. In addition, TP expression is often reported to be in either cancer cells or other tumor stromal cells such as fibroblasts. The angiogenic role of TP is possibly more related to the secretion of angiogenic factors (i.e. dR or chemoattractants such as IL-8) by cancer cells, rather than an autocrine stimulation by endothelial cells themselves. Therefore, the model that we used in the present study was intended to reflect the stimulation of angiogenesis by cancer cells.

Although TP is clearly associated with angiogenesis, the exact link between TdR-related sugars and angiogenesis is still missing. Cells with a high TP can secrete angiogenic factors at high levels, including the vascular endothelial growth factor (VEGF) and interleukin-8 (IL-8), but the role of sugars in the activation of these factors remains unclear [6,13]. TdR-derived sugars accumulated in the cell fractions of the membrane and cytoskeleton, which may be related to glycation of proteins by the formation of AGE. This may lead to the hypothesis that protein glycation plays a role in the activation of the transcription of angiogenic factors, such as IL-8 and VEGF [13] which can subsequently be secreted. The secreted fraction consisted of non-proteins. This indicates that dR may be the main TdR-sugar-product that is secreted by TP-expressing cells. Possibly, protein-glycation of secreted proteins is not the main regulator of angiogenesis. Thus, glycation of proteins that remain intracellularly are possibly important for angiogenesis, by activating transcription of angiogenic enzymes. Since dR can be secreted by the cancer cells, it can be taken up by the endothelial cells, in which it can also undergo Schiff base reactions, forming AGE inside the endothelial cells, stimulating their angiogenic properties (Fig. 1).

A possible role of sugars in the angiogenic potential of TP was studied through its metabolite dR, which can stimulate endothelial cell migration and angiogenesis. dR is a strongly reducing sugar that can generate oxygen radical species during the early stages of protein glycation [13]. dR can bind to an amino group of proteins by a non-enzymatic reaction. This leads to the formation of a Schiff base, which can subsequently rearrange to an α -hydroxyketone. During this reaction, free oxygen radicals are produced. Thus, through the formation of dR, TP induces oxidative stress in TP-overexpressing tumor cells. This may result in the secretion of angiogenic factors, such as VEGF or IL-8 [13]. dR secreted by the cancer cells may in this way directly or indirectly (by its conversion to other sugars) stimulate angiogenesis. dR was secreted by the cancer cells in our study, indicating that dR can directly play a role in the angiogenic potential of TP.

In conclusion, dR-1-P and subsequently dR-5-P and dR are formed from TdR conversion by TP. Both metabolites may stimulate angiogenesis, e.g. by the stimulation of angiogenic factors, while secreted dR may attract endothelial cells to form new blood vessels in the tumor tissue. The sugars formed from TdR will accumulate in the membrane and cytoskeleton of the cells, indicating that the sugars are used for further cellular metabolism, possibly mediating angiogenesis in a more indirect manner.

References

- [1] de Bruin M, Temmink OH, Hoekman K, Pinedo HM, Peters GJ. Role of platelet derived endothelial cell growth factor/thymidine phosphorylase in health and disease. *Cancer Ther* 2006;4:99–124.
- [2] Bronckaers A, Gago F, Balzarini J, Liekens S. The dual role of thymidine phosphorylase in cancer development and chemotherapy. *Med Res Rev* 2009;29:903–53.
- [3] Liekens S, Bronckaers A, Pérez-Pérez MJ, Balzarini J. Targeting platelet-derived endothelial cell growth factor/thymidine phosphorylase for cancer therapy. *Biochem Pharmacol* 2007;74:1555–67.
- [4] Koukourakis MI, Giatromanolaki A, Kakolyris S, O'Byrne KJ, Apostolikas N, Skarlatos J, et al. Different patterns of stromal and cancer cell thymidine phosphorylase reactivity in non-small-cell lung cancer: impact on tumour neoangiogenesis and survival. *Br J Cancer* 1998;77:1696–703.
- [5] Ishikawa F, Miyazono K, Hellman U, Drexler H, Wernstedt C, Hagiwara K, et al. Identification of angiogenic activity and the cloning and expression of platelet-derived endothelial cell growth factor. *Nature* 1989;338:557–62.
- [6] Hotchkiss KA, Ashton AW, Schwartz EL. Thymidine phosphorylase and 2-deoxyribose stimulate human endothelial cell migration by specific activation of the integrins alpha 5 beta 1 and alpha V beta 3. *J Biol Chem* 2003;278:19272–9.
- [7] Hotchkiss KA, Ashton AW, Klein RS, Lenzi ML, Zhu GH, Schwartz EL. Mechanisms by which tumor cells and monocytes expressing the angiogenic factor thymidine phosphorylase mediate human endothelial cell migration. *Cancer Res* 2003;63:527–33.
- [8] Seeliger H, Guba M, Koehl GE, Doenecke A, Steinbauer M, Bruns CJ, et al. Blockage of 2-deoxy-D-ribose-induced angiogenesis with rapamycin counteracts a thymidine phosphorylase-based escape mechanism available for colon cancer under 5-fluorouracil therapy. *Clin Cancer Res* 2004;10:1843–52.
- [9] Moghaddam A, Zhang HT, Fan TP, Hu DE, Lees VC, Turley H, et al. Thymidine phosphorylase is angiogenic and promotes tumor growth. *Proc Natl Acad Sci USA* 1995;92:998–1002.
- [10] Sengupta S, Sellers LA, Matheson HB, Fan TP. Thymidine phosphorylase induces angiogenesis in vivo and in vitro: an evaluation of possible mechanisms. *Br J Pharmacol* 2003;139:219–31.
- [11] de Bruin M, Smid K, Laan AC, Noordhuis P, Fukushima M, Hoekman K, et al. Rapid disappearance of deoxyribose-1-phosphate in platelet derived endothelial cell growth factor/thymidine phosphorylase overexpressing cells. *Biochem Biophys Res Commun* 2003;301:675–9.
- [12] Carta MC, Mattana A, Camici M, Allegrini S, Tozzi MG, Sgarrella F. Catabolism of exogenous deoxyinosine in cultured epithelial amniotic cells. *Biochim Biophys Acta* 2001;1528:74–80.
- [13] Brown NS, Jones A, Fujiyama C, Harris AL, Bicknell R. Thymidine phosphorylase induces carcinoma cell oxidative stress and promotes secretion of angiogenic factors. *Cancer Res* 2000;60:6298–302.
- [14] Wolff SP, Crabbe MJC, Thornalley PJ. The autoxidation of glyceraldehyde and other simple monosaccharides. *Experientia (Basel)* 1984;40:244–6.
- [15] Wolff SP, Dean RT. Glucose autoxidation and protein modification. The potential role of “autoxidative glycosylation” in diabetes. *Biochem J* 1987;245:243–50.
- [16] Monnier VM. Nonenzymatic glycosylation, the Maillard reaction and the aging process. *J Gerontol* 1990;45:B105–11.
- [17] Bierhaus A, Hofmann MA, Ziegler R, Nawroth PP. AGEs and their interaction with AGE-receptors in vascular disease and diabetes mellitus. I. The AGE concept. *Cardiovasc Res* 1998;37:586–600.
- [18] Haraguchi M, Miyadera K, Uemura K, Sumizawa T, Furukawa T, Yamada K, et al. Angiogenic activity of enzymes. *Nature* 1994;368:198.
- [19] Stevenson DP, Milligan SR, Collins WP. Effects of platelet-derived endothelial cell growth factor/thymidine phosphorylase, substrate, and products in a three-dimensional model of angiogenesis. *Am J Pathol* 1998;152:1641–6.
- [20] de Bruin M, van Capel T, Van der Born K, Kruyt FA, Fukushima M, Hoekman K, et al. Role of platelet-derived endothelial cell growth factor/thymidine phosphorylase in fluoropyrimidine sensitivity. *Br J Cancer* 2003;88:957–64.
- [21] Huck JH, Struys EA, Verhoeven NM, Jakobs C, van der Knaap MS. Profiling of pentose phosphate pathway intermediates in blood spots by tandem mass spectrometry: application to transaldolase deficiency. *Clin Chem* 2003;49:1375–80.
- [22] de Bruin M, Peters GJ, Oerlemans R, Assaraf YG, Masterson AJ, Adema AD, et al. Sulfasalazine down-regulates the expression of the angiogenic factors platelet-derived endothelial cell growth factor/thymidine phosphorylase and interleukin-8 in human monocytic-macrophage THP1 and U937 cells. *Mol Pharmacol* 2004;66:1054–60.
- [23] Laurensse EJ, Pinedo HM, Peters GJ. A sensitive non-radioactive assay for pyrimidine nucleoside phosphorylase using reversed-phase high performance liquid chromatography. *Clin Chim Acta* 1988;178:71–8.
- [24] Wamelink MM, Struys EA, Huck JH, Roos B, van der Knaap MS, Jakobs C, et al. Quantification of sugar phosphate intermediates of the pentose phosphate pathway by LC–MS/MS: application to two new inherited defects of metabolism. *J Chromatogr B Anal Technol Biomed Life Sci* 2005;823:18–25.
- [25] Mathews ST, Plaisance EP, Kim T. Imaging systems for westerns: chemiluminescence vs. infrared detection. *Methods Mol Biol* 2009;536:499–513.
- [26] Sgarrella F, Poddie FP, Meloni MA, Sciola L, Pippa P, Tozzi MG. Channelling of deoxyribose moiety of exogenous DNA into carbohydrate metabolism: role of deoxyriboaldolase. *Comp Biochem Physiol B Biochem Mol Biol* 1997;117:253–7.
- [27] Moriaki Y, Yamamoto T, Yamakita J, Takahashi S, Higashino K. Comparative localization of aldehyde oxidase and xanthine oxidoreductase activity in rat tissues. *Histochem J* 1998;30:69–74.
- [28] Sgarrella F, Del Corso A, Tozzi MG, Camici M. Deoxyribose 5-phosphate aldolase of *Bacillus cereus*: purification and properties. *Biochim Biophys Acta* 1992;1118:130–3.
- [29] Cho SJ, Roman G, Yeboah F, Konishi Y. The road to advanced glycation end products: a mechanistic perspective. *Curr Med Chem* 2007;14:1653–71.
- [30] Brown NS, Streeter EH, Jones A, Harris AL, Bicknell R. Cooperative stimulation of vascular endothelial growth factor expression by hypoxia and reactive oxygen species: the effect of targeting vascular endothelial growth factor and oxidative stress in an orthotopic xenograft model of bladder carcinoma. *Br J Cancer* 2005;92:1696–701.
- [31] Tozzi MG, Camici M, Mascia L, Sgarrella F, Ipata PL. Pentose phosphates in nucleoside interconversion and catabolism. *FEBS J* 2006;273:1089–101.
- [32] Pieters R, Huismans DR, Loonen AH, Peters GJ, Hählen K, van der Does-van den Berg A, et al. Adenosine deaminase and purine nucleoside phosphorylase in childhood lymphoblastic leukemia: relation with differentiation stage, in vitro drug resistance and clinical prognosis. *Leukemia* 1992;6:375–80.
- [33] van Laarhoven JP, de Bruyn CH. Purine metabolism in relation to leukemia and lymphoid cell differentiation. *Leuk Res* 1983;7(4):451–80.
- [34] Bardot V, Dutrillaux AM, Delattre JY, Vega F, Poisson M, Dutrillaux B, et al. Purine and pyrimidine metabolism in human gliomas: relation to chromosomal aberrations. *Br J Cancer* 1994;70:212–8.
- [35] Westra JL, Hollema H, Schaapveld M, Platteel I, Oien KA, Keith WN, et al. Predictive value of thymidylate synthase and dihydropyrimidine dehydrogenase protein expression on survival in adjuvantly treated stage III colon cancer patients. *Ann Oncol* 2005;16:1646–53.
- [36] Johnson MR, Diasio RB. Importance of dihydropyrimidine dehydrogenase (DPD) deficiency in patients exhibiting toxicity following treatment with 5-fluorouracil. *Adv Enzyme Regul* 2001;41:151–7.
- [37] Thomas HR, Ezzeldin HH, Guarcello V, Mattison LK, Fridley BL, Diasio RB. Genetic regulation of dihydropyrimidinase and its possible implication in altered uracil catabolism. *Pharmacogenet Genomics* 2007;17:973–87.
- [38] Peters GJ, van Groenigen CJ, Pinedo HM. Dihydropyrimidine dehydrogenase in livers from mouse and rat, and in human colon tumors, and mucosa in relation to anabolism of 5-fluorouracil. *Adv Exp Med Biol* 1998;431:633–6.
- [39] Beck A, Etienne MC, Chéradame S, Fischel JL, Formento P, Renée N, et al. A role for dihydropyrimidine dehydrogenase and thymidylate synthase in tumour sensitivity to fluorouracil. *Eur J Cancer* 1994;30A:1517–22.
- [40] Shimoda M, Sawada T, Kubota K. Thymidylate synthase and dihydropyrimidine dehydrogenase are upregulated in pancreatic and biliary tract cancers. *Pathobiology* 2009;76:193–8.
- [41] Anan K, Mitsuyama S, Tamae K, Suehara N, Nishihara K, Ogawa Y, et al. Increased dihydropyrimidine dehydrogenase activity in breast cancer. *J Surg Oncol* 2003;82:174–9.
- [42] Bijnsdorp IV, Vrijland K, Vrolijk L, Peters GJ. Increased migration by stimulation of thymidine phosphorylase in endothelial cells of different origin. *Nucleos Nucleot Nucl Acids* 2010;29:482–7.
- [43] Pula G, Mayr U, Evans C, Prokopi M, Vara DS, Yin X, et al. Proteomics identifies thymidine phosphorylase as a key regulator of the angiogenic potential of colony-forming units and endothelial progenitor cell cultures. *Circ Res* 2009;104:32–40.