

## ERRATA

Volume 44, No. 3, 1971, in the Communication, "Crystalline Kynureninase from *Pseudomonas marginalis*" by M. Moriguchi, T. Yamamoto and K. Soda; Pages 752-757, Page 754, Table 1. The figures in Total Units and Specific Activity should read:

| Step | Total Units | Specific Activity |
|------|-------------|-------------------|
| 1    | 6,903.7     | 0.069             |
| 2    | 6,795.2     | 0.101             |
| 3    | 6,258.9     | 0.141             |
| 4    | 1,488.0     | 0.458             |
| 5    | 1,032.3     | 1.009             |
| 6    | 787.4       | 1.321             |
| 7    | 393.7       | 9.145             |
| 8    | 238.7       | 9.520             |

Page 755, Line 5 of the first paragraph (Step 7) should read 6.2 instead of 2. Page 757, References, 10. The name of Journal should be added: Biochem. J., 114, 629 (1969).

---

Volume 44, No. 4, 1971, in the Communication, "Crystalline L-Leucine Dehydrogenase" by K. Soda, H. Misono, K. Mori and H. Sakato; Pages 931-935, Page 934, Line 8 of the explanatory material of Table 1 should read 1.0 instead of 0.001.

---

## Addendum

Addendum to the article entitled, "Isolation of a Glycoprotein - Glycolipid Fraction from Human Erythrocyte Membranes," by B. Zvilichovsky, P. M. Gallop, and O. O. Blumenfeld, Vol. 44 (1971), page 1234. The following was inadvertently omitted:

We have recently determined that commercial phospholipids, primarily sphingomyelin or phosphatidyl serine, when subjected to electrophoresis on 5% polyacrylamide gels in presence of sodium dodecyl sulfate give a Schiff-positive band (with or without previous treatment with periodic acid) which moves with the front. Thus phospholipids are also probably present in the isolated fraction and its ethanol-ether extract, and the labeling of this as a "glycolipid fraction" is not meant to indicate that only glycolipid is present.