



# Corrigendum

## Corrigendum to “Effect of human umbilical cord blood cells on Ang-II-induced hypertrophy in mice” [Biochem. Biophys. Res. Commun. 386 (2009) 386–391]

Pravan K. Vanamala<sup>a</sup>, Sreelatha Gopinath<sup>a</sup>, Christopher S. Gondi<sup>a</sup>, Jasti S. Rao<sup>a,b,\*</sup>

<sup>a</sup> Department of Cancer Biology & Pharmacology, University of Illinois, College of Medicine at Peoria, Peoria, IL 61656, USA

<sup>b</sup> Department of Neurosurgery, University of Illinois, College of Medicine at Peoria, Peoria, IL 61656, USA

The authors regret errors in their published Fig. 2. The corrected Fig. 2 and its caption appear below:




Fig. 2. hUCB reverse Ang-II-induced hypertrophy and apoptosis in NRCM. Cardiomyocytes isolated from neonatal rats treated with Ang-II (100 nM) were co-cultured with or without hUCB. (A) Cells were immunostained for  $\alpha$ -sarcomeric actinin. Induction of hypertrophy was determined as function of striated vs. punctate sarcomeric organization. (B) Cells stained for ANF polypeptide. Hypertrophy was determined as function of perinuclear staining.

DOI of original article: <http://dx.doi.org/10.1016/j.bbrc.2009.05.151>

\* Corresponding author at: Department of Cancer Biology & Pharmacology, University of Illinois, College of Medicine at Peoria, Peoria, IL 61656, USA. Fax: +309 671 3442.

E-mail addresses: [jsrao@uic.edu](mailto:jsrao@uic.edu), [shellee@uic.edu](mailto:shellee@uic.edu) (J.S. Rao).