



V α 14iNKT cell deficiency prevents acetaminophen-induced acute liver failure by enhancing hepatic glutathione and altering APAP metabolism

Isaac Downs^a, Tak Yee Aw^a, Jianfeng Liu^a, Patrick Adegboyega^b, Maureen N. Ajuebor^{a,*}

^a Department of Molecular & Cellular Physiology, Louisiana State University Health Sciences Center, Shreveport, LA, USA

^b Department of Pathology, Louisiana State University Health Sciences Center, Shreveport, LA, USA

ARTICLE INFO

Article history:

Received 2 October 2012

Available online 15 October 2012

Keywords:

Glutathione

APAP

V α 14iNKT cells

ROS

Liver

ABSTRACT

Acetaminophen (APAP) overdose is widely regarded as a major cause of acute liver failure in the United States. Intentional or accidental overdose of APAP in man or rodent elicits direct hepatocellular injury that is accompanied by hepatic depletion of the antioxidant, glutathione (GSH). In recent years, the innate immune response has also been shown to promote the development of APAP hepatotoxicity via indirect liver damage. In the present study, we demonstrate that J α 18^{−/−} mice, which are selectively deficient in the innate immune T cell, V α 14iNKT cells, were resistant to APAP hepatotoxicity relative to WT mice as reflected by biochemical and histological liver injury markers. In parallel, improvement in the biochemical and histological parameters of liver injury in J α 18^{−/−} mice was associated with a significant increase in hepatic levels of GSH, which detoxified APAP metabolites to attenuate hepatic oxidative stress, liver injury and necrosis. Notably, the protective effect of hepatic GSH during V α 14iNKT cells deficiency was demonstrated by its depletion in J α 18^{−/−} mice using DL-buthionine-[S,R]-sulfoximine which exacerbated hepatic oxidative and nitrosative stress as well as liver necrosis and caused mice mortality. Extraordinarily, APAP metabolism in J α 18^{−/−} mice was altered in favor of hepatic GSH conjugates and decreased glucuronide conjugates. In summary, we reveal a novel finding establishing a unique association between hepatic innate immunity and GSH levels in altering APAP metabolism to suppress liver injury and necrosis during V α 14iNKT cells deficiency in J α 18^{−/−} mice.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Globally, acetaminophen (APAP; *N*-acetyl-*p*-aminophenol) is a commonly used analgesic and antipyretic that is generally safe at the recommended dose. Widespread use of APAP in hundreds of prescription and over-the counter drugs has increased the prevalence of APAP hepatotoxicity [1,2]. Pre-existing liver disease, nutrition, age, genetic polymorphisms, alcohol, tobacco, or interactions with other drugs have been reported to enhance the toxic effects of APAP [1]. For this reason, APAP overdoses are the most common cause of acute liver failure (ALF) in the US [1,2]. APAP intoxication can initiate liver injury by direct and immune-mediated mechanisms [3].

Direct mechanism of APAP induced cell death occurs through the metabolic or oxidative phase [1,2]. The metabolic phase involves glucuronidation, sulfation, and oxidation pathways. These APAP biotransformation pathways form glucuronide, sulfate, and glutathione conjugates that increase the polarity and solubility of the xenobiotic contributing to excretion. For example, the glucu-

ronidation pathway uses glucuronyl transferases to catalyze UDP-glucuronic acid conjugation to APAP forming APAP-glucuronide whereas the sulfation pathway uses sulfotransferases and the cofactor, 3'-phosphoadenosine-5'-phosphosulfate (PAPS), to add a sulfate group to APAP forming APAP-sulfate [4]. During the oxidation pathway, APAP is oxidized to *N*-acetyl-*p*-benzoquinone imine (NAPQI) catalyzed by cytochrome P450 enzymes (e.g. CYP2E1; an enzyme found in high levels in centrilobular hepatocytes) [2,5]. NAPQI is an electrophile that forms protein adducts (e.g. mitochondrial proteins) during the metabolic phase of APAP toxicity [2]. Glutathione-S-transferase detoxifies NAPQI by catalyzing the conjugation of the antioxidant, glutathione (GSH), to NAPQI to form 3-[glutathione-S-yl]-APAP (APAP-GSH) [1,2]. Therefore, APAP intoxication increases CYP2E1 activity promoting NAPQI accumulation and subsequent hepatic GSH depletion [2,6]. In summary, the oxidative phase involves mitochondrial dysfunction, which increases reactive oxygen species (ROS) production leading to oxidative/nitrosative stress, ATP depletion, hepatocyte injury, and subsequent centrilobular necrosis.

Immune-mediated liver damage is also evident during APAP overdose. Emerging studies have demonstrated pathophysiological roles for innate immune cells such as peroxynitrite-producing Kupffer cells and neutrophils during APAP hepatotoxicity [2,7,8].

* Corresponding author. Address: Department of Molecular & Cellular Physiology, LSUHSC-Shreveport, 1501 Kings Highway, LA 71130-3932, USA.

E-mail address: majueb@gmail.com (M.N. Ajuebor).

Lymphocytes have also been reported to promote the development of APAP liver toxicity as RAG2^{-/-} mice, which are deficient in T and B cells, were found to be resistant to liver injury [9]. Conversely, the adoptive transfer of CD4⁺ T cells into RAG2^{-/-} mice increased APAP-induced liver damage. More recently, both NKT and NK cells were reported to participate in the development of APAP hepatotoxicity since simultaneous depletion of both cell types suppressed hepatic injury [9–11]. Specifically, CD1d^{-/-} mice which lack both Type I NKT cells (i.e. V α 14iNKT cells) and Type II NKT cells were resistant to APAP hepatotoxicity only if NK cells were also depleted [10].

V α 14iNKT cells are a major class of NKT cells that constitute 20–40% of the hepatic lymphocyte population [12]. These innate T cells swiftly produce pro-inflammatory cytokines and chemokines on activation [13]. V α 14iNKT cells have an invariant T cell receptor with a V α 14–J α 18 chain associated with a V β 2, V β 7, or V β 8.2 chain [13,14] and are directly stimulated by TCR and TCR-independent mechanisms. The specific role of V α 14iNKT cells in the development of APAP hepatotoxicity is incompletely defined. In the present study, we demonstrate that V α 14iNKT cells induce pro-inflammatory effects during APAP hepatotoxicity by regulating hepatic GSH production and APAP metabolism.

2. Materials and methods

2.1. Mice

Wildtype (WT) male C57BL/6 mice (5–7 weeks) were purchased from the Jackson Laboratory (Bar Harbor, ME). Breeding pairs of J α 18^{-/-} mice (on C57BL/6 background) provided by Dr. M. Taniguchi (RIKEN Research Center for Allergy & Immunology, Yokohama, Japan) [15] were bred in a pathogen-free breeding facility at LSUHSC-Shreveport [16]. All mice used in this study were continuously fed standard chow pellet diet, drinking water *ad libitum* and were maintained on a 12 h light/dark cycle. All experiments were performed in accordance with NIH guidelines and approved by LSUHSC-Shreveport Animal Care and User Committee.

2.2. Reagents

Acetaminophen (APAP; *N*-acetyl-*p*-aminophenol), DL-buthionine-[S,R]-sulfoximine (BSO), PBS, perchloric acid and protease cocktail inhibitor were all purchased from Sigma-Aldrich (St. Louis, MO). PVDF membrane and ECL western blotting reagent were purchased from Thermo Fisher Scientific (Rockford, IL) whereas ALT commercial kit was supplied by Thermo Electron (Waltham). Primary antibodies, malondialdehyde pAb and 3-nitrotyrosine mAb, were purchased from EMD (Darmstadt, DE) and Invitrogen (Camarillo, CA), respectively. GAPDH mAb was procured from Santa-Cruz Biotech (Santa Cruz, CA) and anti-mouse horseradish peroxidase secondary antibody was acquired from BD Biosciences (San Diego, CA). *In Situ* Cell Death Detection and Protein Assay kits were obtained from Roche Applied Science (Indianapolis, IN) and Bio-Rad Laboratories (Hercules, CA), respectively.

2.3. APAP hepatotoxicity and in vivo treatment protocol

Freshly prepared APAP (600 mg/kg, i.p.) in warm sterile PBS was administered to fed mice. Control mice received an equivalent volume of warm sterile PBS. At indicated time-points, mice were anesthetized with a mixture of xylazine and ketamine hydrochloride and blood serum were collected. Livers were then perfused with ice-cold sterile PBS (to remove blood elements) and harvested for the experimental assays described below. For hepatic GSH depletion, mice were administered BSO (500 mg/kg, i.p.) or sterile PBS,

2 h before APAP and 3 h thereafter until termination of the experiment to sustain GSH depletion [17].

2.4. Biochemical and histological liver injury

Acute liver injury was determined biochemically by measuring serum levels of the liver enzyme, alanine aminotransferase (ALT), using a commercial kit [16,18]. For histological evaluation, paraffin embedded liver sections (5 μ m thick) were deparaffinized, stained with H & E according to standard protocols and then analyzed by light microscopy in a blinded fashion by a pathologist (PAA). The degree of inflammation in the liver and hepatocyte damage was graded as mild, moderate or severe using a combination of the severity of the inflammation, and the degree of hepatocyte degenerative changes including ballooning degeneration, hepatocyte necrosis and frequency of acidophilic bodies [18].

2.5. GSH/GSSG analysis

Perfused livers were snap-frozen in liquid nitrogen immediately after excision from mice. Total hepatic GSH was determined by HPLC using a modified protocol of Reed et al. [19] as we previously described [18,19].

2.6. APAP metabolites analysis

Perfused liver samples were briefly homogenized in cold sterile PBS containing protease cocktail inhibitor and then centrifuged twice at 4 °C for 15 min at 1800g. Supernatants were filtered using a 0.45 μ m pore syringe filter and stored in –80 °C until use. Proteins in liver and serum supernatants were precipitated in 20% perchloric acid at 4 °C by centrifugation for 10 min at 10,000g. Next, APAP metabolites were separated using a Beckman Ultrasphere ODS C18 Reverse Phase column (80A) on a Shimadzu HPLC system as described by Howie et al. [20]. The concentration of APAP metabolites in liver and serum samples were measured based on the APAP standard phenolic ring absorbance at the wavelength of 195 nm [20]. Liver protein concentration was determined using preceding protocol.

2.7. Western Blot analysis

Perfused liver samples were processed and 30 μ g of protein were assayed in accordance with protocol previously described [18]. Primary antibodies were diluted in 5% milk at the following dilutions: malondialdehyde (1:1000) or nitrotyrosine (1:1000), incubated overnight in a cold room. Next, membranes were washed three times with PBS in Tween-20 and counterstained with corresponding secondary antibodies conjugated to horseradish peroxidase (1:1000). Membranes were visualized using Pierce ECL western blotting reagent and chemiluminescence film. Subsequently, all membranes were stripped in stripping buffer (0.08% mercaptoethanol, 0.5 mM Tris–HCl pH 6.8, 10% SDS) and reprobed with GAPDH mAb (1:1000) to verify equal protein loading in samples.

2.8. In situ analysis of liver apoptosis using TUNEL

Paraffin-embedded liver sections were dewaxed in xylene and rehydrated by passage through a graded series of ethanol solutions, and then PBS. Sections were treated with proteinase K (20 μ g/ml in 10 mM Tris–HCl, pH 7.4–8.0) at 37 °C for 15 min, washed and then stained with fluorescein nucleotide mixture with terminal deoxynucleotidyl transferase (TdT) from *In Situ* Cell Death Detection kit. Sections were viewed and photographed using standard fluorescent microscopic techniques [18].

2.9. Statistical analysis

Student unpaired *t* test was used for the comparison of means between two experimental groups. Comparison among three or more experimental groups was performed using a one-way ANOVA, followed by Newman–Keuls post hoc test. A value of $p < 0.05$ was considered significant. Densitometric image analysis was performed using ImageJ 1.43u program (NIH). All data are shown as mean \pm SEM.

3. Results

3.1. Resistance of $J\alpha 18^{-/-}$ mice to APAP liver toxicity

In preliminary experiments, we found that administration of APAP (600 mg/kg) to fed mice significantly increased liver injury as reflected by elevated serum ALT without causing mice mortality (data not shown). This dose was used in all experiments. Next, we assessed whether the presence of hepatic $V\alpha 14iNKT$ cells contribute to the development of APAP hepatotoxicity. In Fig. 1A, we show that APAP administration into WT mice caused a time-dependent significant increase in serum ALT levels at 8 and 24 h relative to naïve. In contrast, $J\alpha 18^{-/-}$ mice, which are selectively deficient in $V\alpha 14iNKT$ cells [15,21], were highly resistant to APAP hepatotoxicity as shown by almost complete significant suppression ($>80\%$ reduction) of serum ALT at 8 and 24 h (Fig. 1A). As expected, normal levels of serum ALT were observed in both naïve WT and $J\alpha 18^{-/-}$ mice (Fig. 1A). A representative histological liver section from WT mice at 24 h following APAP treatment revealed extensive hepatocyte apoptosis and necrotic damage compared to livers from $J\alpha 18^{-/-}$ mice which appeared almost normal (Fig. 1B). Interestingly, the absence of $V\alpha 14iNKT$ cells in $J\alpha 18^{-/-}$ mice was also associated with significantly higher GSH production in the liver at 8 and 24 h post-APAP contrary to corresponding WT mice (Fig. 1C). It is notable that GSH levels in naïve $J\alpha 18^{-/-}$ mice did not significantly differ from naïve WT mice. In summary, the resistance of $J\alpha 18^{-/-}$ mice to APAP hepatotoxicity is associated with elevated hepatic GSH.

3.2. GSH depletion enhances APAP-induced liver damage in $V\alpha 14iNKT$ cells deficient mice

In view of preceding results, we depleted hepatic GSH in $J\alpha 18^{-/-}$ mice prior to APAP administration to see if it enhanced liver injury. BSO, a specific inhibitor targeting the rate determining step in GSH synthesis, was used to deplete hepatic GSH. Hepatic GSH depletion in $J\alpha 18^{-/-}$ mice resulted in enhanced liver injury and mice mortality (60%) following APAP injection relative to any treatment group (Fig. 1D). In parallel, liver sections from these mice exhibited widespread necrosis, severe hemorrhage, and focal steatosis (Fig. 1E). These results suggest a protective role for GSH in the liver of $J\alpha 18^{-/-}$ mice during APAP hepatotoxicity.

3.3. Effect of $V\alpha 14iNKT$ cell deficiency on APAP metabolism in the liver

Considering the observed changes in hepatic GSH levels in $J\alpha 18^{-/-}$ mice during APAP hepatotoxicity, we also quantified hepatic APAP and its metabolites. Following APAP administration, we observed a time-dependent decrease in APAP content in the livers of WT and $J\alpha 18^{-/-}$ mice (Fig. 2A). Notably, APAP content in the liver of $J\alpha 18^{-/-}$ mice was significantly lower at 2 h compared to corresponding WT mice (Fig. 2A). In parallel, we observed augmented hepatic APAP-GSH levels in $J\alpha 18^{-/-}$ mice at 2 h in contrast to corresponding WT mice (Fig. 2B). In Fig. 2C, we also demonstrate that the level of the APAP metabolite, APAP-glucuronide, in the liver of

$J\alpha 18^{-/-}$ mice was significantly reduced at both the 1 and 2 h time-points relative to corresponding WT mice. No significant changes in hepatic APAP-sulfate levels were observed in the liver of WT and $J\alpha 18^{-/-}$ mice throughout the time-course (Fig. 2D).

3.4. Effect of $V\alpha 14iNKT$ cell deficiency on serum APAP and metabolite levels

$J\alpha 18^{-/-}$ mice had significantly lower serum APAP levels at 0.5 and 2 h when compared to corresponding WT mice (Fig. 3A). In contrast to our finding in the liver, no significant change in serum APAP-GSH levels was observed in $J\alpha 18^{-/-}$ mice relative to WT mice throughout the time-course (Fig. 3B). When APAP-GSH is transported outside the cell, it undergoes further metabolism and can form APAP-mercapturate at principal sites of acetylation like the kidneys [22]. At 4 h post APAP administration, we identified significantly high APAP-mercapturate serum levels in $J\alpha 18^{-/-}$ mice vs. corresponding WT mice (Fig. 3C). Serum APAP-glucuronide levels significantly decreased at 0.5–2 h in $J\alpha 18^{-/-}$ mice vs. WT mice (Fig. 3D) whereas serum APAP-sulfate levels were significantly reduced in $J\alpha 18^{-/-}$ mice over WT mice at 2 h post-APAP administration (Fig. 3E).

4. Discussion

$V\alpha 14iNKT$ cells are the most abundant innate T lymphocytes in the liver representing 80–90% of the NKT cell population [12]. During inflammatory responses, $V\alpha 14iNKT$ cells are rapidly activated by TCR-dependent and independent mechanisms to release pro-inflammatory cytokines (e.g. IFN- γ , TNF- α) and chemokines (including CCL5, CCL3) [13]. These pathophysiological responses recruit and/or stimulate intrahepatic leukocytes to produce reactive species, ROS and RNS [13,23,24]. Consequently, $V\alpha 14iNKT$ cells have an important immunoregulatory role representing a critical link between the innate and adaptive immune systems. $V\alpha 14iNKT$ cells have been implicated in the pathophysiology of many models of liver inflammation and injury mediated by ischemia/reperfusion [21], microbes [24], alcohol [25] and adenovirus [16]. The specific contribution of $V\alpha 14iNKT$ cells to the development of APAP-mediated liver injury is incompletely defined.

In the current study, we provide proof-of-principle that highlights a central role for $V\alpha 14iNKT$ cells in suppressing GSH formation in the liver to promote the development of APAP hepatotoxicity. We initially demonstrated that mice deficient in $V\alpha 14iNKT$ cells were resistant to liver injury after APAP administration since biochemical and histological markers of liver injury were significantly reduced ($>80\%$) relative to the levels in WT mice. This observation is in contrast to a previous report [10] using the $CD1d^{-/-}$ mice, which are deficient in both $V14iNKT$ cells and Type II NKT cells. Specifically, $CD1d^{-/-}$ mice were found to be equally susceptible to liver injury as the corresponding WT mice following APAP administration [10]. Interestingly, liver injury in these mice were only alleviated if NK cells were also depleted [10]. NKT and NK cells are known to share similar phenotypic and functional characteristics including cell surface markers, cytokine production and cytotoxicity [10,13]. Therefore, it has been suggested that the deficiency of either NKT or NK cells could be compensated by the cell type present to induce hepatocyte death during APAP intoxication. However, our study undeniably revealed a novel and essential pathophysiological role for $V\alpha 14iNKT$ cells in promoting APAP hepatotoxicity. Next, we evaluated why $V\alpha 14iNKT$ cell deficient mice, $J\alpha 18^{-/-}$ mice, were resistant to APAP liver toxicity. We speculated that high GSH levels in $J\alpha 18^{-/-}$ mice relative to WT mice may underlie the hepatoprotective effects documented in $J\alpha 18^{-/-}$ mice following APAP administration. It is notable that GSH is an




Fig. 1. Time course for APAP hepatotoxicity in Jα18^{-/-} mice and WT mice. (A) Serum ALT levels, (B) Representative H & E staining of liver sections from WT and Jα18^{-/-} mice and (C) HPLC measurement of hepatic GSH levels in WT and Jα18^{-/-} mice before and after APAP treatment. Data in (A) and (C) are presented as mean ± SEM with $n = 6-12$ mice/group with * $p < 0.05$ vs. naïve WT; # $p < 0.05$ vs. corresponding WT as assessed by ANOVA/Newman-Keuls post hoc test. (D, E) Effect of GSH depletion on APAP liver toxicity in Jα18^{-/-} mice. BSO or PBS was injected in WT and Jα18^{-/-} mice (see methods) for assessment of serum ALT levels (D) and Histological injury by H & E staining of liver sections (E). Data in D is shown as mean ± SEM with $n = 7-19$ mice/group with * $p < 0.05$ vs. WT/PBS; # $p < 0.05$ vs. WT/PBS/APAP; $p < 0.05$ vs. Jα18^{-/-}/PBS/APAP.

effective and potent ROS and RNS scavenger shown to suppress/prevent APAP hepatotoxicity [2,26]. Indeed, we documented elevated hepatic levels of GSH in Jα18^{-/-} mice vs. WT mice post-APAP administration. Additionally, we specifically established a hepato-

protective role for hepatic GSH in Jα18^{-/-} mice post-APAP treatment since depletion of hepatic GSH in Jα18^{-/-} mice using BSO significantly exacerbated liver damage (~8-fold) and caused mortality (60%) over APAP-treated Jα18^{-/-} mice administered vehicle.




Fig. 2. Metabolic shift of hepatic APAP and metabolites levels from Jα18^{-/-} mice vs. WT mice during APAP liver toxicity. (a) APAP levels, (b) APAP-GSH levels, (c) APAP-Glucuronide levels and (d) APAP-Sulfate levels. All results are presented as mean ± SEM with *n* = 6–12 mice/group; **p* < 0.05 vs. naive WT; #*p* < 0.05 vs. corresponding WT.

Taken together, we propose that Vα14iNKT cells deficiency positively regulate GSH production in the liver to dampen APAP liver toxicity in Jα18^{-/-} mice.

Given that hepatic GSH induced a hepatoprotective in Jα18^{-/-} mice, we next measured APAP and its metabolites to assess any changes in APAP metabolism. We observed a significant increase in hepatic APAP-GSH and serum APAP-mercapturate levels during Vα14iNKT cells deficiency that correlated with a significant decrease in APAP levels in Jα18^{-/-} mice compared to the WT mice. Our results indicate that hepatic GSH detoxify APAP metabolites during Vα14iNKT cells deficiency thereby reducing liver damage in Jα18^{-/-} mice. Differences between GSH and APAP-related conjugate levels at early time-points as seen in WT and Jα18^{-/-} mice can result from hepatic protein glutathionylation in centrilobular region and biliary excretion [27,28]. For this reason, we also assessed APAP-glucuronide and APAP-sulfate levels. Glucuronyl transferase activity is known to be regulated by signaling pathways (e.g. Nrf2, PKC, MAPK) during APAP intoxication [29,30]. We observed a significant decrease in serum and hepatic APAP-glucuronide levels Jα18^{-/-} mice compared to WT mice. In addition, serum but not hepatic APAP-sulfate levels were significantly decreased in Jα18^{-/-} mice vs. WT mice. Our data suggests that Jα18^{-/-} mice have a decreased dependency to form APAP-sulfate in extrahepatic tissues like the kidneys [4,22]. On the basis of these findings, we propose that the enrichment of GSH in the liver of Jα18^{-/-} mice due to a deficiency in Vα14iNKT cells was related

to a shift in APAP metabolism in favor of GSH conjugation to NAPQI that suppressed liver damage.

GSH is known to effectively scavenge ROS and RNS through enzymatic (e.g. GSH peroxidase) and non-enzymatic reactions that minimizes oxidative and nitrosative stress as well as liver toxicity in response to APAP administration [2,26]. In agreement, GSH depletion in Jα18^{-/-} mice using BSO during APAP hepatotoxicity enhanced oxidative stress (i.e. MDA-protein adducts), nitrosative stress (i.e. nitrotyrosine) and liver necrosis. Conversely, high GSH levels in Jα18^{-/-} mice due to Vα14iNKT cells deficiency was associated with significantly reduced oxidative stress and DNA fragmentation linked to a decrease in liver damage relative to WT mice with similar treatments. All APAP treatments had high levels of nitrotyrosine staining, which is a reliable footprint of peroxynitrite [2,8,31]. Intriguingly, MDA and nitrotyrosine-protein adducts (~60–200 kDa) have been reported to resemble albumin monomers and aggregates suggesting that the blood components form adducts with products of reactive species [32,33]. RNS can elicit DNA fragmentation and is known to inhibit P450 enzymes, caspases, and cytochrome oxidase [34]. This ability of RNS to inhibit caspase activity associates with the lack of DNA fragmentation during GSH depletion accompanying APAP hepatotoxicity.

In summary, our results revealed that Vα14iNKT cells exert pro-inflammatory and pathophysiological responses during APAP hepatotoxicity by suppressing hepatic GSH formation to alter APAP metabolism. Specifically, mice deficient in Vα14iNKT cells had high




Fig. 3. Evaluation of serum APAP and metabolites levels in WT and Jα18^{-/-} mice by HPLC during APAP hepatotoxicity. (A) APAP, (B) APAP-GSH, (C) APAP-Mercapturate, (D) APAP-Glucuronide levels and (E) APAP-Sulfate levels. All data are presented as mean ± SEM with $n = 3-9$ mice/group with * $p < 0.05$ vs. naive WT; # $p < 0.05$ vs. corresponding WT.

hepatic GSH levels that detoxified NAPQI to attenuate oxidative stress and the development of liver necrosis.

Acknowledgment

This study was supported by NIH grants to MNA (NIAID R56AI085150) and TYA (NIDDK DK44510).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbrc.2012.10.029>.

References

- [1] A.M. Larson, Acetaminophen hepatotoxicity, *Clin. Liver Dis.* (11) (2007) 525–548.
- [2] J.A. Hinson, D.W. Roberts, L.P. James, Mechanisms of acetaminophen-induced liver necrosis, *Handb. Exp. Pharmacol.* (196) (2010) 369–405.
- [3] Z. Wu, M. Han, T. Chen, W. Yan, Q. Ning, Acute liver failure: mechanisms of immune-mediated liver injury, *Liver Int.* (30) (2010) 782–794.
- [4] L. Liu, C.D. Klaassen, Different mechanism of saturation of acetaminophen sulfate conjugation in mice and rats, *Toxicol. Appl. Pharm.* (139) (1996) 128–134.
- [5] X. Wang, Y. Lu, A.I. Cederbaum, Induction of cytochrome P450 2E1 increases hepatotoxicity caused by fas agonistic Jo2 antibody in mice, *Hepatology* (42) (2005) 400–410.
- [6] N. Hanawa, N. Shinohara, B. Saberi, W.A. Gaarde, D. Han, N. Kaplowitz, Role of JNK translocation to mitochondria leading to inhibition of mitochondria bioenergetics in acetaminophen-induced liver injury, *J. Biol. Chem.* (283) (2008) 13565–13577.
- [7] Y. Ishida, T. Kondo, T. Ohshima, H. Fujiwara, Y. Iwakura, N. Mukaida, A pivotal involvement of IFN-γ in the pathogenesis of acetaminophen-induced acute liver injury, *FASEB J.* (16) (2002) 1227–1236.
- [8] S.L. Michael, N.R. Pumford, P.R. Mayeux, M.R. Niesman, J.A. Hinson, Pretreatment of mice with macrophage inactivators decreases acetaminophen hepatotoxicity and the formation of reactive oxygen and nitrogen species, *Hepatology* (30) (1999) 186–195.
- [9] K. Numata, M. Kubo, H. Watanabe, K. Takagi, H. Mizuta, S. Okada, et al., Overexpression of suppressor of cytokine signaling-3 in T cells exacerbates acetaminophen-induced hepatotoxicity, *J. Immunol.* (178) (2007) 3777–3785.
- [10] Z.X. Liu, S. Govindarajan, N. Kaplowitz, Innate immune system plays a critical role in determining the progression and severity of acetaminophen hepatotoxicity, *Gastroenterology* (127) (2007) 1760–1774.
- [11] M.J. Masson, L.D. Carpenter, M.L. Graf, L.R. Pohl, Pathogenic role of natural killer T and natural killer cells in acetaminophen-induced liver injury in mice is dependent on the presence of dimethyl sulfoxide, *Hepatology* (48) (2008) 889–897.
- [12] G. Eberl, R. Lees, S.T. Smiley, M. Taniguchi, M.J. Grusby, H.R. MacDonald, Tissue-specific segregation of CD1d-dependent and CD1d-independent NK T cells, *J. Immunol.* (162) (1999) 6410–6419.
- [13] J.L. Matsuda, T. Mallevaey, J. Scott-Browne, L. Gapin, CD1d-restricted iNKT cells the ‘Swiss-Army knife’ of the immune system, *Curr. Opin. Immunol.* (20) (2008) 358–368.
- [14] D.I. Godfrey, H.R. MacDonald, M. Kronenberg, M.J. Smyth, L. Van Kaer, NKT cells: what’s in a name?, *Nature Rev.* (4) (2004) 231–237.
- [15] J. Cui, T. Shin, T. Kawano, H. Sato, E. Kondo, I. Toura, et al., Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors, *Science* (278) (1997) 1623–1626.
- [16] M.N. Ajuebor, Q. Chen, R.M. Strieter, P.A. Adegboyega, T.Y. Aw, V(α)14iNKT cells promote liver pathology during adenovirus infection by inducing CCL5 production: implications for gene therapy, *J. Virol.* (84) (2010) 8520–8529.
- [17] H.M. Shen, W.X. Ding, C.N. Ong, Intracellular glutathione is a cofactor in methylseleninic acid-induced apoptotic cell death of human hepatoma HEPG(2) cells, *Free Radical Biol. Med.* (33) (2002) 552–561.
- [18] I. Downs, J. Liu, P.A. Adegboyega, T.Y. Aw, M.N. Ajuebor, The ROS scavenger NAC regulates hepatic Vα14iNKT cells signaling during Fas mAb-dependent fulminant liver failure, *PLoS ONE* (7) (2012) 1–11.
- [19] D.J. Reed, J.R. Babson, P.W. Beatty, A.E. Brodie, W.W. Ellis, D.W. Potter, High-performance liquid chromatography analysis of nanomole levels of glutathione glutathione disulfide and related thiols and disulfides, *Anal. Biochem.* (106) (1980) 55–62.
- [20] D. Howie, P.I. Adriaenssens, L.F. Prescott, Paracetamol metabolism following overdose: application of high performance liquid chromatography, *J. Pharm. Pharmacol.* (29) (1977) 235–237.
- [21] P. Arrenberg, I. Maricic, V. Kumar, Sulfatide-mediated activation of type II natural killer T cells prevents hepatic ischemic reperfusion injury in mice, *Gastroenterology* (140) (2011) 646–655.
- [22] C.A. Hinchman, H. Matsumoto, T.W. Simmons, N. Ballatori, Intrahepatic conversion of a glutathione conjugate to its mercapturic acid. Metabolism of

- 1-chloro-2,4-dinitrobenzene in isolated perfused rat and guinea pig livers, *J. Biol. Chem.* (266) (1991) 22179–22185.
- [23] H. Nakashima, M. Kinoshita, M. Nakashima, Y. Habu, S. Shono, T. Uchida, et al., Superoxide produced by kupffer cells is an essential effector in concanavalin A-induced hepatitis in mice, *Hepatology* (48) (2008) 1979–1988.
- [24] H. Ohtaki, H. Ito, K. Ando, T. Ishikawa, K. Saito, M. Imawari, et al., Valpha14 NKT cells activated by alpha-galactosylceramide augment lipopolysaccharide-induced nitric oxide production in mouse intra-hepatic lymphocytes, *Biochem. Biophys. Res. Commun.* (378) (2009) 579–583.
- [25] M. Minagawa, Q. Deng, Z.X. Liu, H. Tsukamoto, G. Dennert, Activated natural killer T cells induce liver injury by Fas and tumor necrosis factor-alpha during alcohol consumption, *Gastroenterology* (126) (2004).
- [26] C. Saito, C. Zwingmann, H. Jaeschke, Novel mechanisms of protection against acetaminophen hepatotoxicity in mice by glutathione and N-acetylcysteine, *Hepatology* (51) (2010) 246–254.
- [27] C. Chen, G.E. Hennig, J.E. Manautou, Hepatobiliary excretion of acetaminophen glutathione conjugate and its derivatives in transport-deficient (TR-) hyperbilirubinemic rats, *Drug Metab. Dispos.: The Biol. Fate Chem.* (31) (2003) 798–804.
- [28] X.G.J.A. Yang, A. Shi, Q. Roberts, D.W. Hinson, J.A. Muskheishvili, L. Beger, R. Pence, L.M. Ando, Y. Sun, J. Davis, K. Salminen, Changes in mouse liver protein glutathionylation after acetaminophen exposure, *J. Pharmacol. Exp. Ther.* (340) (2011) 360–369.
- [29] A. Enomoto, K. Itoh, E. Nagayoshi, J. Haruta, T. Kimura, T. O'Connor, et al., High sensitivity of Nrf2 knockout mice to acetaminophen hepatotoxicity associated with decreased expression of ARE-regulated drug metabolizing enzymes and antioxidant genes, *Toxicol. Sci.* (59) (2001) 169–177.
- [30] L. Volak, M. Court, Role for protein kinase C delta in the functional activity of human UGT1A6: implications for drug-drug interactions between PKC inhibitors and UGT1A6, *Xenobiotica* (40) (2010) 306–318.
- [31] S. Burney, J.L. Caulfield, J.C. Niles, J.S. Wishnok, S.R. Tannenbaum, The chemistry of DNA damage from nitric oxide and peroxynitrite, *Mutat. Res.* (424) (1999) 37–49.
- [32] K. Houghlum, M. Filip, J.L. Witztum, M. Chojkier, Malondialdehyde and 4-hydroxynonenal protein adducts in plasma and liver of rats with iron overload, *J. Clin. Invest.* (86) (1990) 1991–1998.
- [33] J.L. Wayenberg, V. Ransy, D. Vermeylen, E. Damis, S.P. Bottari, Nitrated plasma albumin as a marker of nitrative stress and neonatal encephalopathy in perinatal asphyxia, *Free Radical Biol. Med.* (1) (2009) 975–982.
- [34] T. Chen, R. Zamora, B. Zuckerbraun, T.R. Billiar, Role of nitric oxide in liver injury, *Curr. Mol. Med.* (3) (2003) 519–526.