



Construction and direct electrochemistry of orientation controlled laccase electrode



Ying Li^a, Jiwei Zhang^b, Xirong Huang^{a,b,*}, Tianhong Wang^{b,*}

^a Key Laboratory of Colloid & Interface Chemistry of the Education Ministry of China, Shandong University, Jinan 250100, PR China

^b State Key Laboratory of Microbial Technology of China, Shandong University, Jinan 250100, PR China

ARTICLE INFO

Article history:

Received 10 February 2014

Available online 26 February 2014

Keywords:

Recombinant laccase

Gold electrode

Oriented immobilization

Direct electron transfer

Electron transfer pathway

ABSTRACT

A laccase has multiple redox centres. Chemisorption of laccases on a gold electrode through a polypeptide tag introduced at the protein surface provides an isotropic orientation of laccases on the Au surface, which allows the orientation dependent study of the direct electrochemistry of laccase. In this paper, using genetic engineering technology, two forms of recombinant laccase which has Cys-6×His tag at the N or C terminus were generated. Via the Au-S linkage, the recombinant laccase was assembled orientationally on gold electrode. A direct electron transfer and a bioelectrocatalytic activity toward oxygen reduction were observed on the two orientation controlled laccase electrodes, but their electrochemical behaviors were found to be quite different. The orientation of laccase on the gold electrode affects both the electron transfer pathway and the electron transfer efficiency of O₂ reduction. The present study is helpful not only to the in-depth understanding of the direct electrochemistry of laccase, but also to the development of laccase-based biofuel cells.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

Laccase is a copper-containing oxidase [1,2]. Each laccase protein contains four copper ions, which are divided into T1 (one), T2 (one) and T3 (two) categories according to their spectral characteristics. The T1 site is for aromatic substrate oxidation and T2 and T3 copper ions form a trinuclear copper cluster active site for oxygen reduction. During the catalytic oxidation of substrates, one electron from the substrate is transferred to the T1 site, and subsequently it passes through the intramolecular His-Cys-His bridge to the T2/T3 cluster where molecular oxygen is reduced to water instead of the intermediate H₂O₂. Due to these unique properties, laccase has been tried to construct biofuel cells [3–5]. However, the direct electron transfer (DET) of laccase on conventional electrodes is found to be difficult. It depends upon the electrode materials and laccase immobilization strategies [3,6–10]. It has been found that the electrocatalytic reduction of oxygen at non-oriented laccase electrode is accompanied by the production of the intermediate H₂O₂, which decreases the efficiency of the O₂ reduction [11]. It follows that the oriented immobilization of

laccase is a prerequisite for understanding the above phenomena and also it is the difficult point for studying the DET of laccase.

The site-directed biological modification of functional proteins is an ideal approach to achieving an oriented assembly of the proteins [12–17]. The basic strategy is to use genetic engineering technology to modify functional proteins with linkers which provide oriented attachment of the proteins on a specific support. The Au-S bond is usually used to fabricate an assembled film on Au electrode. Therefore, the introduction of cysteine at the N or C-terminus of laccase peptide chain facilitates the oriented immobilization of laccase. To obtain a recombinant laccase with Cysteine-6×Histidine (Cys-6×His) tag at the N or C terminus (named rLacA-N and rLacA-C, respectively), a gene sequence encoding the Cys-6×His tag was fused in the present study to the 3'/5' end of the *lacA* gene from *Trametes* sp. AH 28-2. The obtained recombinant laccase was assembled orientationally on a gold electrode via the Au-S bond. The orientation dependent direct electron transfer of laccase and its electrocatalysis toward oxygen reduction were then investigated and compared. The present study is helpful to the in-depth understanding of the direct electrochemistry of laccase and to the development of laccase-based biofuel cells.

2. Materials and methods

The construction of the expression vector of rLacA with Cys-6×His tagged at the N or C terminus (named pPlacA-N_{Cys-His} and

* Corresponding authors. Address: Key Laboratory of Colloid & Interface Chemistry of the Education Ministry of China, Shandong University, Jinan 250100, PR China. Fax: +86 531 88365433 (X. Huang). Fax: +86 531 88366118 (T. Wang).

E-mail addresses: xrhuang@sdu.edu.cn (X. Huang), wangtianhong@sdu.edu.cn (T. Wang).

pLacA-C_{Cys-His}, respectively) and its heterologous expression in *Pichia pastoris* strain GS115 (Invitrogen, Carlsbad, CA) were performed according to the Xiao's method [18]. The PCR with designed primers that contained the sequence of the tag (the underlined) was carried on the plasmid pLacA (kindly provided by Dr. Xiao et al. [19]) by standard procedures. For the construction of pLacA-N_{Cys-His}, the upstream and downstream primers were 5'-CCGgaattcTGCCATCATCACCATCATCACGCCATTGGGCCCCACCGCTGACCTCA-3' and 5'-AATAgcggccgcCTGGTCGTTGACATCGAGCGCG-3', respectively. For the construction of pLacA-C_{Cys-His}, the two primers were 5'-CCGgaattcGCCATTGGGCCCCACCGCTGACCTCA-3' and 5'-AATAgcggccgcGCAGTGATGATGGTGATGATGCTGGTCGTTGACATCGAGCGCG-3', respectively. Sequence analyses of the target expression vectors (pLacA-N_{Cys-His} and pLacA-C_{Cys-His}) were performed.

The laccase producing transformants were cultured under the conditions described in the manual of the *Pichia* Expression Kit (Invitrogen). To harvest the expressed rLacA, the cells were first removed by centrifugation and the supernatant was then concentrated with a 10 kDa cut-off centrifugal filtering device (Amicon Ultra-15, Millipore Corp. MA). The resulting rLacA was purified with a His-Trap FF column (GE Healthcare) at 4 °C. The purification was monitored by SDS-PAGE. The activity of purified laccases (both free and immobilized) were determined spectrophotometrically [18] and one activity unit (U) was defined as the amount of the enzyme required to oxidize 1 μmol 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) per minute at pH 4.0 at 30 °C.

The rLacA modified electrode was prepared by immersing a pre-treated gold disc electrode [20] (3 mm in diameter) into the enzyme solution at 4 °C for 24 h, followed by rinsing with phosphate buffer (100 mM, pH 6.0). The immobilized laccase was determined using quartz crystal microbalance (ANT Q300 affinity sensor, Taiwan). Cyclic voltammetry was performed using a CHI 630 Electrochemical Analyzer (Shanghai). A bare or laccase modified gold electrode was used as a working electrode, a platinum wire and a saturated calomel electrode (SCE) were used as counter electrode and reference electrode, respectively. Unless otherwise specified, the electrolyte phosphate buffer (100 mM, pH 4.6) was thoroughly deoxygenated with pure N₂ for ca. 30 min prior to experiments and the N₂ atmosphere was kept during the experiments. The CVs were recorded over a broad potential range and with different scan rates. The content of H₂O₂ formed during the electrocatalytic reduction of O₂ was measured according to the method described elsewhere [11].

3. Results and discussion

Sequence analysis indicated that the cDNA fragment of *lacA* with Cys-6×His tag in N or C-terminus was correctly inserted into pPIC9K, resulting in two 15.4 kb plasmids without gene mutation. The gene was expressed in *P. pastoris* with an expression level of 35.9 mg L⁻¹ (3.04×10^4 U L⁻¹) for rLacA-N and 19.9 mg L⁻¹ (0.83×10^4 U L⁻¹) for rLacA-C, indicating that the heterologous expression of the rLacA was successful.

SDS-PAGE experiments indicated that the separated rLacA-N and rLacA-C were both pure with molecular weight estimated to be both 64 kDa, which is very close to that of the native LacA (nLacA) from *Trametes* sp. AH28-2 (62 kDa) [19]. The apparent kinetic parameters of K_m and k_{cat} , determined with ABTS as substrate, are 16.7 μM and 68.6 s⁻¹ for rLacA-N, and 17.2 μM and 65.3 s⁻¹ for rLacA-C, respectively. These data suggest that there is little difference between rLacA-N and rLacA-C in the catalytic performance and the insertion of Cys-6×His tag has no notable effect on the conformation of laccase.

The structure of LacA from *Trametes* sp. AH 28-2 is not reported, but that of LacB from the same strain has been released (Protein Data Bank code 3KW7) [21]. The structure of rLacA was built via homology modelling using the SWISS-MODEL function in Swiss Pdb viewer version 3.7 [22]. Its structure and probable orientation on gold surface (the rLacA was immobilized via the Au-S bond) is schematically represented in Fig. 1.

In rLacA-N, the T1 site is away from the Au electrode surface, while the T2/T3 site is near the electrode surface; in rLacA-C, however, the distance between the T1 and the electrode surface is nearly the same as that between the T2/T3 and the electrode surface. It follows that the difference in site for tag modification results in different orientations of laccase on the gold surface.

The amount of immobilized laccase on the gold surface shows that the unmodified laccase is not readily adsorbed on the gold surface, while the modified laccase is easily immobilized on the gold surface due to the thiol from the Cys-6×His tag. The surface coverages of the two modified laccases are $(5.41 \pm 0.28) \times 10^{-12}$ mol cm⁻² for rLacA-N and $(4.99 \pm 0.19) \times 10^{-12}$ mol cm⁻² for rLacA-C, respectively. These data are close to the theoretical value of 4.65×10^{-12} mol cm⁻² calculated based on the monolayer assembly of laccase [7]. The slight difference in the coverage may be due to differences in orientation, transverse sectional area and steric hindrance, etc.

The measurements of the activity of the immobilized laccase shows that after chemisorption of rLacA on the Au surface, the




Fig. 1. Schematic representation of rLacA immobilized orientationally on bare gold surface with Cys-6×His tag at its N-terminus (A) or C-terminus (B).

two modified laccases are still active with specific activities of 174.4 U mg^{-1} for rLacA-N and 162.8 U mg^{-1} for rLacA-C, respectively. The activity maintenance should be attributed to the linker, which is projecting on the outer surface and away from the active centre.

Fig. 2 shows the cyclic voltammograms (CVs) of different laccase modified gold electrodes in phosphate buffer (pH 4.6). Under the same conditions, no peak was observed at the native laccase modified electrode over the potential range, but a pair of well-defined, quasi-reversible redox peaks were observed for rLacA-N modified gold electrode with $E_{pc} = 289 \text{ mV vs. NHE}$ and $E_{pa} = 465 \text{ mV vs. NHE}$. The obtained formal potential ($E_0 = 377 \text{ mV vs. NHE}$) at rLacA-N modified gold electrode is very close to the potential at T2/T3 cluster reported elsewhere [11], i.e., this pair of redox peaks could be attributed to the direct electron transfer between the T2/T3 cluster of laccase and the electrode. The homology modeling supports this inference (see Fig. 1A). For rLacA-C modified gold electrode, a pair of well-defined but irreversible peaks were observed with $E_{pc} = 316 \text{ mV vs. NHE}$ and $E_{pa} = 624 \text{ mV vs. NHE}$. Its formal potential (470 mV vs. NHE) is higher than that of rLacA-N. The broad peaks may be ascribed to the combined effects of the T1 and T2/T3 copper ions. Based on the presumed orientation of rLacA-C on the electrode surface, there should exist two possible pathways of the electron transfer between the electrode and rLacA-C (T1 and T2/T3 cluster). Since the potential at T1 is higher than that at the T2/T3 cluster, the half-peak widths of both the overlapped anodic peak and the cathodic one increase (the corresponding onset potentials support the inference). The irreversibility of the redox peaks at the rLacA-C modified electrode may also correlate with the unidirectional intramolecular electron transfer from T1 to T2/T3 cluster [11]. It is worth mentioning that no signal at the native laccase/Au electrode is due to small amounts of immobilized laccase on the electrode.

For rLacA-N and rLacA-C modified gold disk electrodes, their peak currents were found to vary linearly with the scan rate over a scan rate range of $5\text{--}100 \text{ mV s}^{-1}$, indicating that the electron transfer process is surface-confined. For a surface-confined process, the following equation can be used to estimate the number of electrons transferred:

$$I_p = n^2 F^2 v A \Gamma / 4RT$$

where the symbols have their usual meanings. For rLacA-C and rLacA-N, the calculated n values are 1.82 and 0.89, respectively. That means that for each laccase immobilized orientationally on the Au surface, the number of electrons transferred as well as the DET pathway is closely related with the orientation of the recombinant laccase. Different orientation results in different pathway and different DET efficiency.

Fig. 3 shows the cyclic voltammograms of rLacA modified gold electrodes under aerobic condition. Compared with CVs recorded under anaerobic condition, the cathodic peak currents in the presence of O_2 are increased, especially for rLacA-C. It was also observed that the onset potential for oxygen reduction at the rLacA-C modified electrode (495 mV vs. NHE) was more positive than that at the rLacA-N modified electrode (340 mV vs. NHE). These results indicate that both recombinant laccases have electrocatalytic activities for oxygen reduction, and that their difference in electrocatalytic activity should be related to the orientation of laccase on the electrode and the active sites of laccase involved in the electron transfer.

It has been reported that the electrocatalytic reduction of O_2 at non-oriented laccase electrode is accompanied by the production of intermediate H_2O_2 [23]. To explore the relationship between the orientation of laccase on Au electrode and the DET pathway, the amount of H_2O_2 formed in the electrolyte was determined after repeated cycles of CV scan over a potential range of $142\text{--}742 \text{ mV vs. NHE}$. The obtained H_2O_2 concentration was determined to be $2.75 \mu\text{M}$ and $7.53 \mu\text{M}$ for rLacA-C and rLacA-N modified gold electrodes, respectively. This means that the amount of H_2O_2 produced at the rLacA-C modified electrode is much less than that produced at the rLacA-N modified electrode although the maximum catalytic current at rLacA-C modified electrode is approximately two times as large as that at rLacA-N modified electrode. Because the specific activities of the immobilized laccases are close to each other, the significant difference in the electrochemical behavior must have been caused by their orientation on the Au surface. Based on the simulated orientation of the two laccases on the Au surface, the T1 site of rLacA-N is far from the electrode and the electron transfer takes place mostly between the Au electrode and the T2/T3 site of rLacA-N. For rLacA-C, however, its T1 site is much closer to the Au surface than that of rLacA-N; moreover, as predicted, the T1 site of rLacA-C is closer than its T2/T3 site to the electrode




Fig. 2. Cyclic voltammograms (second cycle) of recombinant laccase (rLacA-C (a) and rLacA-N (b)) and nonrecombinant laccase (c) modified gold electrodes in 0.1 M phosphate buffer (pH 4.6) under anaerobic conditions at a scan rate of 20 mV s^{-1} .




Fig. 3. Cyclic voltammograms (second cycle) of rLacA-N (dotted line) or rLacA-C (solid line) immobilized gold electrode under air saturated conditions in 0.1 M pH 4.6 phosphate buffer. Scan rate: 20 mV s⁻¹.




Fig. 4. Schematic representation of the electron transfer pathway for the electrocatalytic reduction of O₂ by rLacA-N (A) and rLacA-C (B) immobilized orientationally on gold surface.

surface. So rLacA-C has two possible pathways for its communication with gold electrode (see Fig. 4). Based on the present electrocatalytic behavior of the two recombinant laccases on the Au surface, it could be concluded that for rLacA-N, the electron from the electrode is transferred directly to the T2/T3 site where O₂ is mostly reduced to H₂O₂; for rLacA-C, however, there exists another route, i.e., the electron from the electrode transfers via the T1 site to the T2/T3 site where O₂ is reduced to H₂O. It follows that the orientation of laccase on Au electrode affects both the electron transfer pathway and the electron transfer efficiency of O₂ reduction.

The two recombinant laccases, rLacA-N and rLacA-C, obtained using genetic engineering technology, were demonstrated to have similar enzymatic properties, but their electrochemical behaviors were shown to be quite different when they were orientationally immobilized on a bare gold surface. The orientation of laccase on Au electrode affects both the electron transfer pathway and the electron transfer efficiency of O₂ reduction.

Acknowledgments

The authors gratefully acknowledge the financial support from the State Key Laboratory of Microbial Technology of China, the National Natural Science Foundation of China (20973103 and

21173133), and the National Basic Research Program of China (2011CB707400).

References

- [1] E.I. Solomon, U.M. Sundaram, T.E. Machonkin, Multicopper oxidases and oxygenases, *Chem. Rev.* 96 (1996) 2563–2605.
- [2] S. Shleev, J. Tkac, A. Christenson, T. Ruzgas, A.I. Yaropolov, J.W. Whittaker, L. Gorton, Direct electron transfer between copper-containing proteins and electrodes, *Biosens. Bioelectron.* 20 (2005) 2517–2554.
- [3] C. Gutiérrez-Sánchez, M. Pita, C. Vaz-Dominguez, S. Shleev, A.L. De Lacey, Gold nanoparticles as electronic bridges for laccase-based biocathodes, *J. Am. Chem. Soc.* 134 (2012) 17212–17220.
- [4] J. Martínez-Ortiz, R. Flores, R. Vazquez-Duhalt, Molecular design of laccase cathode for direct electron transfer in a biofuel cell, *Biosens. Bioelectron.* 26 (2011) 2626–2631.
- [5] M. Pita, C. Gutiérrez-Sánchez, D. Olea, M. Velez, C. García-Diego, S. Shleev, V.M. Fernández, A.L. De Lacey, High redox potential cathode based on laccase covalently attached to gold electrode, *J. Phys. Chem. C* 115 (2011) 13420–13428.
- [6] V. Balland, C. Hureau, A.M. Cusano, Y. Liu, T. Tron, B. Limoges, Oriented immobilization of a fully active monolayer of histidine-tagged recombinant laccase on modified gold electrodes, *Chem. Eur. J.* 14 (2008) 7186–7192.
- [7] H. Qiu, C. Xu, X. Huang, Y. Ding, Y. Qu, P. Gao, Adsorption of laccase on the surface of nanoporous gold and the direct electron transfer between them, *J. Phys. Chem. C* 112 (2008) 14781–14785.
- [8] C. Vaz-Dominguez, S. Campuzano, O. Rüdiger, M. Pita, M. Gorbacheva, S. Shleev, V.M. Fernandez, A.L. De Lacey, Laccase electrode for direct electrocatalytic reduction of O₂ to H₂O with high-operational stability and resistance to chloride inhibition, *Biosens. Bioelectron.* 24 (2008) 531–537.

- [9] H. Qiu, C. Xu, X. Huang, Y. Ding, Y. Qu, P. Gao, Immobilization of laccase on nanoporous gold: comparative studies on the immobilization strategies and the particle size effects, *J. Phys. Chem. C* 113 (2009) 2521–2525.
- [10] M.S. Thorum, C.A. Anderson, J.J. Hatch, A.S. Campbell, N.M. Marshall, S.C. Zimmerman, Y. Lu, A.A. Gewirth, Direct, electrocatalytic oxygen reduction by laccase on anthracene-2-methanethiol-modified gold, *J. Phys. Chem. Lett.* 1 (2010) 2251–2254.
- [11] M. Pita, S. Shleev, T. Ruzgas, V.M. Fernández, A.I. Yaropolov, L. Gorton, Direct heterogeneous electron transfer reactions of fungal laccases at bare and thiol-modified gold electrodes, *Electrochem. Commun.* 8 (2006) 747–753.
- [12] K. Ataka, F. Giess, W. Knoll, R. Naumann, S. Haber-Pohlmeier, B. Richter, J. Heberle, Oriented attachment and membrane reconstitution of His-tagged cytochrome c oxidase to a gold electrode: in situ monitoring by surface-enhanced infrared absorption spectroscopy, *J. Am. Chem. Soc.* 126 (2004) 16199–16206.
- [13] D.E. Benson, D.W. Conrad, S.A. Trammell, H.W. Hellinga, Design of bioelectronic interfaces by exploiting hinge-bending motions in proteins, *Science* 293 (2001) 1641–1644.
- [14] C. Gutiérrez-Sánchez, D. Olea, M. Marques, V.M. Fernández, I.A. Pereira, M. Vélez, A.L. De Lacey, Oriented immobilization of a membrane-bound hydrogenase onto an electrode for direct electron transfer, *Langmuir* 27 (2011) 6449–6457.
- [15] C. Gwenin, M. Kalaji, P. Williams, R. Jones, The orientationally controlled assembly of genetically modified enzymes in an amperometric biosensor, *Biosens. Bioelectron.* 22 (2007) 2869–2875.
- [16] C. Ley, D. Holtmann, K.M. Mangold, J. Schrader, Immobilization of histidine-tagged proteins on electrodes, *Colloids Surf. B* 88 (2011) 539–551.
- [17] F. Schröper, A. Baumann, A. Offenhäusser, D. Mayer, Direct electrochemistry of novel affinity-tag immobilized recombinant horse heart cytochrome c, *Biosens. Bioelectron.* 34 (2012) 171–177.
- [18] Y. Hong, Y. Xiao, H. Zhou, W. Fang, M. Zhang, J. Wang, L. Wu, Z. Yu, Expression of a laccase cDNA from *Trametes* sp. AH28-2 in *Pichia pastoris* and mutagenesis of transformants by nitrogen ion implantation, *FEMS Microbiol. Lett.* 258 (2006) 96–101.
- [19] Y. Xiao, X. Tu, J. Wang, M. Zhang, Q. Cheng, W. Zeng, Y. Shi, Purification, molecular characterization and reactivity with aromatic compounds of a laccase from basidiomycete *Trametes* sp. strain AH28-2, *Appl. Microbiol. Biotechnol.* 60 (2003) 700–707.
- [20] H. Qiu, Y. Sun, X. Huang, Y. Qu, A sensitive nanoporous gold-based electrochemical aptasensor for thrombin detection, *Colloids Surf. B* 79 (2010) 304–308.
- [21] H. Ge, Y. Gao, Y. Hong, M. Zhang, Y. Xiao, M. Teng, L. Niu, Structure of Native Laccase B from *Trametes* sp. AH28-2, *Acta Crystallogr. Sect. F* 66 (2010) 254–258.
- [22] N. Guex, M.C. Peitsch, SWISS-MODEL and the Swiss-Pdb viewer: an environment for comparative protein modeling, *Electrophoresis* 18 (1997) 2714–2723.
- [23] S. Shleev, M. Pita, A.I. Yaropolov, T. Ruzgas, L. Gorton, Direct heterogeneous electron transfer reactions of *Trametes hirsuta* laccase at bare and thiol-modified gold electrodes, *Electroanalysis* 18 (2006) 1901–1908.