

A Hotspot of Inactivation: The A22S and V108M Polymorphisms Individually Destabilize the Active Site Structure of Catechol *O*-Methyltransferase[†]

Karen Rutherford^{‡,||} and Valerie Daggett^{*,‡,§}

[‡]Departments of Biochemistry and [§]Bioengineering, Box 355013, University of Washington, Seattle, Washington 98195-5013.

^{||}Present address: Department of Biochemistry and Biophysics, Howard Hughes Medical Institute, University of Pennsylvania School of Medicine, Philadelphia, PA 19104

Received February 3, 2009; Revised Manuscript Received May 12, 2009

ABSTRACT: Human catechol *O*-methyltransferase (COMT) contains three common polymorphisms (A22S, A52T, and V108M), two of which (A22S and V108M) render the protein susceptible to deactivation by temperature or oxidation. We have performed multiple molecular dynamics simulations of the wild-type, A22S, A52T, and V108M COMT proteins to explore the structural consequences of these mutations. In total, we have amassed more than 1.4 μ s of simulation time, representing the largest set of simulations detailing the effects of polymorphisms on a protein system to date. The A52T mutation had no significant effect on COMT structure in accord with experiment, thereby serving as a good negative control for the simulation set. Residues 22 (α 2) and 108 (α 5) interact with each other throughout the simulations and are located in a polymorphic hotspot \sim 20 Å from the active site. Introduction of either the larger Ser (22) or Met (108) tightens this interaction, pulling α 2 and α 5 toward each other and away from the protein core. The V108M polymorphism rearranges active-site residues in α 5, β 3, and α 6, increasing the *S*-adenosylmethionine site solvent exposure. The A22S mutation reorients α 2, moving critical catechol-binding residues away from the substrate-binding pocket. The A22S and V108M polymorphisms evolved independently in Northern European and Asian populations. While the decreased activities of both A22S and V108M COMT are associated with an increased risk for schizophrenia, the V108M-induced destabilization is also linked with improved cognitive function. These results suggest that polymorphisms within this hotspot may have evolved to regulate COMT activity and that heterozygosity for either mutation may be advantageous.

Catechol *O*-methyltransferase (COMT,¹ EC 2.1.16) catalyzes the transfer of a methyl group from cosubstrate *S*-adenosylmethionine (SAM) to a variety of catechol substrates, thus playing an important role in the metabolism and inactivation of catecholamine neurotransmitters and catechol estrogens (1–3). COMT exists as two isozymes: a ubiquitous 221-residue soluble form (termed COMT in this paper) expressed primarily in the liver and kidneys and a 271-residue membrane-bound form (*mb*-COMT) found predominantly in the brain and adrenal medulla (4–8). COMT and *mb*-COMT have different substrate specificities, with *mb*-COMT methylating neurotransmitters much more efficiently than the cytosolic protein does (9, 10).

[†]Financial support for this work was provided by the National Institutes of Health (Grant GM50789 to V.D.) and the Canadian Institutes of Health Research (DRA MD-75910 to K.R.).

^{*}To whom correspondence should be addressed. Phone: (206) 685-7420. Fax: (206) 685-3300. E-mail: daggett@u.washington.edu.

¹Abbreviations: COMT, catechol *O*-methyltransferase; C α -rmsd, C α root-mean-square deviation from the starting structure; C α -rmsf, C α root-mean-square fluctuation about the mean structure; DNC, 3,5-dinitrocatechol; MD, molecular dynamics; PDB, Protein Data Bank; SAH, *S*-adenosylhomocysteine; SAM, *S*-adenosylmethionine; SASA, solvent-accessible surface area; *T*_m, temperature resulting in 50% unfolding; *T*₅₀, temperature resulting in 50% loss of enzymatic activity; WT, wild-type (22A, 52A, 108V) protein.

Inhibitors of COMT are currently included in the primary treatment in the management of Parkinson's disease (11–13).

The human *COMT* gene contains three common coding polymorphisms: A22S (rs6267), A52T (rs5031015), and V108M (rs4680) (14–16) (Figure 1). Twenty-five percent of U.S. and Northern European Caucasians are homozygous for the 108M allele (17, 18), while the 22S allele occurs in \sim 10% of Asian populations (www.ncbi.nlm.nih.gov/SNP). There are only limited data regarding the population distribution of the 52T allele, although it appears to occur most frequently in Sub-Saharan African and African American populations (www.ncbi.nlm.nih.gov/SNP). Of the three polymorphisms, the structural and epidemiological effects of the V108M mutation are best characterized. The side chain of residue 108 is buried within a hydrophobic pocket in a loop between α 5 and β 3, approximately 16 Å from the protein's active site (Figure 1). While the wild-type and V108M proteins display similar kinetic properties in vitro (10, 19, 20), the V108M polymorphism destabilizes the protein, increasing its susceptibility to thermal (14, 21) and chemical denaturation (21), as well as to oxidative stress (22–24). This destabilization results in decreased protein levels (19, 25, 26), and therefore activity (27–32), in vivo relative to the wild-type protein. The association of the 108M allele with an increased




FIGURE 1: A22S, A52T, and V108M COMT polymorphisms. Ribbon diagram of wild-type COMT [PDB entry 3BWM (54)] colored from blue (N-terminus) to red (C-terminus). *S*-Adenosylmethionine (SAM) and 3,5-dinitrocatechol (DNC) are shown in stick representation and colored by atom type. Polymorphic residues 22, 52, and 108 are shown in space-filling representation and colored blue, cyan, and green, respectively.

risk for breast cancer (20, 33–35) and neuropsychiatric diseases (36–40) has been well documented. Interestingly, recent studies have linked the 108M allele with increased sensitivity to pain (41–43) and with improved prefrontal cognition, especially in working memory (44–46).

Residue 22 is positioned in a surface loop between $\alpha 1$ and $\alpha 2$, 13 Å from the SAM-binding site and ~20 Å from the catechol-binding site (Figure 1). The A22S mutation decreases COMT activity by 30% relative to that of the wild-type protein, and incubating 22S COMT for 1 h at 45 °C lowers enzymatic activity by 80% compared with that of the wild-type protein (47). However, the A22S polymorphism significantly increases COMT's affinity for SAM (47). Three studies have been published linking the S22 allele with increased risk for schizophrenia (48, 49) and estrogen-associated cancers (47) in Koreans. However, two studies find no association between the A22S polymorphism and either renal cell (50) or prostate cancer (51) in Japanese men.

Residue 52 is positioned on the protein surface in helix $\alpha 3$ which contains residues critical for binding both SAM and catechol substrates (Figure 1). The wild-type and 52T proteins have T_{50} (temperatures resulting in 50% inactivation) values of 53.7 ± 0.10 and 52.7 ± 0.34 °C, respectively (14). In addition, Li et al. showed that while the A52T mutation slightly decreases COMT's affinity for SAM, it has no significant effect on COMT activity (47). No data associating the A52T polymorphism with disease have been published, to the best of our knowledge.

In 2006, we published a molecular dynamics (MD) study describing the destabilizing effects of the V108M polymorphism on the structure of COMT at atomic resolution (52). The mutation distorted the overall structure of COMT, increasing the solvent exposure of both the SAM- and catechol-binding sites. These MD studies were performed using a homology model of human COMT generated from a 2 Å crystal structure of the highly homologous rat protein [PDB entry 1VID (53)] which shares 81% sequence identity with the human protein. Crystal structures of human wild-type and V108M COMT bound with SAM and a substrate analogue are now available (54). Here we describe MD simulations of the wild-type and V108M COMT crystal structures to validate the results of our previous simulations based on the homology model of human COMT (52). We also expand upon these studies to investigate the effects of the

A22S and A52T polymorphisms on COMT structure and dynamics via multiple MD simulations. In total, we have amassed more than 1.4 μ s of simulation time describing the structural effects of COMT polymorphisms, representing the largest set of simulations detailing the effects of polymorphisms on a protein system to date.

MATERIALS AND METHODS

Homology Model. An initial homology model of human 108 V COMT was generated from a 2 Å crystal structure of rat COMT [PDB entry 1VID (53), residues 4–216], which shares 81% sequence identity with the human protein. The side chains of the nonidentical rat residues were replaced sequentially using a Monte Carlo procedure that minimized the torsional, electrostatic, and van der Waals energies of each residue (55). Two passes through the sequence were completed so that residues altered early in the sequence could rearrange in response to later modifications. Altogether, 40 residues were changed, including 30 conservative replacements. Models of A22S, A52T, and V108M COMT were generated from the homology model by replacing each wild-type residue with its polymorphic variant and minimizing the torsional, electrostatic, and van der Waals energies of the resultant structure in vacuo (56). The C α -rmsd between the starting structures of wild-type COMT and all of the variant proteins was ~0.1 Å.

Protein Preparation. The crystal structures of human 108V and 108M COMT were determined after all of the simulations of the wild-type, A22S, A52T, and V108M COMT homology models described here were completed. We therefore performed simulations using the 108V and 108M COMT crystal structures to validate the simulations of the human homology model described here and in 2006 (52). The 1.98 Å crystal structure of human catechol *O*-methyltransferase [PDB entry 3BWM (54), residues 2–215] bound with *S*-adenosylmethionine (SAM), 3,5-dinitrocatechol (DNC), Mg^{2+} , and K^+ was the starting structure for the simulation of wild-type COMT. The 1.30 Å crystal structure of human 108M COMT [PDB entry 3BWY (54), residues 2–215] bound with SAM, DNC, Mg^{2+} , and 2-methyl-2,4-pentanediol (MPD) was used as the starting structure for the simulation of 108M COMT. SAM, DNC, Mg^{2+} , K^+ , and MPD were removed, and the resultant structures underwent 1000 steps of minimization to minimize the torsional, electrostatic, and van der Waals energies of the resultant structures in vacuo (56). One 21 ns control simulation of each of the wild-type and V108M COMT crystal structures was completed at 37 °C using the protocol described below.

Molecular Dynamics Simulations. Molecular dynamics studies of the wild-type and polymorphic COMT proteins were performed with the *in lucem* molecular mechanics (*imm*) simulation package (57) using protocols and the potential energy function that have been previously described (58–60). The simulations included all hydrogen atoms and explicit flexible three-centered waters (60). The proteins were solvated in a periodic rectangular box with walls extending at least 10 Å from all protein atoms. Solvent densities were set to 0.997 and 0.933 g/mL for simulations performed at 25 and 37 °C, respectively (61). Once the densities were set, the box volume was held fixed and the NVE microcanonical ensemble employed. A 10 Å force-shifted, nonbonded cutoff was used for all nonbonded interactions, and the interaction pair lists were updated every two steps (62). Nonbonded interactions between atoms in charge groups separated by three bonds were included and scaled

Table 1: Comparison of General Properties from MD Simulations Based on Homology Models and Crystal Structures of Human Wild-Type and V108M COMT

property ^a	homology model				crystal structure ^d	
	2006 ^b		2008 ^c		WT	V108M
	WT	V108M	WT	V108M		
C α -rmsd (Å)	2.6 ± 0.3	3.3 ± 0.6	3.1 ± 0.2	3.7 ± 0.4 ⁱ	2.4 ± 0.2	3.9 ± 0.1 ^j
total SASA (Å ²) ^c	10948 ± 208	11148 ± 248	11344 ± 245	11850 ± 263 ⁱ	10944 ± 187	12145 ± 143 ^k
Active Site						
SAM-binding site SASA (Å ²) ^f	1016 ± 56	1146 ± 37 ^l	989 ± 105	1150 ± 79	983	1264
E90 C γ –Q120 C γ (Å)	14.7 ± 1.0	15.7 ± 0.9	14.0 ± 2.3	15.4 ± 1.0	16.8 ± 1.4	9.2 ± 0.7
Q120 C γ –W143 C β (Å)	14.9 ± 4.0	18.6 ± 5.2	13.3 ± 1.8	18.1 ± 1.7 ^l	14.9 ± 1.7	15.0 ± 2.2
W143 C β –E90 C γ (Å)	10.8 ± 0.7	12.5 ± 1.8	11.5 ± 2.1	14.0 ± 3.3	11.9 ± 0.5	13.8 ± 1.4 ⁱ
catechol-binding site SASA (Å ²) ^g	723 ± 44	812 ± 107	937 ± 77	851 ± 96	754	999
W38 C γ –P174 C γ (Å)	20.0 ± 2.3	23.7 ± 4.3	19.0 ± 1.8	23.7 ± 4.8	18.8 ± 1.2	29.0 ± 1.6 ^k
Polymorphic Site						
polymorphic site SASA (Å ²) ^h	322 ± 30	366 ± 29	384 ± 95	452 ± 38	376	362
V108 C γ 1/M108 S δ –A22 C β (Å)	4.1 ± 0.3	6.4 ± 3.4	6.3 ± 3.0	4.9 ± 0.7	4.7 ± 0.5	6.0 ± 0.6 ⁱ
V108 C γ 2/M108 C ϵ –V74 C γ 1 (Å)	4.4 ± 0.6	8.0 ± 3.3	7.6 ± 2.0	6.0 ± 1.3	5.6 ± 0.8	4.8 ± 0.6
V108 C γ 1/M108 S δ –R78 C β (Å)	6.5 ± 0.2	6.8 ± 3.3	8.4 ± 3.1	8.5 ± 2.8	8.5 ± 0.6	8.6 ± 0.5
V108 C γ 2/M108 C ϵ –V103 C γ 1 (Å)	4.2 ± 0.3	7.5 ± 0.7 ^k	4.9 ± 0.9	6.4 ± 2.0	4.5 ± 0.4	5.4 ± 0.9

^a All values were calculated using structures from the last 5 ns of each simulation. ^b Data from simulations published in ref (52). Three independent simulations of each protein were run with a *C-scale* value of 0.0. ^c Three independent simulations of each protein were run using a *C-scale* value of 0.4 (see Materials and Methods). ^d Control simulations were run using either 3BWM (WT COMT) or 3BWY (108 MCOMT) as the starting structure (see Materials and Methods) (54). ^e The total solvent-accessible surface area (SASA) was determined using the NACCESS algorithm (65). ^f The following residues were used to calculate the SASA of the SAM-binding site: A39, M40, N41, V42, G66, A67, Y68, S72, E90, I91, N92, G117, A118, S119, Q120, D141, H142, and W143. ^g The following residues were used to calculate the catechol-binding site SASA: W38, M40, K46, D141, K144, D169, N170, C173, P174, L198, E199, Y200, and D205. ^h The following residues were used to calculate the polymorphic site SASA: A22, A73, V74, R78, V103, A106, G107, K109, K111, V112, and residue 108. ⁱ $p < 0.10$. ^j $p < 0.005$. ^k $p < 0.01$. ^l $p < 0.05$.

(*C-scale*) by 0.4 (63, 64); while the inclusion of these interactions is the current working protocol in the group, these interactions were not included in the past (*C-scale* = 0). Very few protein systems are affected by this change, but some proteins are more stable with *C-scale* = 0.4. To facilitate comparisons with our earlier published work (*C-scale* = 0) (52), we present simulations here using both approaches. COMT is unaffected by the change in *C-scale*. A time step of 2 fs was used in all calculations, and structures were saved every 1 ps for analysis. One control simulation of each system was performed at 25 °C to ensure that the protein fold was maintained at low temperatures and that the protein was suitable for study using MD (Figure S1A of the Supporting Information). Three independent simulations at 37 °C of 41–80 ns were performed for the wild-type, A22S, A52T, and V108M proteins for a total of 807 ns of simulation time. In combination with the 650 ns of simulation time published in 2006 (52), we have amassed more than 1.4 μ s of simulation time describing the structural effects of polymorphisms in COMT.

Data Analyses. Statistical analyses was performed using data from the final 5–10 ns of each simulation to ensure that the analyses included only structures from equilibrated systems in which the energies and properties had converged. Average values for C α -rmsd, solvent-accessible surface area (SASA), and contact distances were calculated using structures from the last 5 ns (5000 structures) of each simulation. A contact was defined as a C–C atom distance of ≤ 5.4 Å, or a heavy atom (O, N, or S) distance of ≤ 4.6 Å between two noncontiguous residues. SASA was calculated using in-house software implementing the NACCESS algorithm (65). C α -rmsf values were calculated relative to the average structure over the last 10 ns of each simulation at 37 °C. C α -rmsf values for the crystal structures of the rat

apoprotein [PDB entry 2ZLB (66)] and holoprotein [PDB entry 1VID (53)] were calculated using the crystallographic *B*-factors (*B*) using the relationship $\text{C}\alpha\text{-rmsf} = (3B/8\pi^2)^{1/2}$ (Figure 3B).

The data described in Tables 1 and 2 are based on sets of three independent simulations at 37 °C for each wild-type and variant protein. The errors are the standard deviations in the average values of the ensembles for each reported property. The statistical significance of the data was determined using the Student's *t* test (67).

RESULTS AND DISCUSSION

V108M Polymorphism: Homology Model versus Crystal Structure. Our earlier MD study examining the destabilizing effects of the V108M polymorphism utilized a homology model of human COMT generated from a crystal structure of the highly homologous rat protein [1VID (53)], which shares 81% sequence identity with the human protein (52). Crystal structures of the human wild-type [3BWM (54)] and V108M [3BWY (54)] COMT proteins bound with both SAM and dinitrocatechol (DNC) are now available. Following energy minimization, the root-mean-square deviation of the C α atoms (C α -rmsd) of the human crystal structures and homology models differed by only 0.2 Å, and the positions of residues within the SAM- and catechol-binding sites were nearly identical, suggesting that our homology model was appropriate. It is interesting to note that the crystal structures of wild-type and V108M COMT are virtually identical, probably due to the presence of SAM and DNC. The lack of information from these crystal structures regarding how the V108M polymorphism destabilizes COMT makes MD simulations of this system detailing the structural and dynamic consequences of this mutation all the more important.

Table 2: Properties from the MD Simulations of WT, A22S, A52T, and V108M COMT

property ^a	WT COMT	A52T COMT	A22S COMT	V108M COMT
Cα-rmsd (Å)	3.1 ± 0.2	2.9 ± 0.3	4.0 ± 0.5 ^h	3.7 ± 0.4 ^h
total SASA (Å ²) ^b	11344 ± 245	11290 ± 329	11759 ± 277 ^h	11850 ± 263 ^h
SAM-binding site SASA (Å ²) ^c	989 ± 105	968 ± 180	965 ± 180	1150 ± 79 ^h
catechol-binding site SASA (Å ²) ^d	937 ± 77	924 ± 30	1088 ± 30 ⁱ	851 ± 96
W38 Cγ–P174 Cγ (Å)	19.0 ± 1.8	21.2 ± 2.3	22.8 ± 1.6 ^h	23.7 ± 4.8
A52T Polymorphic Site				
polymorphic site SASA (Å ²) ^e	427 ± 49	434 ± 41	393 ± 15	392 ± 20
A/T52 Cβ–I49 Cγ2 (Å)	5.8 ± 0.5	6.2 ± 0.4	6.0 ± 0.4	6.0 ± 0.4
A/T52 Cβ–I54 Cβ (Å)	7.4 ± 0.2	7.4 ± 0.2	7.5 ± 0.2	7.4 ± 0.2
A/T52 Cβ–Q55 Cβ (Å)	5.6 ± 0.4	5.7 ± 0.4	5.6 ± 0.4	6.0 ± 0.5
A52 Cβ/T52 Cγ2–E56 Cγ (Å)	6.0 ± 0.7	5.2 ± 0.9	6.5 ± 0.8	5.8 ± 0.9
A/T52 Cβ–Y194 Cε2 (Å)	7.5 ± 0.9	7.8 ± 1.1	6.7 ± 1.1	8.1 ± 1.0
A22S Polymorphic Site				
polymorphic site SASA (Å ²) ^f	318 ± 86	299 ± 33	386 ± 37	392 ± 63
helix α2 Cα-rmsd (Å)	0.5 ± 0.1	0.4 ± 0.1	0.5 ± 0.1	0.4 ± 0.1
helix α4 Cα-rmsd (Å)	0.9 ± 0.3	0.7 ± 0.1	1.0 ± 0.3	0.6 ± 0.2
A/S22 Cβ–V74 γ1 (Å)	4.9 ± 0.7	4.5 ± 0.5	7.0 ± 3.1	7.2 ± 1.3 ^h
A/S22 Cβ–R78 Cγ (Å)	4.5 ± 0.6	5.4 ± 1.3	10.5 ± 3.8 ^h	8.2 ± 2.5 ^h
A/S22 Cβ–A106 Cβ (Å)	6.4 ± 2.4	4.9 ± 0.6	5.3 ± 0.9	4.9 ± 0.8
A22 Cβ/S22 Oγ–G107 Cα (Å)	8.9 ± 3.0	7.2 ± 0.8	5.7 ± 1.0	6.5 ± 0.7
A/S22 Cβ–V108 Cγ1/M108 Sδ (Å)	6.3 ± 3.0	4.6 ± 0.7	5.3 ± 1.4	4.8 ± 0.8
V108M Polymorphic Site				
polymorphic site SASA (Å ²) ^g	384 ± 95	369 ± 31	425 ± 41	452 ± 38
V108 Cγ1/M108 Sδ–A22 Cβ (Å)	6.3 ± 3.0	4.6 ± 0.7	5.3 ± 1.4	4.9 ± 0.7
V108 Cγ1–L26 Cδ1/M108 Cε–L26 Cδ2 (Å)	11.7 ± 1.7	10.2 ± 0.8	7.2 ± 2.0 ⁱ	7.0 ± 1.5 ⁱ
V108 Cγ2/M108 Sδ–A73 Cβ (Å)	5.0 ± 1.1	4.6 ± 0.5	6.5 ± 2.0	6.6 ± 2.2
V108 Cγ2/M108 Cε–V74 Cγ1 (Å)	7.6 ± 2.5	5.2 ± 0.8	5.2 ± 0.9	6.0 ± 1.3
V108 Cγ1/M108 Sδ–A77 Cβ (Å)	5.2 ± 1.3	4.2 ± 0.6	6.6 ± 3.1	5.0 ± 1.4
V108 Cγ1/M108 Sδ–R78 Cβ (Å)	8.4 ± 3.1	7.6 ± 1.4	9.3 ± 1.3	8.5 ± 2.0
V108 Cγ2/M108 Cε–V103 Cγ1 (Å)	4.9 ± 0.9	4.3 ± 0.3	4.3 ± 0.3	6.4 ± 2.0
V108 Cγ2/M108 Cε–A106 Cβ (Å)	4.5 ± 0.4	4.2 ± 0.4	4.2 ± 0.3	4.8 ± 0.8

^a All values were calculated using structures from the last 5 ns of each simulation. All values are expressed as means and standard deviations of the means from three independent simulations at 37 °C. ^b The total solvent-accessible surface area (SASA) was determined using the NACCESS algorithm (65). ^c The following residues were used to calculate the SASA of the SAM-binding site: A39, M40, N41, V42, G66, A67, Y68, S72, E90, I91, N92, G117, A118, S119, Q120, D141, H142, and W143. ^d The following residues were used to calculate the catechol-binding site SASA: W38, M40, K46, D141, K144, D169, N170, C173, P174, L198, E199, Y200, and D205. ^e The following residues were used to calculate the A52T polymorphic site SASA: I49, V53, Q55, E56, Y194, and residue 52. ^f The following residues were used to calculate the A22S polymorphic site SASA: Q23, V25, L26, V74, A77, R78, A106, residue 108, and residue 22. ^g The following residues were used to calculate the V108M polymorphic site SASA: residue 22, A73, V74, R78, V103, A106, G107, K109, K111, V112, and residue 108. ^h $p < 0.10$. ⁱ $p < 0.05$.

We performed simulations of human wild-type and V108M COMT beginning with the human crystal structures to validate the results obtained from simulations using the homology models. Since the original results were published (52), our simulation package, *iimm*, has been updated to include a scaling factor (*C-scale*) of 0.4 for nonbonded interactions between atoms in charge groups separated by three bonds (63, 64). Therefore, we have performed simulations of both COMT variants using a scaling factor of 0.4 for nonbonded interactions to verify that this change does not affect the previously published results (52) and to ensure consistency with our current simulation protocols, particularly for comparison with other proteins in our Dyanemomics database (63, 68).

The overall methyltransferase fold was maintained in all of the simulations (Figure 2). Cα-rmsd plots of structures from the simulations using the human crystal structures and homology models at 25 °C are very similar (Figure S1 of the Supporting Information), consistent with experimental data that show that WT and V108M behave similarly at low temperatures (21). Our focus here is the behavior of the proteins at physiological pH.

Our previous work (52) looked extensively at how the V108M polymorphism affected COMT structure at 25, 37, and 50 °C. Those results indicated that the proteins behave similarly at both 25 and 50 °C and that differences in structure and stability are most prevalent at 37 °C. Experimental studies describing protein levels in vivo (25, 26) as well as protein stability and aggregation in vitro (21) also show larger differences between the WT and V108M proteins at 37 °C.

At 37 °C, substituting Met for Val at position 108 expands the protein, increasing both the Cα-rmsd and the total solvent-accessible surface area of the 108M protein relative to those of 108V COMT in all of the simulations (Table 1). The larger Met interacted differently than the Val with residues in helices α2 (A22), α4 (V74), and α5 (V103), altering the orientation of secondary structure elements neighboring the polymorphic site (Table 1 and Figure 2). 108M COMT was also more prone to distortion in helices α6 (Q120) and α8 (P174), both of which contain active site residues. These changes resulted in a significant increase in the solvent exposure of the SAM-binding sites of the 108M proteins relative to wild-type COMT




FIGURE 2: Comparison of MD simulations performed at 37 °C using crystal structures and homology models of human wild-type and V108M COMT. Snapshots of structures from the final nanosecond of MD simulations of (left) the crystal structures of human wild-type [3BWM (54)] and V108M [3BWY (54)] COMT and (center and right) homology models of human COMT based on the crystal structure of rat COMT [1VID (53)] performed with *C-scale* values of either 0.0 (center) or 0.4 (right) (see Materials and Methods). (A) The 108M COMT structure is prone to disruption at 37 °C. Altered packing around residue M108 reorients helix $\alpha 6$, pulling Q120 away from the SAM-binding site. Helix $\alpha 8$ (and P174) pulls away from the protein core, resulting in an expanded conformation. These motions occur during simulations of both the crystal structure and homology models of the V108M COMT protein but are not observed during the simulations of wild-type COMT. SAM-binding residues (E90, Q120, and W143), catechol-binding residues (W38 and P174), and residue 108 are shown in space-filling representation and colored green, red, and magenta, respectively. (B) The 108 MCOMT active site is more exposed to solvent and distorted at 37 °C than that of the wild-type protein. Residues in the SAM- and catechol-binding sites are colored green and red, respectively.

(Table 1 and Figure 2). While the magnitude of the differences between the 108V and 108M structures varied between simulations, the overall trends describing the destabilizing effects of the V108M polymorphism on COMT structure were consistent under all conditions tested. The results indicate that simulations employing the COMT homology models and the crystal structures provide consistent descriptions of the structural and dynamic effects of the V108M mutation. Also, the addition of a scaling factor for nonbonded interactions does not significantly alter the results of the COMT simulations at 37 °C.

A crystal structure of the rat COMT apoprotein [2ZLB (66)] is now available. The overall structures of the apo and holo rat proteins are very similar, differing only by a $C\alpha$ -rmsd of 0.72 Å with all of the secondary structure elements (helices and

β -strands) superposable (Figure 3A). The major differences between the apo and holo structures lie in the loop regions that define the catechol-binding site. The $\alpha 2$ – $\alpha 3$ loop (W38 and M40), the $\beta 5$ – $\alpha 8$ loop (P174), and the catalytic loop (residues 197–202) all pull away from the protein core in the COMT apoprotein, exposing the catechol-binding site and a portion of the SAM-binding site (Figure 3A,B). Interestingly, the adenosine-binding pockets of the SAM sites are very similar in both the apo- and holoproteins, showing only slight displacements in the positions of key active site residues E90, Q120, and W143. COMT first binds SAM and then the catechol substrate in an ordered mechanism (7). The apoprotein structure suggests that the adenosine moiety of SAM directs binding of the cosubstrate into a partially formed active site pocket. It is probable that upon SAM docking, the $\alpha 2$ – $\alpha 3$ and $\beta 5$ – $\alpha 8$ loops move toward the protein core, enabling active site residues W38, M40, and N170 to contact SAM. This rearrangement of loops would also bring “gatekeeper” residues W38 and P174 closer together to form a binding pocket for the catechol substrate (Figure 3A). Because the rat and human COMT proteins are 81% identical in sequence and the crystal structures of the rat and human COMT holoproteins are virtually identical (Figure 3A), it is reasonable to believe that the structure of the wild-type human apoprotein is similar to that of rat COMT.

The starting structure for all of the simulations of human COMT was a COMT holoprotein stripped of both SAM and the catechol substrate. Figure 3C shows the crystal structure of the rat COMT apoprotein overlaid with structures from the final nanosecond of three independent simulations of the human wild-type and V108M COMT apoproteins. The structures from the simulations of human COMT are similar to that of the rat apoprotein structure in that the protein expands while the overall methyltransferase fold and the structure of the binding pocket for the adenosine moiety of SAM are maintained. In addition, the $\alpha 2$ – $\alpha 3$, $\beta 5$ – $\alpha 8$, and catalytic loops become highly flexible, revealing the active site to facilitate substrate binding. These similarities suggest that our descriptions of the human apoprotein structures published in 2006 (52) and reported here are correct. These characteristic apoprotein features are also seen during simulations of the V108M variant (Figure 3C). Interestingly, the V108M apoprotein demonstrates an additional large displacement of $\beta 3$ and $\alpha 6$ that opens and disorders the SAM-binding pocket (Figure 3C). This displacement is a direct result of altered side chain packing around the larger Met 108 in the nearby $\alpha 5$ – $\beta 3$ loop (Figures 3C and 5E,F). The $\beta 3$ – $\alpha 6$ loop motion is quite dynamic at 37 °C (Figure 4C), and SAM is still able to bind to the V108M protein. Indeed, several studies have shown that the presence of SAM, which would stabilize $\alpha 6$ through interactions with S119 and Q120, protects both the helical structure and activity of COMT at elevated temperatures (10, 21). However, it is possible that the reorientation of $\alpha 6$ nucleates the destabilization of V108M COMT, resulting in the degradation of a portion of the protein pool.

Competitive substrate-based COMT inhibitors have long been used in the treatment of Parkinson’s disease (11, 12, 69–71). Recent studies linking the destabilizing V108M polymorphism with improved prefrontal cognition (45, 46) have led to the use of several of these COMT inhibitors to improve cortical processing in individuals carrying the wild-type COMT allele (72, 73). However, the 108M allele has also been associated with an increased risk for breast cancer (20, 34, 35) and neuropsychiatric disease (36–40). Therefore, the development of stabilizing agents




FIGURE 3: Structural characterization of the COMT apoprotein. (A) Superposed ribbon diagrams of the rat COMT apoprotein [gray, 2ZLB (66)] and the rat [1VID (53)] and human [3BWM (54)] holoproteins bound with SAM and DNC and colored from blue (N-terminus) to red (C-terminus). The overall structures of the rat and human holoproteins are virtually identical. The rat apo- and holoproteins differ mainly in the loop regions that define the catechol-binding site. The $\alpha 2$ – $\alpha 3$ loop, the $\beta 5$ – $\alpha 8$ loop, and the catalytic loop (residues 197–202) pull away from the protein core in the apoprotein, exposing the SAM- and catechol-binding sites. This open configuration may facilitate substrate binding. Interestingly, the adenosine pockets of the apo- and holoprotein SAM-binding sites are very similar, showing only a slight displacement of key active site residues (E90, Q120, and W143). Active site residues are shown in stick representation and colored or labeled in either gray (apoprotein) or to match the structure (holoprotein). (B) C α -rmsf values (in angstroms) per residue for the rat apoprotein [black, 2ZLB (66)] and holoprotein [green, 1VID (53)]. (C) Structural overlay of the rat apoprotein crystal structure [gray, 2ZLB (66)] with structures from the final nanosecond of three independent simulations of human WT (gold) and V108M (pink) COMT. The human WT and rat apoproteins behave similarly in that the structure of the adenosine pocket in the SAM-binding site ($\alpha 5$, $\alpha 6$, and $\alpha 7$) is maintained, with some flexibility in the orientations of active site residues E90, Q120, and W143, while the $\alpha 2$ – $\alpha 3$, $\beta 5$ – $\alpha 8$, and catalytic loops fluctuate greatly, opening the active site core and moving W38, M40, P174, and E199 out of the substrate-binding pocket. In addition to these motions, both $\beta 3$ and $\alpha 6$ pull away from the protein core during simulations of V108M COMT. SAM- and catechol-binding residues are shown in stick representation and colored green and red, respectively. Residue 108 is shown in space-filling representation and colored blue. (D) Ribbon (left) and van der Waals surface (right) representation of the $\beta 3$ – $\alpha 6$ pocket from human WT COMT. The V108M polymorphism alters the orientations of $\beta 3$ and $\alpha 6$, possibly destabilizing the protein. Therefore, the $\beta 3$ – $\alpha 6$ pocket may serve as a target site for docking small molecules to stabilize the V108M COMT structure. Positively and negatively charged residues in the surface representation are colored blue and red, respectively.




FIGURE 4: Mobility of the wild-type, A22S, A52T, and V108M COMT variants at 37 °C. C α -rmsf values (in angstroms) per residue for the wild-type protein (blue) compared with those of the (A) A52T, (B) A22S, and (C) V108M COMT variants (all in red). C α -rmsf values were calculated relative to the average structure over the last 10 ns of each simulation. Secondary structural elements are depicted as cylinders for α -helices and arrows for β -strands and are colored to match the structure in Figure 1. The polymorphic residues are marked with asterisks.

to supplement the treatment of schizophrenia and obsessive-compulsive disorder is also desirable. Because $\beta 3$ and $\alpha 6$ are most readily affected by the V108M polymorphism in simulation and because SAM stabilizes COMT through interactions with residues in the $\beta 3$ – $\alpha 6$ loop, they represent a logical target site for the docking of a stabilizing small molecule that is not competitive with substrate binding. $\alpha 5$, $\beta 3$, and $\alpha 6$ form a pocket, ~ 12 Å in diameter and 6 Å deep, on the protein surface (Figure 3D). The pocket has a hydrophobic core and is bordered by several positively charged residues on $\alpha 4$ (R85), $\beta 3$ (K109 and K111) and $\alpha 6$




FIGURE 5: Effects of the A52T, A22S, and V108M COMT polymorphisms on local tertiary structure. (A and B) A52 ($\alpha 3$) forms main chain hydrogen bonds with K48, I49, and E56 of $\alpha 3$ and side chain contacts with residues in $\alpha 3$ (I49, V53, Q55, and E56) and $\beta 6$ (Y194). T52 maintains all of these contacts, while T52 OG1 forms additional hydrogen bonds with the backbone carbonyl groups of K48 and I49. (C and D) A22 is positioned in a loop between $\alpha 1$ and $\alpha 2$ where it interacts with residues in $\alpha 2$ (Q23, V25, and L26), $\alpha 4$ (V74, A77, and R78), and $\alpha 5$ (A106 and V108). The larger S22 forms an additional S22 OG—G107 O hydrogen bond, enabling closer contact between S22 and residues in $\alpha 5$ (A106, G107, and V108). This altered packing moves S22 away from residues V74 and R78 of $\alpha 4$, affecting the orientations of both $\alpha 2$ and $\alpha 4$ relative to their positions in the wild-type protein. (E and F) Residue 108 is located in a loop between $\alpha 4$ and $\alpha 5$. Both V108 and M108 contact residues in $\alpha 2$ (A22), $\alpha 4$ (A73, V74, and R78), and $\alpha 5$ (V103 and A106). The larger methionine forms additional contacts with V25 and L26 of $\alpha 2$ and M102 of $\alpha 5$, effectively moving helices $\alpha 2$ and $\alpha 5$ closer together. Structures from the final nanosecond of three independent simulations are superimposed in panels A–F and colored to match Figure 1. Side chains of the polymorphic residues and their contacts are shown in space-filling and stick representations, respectively.

(K128, K129). In addition, T113 and Y130 could provide contacts for hydrogen bonding interactions with a stabilizing molecule (Figure 3D). It will be interesting to see if small molecules docked into this site do protect the structure of 108M COMT.

A52T Polymorphism. The A52T polymorphism does not significantly affect either the catalytic activity or thermal stability of COMT (14, 47). The kinetic similarities between the wild-type and A52T proteins are reflected in the simulations. Both proteins reach an overall C α -rmsd of 3.0 Å and a total solvent-accessible surface area of 11300 Å² (Table 2 and Figure S1). The average C α -atom root-mean-square fluctuations (C α -rmsf) values reflecting the degree of motion about the mean structure over all three simulations are shown in Figure 4A. The C α -rmsf values for both the wild-type and A52T proteins follow similar patterns. The wild-type protein has higher C α -rmsf values in the catalytic loop, while the 52T variant shows slight spikes in flexibility at residue 52 and in the loop regions containing both SAM-binding




FIGURE 6: A22S polymorphism disrupts the catechol-binding site. (A) The altered packing around residue S22 causes $\alpha 2$ and $\alpha 5$ to move toward each other while the last turn of $\alpha 4$ pulls away from the polymorphic site. This rearrangement results in the formation of new salt bridges between D30 ($\alpha 2$) and R75 ($\alpha 4$), E34 ($\alpha 2$) and R75 ($\alpha 4$), and E34 ($\alpha 2$) and K48 ($\alpha 3$). The altered packing around the polymorphic residue is propagated through $\alpha 3$ toward the protein's C-terminus, leading to an overall expansion of the protein and the separation of key catechol-binding residues W38, P174, and E199. Residue 22 and residues involved in salt bridges are shown in space-filling representation and colored by atom type. V108 and catechol-binding residues are colored green and magenta, respectively. Structures are colored as in Figure 1. (B) The rearrangement of $\alpha 2$ and $\alpha 3$ both increases the mobility of several residues in the $\alpha 2$ – $\alpha 3$ loop, including W38, a gatekeeping residue involved in catechol binding, and pulls these residues away from the protein's C-terminus. This movement opens a large cleft in the protein, which greatly increases the solvent exposure of the catechol-binding site, relative to that of the wild-type protein. Residue 22 and the catechol-binding residues (W38, P174, and E199) are shown in space-filling representation and colored magenta. $\alpha 2$, $\alpha 3$, and $\alpha 4$ are colored as in Figure 1.

(E90, I91, S119, and Q120) and catechol-binding (P174) residues. However, these differences in C α -rmsf values do not translate to altered active site structures. The solvent exposures of the SAM- and catechol-binding sites are ~975 and 930 Å², respectively, in both the wild-type and A52T proteins (Table 2).

Residue 52 is a surface residue found in the third turn of helix $\alpha 3$ (Figure 1). A52 forms main chain hydrogen bonds with K48, I49, and E56 of $\alpha 3$ and hydrophobic contacts with residues in $\alpha 3$ (I49, V53, Q55, and E56) and $\beta 6$ (Y194) (Table 2 and Figure 5A). The larger Thr maintains all of these contacts, and the CB atoms of A52 and T52 are positioned similarly throughout the simulations and oriented toward the solvent (Table 2 and Figure 5A,B). Instead of interacting with any new nearby residues or solvent molecules, the OG1 atom of T52 forms additional hydrogen bonds with the backbone carbonyl groups of K48 and I49. Phylogenetic analyses show conservation of an alanine residue at position 52 in most species. However, *Bos taurus* and *Sus scrofa*




FIGURE 7: Total C α -rmsd distributions of structures from MD simulations of wild-type, A22S, A52T, and V108M COMT. Structures from the WT COMT simulations fall into a narrow distribution centered at a 3.3 Å C α -rmsd from the starting structure. The 52T distribution is centered at a 2.6 Å C α -rmsd, indicating that the A52T polymorphism somewhat stabilizes the protein structure. However, structures from the 52T COMT simulations fall into a broader distribution than those of the WT protein. Both the 22S and 108M COMT variants exist as ensembles of structures that are more distorted than those of the WT protein. The vertical scale is arbitrary.

have an Arg and a Thr, respectively, at this position, suggesting that COMT structure and function are tolerant of several substitutions at this location. The observation that the Ala to Thr polymorphism at residue 52 only minimally affects COMT structure in both simulation and experiment makes the simulations of A52T COMT good negative controls for the effects of computationally introducing mutations into wild-type crystal structures.

A22S Polymorphism. The A22S polymorphism significantly decreases the overall activity of COMT (47). The A22S variant loses ~80% of its activity upon incubation at 45 °C (47). The mechanism for this deactivation is unclear as the A22S mutation decreases the protein's K_m for SAM while increasing its K_m for hydroxyestradiol relative to the wild-type protein. We performed multiple MD simulations of the A22S COMT variant to investigate the molecular mechanisms underlying its inactivation.

The methyltransferase fold was maintained throughout all of the simulations. Structures of the wild-type protein and A22S variant reached C α -rmsd values of 3.1 ± 0.2 and 4.0 ± 0.5 Å, respectively, during the simulations at 25 and 37 °C (Table 2 and Figure S1). The structure of 22S COMT expanded relative to the wild-type protein, with a 400 Å² increase in total solvent-accessible surface area (Table 2). Figure 4 shows C α -rmsf values for both the wild-type and A22S COMT proteins at 37 °C. The A22S variant exhibited increased flexibility throughout the protein, with large peaks in C α -rmsf values for residues in α 1, α 2 (S22), α 5, α 6, α 7, and the catalytic loop, including SAM- and catechol-binding residues Q120 and E202.

Residue 22 is located in the loop between helices α 1 and α 2, ~13 Å from the SAM-binding site and ~20 Å from the catechol-binding site (Figure 1). No active site residues are located within α 1 or α 2. However, the surface loop between α 2 and α 3 does contain W38, a gatekeeper residue that forms a hydrophobic pocket with P174 of α 8 to orient the catechol substrate for

methylation. Ala 22 forms hydrogen bonds with Q23 OE1 and the backbone amides of V25 and L26. Hydrophobic contacts between A22 and residues in α 2 (Q23, V25, and L26), α 4 (V74, A77, and R78), and α 5 (A106 and V108) are maintained throughout the simulations (Table 2 and Figure 5C). S22 maintains many of its contacts with α 2 and α 5 while forming an additional hydrogen bond with G107 (Table 2 and Figure 5D). The S22–G107 interaction packs α 2 and α 5 more closely together, moving both S22 and α 2 away from their contacts with V74, A77, and R78 in α 4 (Table 2) and moving W38 away from residues P174 and E199 in the protein's C-terminus (Table 2 and Figure 6). This motion moves the opposite (N-terminal) ends of α 2 and α 4 toward each other, enabling salt bridges among D30, R75, and E34 to form. The formation of this network pulls α 4 and α 3 away from each other, and away from the protein's C-terminus, leading to an expansion of the overall structure and the opening of a large cleft in the protein (Figure 6B). This repacking around the polymorphic site is translated throughout the protein, significantly increasing the solvent exposure of the catechol-binding site ~20 Å away, in agreement with the A22S mutation's detrimental effect on substrate binding (47) (Table 2).

A22S and V108M Polymorphic Hotspot. The A22S and V108M polymorphisms have been shown experimentally to decrease COMT stability and activity. The A22S mutation alters catechol binding (47), while the V108M mutation distorts the SAM-binding site (10, 21, 52). Both A22 (α 1– α 2) and V108 (α 5– β 3) are located in surface loops 16–20 Å from the active sites. However, the polymorphic residues pack against secondary structure elements that contain substrate-binding residues on their distal ends. Interestingly, polymorphic residues 22 and 108 contact each other throughout all of the simulations (Figure 5 and Table 2), marking this region as a hotspot for mutations that regulate but do not abolish COMT activity. Substitutions of the

larger Ser and Met residues for Ala and Val similarly distort interactions and packing within the polymorphic sites of both residues 22 and 108 (Table 2). These packing differences are translated throughout both proteins, resulting in large increases in the overall solvent-accessible surface area relative to that of the wild-type protein. However, the two polymorphisms have very different effects on the active sites. The A22S mutation results in the formation of a large cleft in the protein, distorting and significantly increasing the solvent exposure of the catechol-binding site. The introduction of S22 has little effect on the solvent exposure or structure of the SAM-binding site. The opposite is true for the V108M polymorphism, which distorts the SAM-binding site through the rearrangement of active site residues positioned on $\alpha 5$ (E90 and N92), $\beta 3$ (A116), and $\alpha 6$ (S119 and Q120). This difference could depend simply on whether SAM- or catechol-binding residues are on the distal ends of the reoriented helices ($\alpha 2$, $\alpha 5$, and $\alpha 6$). The overall effects of mutation can be seen in the C α RMSD distribution in Figure 7. Both A22S and V108M distort the protein relative to wild type while A52T stabilizes it.

It is interesting to note that the A22S and V108M polymorphisms developed independently in Northern European and Asian populations, possibly as different ways of destabilizing and deactivating the COMT protein. Both the 22S and 108M alleles have been associated with increased risk of cancer (20, 33–35, 47) and neuropsychiatric dysfunction (36–40, 48, 49). However, the 108M allele also confers improved prefrontal cognitive function, especially in working memory (44–46). These results suggest that individuals heterozygous for the V108M polymorphism may have an advantage in that intermediate COMT activity may minimize the individual's risk for neuropsychiatric disease while slightly improving cortical processing. A similar phenomenon is seen with protein isoaspartate O-methyltransferase (PIMT), an enzyme critical for protein repair, where heterozygosity for a V119I polymorphism promotes successful aging, due in part to the complementary substrate affinities of the 119V and 119I proteins (74). Both the V108M and V119I mutations in COMT and PIMT, respectively, are present at high frequencies in Caucasian populations, with ~50% of individuals heterozygous for either polymorphism (17, 74, 75). Further studies are necessary to determine if the A22S polymorphism is associated with any "beneficial" phenotypes (Figure 7).

ACKNOWLEDGMENT

Most simulations in this study were performed using processor time from Microsoft Research through Computing @ Microsoft (www.microsoft.com/science). Figures were produced using the UCSF Chimera package from the Computer Graphics Laboratory, University of California, San Francisco (supported by NIH Grant P41 RR-01081) (76). K.R. thanks Dr. William Parson for helpful discussions and Amanda Jonsson for running the simulations of the COMT homology models.

SUPPORTING INFORMATION AVAILABLE

Evolution of C α -rmsd (in angstroms) with time for simulations of the WT, A22S, A52T, and V108M COMT proteins performed at 25 and 37 °C (Figure S1). This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

1. Axelrod, J., and Szara, S. (1958) Enzymic conversion of metanephrine to epinephrine. *Biochim. Biophys. Acta* 30, 188–189.

2. Mannisto, P. T., and Kaakkola, S. (1999) Catechol-O-methyltransferase (COMT): Biochemistry, molecular biology, pharmacology, and clinical efficacy of the new selective COMT inhibitors. *Pharmacol. Rev.* 51, 593–628.
3. Weinshilboum, R. M., Otterness, D. M., and Szumlanski, C. L. (1999) Methylation pharmacogenetics: Catechol O-methyltransferase, thio-purine methyltransferase, and histamine N-methyltransferase. *Annu. Rev. Pharmacol. Toxicol.* 39, 19–52.
4. Tenhunen, J., Salminen, M., Jalanko, A., Ukkonen, S., and Ulmanen, I. (1993) Structure of the rat catechol-O-methyltransferase gene: Separate promoters are used to produce mRNAs for soluble and membrane-bound forms of the enzyme. *DNA Cell Biol.* 12, 253–263.
5. Tenhunen, J., Salminen, M., Lundstrom, K., Kiviluoto, T., Savolainen, R., and Ulmanen, I. (1994) Genomic organization of the human catechol O-methyltransferase gene and its expression from two distinct promoters. *Eur. J. Biochem.* 223, 1049–1059.
6. Rivett, A. J., Francis, A., and Roth, J. A. (1983) Localization of membrane-bound catechol-O-methyltransferase. *J. Neurochem.* 40, 1494–1496.
7. Rivett, A. J., and Roth, J. A. (1982) Kinetic studies on the O-methylation of dopamine by human brain membrane-bound catechol O-methyltransferase. *Biochemistry* 21, 1740–1742.
8. Huh, M. M., and Friedhoff, A. J. (1979) Multiple molecular forms of catechol-O-methyltransferase. Evidence for two distinct forms, and their purification and physical characterization. *J. Biol. Chem.* 254, 299–308.
9. Bonifati, V., and Meco, G. (1999) New, selective catechol-O-methyltransferase inhibitors as therapeutic agents in Parkinson's disease. *Pharmacol. Ther.* 81, 1–36.
10. Lotta, T., Vidgren, J., Tilgmann, C., Ulmanen, I., Melen, K., Julkunen, I., and Taskinen, J. (1995) Kinetics of human soluble and membrane-bound catechol O-methyltransferase: A revised mechanism and description of the thermolabile variant of the enzyme. *Biochemistry* 34, 4202–4210.
11. Lees, A. J. (2008) Evidence-based efficacy comparison of tolcapone and entacapone as adjunctive therapy in Parkinson's disease. *CNS Neurosci. Ther.* 14, 83–93.
12. Chan, D. K., Cordato, D. J., and O'Rourke, F. (2008) Management for motor and non-motor complications in late Parkinson's disease. *Geriatrics* 63, 22–27.
13. Factor, S. A. (2008) Current status of symptomatic medical therapy in Parkinson's disease. *Neurotherapeutics* 5, 164–180.
14. Shield, A. J., Thomae, B. A., Eckloff, B. W., Wieben, E. D., and Weinshilboum, R. M. (2004) Human catechol O-methyltransferase genetic variation: Gene resequencing and functional characterization of variant allozymes. *Mol. Psychiatry* 9, 151–160.
15. Saito, S., Iida, A., Sekine, A., Miura, Y., Sakamoto, T., Ogawa, C., Kawauchi, S., Higuchi, S., and Nakamura, Y. (2001) Identification of 197 genetic variations in six human methyltransferase genes in the Japanese population. *J. Hum. Genet.* 46, 529–537.
16. Cargill, M., Altshuler, D., Ireland, J., Sklar, P., Ardlie, K., Patil, N., Shaw, N., Lane, C. R., Lim, E. P., Kalyanaraman, N., Nemesh, J., Ziaugra, L., Friedland, L., Rolfe, A., Warrington, J., Lipshutz, R., Daley, G. Q., and Lander, E. S. (1999) Characterization of single-nucleotide polymorphisms in coding regions of human genes. *Nat. Genet.* 22, 231–238.
17. Palmatier, M. A., Kang, A. M., and Kidd, K. K. (1999) Global variation in the frequencies of functionally different catechol-O-methyltransferase alleles. *Biol. Psychiatry* 46, 557–567.
18. Weinshilboum, R. M., and Raymond, F. A. (1977) Inheritance of low erythrocyte catechol-o-methyltransferase activity in man. *Am. J. Hum. Genet.* 29, 125–135.
19. Chen, J., Lipska, B. K., Halim, N., Ma, Q. D., Matsumoto, M., Melhem, S., Kolachana, B. S., Hyde, T. M., Herman, M. M., Apud, J., Egan, M. F., Kleinman, J. E., and Weinberger, D. R. (2004) Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): Effects on mRNA, protein, and enzyme activity in postmortem human brain. *Am. J. Hum. Genet.* 75, 807–821.
20. Goodman, J. E., Jensen, L. T., He, P., and Yager, J. D. (2002) Characterization of human soluble high and low activity catechol-O-methyltransferase catalyzed catechol estrogen methylation. *Pharmacogenetics* 12, 517–528.
21. Rutherford, K., Alphonso, E., McMillan, A., Daggett, V., and Parson, W. W. (2008) The V108M mutation decreases the structural stability of catechol O-methyltransferase. *Biochim. Biophys. Acta* 1784, 1098–1105.
22. Cotton, N. J., Stoddard, B., and Parson, W. W. (2004) Oxidative inhibition of human soluble catechol-O-methyltransferase. *J. Biol. Chem.* 279, 23710–23718.

23. Li, Y., Yao, J., Chang, M., Nikolic, D., Yu, L., Yager, J. D., Mesecar, A. D., van Breemen, R. B., and Bolton, J. L. (2004) Equine catechol estrogen 4-hydroxyequilenin is a more potent inhibitor of the variant form of catechol-O-methyltransferase. *Chem. Res. Toxicol.* 17, 512–520.
24. Li, Y., Yang, X., Chang, M., Yager, J. D., van Breemen, R. B., and Bolton, J. L. (2005) Functional and structural comparisons of cysteine residues in the Val108 wild type and Met108 variant of human soluble catechol O-methyltransferase. *Chem.-Biol. Interact.* 152, 151–163.
25. Doyle, A. E., Goodman, J. E., Silber, P. M., and Yager, J. D. (2004) Catechol-O-methyltransferase low activity genotype (COMTLL) is associated with low levels of COMT protein in human hepatocytes. *Cancer Lett.* 214, 189–195.
26. Doyle, A. E., and Yager, J. D. (2008) Catechol-O-methyltransferase: Effects of the val108met polymorphism on protein turnover in human cells. *Biochim. Biophys. Acta* 1780, 27–33.
27. Aksoy, S., Klenner, J., and Weinshilboum, R. M. (1993) Catechol O-methyltransferase pharmacogenetics: Photoaffinity labelling and western blot analysis of human liver samples. *Pharmacogenetics* 3, 116–122.
28. Boudikova, B., Szumlanski, C., Maidak, B., and Weinshilboum, R. (1990) Human liver catechol-O-methyltransferase pharmacogenetics. *Clin. Pharmacol. Ther.* 48, 381–389.
29. Grossman, M. H., Szumlanski, C., Littrell, J. B., Weinstein, R., and Weinshilboum, R. M. (1992) Electrophoretic analysis of low and high activity forms of catechol-O-methyltransferase in human erythrocytes. *Life Sci.* 50, 473–480.
30. Scanlon, P. D., Raymond, F. A., and Weinshilboum, R. M. (1979) Catechol-O-methyltransferase: Thermolabile enzyme in erythrocytes of subjects homozygous for allele for low activity. *Science* 203, 63–65.
31. Spielman, R. S., and Weinshilboum, R. M. (1981) Genetics of red cell COMT activity: Analysis of thermal stability and family data. *Am. J. Med. Genet.* 10, 279–290.
32. Weinshilboum, R. M., and Dunnette, J. (1981) Thermal stability and the biochemical genetics of erythrocyte catechol-O-methyltransferase and plasma dopamine- β -hydroxylase. *Clin. Genet.* 19, 426–437.
33. Dawling, S., Roodi, N., Mernaugh, R. L., Wang, X., and Parl, F. F. (2001) Catechol-O-methyltransferase (COMT)-mediated metabolism of catechol estrogens: Comparison of wild-type and variant COMT isoforms. *Cancer Res.* 61, 6716–6722.
34. Wedren, S., Rudqvist, T. R., Granath, F., Weiderpass, E., Ingelman-Sundberg, M., Persson, I., and Magnusson, C. (2003) Catechol-O-methyltransferase gene polymorphism and post-menopausal breast cancer risk. *Carcinogenesis* 24, 681–687.
35. Yim, D. S., Park, S. K., Yoo, K. Y., Yoon, K. S., Chung, H. H., Kang, H. L., Ahn, S. H., Noh, D. Y., Choe, K. J., Jang, I. J., Shin, S. G., Strickland, P. T., Hirvonen, A., and Kang, D. (2001) Relationship between the Val158Met polymorphism of catechol O-methyl transferase and breast cancer. *Pharmacogenetics* 11, 279–286.
36. Lachman, H. M., Morrow, B., Shprintzen, R., Veit, S., Parsia, S. S., Faedda, G., Goldberg, R., Kucherlapati, R., and Papolos, D. F. (1996) Association of codon 108/158 catechol-O-methyltransferase gene polymorphism with the psychiatric manifestations of velo-cardio-facial syndrome. *Am. J. Med. Genet.* 67, 468–472.
37. Lachman, H. M., Nolan, K. A., Mohr, P., Saito, T., and Volavka, J. (1998) Association between catechol O-methyltransferase genotype and violence in schizophrenia and schizoaffective disorder. *Am. J. Psychiatry* 155, 835–837.
38. Nolan, K. A., Volavka, J., Czobor, P., Cseh, A., Lachman, H., Saito, T., Tiihonen, J., Putkonen, A., Hallikainen, T., Kotilainen, I., Rasanen, P., Isohanni, M., Jarvelin, M. R., and Karvonen, M. K. (2000) Suicidal behavior in patients with schizophrenia is related to COMT polymorphism. *Psychiatr. Genet.* 10, 117–124.
39. Strous, R. D., Bark, N., Parsia, S. S., Volavka, J., and Lachman, H. M. (1997) Analysis of a functional catechol-O-methyltransferase gene polymorphism in schizophrenia: Evidence for association with aggressive and antisocial behavior. *Psychiatry Res.* 69, 71–77.
40. Strous, R. D., Nolan, K. A., Lapidus, R., Diaz, L., Saito, T., and Lachman, H. M. (2003) Aggressive behavior in schizophrenia is associated with the low enzyme activity COMT polymorphism: A replication study. *Am. J. Med. Genet., Part B* 120, 29–34.
41. Diatchenko, L., Nackley, A. G., Slade, G. D., Bhalang, K., Belfer, I., Max, M. B., Goldman, D., and Maixner, W. (2006) Catechol-O-methyltransferase gene polymorphisms are associated with multiple pain-evoking stimuli. *Pain* 125, 216–224.
42. Zubieta, J. K., Heitzeg, M. M., Smith, Y. R., Bueller, J. A., Xu, K., Xu, Y., Koeppe, R. A., Stohler, C. S., and Goldman, D. (2003) COMT val158met genotype affects μ -opioid neurotransmitter responses to a pain stressor. *Science* 299, 1240–1243.
43. Nackley, A. G., Tan, K. S., Fecho, K., Flood, P., Diatchenko, L., and Maixner, W. (2007) Catechol-O-methyltransferase inhibition increases pain sensitivity through activation of both β 2- and β 3-adrenergic receptors. *Pain* 128, 199–208.
44. Sheldrick, A. J., Krug, A., Markov, V., Leube, D., Michel, T. M., Zerres, K., Eggermann, T., and Kircher, T. (2008) Effect of COMT val158met genotype on cognition and personality. *Eur. Psychiatry* 23, 385–389.
45. Goldberg, T. E., Egan, M. F., Gscheidle, T., Coppola, R., Weickert, T., Kolachana, B. S., Goldman, D., and Weinberger, D. R. (2003) Executive subprocesses in working memory: Relationship to catechol-O-methyltransferase Val158Met genotype and schizophrenia. *Arch. Gen. Psychiatry* 60, 889–896.
46. Meyer-Lindenberg, A., Kohn, P. D., Kolachana, B., Kippenhan, S., McInerney-Leo, A., Nussbaum, R., Weinberger, D. R., and Berman, K. F. (2005) Midbrain dopamine and prefrontal function in humans: interaction and modulation by COMT genotype. *Nat. Neurosci.* 8, 594–596.
47. Li, Y., Yang, X., van Breemen, R. B., and Bolton, J. L. (2005) Characterization of two new variants of human catechol O-methyltransferase in vitro. *Cancer Lett.* 230, 81–89.
48. Hong, J. P., Lee, J. S., Chung, S., Jung, J., Yoo, H. K., Chang, S. M., and Kim, C. Y. (2008) New functional single nucleotide polymorphism (Ala72Ser) in the COMT gene is associated with aggressive behavior in male schizophrenia. *Am. J. Med. Genet., Part B* 147, 658–660.
49. Lee, S. G., Joo, Y., Kim, B., Chung, S., Kim, H. L., Lee, I., Choi, B., Kim, C., and Song, K. (2005) Association of Ala72Ser polymorphism with COMT enzyme activity and the risk of schizophrenia in Koreans. *Hum. Genet.* 116, 319–328.
50. Tanaka, Y., Hirata, H., Chen, Z., Kikuno, N., Kawamoto, K., Majid, S., Tokizane, T., Urakami, S., Shiina, H., Nakajima, K., Dhir, R., and Dahiya, R. (2007) Polymorphisms of catechol-O-methyltransferase in men with renal cell cancer. *Cancer Epidemiol., Biomarkers Prev.* 16, 92–97.
51. Tanaka, Y., Sasaki, M., Shiina, H., Tokizane, T., Deguchi, M., Hirata, H., Hinoda, Y., Okayama, N., Suehiro, Y., Urakami, S., Kawakami, T., Kaneuchi, M., Pookot, D., Igawa, M., Okuyama, A., Ishii, N., and Dahiya, R. (2006) Catechol-O-methyltransferase gene polymorphisms in benign prostatic hyperplasia and sporadic prostate cancer. *Cancer Epidemiol., Biomarkers Prev.* 15, 238–244.
52. Rutherford, K., Bennion, B. J., Parson, W. W., and Daggett, V. (2006) The 108M polymorph of human catechol O-methyltransferase is prone to deformation at physiological temperatures. *Biochemistry* 45, 2178–2188.
53. Vidgren, J., Svensson, L. A., and Liljas, A. (1994) Crystal structure of catechol O-methyltransferase. *Nature* 368, 354–358.
54. Rutherford, K., Le Trong, I., Stenkamp, R. E., and Parson, W. W. (2008) Crystal structures of human 108V and 108M catechol O-methyltransferase. *J. Mol. Biol.* 380, 120–130.
55. Alden, R. G., Parson, W. W., Chu, Z. T., and Warshel, A. (1996) Orientation of the OH dipole of tyrosine (M210) and its effect on electrostatic energies in photosynthetic bacterial reaction centers. *J. Phys. Chem.* 100, 16761–16770.
56. Levitt, M. (1990) ENCAD, Energy Calculations and Dynamics, Molecular Applications Group, Palo Alto, CA, and Yeda, Rehovot, Israel.
57. Beck, D. A. C., Alonso, D. O. V., and Daggett, V. (2000–2009) ilmm, in *Lucem Molecular Mechanics*, University of Washington, Seattle.
58. Beck, D. A., and Daggett, V. (2004) Methods for molecular dynamics simulations of protein folding/unfolding in solution. *Methods* 34, 112–120.
59. Levitt, M., Hirshberg, M., Sharon, R., and Daggett, V. (1995) Potential energy function and parameters for simulations of the molecular dynamics of proteins and nucleic acids in solution. *Comput. Phys. Commun.* 91, 215–231.
60. Levitt, M., Hirshberg, M., Sharon, R., Laidig, K. E., and Daggett, V. (1997) Calibration and testing of a water model for simulation of the molecular dynamics of proteins and nucleic acids in solution. *J. Phys. Chem. B* 101, 5051–5061.
61. Kell, G. S. (1967) Precise representation of volume properties of water at one atmosphere. *J. Chem. Eng. Data* 12, 66–69.
62. Beck, D. A., Armen, R. S., and Daggett, V. (2005) Cutoff size need not strongly influence molecular dynamics results for solvated polypeptides. *Biochemistry* 44, 609–616.
63. Beck, D. A., Jonsson, A. L., Schaeffer, R. D., Scott, K. A., Day, R., Toofanny, R. D., Alonso, D. O., and Daggett, V. (2008) Dynamematics: Mass annotation of protein dynamics and unfolding in water by high-throughput atomistic molecular dynamics simulations. *Protein Eng., Des. Sel.* 21, 353–368.
64. Armen, R. S., Bernard, B. M., Day, R., Alonso, D. O., and Daggett, V. (2005) Characterization of a possible amyloidogenic precursor in

- glutamine-repeat neurodegenerative diseases. *Proc. Natl. Acad. Sci. U.S.A.* 102, 13433–13438.
65. Hubbard, S. J., and Thornton, J. M. (1993) NACCESS, Department of Biochemistry and Molecular Biology, University College, London.
66. Tsuji, E., Okazaki, K., Isaji, M., and Takeda, K. (2009) Crystal structures of the Apo and Holo form of rat catechol-O-methyltransferase. *J. Struct. Biol.* 165, 133–139.
67. Hoel, P. G. (1954) Introduction to Mathematical Statistics, 2nd ed., John Wiley, New York.
68. Benson, N., and Daggett, V. (2008) Dynameomics: Large-Scale Assessment of Native Protein Flexibility. *Protein Sci.* 17, 2038–2050.
69. Masjost, B., Ballmer, P., Borroni, E., Zurcher, G., Winkler, F. K., Jakob-Roetne, R., and Diederich, F. (2000) Structure-based design, synthesis, and in vitro evaluation of bisubstrate inhibitors for catechol O-methyltransferase (COMT). *Chemistry* 6, 971–982.
70. Kaakkola, S., Gordin, A., and Mannisto, P. T. (1994) General properties and clinical possibilities of new selective inhibitors of catechol O-methyltransferase. *Gen. Pharmacol.* 25, 813–824.
71. Kaakkola, S., Teravainen, H., Ahtila, S., Rita, H., and Gordin, A. (1994) Effect of entacapone, a COMT inhibitor, on clinical disability and levodopa metabolism in parkinsonian patients. *Neurology* 44, 77–80.
72. Apud, J. A., Mattay, V., Chen, J., Kolachana, B. S., Callicott, J. H., Rasetti, R., Alce, G., Iudicello, J. E., Akbar, N., Egan, M. F., Goldberg, T. E., and Weinberger, D. R. (2007) Tolcapone improves cognition and cortical information processing in normal human subjects. *Neuropsychopharmacology* 32, 1011–1020.
73. Apud, J. A., and Weinberger, D. R. (2007) Treatment of cognitive deficits associated with schizophrenia: Potential role of catechol-O-methyltransferase inhibitors. *CNS Drugs* 21, 535–557.
74. DeVry, C. G., and Clarke, S. (1999) Polymorphic forms of the protein L-isoaspartate (D-aspartate) O-methyltransferase involved in the repair of age-damaged proteins. *J. Hum. Genet.* 44, 275–288.
75. Tsai, W., and Clarke, S. (1994) Amino acid polymorphisms of the human L-isoaspartyl/D-aspartyl methyltransferase involved in protein repair. *Biochem. Biophys. Res. Commun.* 203, 491–497.
76. Pettersen, E. F., Goddard, T. D., Huang, C. C., Couch, G. S., Greenblatt, D. M., Meng, E. C., and Ferrin, T. E. (2004) UCSF Chimera: A visualization system for exploratory research and analysis. *J. Comput. Chem.* 25, 1605–1612.