

## CORRECTIONS

3 $\beta$ -Hydroxy- $\Delta^5$ -steroid Dehydrogenase/3-Keto- $\Delta^5$ -steroid Isomerase from Bovine Adrenals: Mechanism of Inhibition by 3-Oxo-4-aza Steroids and Kinetic Mechanism of the Dehydrogenase, by Martin Brandt and Mark A. Levy\*, Volume 28, Number 1, January 10, 1989, pages 140–148.

Page 144. In Table II, the R<sub>1</sub> group of compound **1c** should be  $-\text{C}(\text{O})\text{N}(\text{CH}_2\text{CH}_3)_2$ ; this properly designates that compound **1c** is equivalent to 4-MA as indicated under Discussion.

Cooperativity in A-Tract Structure and Bending Properties of Composite T<sub>n</sub>A<sub>n</sub> Blocks, by Tali E. Haran and Donald M. Crothers\*, Volume 28, Number 7, April 4, 1989, pages 2763–2767.

Page 2764. In Table I, the correct identification of oligomers A<sub>8</sub>N<sub>13</sub> and A<sub>10</sub>N<sub>11</sub> is

|                                 |                       |
|---------------------------------|-----------------------|
| A <sub>8</sub> N <sub>13</sub>  | CCGGCCAAAAAAAACGCGCGC |
| A <sub>10</sub> N <sub>11</sub> | CCGGCCAAAAAAAACGCGCGC |

Platelet Receptor Recognition Domain on the  $\gamma$  Chain of Human Fibrinogen and Its Synthetic Peptide Analogues, by Marek Kloczewiak, Sheila Timmons, Maria A. Bednarek, Masato Sakon, and Jacek Hawiger\*, Volume 28, Number 7, April 4, 1989, pages 2915–2919.

Page 2916. In Table II, column 2, line 4, 0 should read 5.

Structural Characterization of the Interactions between Calmodulin and Skeletal Muscle Myosin Light Chain Kinase: Effect of Peptide (576–594)G Binding on the Ca<sup>2+</sup>-Binding Domains, by Steven H. Seeholzer and A. Joshua Wand\*, Volume 28, Number 9, May 2, 1989, pages 4011–4020.

Page 4012. In column 1, the sequence of the peptide should read as follows: Ac-M-K-R-R-W-K-K-N-F-I-A-V-S-A-A-N-R-F-G-NH<sub>2</sub>.

DNA Bending by the Bulge Defect, by Janet A. Rice and Donald M. Crothers\*, Volume 28, Number 10, May 16, 1989, pages 4512–4516.

Page 4514. The caption to Figure 4 should read as follows: Mobilities of 19–22 bp sequences containing a bulged adenine and an A<sub>6</sub> tract (see Figure 2) measured relative to *Bam*HI. (□) 19 base pair monomer fragment; (▽) 20 base pair monomer fragment; (▲) 21 base pair monomer fragment; (○) 22 base pair monomer fragment. The mobilities of the (20)<sub>n</sub> and (21)<sub>n</sub> fragments are almost identical.