

- Steigemann, W. (1974) Ph.D. Dissertation, Technische Universität, München.
- Taylor, S. S. (1989) *J. Biol. Chem.* 263, 8443–8446.
- Toda, T., Cameron, S., Sass, P., Zoller, M., Scott, J. D., McMullen, B., Hurwitz, M., Krebs, E. G., & Wigler, M. (1987a) *Mol. Cell. Biol.* 7, 1371–1377.
- Toda, T., Cameron, S., Sass, P., Zoller, M., & Wigler, M. (1987b) *Cell* 50, 277–285.
- Vieira, J., & Messing, J. (1987) *Methods Enzymol.* 153, 3–11.
- Westbrook, E. M. (1985) *Methods Enzymol.* 114, 187–196.
- Wierenga, R. K., & Hol, W. G. J. (1983) *Nature* 302, 842–844.
- Yamano, S., Tanaka, K., Matsumoto, K., & Toh-e, A. (1987) *Mol. Gen. Genet.* 210, 413–418.
- Zoller, M. J., Kuret, J., Cameron, S., Levin, L., & Johnson, K. E. (1988) *J. Biol. Chem.* 263, 9142–9148.

CORRECTION

Secondary Structure Analysis of the Scrapie-Associated Protein PrP 27–30 in Water by Infrared Spectroscopy, by Byron W. Caughey,* Aichun Dong, Kolari S. Bhat, Darwin Ernst, Stanley F. Hayes, and Winslow S. Caughey, Volume 30, Number 31, August 6, 1991, pages 7672–7680.

Page 7674. Due to a printing error, the detail in Figure 2 was lost. An appropriate representation is shown below.



FIGURE 2: Electron microscopic analysis of PrP-res 27–30. Appearance of PrP-res 27–30 aggregates negatively stained with 0.5% ammonium molybdate. Magnification = 78750 $\times$. Bar = 100 nm.