

Mutation of R116C Results in Highly Oligomerized α A-Crystallin with Modified Structure and Defective Chaperone-like Function[†]

Nilufer P. Shroff, Mary Cherian-Shaw, Sibes Bera, and Edathara C. Abraham*

Department of Biochemistry and Molecular Biology, Medical College of Georgia, Augusta, Georgia 30912-2100

Received July 19, 1999; Revised Manuscript Received October 6, 1999

ABSTRACT: An autosomal dominant congenital cataract in human is associated with mutation of Arg-116 to Cys (R116C) in α A-crystallin. To investigate the molecular basis of cataract formation, rat α A-crystallin cDNA was cloned into pET-23d(+), and the site-directed mutants S142C (similar to wild-type human α A) and R116C/S142C or R116C (matrix to human R116C variant) were generated. These were expressed in *E. coli* and the recombinant α A-crystallins purified by Sephacryl size-exclusion chromatography. The chaperone-like function of mutant R116C determined at 37 °C with insulin and alcohol dehydrogenase as target proteins was about 40% lower than those of wild-type and mutant S142C. Based on size-exclusion chromatography data, the oligomeric size of the R116C mutant was about 2000 kDa at 25 °C, 1400 kDa at 37 °C, and 900 kDa at 45 °C. In comparison, α A-wild-type and α A-S142C ranged from 477 to 581 kDa. Heat stability studies corroborated the effect of temperature on the dynamic quaternary structure of the R116C mutant. Circular dichroism spectra showed secondary and tertiary structural changes, and ANS fluorescence spectra showed loss of surface hydrophobicity in the R116C mutant. These findings suggest that the molecular basis for the congenital cataract with the α A-R116C mutation is due to the generation of a highly oligomerized α A-crystallin having a modified structure and decreased chaperone-like function.

α -Crystallin constitutes about 25–50% of the total soluble protein in the lens and is believed to be a major structural element in the highly concentrated protein matrix which is essential for maintenance of transparency. Native α -crystallin is composed of α A- and α B-subunit polypeptides with molecular weights of approximately 20 000 each (1, 2). α A and α B share 57% identity in their polypeptide chains (1). The human α A and α B genes have been localized to chromosomes 21 (17 in rat) and 11, respectively (3, 4). α A-Crystallin is expressed extralenticularly in small amounts in the brain, liver, spleen, retina, and thymus (5, 6). α B-Crystallin is most abundant in the lenticular tissue, and is also expressed in cardiac muscle, skeletal muscle, brain, retina, and lung (7, 8). There is involvement of hsp and α B-crystallin in major cytomorphological organization during normal growth and under pathological conditions (9–13). α -Crystallin oligomers have a molecular mass in the 300–800 kDa range (14, 15). Both subunits of α -crystallin share sequence similarity with small heat-shock protein, and this sequence homology is confined to a stretch of 100 amino acids in the C-terminal domain (16). In addition to its structural function, Horwitz (17) demonstrated that α -crystallin in vitro has chaperone-like activity, a feature which allows it to suppress aggregation of other proteins. The presence of an active molecular chaperone α -crystallin may be essential for maintaining lens transparency. Brady et al. (18) generated a knock-out mouse homologous for the α A-crystallin gene. The α A-crystallin $-/-$ mice had no detect-

able α A-crystallin in their lens, and the lens size was slightly smaller than the wild-type lenses but had normal gross structure. However, several weeks after birth, lens opacification and dense inclusion bodies that contained insoluble α B-crystallin in the lens fiber cells were seen. This was the first in vivo report showing the necessity of α A-crystallin for maintenance of lens clarity and solubility of α B-crystallin.

Due to its large oligomeric size and its microheterogeneity, the quaternary structure has eluded both NMR and crystallographic studies. Haley et al. (19), using cryo-electron microscopy (cryo-EM) and image processing, analyzed the structure of recombinant human α B-crystallin. They proposed that α B-crystallin has an asymmetric quaternary structure with variable monomer binding and it forms roughly spherical particles with a large central cavity and a protein shell which has a lot of structural divergence. The proposed structure of the recombinant α B-crystallin is similar to the crystal structure of a small heat-shock protein from *Methanococcus jannaschii*. It is composed of 24 monomers and assumes a hollow spherical complex with octahedral symmetry having 8 trigonal and 6 square windows (20). In α A-crystallin, site-directed spin-label studies of the amino acid residues between 109 and 120 revealed the presence of a β -strand containing the buried residues R112 and R116 (21). Further studies demonstrated that residues between 84 and 120 had antiparallel β -sheets consisting of three strands arranged in consecutive β -hairpins (22).

Congenital cataracts are a common cause of blindness in infants (23). Autosomal dominant congenital cataract is the most common familial form (24). Several types of mutations occurring in mammalian crystallin genes are associated with

[†] This work was supported by Grants EY 11352 and EY 07394 from the National Institutes of Health.

* To whom correspondence should be addressed.

hereditary cataracts (25–30). An autosomal dominant cataract associated with a missense mutation in the human α A-crystallin gene has been reported recently (30). In this form of cataract, the α A-crystallin mutation Arg-116 to Cys (R116C) was observed. However, it has not been shown how this mutation in α A-crystallin leads to cataract in the affected persons and whether this mutation causes structural and functional changes in α A-crystallin. Rat α A-crystallin has 98% sequence homology with human α A-crystallin, and residues 109–120 show 100% homology. Due to this strong sequence homology and since it is difficult to obtain a human α A-crystallin clone, we have cloned rat α A-crystallin and generated a rat α A-crystallin containing the same mutation. This study demonstrates that the mutant R116C exists as a highly oligomerized protein with perturbed secondary and tertiary structures and decreased chaperone-like activity.

EXPERIMENTAL PROCEDURES

Cloning and Site-Directed Mutagenesis of Rat α A-Crystallin DNA. Total RNA was isolated using Trizol Reagent (Life Technologies) from a 1-month-old fresh rat lens. RT-PCR was done using the upstream primer 5'-CCC AAG CTT TCA GGA AGG CAG ACT CTT T-3' containing an *EcoRV* site and the downstream primer 5'-CCC GAT ATC ATG GAC GTC ACC ATC CA-3' containing a *HindIII* site. The α A-crystallin DNA was ligated into *EcoRV*-digested pGEM-T vector (Promega) and used to transform DH5 α *E. coli* cells (Stratagene). Plasmid DNA from the resulting clones was sequenced by automated DNA sequencing (ABI Prism 377 DNA sequencer). To subclone into the expression vector, pET-23d(+) (Novagen) was sequentially digested with *NcoI*, mung bean nuclease, and *HindIII*, and the α A-crystallin DNA was excised from pGEM-T using *HindIII* and *EcoRV* and ligated into this vector. The QuickChange Site-Directed Mutagenesis Kit (Stratagene) was used to generate the mutants α A-S142C and α A-R116C/S142C (abbreviated as R116C), and their nucleotide sequences were confirmed by automated DNA sequencing.

Expression and Purification of Recombinant Wild-Type and Mutant α A-Crystallins. BL21(DE3)pLysS *E. coli* cells (Novagen) were used as the expression host. These cells were transformed with the appropriate amplicons and grown in 500 mL of Luria broth at 37 °C with aeration. At an OD_{600 nm} \approx 0.6, the T7 promoter was induced using 0.5 mM isopropyl-1-thio- β -D-galactopyranoside and incubation continued for 4 h. The cells were harvested and resuspended in 10 mL of lysis buffer [20 mM Tris, pH 7.4, 1 mM EDTA, 4 μ g/mL pepstatin A, 4 μ g/mL aprotinin, 0.1 M PMSF (phenylmethylsulfonyl fluoride)]. After 3 freeze–thaw cycles, 800 units of DNase (Sigma), 5 units of RNase (Sigma), and 10 mM MgCl₂ were added to the cell lysate, and incubation was carried out at room temperature on the nutator for 2 h with subsequent centrifugation at 28000g for 45 min. The protein was filtered through a 0.2 μ m filter and loaded onto a Sephacryl S-300HR molecular sieve column (120 \times 2.6 cm). SDS–PAGE was used to examine the fractions; those containing the wild-type and mutant α A-crystallins were pooled and concentrated by ultrafiltration in an Amicon stirred cell.

SDS–PAGE and Western Blot. SDS–PAGE was performed using a 12% separating gel and a 4% stacking gel

under reducing conditions, according to the method of Laemmli (31). Three micrograms was run at 200 V for 45 min, and the proteins were then blotted onto a Millipore poly(vinylidene difluoride) (PVDF) membrane. Rabbit anti- α A/ α B-crystallin peptide polyclonal antibody (Stress Gene) was used as the first antibody, and the crystallins were visualized by incubating with Western Blue stabilized substrate for alkaline phosphatase (Protoblot System, Promega).

Size-Exclusion Chromatography for Determining Molecular Masses. The aggregate sizes were determined by size-exclusion chromatography performed with a water-jacketed Sephacryl-S-300 HR column (1.6 \times 100 cm) at different temperatures (25, 37, and 45 °C). A constant flow-rate (34 mL/h) and temperatures were maintained by a peristaltic pump and water bath, respectively. Protein, 2 mg in 1 mL of PBS buffer, pH 7.4, was incubated for 1 h at the above temperatures and then applied to the column which had been equilibrated to the desired temperatures. The elution was performed with the same buffer, and fractions of 1 mL each were collected and monitored at 280 nm. High molecular mass standards (Sigma) were used to calibrate the column. Molecular masses (*M*) were determined from a plot of log *M* vs *V*_e/*V*_o (*V*_e = elution volume and *V*_o = void volume).

Circular Dichroism Measurements. To investigate secondary and tertiary structural changes of recombinant rat α A-wild-type, α A-S142C, and α A-R116C, their near- and far-UV CD spectra were recorded at 25 and 37 °C using a Jasco 710 spectropolarimeter. Protein concentrations of 0.5 and 0.1 mg/mL in 50 mM potassium phosphate buffer, pH 7.4, were used for recording the near- and far-UV CD spectra, respectively. The reported CD spectra are the average of five scans smoothed. Secondary structure parameters were estimated by the computer program PROSEC derived from Yang et al. (32).

Determination of Molecular Chaperone-like Activity of α A-Crystallin. The aggregation and denaturation of insulin or alcohol dehydrogenase (ADH) in the absence and in the presence of recombinant wild-type and mutant α A-crystallins was measured in a Shimadzu UV160 spectrophotometer equipped with a temperature-regulated cell holder. Ratios of 1:1 and 2:1 α A-crystallin to insulin and α A-crystallin to ADH were used to determine chaperone-like activity. Insulin was denatured using 20 mM DTT and ADH using 10 mM EDTA at 37 \pm 1 °C, and light scattering was monitored at 360 nm.

Determination of Protein Stability of α A-Wild-Type and Its Mutants. The heat stability of 1 mg/mL protein in PBS, pH 7.4, was tested at 25, 37, and 45 °C by monitoring light scattering at 360 nm. Stability studies were also done in the presence of EDTA or DTT.

Tryptophan Fluorescence Measurements. Fluorescence measurements were performed with a Shimadzu RF-540 spectrofluorophotometer. Protein samples of 0.1 mg/mL in 50 mM phosphate buffer, pH 7.4, were used. Excitation was fixed at 295 nm and emission scanned between 310 and 400 nm.

ANS Fluorescence Measurements. The hydrophobic probe ANS binding studies were done as follows: to 1 mL of protein (0.1 mg/mL) solution in 50 mM phosphate buffer, pH 7.4, was added 10 μ L of 20 mM methanolic solution of ANS and incubated for 2 h. Emission spectra were recorded




FIGURE 1: SDS-PAGE and Western blot of bacterial expressed rat α A-wild-type and mutants S142C and R116C/S142C. Lane 2: α L fraction from rat lens. Lanes 3–5: total cell lysate of expressed α A-crystallins. Lanes 6–8: size exclusion chromatography purified α A-crystallins. Lanes 9–11: Western blot of lanes 6–8 with anti- α A/ α B polyclonal antibody.

between 420 and 600 nm while the excitation wavelength was fixed at 390 nm.

RESULTS

Cloning, Mutagenesis, Expression, and Purification of Wild-Type and Mutant Rat α A-Crystallins. The rat α A-crystallin DNA obtained by RT-PCR was cloned into the expression vector pET-23d(+). The nucleotide sequence was confirmed by automated sequencing and found to be identical to the coding region of GenBank sequence accession number U47922. Rat α A-crystallin has only one cysteine (position 131) whereas human α A-crystallin has two cysteines (positions 131 and 142). Therefore, a cysteine was introduced at position 142 (S142C) in the rat α A-crystallin in order to mimic the number of cysteines in the native human α A-crystallin. Then, using the α A-S142C mutant amplicon as the template, a second mutant, R116C/S142C (R116C), was generated. This mutant is structurally similar to the genetic variant previously reported to have caused congenital cataract in humans (30). The expression host BL21(DE3)pLysS *E. coli* cells were transformed with pET-23d(+)- α A amplicons. After induction with IPTG and further growth, the cells were pelleted and lysed, and protein was analyzed by SDS-PAGE (Figure 1, lanes 3–5). The recombinant α A-crystallins and the α A-crystallin from the total rat α L-crystallin fraction corresponded approximately to the 20 kDa band of the marker. The expressed proteins were purified by Sephacryl gel permeation chromatography, which resulted in about 95% pure recombinant α A-crystallin (Figure 1, lanes 6–8). During purification, it was noticed that the α A-R116C mutant eluted ahead of the wild-type and the α A-S142C mutant, indicating that α A-R116C was more oligomerized than the others. After purification, Western blotting with a rabbit anti- α A/ α B-crystallin peptide polyclonal antibody further confirmed the identity of the recombinant α A-crystallin (Figure 1, lanes 9–11).

Molecular Masses of the Recombinant Wild-Type and Mutant α A-Crystallins. To ascertain if mutation of Arg-116 to Cys influenced α A-crystallin oligomerization, size-exclu-




FIGURE 2: Determination of the oligomeric sizes of recombinant α A-crystallins by size exclusion chromatography with a temperature-regulated Sephacryl S-300 HR column at 25, 37, and 45 °C. The numbers on the top refer to the masses of the molecular weight protein standards.

Table 1: Molecular Masses^a (kDa) vs α A-Wild-Type (α A-WT) and Its Mutants

temp (°C)	α A-WT	α A-S142C	α A-R116C
25	581	528	2000
37	581	489	1403
45	516	477	903

^a Molecular masses (*M*) were determined from a plot of log *M* vs *V_e*/*V_o* of molecular mass standards.

sion chromatography was performed after protein purification. α A-Wild-type, α A-S142C, and α A-R116C were analyzed at 25, 37, and 45 °C; representative chromatograms are given in Figure 2 and their molecular masses are given in Table 1. At 25 °C, the size of the R116C mutant was close to 2000 kDa whereas for the wild-type and the S142C mutant the values were 581 and 528 kDa, respectively. At 37 and 45 °C, the size of the R116C mutant decreased to about 1400 and 900 kDa, respectively, while the size of the wild-type and S142C remained nearly the same.

Conformational Studies. The far-UV and near-UV CD spectra of rat α A-wild-type and S142C and R116C mutants, recorded at 25 and 37 °C, are shown in Figure 3. The far-UV profiles indicate the minima to be around 217 nm, which is the characteristic of mainly β -conformation even though their mean residue ellipticity values have been found to be slightly different (Figure 3A,B). Estimates of the secondary structure parameters were done by the computer program PROSEC. At 25 °C, contents of α -helix, β -sheet, β -turn, and random coil for α A-wild-type were 9.3, 44.1, 21.4, and 22.2%, respectively; similar levels of the various structural components were found in the two mutants. These data are in good agreement with the published values for α A-crystallin (33). However, at 37 °C, the data differ markedly. While the α -helix conformation remained the same, the β -sheet content was reduced by 50% with concomitant increase in random coil conformation. Near-UV CD spectra of α A-wild-type, S142C mutant, and R116C mutant (Figure 3C,D) show the maxima at 259 and 265 nm, which are the characteristics of phenylalanine fine structure. The remaining transitions between 270 and 290 nm arise from Tyr and/or Trp residues. The negative vibronic transition at around 293 nm is considered to be due to the contribution of Trp residues. But, for α A-R116C there are no distinct bands due to Tyr and/or Trp. This indicates different Tyr and Trp microenvironments in this mutant when compared with α A-




FIGURE 3: Far-UV CD spectra for α A-wild-type and S142C and R116C/S142C mutants at (A) 25 °C and (B) 37 °C. Near-UV spectra for these α A-crystallins at (C) 25 °C and (D) 37 °C. The protein concentrations for far-UV CD and near-UV CD measurements were 0.1 and 0.5 mg/mL, respectively.




FIGURE 4: Trp fluorescence spectra (A) and ANS fluorescence spectra (B) for α A-wild-type and S142C and R116C/S142C mutants at 37 °C. For Trp fluorescence measurements, the excitation wavelength was 295 nm, and for ANS fluorescence, 390 nm and the protein concentration was 0.1 mg/mL.

wild-type and α A-S142C. Figure 4 represents fluorescence monitoring before and after ANS binding. Intrinsic Trp fluorescence spectra showed similar wavelength maxima and fluorescence intensity at around 336 nm for all three α A-crystallins (Figure 4A). ANS fluoresces at about 512 nm, and in the presence of α -crystallin, it fluoresces significantly intense and blue-shifts to 483 nm (Figure 4B). ANS binding to the R116C mutant was decreased significantly, implying

that the available hydrophobic surface is decreased appreciably.

Molecular Chaperone-like Activity of the Wild-Type and the Mutant S142C and R116C α A-Crystallins. Chaperone-like activity was measured at 37 °C by in vitro assays using insulin and ADH as target proteins with two different proportions of α A:target protein (1:1 and 2:1). Different preparations of the α A-wild-type, mutant α A-S142C, and




FIGURE 5: Chaperone-like activity of α A-wild-type and S142C and R116C/S142C mutants at 37 °C using insulin as the target protein in the presence of DTT (A, B) and ADH as the target protein in the presence of EDTA (C, D). Each curve is the mean of chaperone assays done on three different preparations, and bars represent standard deviations.

mutant α A-R116C were assayed, and the means \pm SD of protein denaturation curves are presented (Figure 5). In the absence of α A-crystallin, insulin is fully aggregated by DTT at 37 °C whereas the presence of α A-crystallin prevents this aggregation (Figure 5A,B). This inhibitory effect on insulin aggregation is taken as the chaperone-like activity (chaperone activity or chaperone function). The chaperone activity of α A-S142C was not significantly different from α A-wild-type. α A-R116C, on the other hand, showed about 40% loss in chaperone activity when the α -crystallin-to-insulin ratio was 1:1 (Figure 5A). A similar, but slightly smaller, decrease of chaperone activity was observed at a α :insulin ratio of 2:1 (Figure 5B). ADH was used as a target protein in the presence of EDTA with no DTT. Chaperone activity assays were done with ADH as the target protein, which gave similar results (Figure 5C,D).

Heat Stability of the Wild-Type and Mutant α A-Crystallins. Heat stability of the wild-type and the R116C and S142C mutants was determined at 25, 37, and 45 °C by monitoring light scattering at these temperatures using the same setup as in chaperone assays in the absence of any target proteins; the data are presented in Figure 6. It is well-known that the absorbance at 360 nm is due to light scattering of the protein molecule which is directly proportional to the size. As expected, the R116C mutant gave higher absorbance than the wild-type and the S142C mutant. At all three temperatures, the wild-type and the S142C mutant showed no evidence of denaturation and aggregation to molecules or disaggregation to smaller size. The R116C mutant, on the other hand, behaved very differently at 37 and 45 °C; at these temperatures, light scattering significantly decreased, indicat-

ing dissociation of the protein into smaller molecules. This observation corroborates with our size exclusion chromatography experiments which also showed dissociation of the R116C super-oligomers into smaller molecules at 37 and 45 °C. The presence of DTT or EDTA did not show any additional effect on the protein stability at all temperatures.

DISCUSSION

Autosomal dominant congenital cataracts are the major form of familial cataract. Mutations in mammalian crystallin genes associated with hereditary cataracts have been reported in human and other species (23–30). The basis for the present study is a recent report by Litt et al. (30) of autosomal dominant congenital cataract in human associated with a missense mutation, R116C, in α A-crystallin. These authors have speculated that the mutant α A-polypeptide might have an increased tendency to aggregate because of its decreased positive charge and this may result in the loss of chaperone activity. We investigated this possibility by generating a rat α A-crystallin with strong structural similarity to the above mutant. In fact, three different recombinant rat α A-crystallins were generated: α A-wild-type, mutant α A-S142C (structurally similar to normal human α A-crystallin), and mutant α A-R116C (structurally similar to the genetic variant causing cataract). By utilizing these α A-crystallins, we were able to conclude that the variant α A-R116C forms a highly oligomerized protein (Figure 2) with modified structure (Figures 3 and 4) and partially impaired chaperone function (Figure 5). It is also noteworthy that the dynamic quaternary structure of the R116C mutant is temperature-dependent (Figure 6);




FIGURE 6: Heat stability curves of α A-wild-type and S142C and R116C/S142C mutants at 25, 37, and 45 °C without the target protein. Light scattering was recorded at 360 nm.

at physiological temperature, it has more than twice the size of α A-wild-type.

The mutant α A-R116C had an oligomeric size 2–4 times larger (depending on the temperature) than either the α A-wild-type or the α A-S142C (Figure 2). Proper protein assembly may depend on a salt bridge involving the positively charged guanidino group of Arg-116, the absence of which probably explains increased oligomeric size. α A-R116C, which is conserved in many species, lies in a buried region of the protein (21). Arginines in general are known to be involved in salt bridges and hydrogen bonds. Before the discovery of the autosomal dominant congenital cataract associated with a R116C mutation in α A-crystallin, Berengian et al. (21) made a series of cysteine mutants extending between amino acid residues 109 and 120 in rat α A-crystallin. The purpose of their study was to apply site-directed spin-labeling in the form of attached nitroxides to investigate the role of the above sequence in the assembly of α A-crystallin quaternary structure. Unlike the mutants used in the present study, they mutated Cys-131 to Ala and did not mutate Ser-142 to Cys. The mutant R116C with a nitroxide attached to the Cys studied by these investigators had a molecular mass 30% higher and chaperone activity similar to the wild-type α A-crystallin. This contradicts our observation, probably because the mutant characterized by the above authors has the mutated cysteine carrying a nitroxide and this additional derivatization seems to suppress excessive protein aggregation. Moreover, the rat α A-crys-

tallin R116C mutant used in our study has cysteines in two other positions, namely, 131 and 142, similar to the human α A-crystallin. Since there are a total of three Cys in R116C, it is possible that aggregation of the R116C mutant involves inter- or intrapolypeptide disulfides. However, the oligomeric sizes and chaperone activity were not influenced by DTT treatment (Figures 2 and 5), indicating that protein disulfides are not formed, or, if formed, do not affect oligomerization and chaperone function.

There is strong evidence that mutation of Arg-116 to Cys causes conformational changes in the α A-crystallin (Figures 3 and 4). Since chaperone assays were done at 37 °C, we performed all the biophysical studies at 25 and 37 °C. CD studies clearly showed structural and conformational differences at 37 °C, in particular. Although intrinsic tryptophan fluorescence spectra (Figure 4A) did not suggest the presence of any significant conformational changes, near-UV CD spectra implicated changes in Trp and Tyr microenvironments. Near-UV CD spectra are sensitive to changes in the fine structure and microenvironment of Trp and Tyr. Interestingly, ANS binding showed significantly reduced surface hydrophobicity in the R116C mutant (Figure 4B). In fact, hydrophobic surfaces are believed to be needed for proper chaperoning function, and their loss is expected to reduce chaperone-like function. However, mutation of R116C by itself is not expected to decrease the availability of hydrophobic surfaces unless it causes changes in protein conformation.

Another most noteworthy example of a single point mutation in one of the α -crystallin subunits is the missense mutation R120G in α B-crystallin which has been shown to cosegregate with a desmin-related myopathy in a French family (34). This mutant α B-crystallin has been characterized recently by Bova et al. (35), who showed irregular quaternary structure and complete loss of chaperone function. Interestingly, Arg-120, in α B-crystallin, lies in the most conserved region of the small hsp family (36). It is noteworthy that the oligomeric size of this mutant was nearly 1400 kDa. Arg-116 of α A-crystallin also lies in a conserved region of residues 109 through 120 (21) and was shown to exist in a buried environment with no access to aqueous solvent (21). Buried charged amino acids are believed to form salt bridges with residues of opposite charges, and the absence of such salt bridges could lead to protein conformational changes. The site-directed D69S mutant of α A-crystallin resulted in reduced chaperone-like function (37). Thus, it appears that α -crystallin may not withstand mutation of a charged residue like Arg to an uncharged residue, within a conserved region, in particular.

To assess the chaperone-like activity at physiological temperature, we have used two independent assay systems with insulin and ADH as target proteins. Both systems showed about 40% loss in chaperone activity when the α A:target protein proportion (w/w) was 1:1 (Figure 5). The physiological significance of this finding comes from the fact that significant loss of chaperone activity was seen when the α -crystallin proportion in the assay system was similar to the proportion in human lenses. It is also noteworthy that loss in chaperone activity, although smaller (21–27%), could be demonstrated even when α A:target protein proportion was raised to 2:1, which is significantly higher than in human lens. It was crucial that we performed all the biophysical

studies at 37 °C to relate structural changes to a decrease in function at this temperature. Structural and conformational changes are present in the R116C mutant at 37 °C (Figures 3 and 4), although the oligomeric size is smaller at 37 °C than at 25 °C due to some dissociation of the super-oligomers. From the present study, it appears that the molecular basis for the development of congenital cataract in the affected individuals is the formation of highly oligomerized α A-crystallin with structural and conformational changes and defective chaperone-like function. However, it can be argued that, since α B-crystallin is a better chaperone than α A-crystallin, the loss of chaperone activity of the R116C mutant may play only a small role in cataractogenesis. On the other hand, it is known that this cataract develops mostly in infancy and late childhood (30) and the proportion of α A-R116C: α B would be 2:1 in these patients (38). So, it is conceivable that 40% loss in chaperone activity contributes to cataractogenesis.

ACKNOWLEDGMENT

We thank Prof. Leon Dure and Ms. Tracy Grant of the University of Georgia for assistance in CD measurements.

REFERENCES

- van der Ouderaa, F. J., de Jong, W. W., Hilderink, A., and Bloemendal, H. (1973) *Eur. J. Biochem.* 39, 207–222.
- van der Ouderaa, F. J., de Jong, W. W., and Bloemendal, H. (1974) *Eur. J. Biochem.* 49, 157–168.
- Hawkins, J. W., van Keuren, M. L., Piatigorsky, J., Law, M. L., Patterson, D., and Kao, F. T. (1987) *Hum. Genet.* 76, 375–380.
- Ngo, J. T., Klisak, I., Dubin, R. A., Piatigorsky, J., Mohandas, T., Sparkes, R. S., and Bateman, J. B. (1989) *Genomics* 5, 665–669.
- Deretic, D., Aebersold, R. H., Morrison, H. D., and Papenmaster, D. S. (1994) *J. Biol. Chem.* 269, 16853–16861.
- Srinivasan, A. N., Nagineni, C. N., and Bhat, S. P. (1992) *J. Biol. Chem.* 267, 23337–23341.
- Bhat, S. P., and Nagineni, C. N. (1989) *Biochem. Biophys. Res. Commun.* 158, 319–325.
- Dubin, R. A., Wawrousek, E. F., and Piatigorsky, J. (1989) *Mol. Cell. Biol.* 9, 1083–1091.
- Lowe, J., McDermot, H., Pike, I., Spendlove, I., Landon, M., and Mayer, R. J. (1992) *J. Pathol.* 166, 61–68.
- Iwaki, T., Wisniewski, T., Iwaki, A., Corbin, E., Tomokane, N., Tateishi, J., and Goldman, J. E. (1992) *Am. J. Pathol.* 140, 345–356.
- Scotting, P., McDermott, H., and Mayer, R. J. (1991) *FEBS Lett.* 285, 75–79.
- Iwaki, T., Iwaki, A., Liem, R. K., and Goldman, J. E. (1991) *Kidney Int.* 40, 52–56.
- Renkawek, K., Bosman, G. J., and de Jong, W. W. (1994) *Acta Neuropathol.* 87, 155–160.
- Harding, J. J., and Crabbe, M. J. C. (1984) The Lens: development, protein, metabolism and cataract. in *The Eye*, pp 207–492, Academic Press, New York.
- Augusteyn, R. C., Parkhill, E. M., and Stevens, A. (1992) *Exp. Eye Res.* 54, 219–228.
- Ingolia, T. D., and Craig, E. A. (1982) *Proc. Natl. Acad. Sci. U.S.A.* 79, 2360–2364.
- Horwitz, J. (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89, 10449–10453.
- Brady, J. P., Garland, D., Douglas-Tabor, Y., Robison, W. G., Groome, A., and Wawrousek, E. F. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94, 884–889.
- Haley, D. A., Horwitz, J., and Stewart, P. L. (1998) *Invest. Ophthalmol. Visual Sci.* 39, S870.
- Kim, K. K., Kim, R., and Kim, S. (1998) *Nature* 394, 595–599.
- Berengian, A. R., Bova, M. P., and Mchaourab, H. S. (1997) *Biochemistry* 36, 9951–9957.
- Koteiche, H. A., Berengian, A. R., and Mchaourab, H. S. (1998) *Biochemistry* 37, 12681–12688.
- Lund, A. M., Eiberg, H., Rosenberg, T., and Warburg, M. (1992) *Clin. Genet.* 41, 65–69.
- Reese, P., Truck-Mullerr, C., and Maumenee, I. (1987) *Arch. Ophthalmol.* 105, 1382–1384.
- Rodriguez, I. R., Gonzalez, P., Zigler, J. S., Jr., and Borrás, T. (1992) *Biochim. Biophys. Acta* 1180, 44–52.
- Cartier, M., Breitman, M. L., and Tsui, L. C. (1992) *Nat. Genet.* 2, 42–45.
- Chambers, C., and Russell, P. (1991) *J. Biol. Chem.* 266, 6742–6746.
- Litt, M., Carrero-Valenzuela, R., LaMorticella, D. M., Schultz, D. W., Mitchell, T. N., Kramer, P., and Maumenee, I. H. (1997) *Hum. Mol. Genet.* 6, 665–668.
- Brakenhoff, R. H., Henskens, H. A., van Rossum, M. W., Lubsen, N. H., and Schoenmakers, J. G. (1994) *Hum. Mol. Genet.* 3, 279–283.
- Litt, M., Kramer, P., LaMorticella, D. M., Murphey, W., Lovrien, E. W., and Weleber, R. G. (1998) *Hum. Mol. Genet.* 7, 471–474.
- Laemmli, U. K. (1970) *Nature* 227, 680–685.
- Yang, J. T., Wu, C. H. C., and Martinez, H. M. (1986) *Methods Enzymol.* 130, 208–269.
- Sun, T., Das, B. K., and Liang, J. N. (1997) *J. Biol. Chem.* 272, 6220–6225.
- Vicart, P., Caurn, A., Guicheney, P., Li, A., Prevost, M. C., Faure, A., Chateau, D., Chapou, F., Tone, F., Dupret, J. M., Paulin, D., and Fardeau, M. (1998) *Nat. Genet.* 20, 92–95.
- Bova, M. P., Yaron, O., Huang, Q., Ding, L., Haley, D., Stewart, P., and Horwitz, J. (1999) *Proc. Natl. Acad. Sci. U.S.A.* 96, 6137–6142.
- de Jong, W. W., Leurissen, J. A. M., and Voorter, C. E. M. (1993) *Mol. Biol. Evol.* 10, 103–126.
- Smulders, R. H. P. H., Merck, K. B., Aendekerk, J., Horwitz, J., Takemoto, L., Slingsby, C., Bloemendal, H., and de Jong, W. W. (1995) *Eur. J. Biochem.* 232, 834–838.
- Swamy, M. S., and Abraham, E. C. (1991) *Curr. Eye Res.* 10, 213–220.

B1991656B