Haloalkane Dehalogenase LinB from *Sphingomonas paucimobilis* UT26: X-ray Crystallographic Studies of Dehalogenation of Brominated Substrates, by Victor A. Streltsov,* Zbyněk Prokop, Jiří Damborský,* Yuji Nagata, Aaron Oakley, and Matthew C. J. Wilce*, Volume 42, Number 34, September 2, 2003, pages 10104–10112.

Page 10110 and page 10111. While the figure legends are correct, the graphical components of Figures 4 and 5 should be switched and reorganized. Panels A and B of Figure 4 are identical to panels C and D of Figure 5, respectively, but should be panels A and B of Figure 5, respectively. Panels A and B of Figure 5 should be panels C and D of Figure 4, respectively. Correct versions of Figures 4 and 5 follow.




FIGURE 4: Dehalogenation of 1,2-dibromopropane (RT = 4.65 min) and formation of two products (RT = 3.71 and 3.92 min) at 20, 80, and 160 min (A). The mass spectrum of the product with a retention time of 3.71 min identified as 1-bromopropane-2-ol (B) and that with a retention time of 3.92 min as 2-bromopropane-1-ol (C) and 1,2-propanediol (D). 1,2-Propanediol was identified in the separate experiment conducted at a high protein concentration (90 μ M) and with a long reaction time (1020 min).




FIGURE 5: Dehalogenation of 2,3-dibromopropane (RT = 4.77 min) and formation of the product (RT = 4.20 min) at 2, 5, and 10 min (A). The mass spectrum of the product identified as 2-bromo-2-propene-1-ol.

BI0330141

10.1021/bi0330141 Published on Web 10/07/2003