

Quiescin Sulphydryl Oxidase from *Trypanosoma brucei*: Catalytic Activity and Mechanism of a QSOX Family Member with a Single Thioredoxin Domain[†]

Vamsi K. Kodali and Colin Thorpe*

Department of Chemistry and Biochemistry, University of Delaware, Newark, Delaware 19716

Received December 29, 2009; Revised Manuscript Received February 1, 2010

ABSTRACT: Quiescin sulphydryl oxidase (QSOX) flavoenzymes catalyze the direct, facile, insertion of disulfide bonds into reduced unfolded proteins with the reduction of oxygen to hydrogen peroxide. To date, only QSOXs from vertebrates have been characterized enzymatically. These metazoan sulphydryl oxidases have four recognizable domains: a redox-active thioredoxin (Trx) domain containing the first of three CxxC motifs (C^I–C^{II}), a second Trx domain with no obvious redox-active disulfide, a helix-rich domain, and then an Erv/ALR domain. This last domain contains the FAD moiety, a proximal C^{III}–C^{IV} disulfide, and a third CxxC of unknown function (C^V–C^{VI}). Plant and protist QSOXs lack the second Trx domain but otherwise appear to contain the same complement of redox centers. This work presents the first characterization of a single-Trx QSOX. *Trypanosoma brucei* QSOX was expressed in *Escherichia coli* using a synthetic gene and found to be a stable, monomeric, FAD-containing protein. Although evidently lacking an entire domain, *Tb*QSOX shows catalytic activity and substrate specificity similar to the vertebrate QSOXs examined previously. Unfolded reduced proteins are more than 200-fold more effective substrates on a per thiol basis than glutathione and some 10-fold better than the parasite bisglutathione analogue, trypanothione. These data are consistent with a role for the protist QSOX in oxidative protein folding. Site-directed mutagenesis of each of the six cysteine residues (to serines) shows that the CxxC motif in the single-Trx domain is crucial for efficient catalysis of the oxidation of both reduced RNase and the model substrate dithiothreitol. As expected, the proximal disulfide C^{III}–C^{IV}, which interacts with the flavin, is catalytically crucial. However, as observed with human QSOX1, the third CxxC motif shows no obvious catalytic role during the *in vitro* oxidation of reduced RNase or dithiothreitol. Pre-steady-state kinetics demonstrates that turnover in *Tb*QSOX is limited by an internal redox step leading to 2-electron reduction of the FAD cofactor. In sum, the single-Trx domain QSOX studied here shows a striking similarity in enzymatic behavior to its double-Trx metazoan counterparts.

The quiescin sulphydryl oxidase (QSOX)¹ family of enzymes catalyze the net generation of disulfide bonds with reduction of oxygen to hydrogen peroxide (1–4):



Reduced unfolded proteins are excellent substrates of these multi-domain flavin-dependent oxidases, suggesting that they contribute to the net generation of disulfide bonds during oxidative protein folding. QSOXs are found in the genomes of almost all eukaryotes, with the notable exception of the fungi (2–4). However, all of our current insights into the substrate specificity and the mechanism of these oxidases have come from study of metazoan QSOXs: first from murine seminal vesicles (5, 6), then from egg white (7, 8) and

bovine milk (9), and, most recently, from a human recombinant enzyme expressed in *Escherichia coli* (10).

The domain organization of a metazoan QSOX is shown in Figure 1A. These enzymes have two N-terminal thioredoxin domains (Trx1 and Trx2) joined to a C-terminal Erv/ALR domain by a helix-rich region (HRR) (1, 3, 4, 10, 11). A transmembrane helix is found at the C-terminus of most QSOX sequences, although alternately spliced transcripts lacking this feature have been identified (1, 4). Collectively, the Trx1 and Trx2 domains are similar to *a* and *b* domains of the protein disulfide isomerase (PDI) (12–15), with only the first containing a redox-active CxxC motif. Following the second thioredoxin domain in metazoan QSOXs is the HRR domain: an apparently unique feature of these oxidases. The Erv/ALR domain was first described as a yeast growth factor, essential for respiration and viability (Erv1) (16), and later found to be a small stand-alone flavin-linked sulphydryl oxidase (17). Crystal structures of yeast Erv2 (18), rat augmenter of liver regeneration (ALR) (19), and other Erv/ALR family members (18, 20–22) show that the flavin ring is bound at the mouth of a compact four-helix bundle. Additionally, the Erv/ALR domain houses a second redox center: a proximal CxxC disulfide that relays reducing equivalents from dithiol substrates, or from distal dithiol motifs, to the isoalloxazine ring. While the roles of thioredoxin and proximal disulfides in QSOX catalysis are established (10), there is a third CxxC motif of unknown function that is also conserved

[†]This work was supported in part by National Institutes of Health Grant GM26643. The content of this work is solely the responsibility of the authors and does not necessarily reflect the official views of the National Institute of General Medical Sciences or the National Institutes of Health.

*To whom correspondence should be addressed. Phone: (302) 831-2689. Fax: (302) 831-6335. E-mail: cthorpe@udel.edu.

¹Abbreviations: ALR, augmenter of liver regeneration; DTNB, 5,5'-dithiobis(2-nitrobenzoate); DTT, dithiothreitol; ERV, a protein essential for respiration and viability in yeast; GSH, reduced glutathione; GSSG, oxidized glutathione; GdnHCl, guanidine hydrochloride; HRR, helix-rich region; IPTG, isopropyl β-D-thiogalactopyranoside; QSOX, quiescin sulphydryl oxidase; RfBP, riboflavin binding protein; RNase, ribonuclease A; SDS, sodium dodecyl sulfate; TCEP, tris(carboxyethyl)phosphine; Trx, thioredoxin domain; VSG, variable surface glycoproteins; WT, wild type.




FIGURE 1: Domain organization of human and trypanosomal QSOX proteins and amino acid sequence of *TbQSOX*. Panels A and B show the domain organization of human and trypanosomal QSOX proteins, respectively. CxxC motifs are shown as red bars, labeled as previously (10) and using the modified nomenclature implemented in this work. The flavin ring system of FAD is shown above the Erv/ALR label. Panel C represents the amino acid sequence (NCBI Accession Number XP_845306) and predicted secondary structure of *TbQSOX*. The predicted signal sequence is underlined. Trx, HRR, and Erv/ALR domains are highlighted in blue, orange, and yellow, respectively. A predicted transmembrane region is highlighted in pink. CxxC motifs are highlighted in red and underlined in boldface. The remaining cysteine residues are shown against a gray background.

in all QSOX sequences available to date (1, 10). It lies at the C-terminus of the Erv/ALR domain and is not catalytically essential for the *in vitro* oxidation of DTT or reduced RNase (rRNase) by the recombinant human QSOX1 (10). It might play a role in the regulation of catalytic activity (10).

Plant and protist QSOXs lack all, or parts, of the second thioredoxin domain found in their metazoan counterparts (Figure 1B) (1, 3, 4). Since all of our current insights into the QSOX family have come from oxidases with two thioredoxin domains, we wanted to investigate a representative example of a structurally simpler QSOX. We chose the enzyme from *Trypanosoma brucei*, a protozoan pathogen (family *Trypanosomatidae*, order *Kinetoplastida*) that causes African sleeping sickness. In its predominant blood-stream phase, the parasite counters the host immune response with the help of a monolayer of variable surface glycoproteins (VSG) anchored to the outer leaflet of the plasma membrane by a glycosylphosphatidylinositol moiety (23). Although this coating is highly immunogenic, at each division cycle a proportion of the daughter cells change their VSG coat and thus evade detection by the human immune system. These highly variable VSGs do however contain multiple conserved disulfide bonds (24–28), and hence the ability of the parasite to

generate these structural disulfides would be expected to be important for pathogenicity.

Trypanosoma are known to have an unusual redox biology, in which a bisglutathione derivative, trypanothione, plays a major role (29, 30). However, we know very little about the ways these parasites catalyze the net generation of disulfide bridges during oxidative protein folding. Three flavin-linked sulfhydryl oxidases can be identified from the *T. brucei* genome. One is homologous to yeast Ero1p and its metazoan counterparts; these flavoproteins are resident in the endoplasmic reticulum and are thought to play significant roles in disulfide bond generation (31–35). A second, *TbALR*, is the counterpart of yeast Erv1p and human ALR and hence would be expected to support oxidative protein folding in the mitochondrial intermembrane space (36–39). The third is a QSOX sequence of the type discussed above with three, not four, recognizable domains.

As part of our effort to understand catalytic strategies in the QSOX family of proteins, we report the first enzymatic characterization of a protist QSOX. The protein was expressed using a synthetic gene, and its behavior was analyzed by a range of static methods, together with steady-state and rapid-reaction studies. These approaches reveal striking commonalities of behavior between these evolutionarily distant and structurally diverse representatives of the QSOX family.

EXPERIMENTAL PROCEDURES

Materials. Chemical reagents were obtained as described previously (40). Primers for subcloning of all constructs and for mutagenesis were from Integrated DNA Technologies. Protein and DNA molecular mass standards and all of the enzymes used for subcloning were from Promega. All bacterial culture media components were from Fisher Scientific.

***TbQSOX* Constructs.** A synthetic gene encoding *TbQSOX* protein (NCBI Accession Number XP_845306) codon-optimized for expression in *E. coli* was designed using Gene Designer software (DNA 2.0, www.dna20.com) and was obtained from DNA 2.0 in a pJ201 vector (see Supporting Information, Figure S1). All protein expression constructs were then obtained by subcloning the desired region into a pET-28a(+) vector (Novagen) using appropriate PCR primers (Supporting Information, Table S1). DNA sequences of all of the inserts were confirmed by sequencing to ensure in-frame ligation and the absence of unwanted mutations. Specific cysteine mutations were introduced by using the QuikChange site-directed mutagenesis kit (Stratagene) with the primers shown in Table S1. All mutant constructs were also sequenced to verify mutations.

Expression of *TbQSOX* Constructs and Mutants. Full-length *TbQSOX* constructs started at either amino acid residues S20 or G32 and ended at G485. Initial attempts at expression and purification from *E. coli* BL21 Star (DE3) (Invitrogen) cells induced at either 15 or 37 °C were unsuccessful. However, a full-length construct starting at residue S20 was expressed in a soluble form in *E. coli* Origami2 (DE3) (Novagen) cells. Thereafter, all *TbQSOX* constructs were expressed in this strain. Starter cultures were grown overnight at 37 °C in LB broth containing 15 µg/mL kanamycin and 12.5 µg/mL tetracycline. A 5 mL volume of starter culture was used to inoculate 500 mL of Terrific broth containing the same levels of antibiotics, and the media were incubated at 37 °C until an absorbance of 0.8–1.0 at 600 nm was reached. Protein expression was induced with 0.5 mM IPTG at 15 °C for 24 h. Cells were then harvested by centrifugation at 5000g at 4 °C and stored at –20 °C.

Purification of *TbQSOX* Constructs and Mutants. Cell pellets from 2 L of culture were resuspended in 20 mL of 50 mM potassium phosphate buffer, pH 7.5, containing 300 mM NaCl, 100 μ M FAD, and two tablets of protease inhibitor cocktail for His-tagged proteins (Complete, Mini, EDTA free; Roche). Cells were lysed by two passages through a French press (at 10000 psi) followed by brief sonication to shear DNA. The resulting lysate was centrifuged at 17000g for 30 min at 4 °C. The supernatant was carefully removed and added to Pro Bond Ni-NTA resin (Invitrogen) previously equilibrated in 50 mM phosphate buffer, pH 7.5, containing 300 mM NaCl. The suspension was rocked at 4 °C for 3 h and poured into an empty column. Flow-through was collected for further analysis, and the column was washed with four column volumes of 50 mM phosphate buffer, pH 7.5, containing 300 mM NaCl followed by four column volumes of 50 mM phosphate buffer, pH 6.0. Protein bound to the resin was eluted using a gradient of imidazole in 50 mM phosphate buffer, pH 6.0, and collected in 1.5 mL aliquots. Fractions containing *TbQSOX* enzyme were pooled and further purified using hydrophobic interaction chromatography. Here, ammonium sulfate was added to 40% saturation at 4 °C and the solution clarified by centrifugation at 17000g for 30 min. The supernatant was then loaded onto a butyl-Sepharose column (6 \times 2 cm) previously equilibrated with 50 mM phosphate buffer, pH 7.5, containing 40% saturated ammonium sulfate and 1 mM EDTA. The column was washed with three column volumes of 50 mM phosphate buffer, pH 7.5, containing 40% saturated ammonium sulfate and 1 mM EDTA. *TbQSOX* was eluted with a decreasing gradient of ammonium sulfate. Fractions (5 mL) were pooled and concentrated using Amicon Ultra centrifugal filter devices (Millipore).

UV/Vis Spectroscopy, Determination of Extinction Coefficients, and Thiol Titer. All absorption spectra were collected on an Agilent 8453 UV/vis spectrophotometer, and the data were analyzed using the ChemStation software (Agilent Technologies). Extinction coefficients for the bound flavin of wild-type *TbQSOX* and the shorter HRR-Erv construct (12250 and 12300 M⁻¹ cm⁻¹, respectively) were determined as described earlier (10). Thiol titer of full-length *TbQSOX* was determined by following the reaction of the native and SDS-denatured enzyme with 0.5 mM DTNB (10).

rRNase Assays. RNase was reduced for 2 h at 37 °C in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA, a 10-fold excess of DTT over total thiols, and 6 M GdnHCl. The mixture (0.6 mL) was applied to a PD10 column (Amersham Biosciences) equilibrated with 10 mM sodium acetate, pH 4.0, giving a baseline separation between rRNase and excess DTT. Thiol titer confirmed complete reduction, and the resulting protein was stored under anaerobic conditions at 4 °C (10). rRNase assays were performed as described earlier (10). Briefly, 50 nM *TbQSOX* was added to a 300 μ L reaction containing varying amounts of rRNase in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA at 25 °C. Aliquots (20 μ L) of the reaction mixture were transferred into 0.5 mM DTNB in phosphate buffer, pH 7.5, and thiol titer was monitored at 412 nm. Control reactions were run under identical conditions in the absence of *TbQSOX*.

Reduced Riboflavin Binding Protein Assays. Chicken riboflavin binding protein (a gift from Dr. Harold B. White III, University of Delaware) was reduced for 3 h at 37 °C in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA, a 10-fold excess of DTT over total thiols, and 6 M GdnHCl. Reduced protein was purified using a PD10 column equilibrated at 4 °C with 10 mM sodium acetate buffer at pH 5.0 containing 1 mM EDTA and 1 M GdnHCl. The thiol content of reduced RfBP

was measured using DTNB to confirm complete reduction, and rRfBP was stored under anaerobic conditions at 4 °C. rRfBP assays were performed essentially as above by adding 125 nM *TbQSOX* to a 300 μ L reaction containing varying amounts of reduced protein in phosphate buffer containing 1 mM EDTA at pH 7.5 at 25 °C. Thiol titers were measured using DTNB to allow calculation of rate of thiol oxidation and of enzymatic turnover numbers.

Trypanothione Assays. A solution of 14 mg/mL oxidized trypanothione (Bachem) in 50 mM phosphate buffer at pH 7.5 containing 1 mM EDTA was incubated with an equal amount of immobilized TCEP disulfide reducing gel (Pierce) for 90 min. Reduced trypanothione was separated from the matrix by centrifugation, and its thiol titer was measured by treatment with DTNB. Trypanothione assays were performed at 25 °C in a 150 μ L reaction volume containing 2.5, 5, or 10 mM reduced trypanothione in phosphate buffer, pH 7.5, containing 1 mM EDTA. Reactions were started by the addition of 1 μ M *TbQSOX* and followed by thiol titers as above. Control incubations lacked enzyme.

Oxygen Electrode Assays. Catalytic activities of all *TbQSOX* constructs were determined using DTT and GSH by using a Clark-type oxygen electrode as described previously (7). DTT and GSH were prepared in 50 mM phosphate buffer, containing 1 mM EDTA. Solutions were readjusted to pH 7.5 as necessary and standardized using DTNB before use.

pH Dependence of Turnover with DTT and rRNase. The following 50 mM buffers containing 1 mM EDTA were used: sodium acetate, potassium phosphate, Tris-HCl, and CAPS, with either 5 mM DTT or 30 μ M rRNase. Assays were initiated by the addition of 50 nM *TbQSOX*, using the oxygen electrode for DTT and discontinuous thiol titer measurements for rRNase.

Dithionite Titrations. Dithionite titrations were conducted under anaerobic conditions as before (7).

Stopped-Flow Spectrophotometry. Data were collected using an SF-61 DX2 double mixing stopped-flow system (Hi-Tech Instruments) at 25 °C in either diode-array or monochromator modes and analyzed by their KinetAsyst 3 software. For experiments under aerobic conditions, *TbQSOX* and DTT were prepared in air-saturated 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA at 25 °C. rRNase was prepared as described earlier. For experiments under anaerobic conditions, wild-type or mutant *TbQSOX* proteins were added to tonometers containing phosphate buffer, pH 7.5, with 1 mM EDTA, 5 mM glucose, 5 nM glucose oxidase, and 1 nM catalase. Separate tonometers contained DTT in the same buffer. The tonometers were made thoroughly anaerobic by rocking for 20 min using alternate cycles of vacuum and flushing with nitrogen. For experiments to determine reactivity toward oxygen, the C¹S *TbQSOX* mutant dissolved in phosphate buffer was made anaerobic as before, except that 1 nM glucose oxidase was used. The mutant enzyme was then reduced by the addition of 2-electron equivalents from a standardized solution of dithionite.

Sequence Analyses. The signalP server was used to predict signal sequence in *TbQSOX* (41). The transmembrane helix and the topology of the protein were predicted by the TMHMM server (41). Secondary structure predictions for all proteins were obtained by submitting sequences to PSIPRED (42). Protein sequence alignments were obtained using ClustalW (43).

RESULTS AND DISCUSSION

***TbQSOX* Gene and Protein.** The single *T. brucei* QSOX gene is located on chromosome 6 between a cluster of snoRNA

Table 1: The Cysteine Residues in *Tb*QSOX

cysteine residue	motif	corresponding cysteines in related proteins
Trx Domain		
58	n/a	not conserved
69, 72	C ^I -C ^{II}	C ⁷⁰ -C ⁷³ (CxxC _{trx}) of human QSOX1b; C ⁶¹ -C ⁶⁴ of yeast Pdi1p
105, 112	C ₆ C	C ¹⁰¹ -C ¹¹⁰ (C ₈ C) of human QSOX1b; C ⁹⁰ -C ⁹⁷ (C ₆ C) of yeast Pdi1p
132, 138	C ₈ C	not conserved
HRR Domain		
180	n/a	not conserved
Erv/ALR Domain		
305, 317	C ₁₁ C	unique to QSOX family; C ³⁹³ -C ⁴⁰⁵ (C ₁₁ C) of human QSOX1b
350, 353	C ^{III} -C ^{IV}	C ⁴⁴⁹ -C ⁴⁵² (CxxC _{prox}) of human QSOX1b; C ¹²¹ -C ¹²⁴ of yeast Erv2p
405, 408	C ^V -C ^{VI}	C ⁵⁰⁹ -C ⁵¹² (CxxC _{tm}) of human QSOX1b; C ¹⁷⁶ -C ¹⁷⁸ of yeast Erv2p

genes and a diacylglycerol kinase gene. *Tb*QSOX is 1581 nucleotides long and encodes a 526 amino acid protein starting with an N-terminal signal sequence with a predicted cleavage site after residue 22 (Figure 1C). The full-length sequence corresponds to a type I membrane protein with a single transmembrane helix (residues 491–513) and a short cytoplasmic C-terminal tail. In higher organisms, alternate splicing generates soluble QSOX forms lacking the membrane span (2, 4, 6). Whether the parasite generates correspondingly truncated QSOX forms is not yet known. As mentioned earlier, the domain structure of *Tb*QSOX is distinctly different from its metazoan orthologues. Analysis of predicted secondary structure (Figure 1C) shows the presence of an alternating arrangement of strands and helices typical of a thioredoxin fold. This feature is supplemented by an N-terminal additional $\beta\alpha$ motif similar to that of an α domain of PDI. Following the HRR domain, encompassing residues 168–304 and dominated by α -helices, is the Erv/ALR domain, another helix-rich domain which serves to bind the FAD moiety and which houses the proximal disulfide that is critical for communication with the Trx1 domain in metazoan QSOXs.

The Cysteine Residues of *Tb*QSOX. Table 1 lists the cysteine residues, and their domain context, in this single-Trx QSOX. Three CxxC motifs are conserved in all of the available QSOX sequences, from the smallest free-living eukaryote (*Ostreococcus tauri*) to humans (1). For simplicity, we refer to these cysteine pairs as C^I–C^{II}, C^{III}–C^{IV}, and C^V–C^{VI}, together with their numbering in the context of the full-length protein (Table 1). C^I–C^{II} cysteine residues comprise the CxxC motif located at the N-terminus of the α_1 helix of the thioredoxin domain. This motif is highly conserved among proteins with thioredoxin domains; it has been shown to be critical for the catalytic activity of, among other proteins, human QSOX1 (10) and PDI (12, 14, 44). This sequence in QSOXs from mammals and birds is typically CGHC and is immediately preceded by a tryptophan residue (1). However, the corresponding sequence in *Tb*QSOX is CGAC (Figure 1C), preceded by a glycine rather than an aromatic residue. In metazoan QSOXs, this disulfide

receives reducing equivalents from dithiol substrates and then transmits them to C^{III}–C^{IV} of the Erv/ALR flavin-binding domain (10, 11). Site-directed mutagenesis with the human QSOX1 confirms that C^{III}–C^{IV} is catalytically essential because it needs to communicate first with C^I–C^{II} in the Trx domain and then with the adjacent flavin moiety (10). The third CxxC motif, C^V–C^{VI}, is also invariant; however, mutation of either, or both, of the C^V–C^{VI} cysteine residues did not result in a major change in activity toward small molecule or protein substrates in QSOX1 (10). The role of this conserved disulfide is currently cryptic.

In addition to the three CxxC disulfides mentioned above, a further eight cysteines, likely in disulfide linkage (see later), are present in full-length *Tb*QSOX. A C₆C motif (C105–C112) is placed comparably to the C₈C motif present in human QSOX1 protein (10), to the C₆C motif present in yeast Pdi1p (13), and to the C_{6–8}C motifs from several other members of the PDI family. In yeast Pdi1p, these residues form a disulfide (13) and may be involved in regulation of catalytic function (45). The C₁₁C motif (C305–C317) lies at the beginning of the Erv/ALR domains. *Tb*QSOX contains an additional four nonconserved cysteine residues (58, 132, 138, and 180) of uncertain role. This report will address mutations of only the three CxxC motifs that have been previously suggested to play catalytic roles in QSOX family members.

Subcloning, Expression, and Purification of *Tb*QSOX. The *Tb*QSOX cDNA sequence (NCBI Accession Number XM_840213) was edited to reflect codon preferences in *E. coli* and synthesized as described in Experimental Procedures. We initially attempted to express *Tb*QSOX with an N-terminal His tag by starting the QSOX sequence at residue G32. This residue lies in the middle of a region with no predicted secondary structure, between the end of the signal sequence and the beginning of a short β -strand that starts the PDI-like Trx domain (Figure 1C). This construct failed to express (see Experimental Procedures). However, the inclusion of an additional 12 residues at the N-terminus of the oxidase sequence yielded a stable,




FIGURE 2: UV/vis spectrum of *TbQSOX*. The spectrum was recorded in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA. The inset shows SDS–PAGE of *TbQSOX* under reducing (Red) and nonreducing (NR) conditions together with protein molecular mass standards in kDa.

soluble protein in ample amounts for the work described in this study (see below). The codon-optimized gene sequence and the amino acid sequence of the wild-type protein, used throughout this work, are included in Supporting Information (Figures S1 and S2).

The number of anticipated catalytic and structural disulfide bonds in *TbQSOX* suggested that expression in *E. coli* BL21 Star (DE3) cells might be problematic because this strain does not favor the cytosolic expression of disulfide-containing proteins. Indeed, expression trials using this strain led to the accumulation of QSOX in inclusion bodies (not shown). However, we could obtain soluble and active protein using a *gor/trx* mutant strain (*E. coli* Origami2 (DE3)) grown until mid log phase at 37 °C and then induced with 0.5 mM IPTG at 15 °C for 24 h. Induction at higher temperatures led to progressively more QSOX appearing in the insoluble fraction (not shown). *TbQSOX* was then purified utilizing the N-terminal His-tag followed by an additional purification step using a butyl-Sepharose column to remove a contaminating flavoprotein (identified as alkyl hydroperoxide reductase, AhpF; data not shown). Typically, we obtained ~7 mg of pure *TbQSOX* per liter of culture for wild type and cysteine mutants. The resulting protein was stable for at least 6 months at 4 °C in 50 mM potassium phosphate buffer containing 1 mM EDTA at pH 7.5. *TbQSOX* could also be stored for at least 1 year at –20 °C without significant loss of enzyme activity. The purified protein showed a typical unresolved flavin absorbance in the visible region with a maximum at 456 nm and an experimentally determined molar extinction coefficient of 12,250 M^{–1} cm^{–1} (Figure 2; see Experimental Procedures). No free thiols were detected when protein denatured with SDS was incubated with DTNB (see Experimental Procedures), suggesting that all 14 cysteine residues in *TbQSOX* comprise 7 disulfide bonds. SDS–PAGE, performed under reducing and nonreducing conditions, showed a single band at the expected molecular mass of 54 kDa (Figure 2, inset). Analytical ultracentrifugation experiments gave a solution molecular mass of 54.3 kDa (Supporting Information, Figure S3).

Catalytic Activity of *TbQSOX*. The divergent redox biology and the evolutionary distance between *T. brucei* and metazoan QSOXs justify an examination of thiol substrate specificity for the parasite oxidase. As for all the other QSOXs investigated (5, 7, 9, 10), DTT proves a useful model substrate of *TbQSOX*. Here, *TbQSOX* shows a pH dependence with a




FIGURE 3: pH dependence for the oxidation of DTT by *TbQSOX*. The ascending and descending limbs for the oxidation of 5 mM DTT were fit to pK values of 5.7 and 8.9, respectively, setting upper and lower limits of 2608 s^{–1} and 0 s^{–1}.

maximal activity at neutral pH values (with ascending and descending limbs fit to pK values of 5.7 and 8.9, respectively; Figure 3). This pH dependence, and the optimal pH value of about 7, is similar to those of the egg white oxidase (7). We utilized 50 mM potassium phosphate buffer, pH 7.5, containing 1 mM EDTA for most of the experiments described herein. Compared to DTT, glutathione is a very poor substrate of *TbQSOX* with an approximately 2000-fold lower k_{cat}/K_m value (reflecting a much larger K_m term for the monothiol; Table 2). While glutathione is present in trypanosomes, the dithiol analogue trypanothione [*N*¹,*N*⁸-bis(glutathionyl)spermidine] substitutes for many of its intracellular functions (29, 46). While trypanothione shows a roughly 20-fold increase in catalytic efficiency over glutathione, this dithiol is approximately 10-fold poorer as a substrate than two reduced proteins selected for study (Table 2).

Both reduced proteins, RNase and RfBP, are substantially unfolded in the absence of denaturant. Reduction of the disulfide bonds in RNase generates a largely unstructured protein (47) in which reaction of all 8 thiols with 2 mM polar reagent DTNB is complete in less than 200 ms (40). Similarly, about 13 of the 18 thiol groups of rRfBP react in 0.5 s under the same conditions (40). The catalytic efficiencies of *TbQSOX* in oxidizing rRNase are some 3–6-fold lower than for the avian, milk, and human enzymes (9, 10, 48). An estimate of the pH dependence for the oxidation of rRNase shows ascending and descending limbs with pK values of 6.8 and 9.8, respectively (Supporting Information, Figure S4). For rRfBP, the trypanosomal enzyme shows a catalytic efficiency that is 2-fold lower than that of the avian enzyme (48). These differences between protist and metazoan enzymes are largely a reflection of the K_m term. In spite of these rather minor decreases in catalytic efficiency between parasite and metazoan QSOXs, the trypanosomal enzyme remains a facile oxidant of reduced proteins. Hence, *TbQSOX* may generate some fraction of secreted protein disulfides in trypanosomes. Finally, *TbQSOX* can cooperate with reduced PDI in the successful oxidative protein folding of RfBP ((49) data not shown).

Roles of CxxC Motifs in *TbQSOX* Catalysis. Recent work with the recombinant human QSOX1 showed that, while C^I–C^{II} and the C^{III}–C^{IV} disulfides were crucial for the efficient oxidation of both DTT and rRNase thiols, the conserved

Table 2: Comparison of Steady-State Parameters for Oxidation of Thiol Substrates between *Tb*QSOX and Three Metazoan QSOXs^a

	<i>Tb</i> QSOX			<i>Gg</i> QSOX1b ^b	<i>Br</i> QSOX1b ^c	<i>Hs</i> QSOX1b ^c
	k_{cat} (s ⁻¹)	K_m (mM)	k_{cat}/K_m (M ⁻¹ s ⁻¹)	k_{cat}/K_m (M ⁻¹ s ⁻¹)	k_{cat}/K_m (M ⁻¹ s ⁻¹)	k_{cat}/K_m (M ⁻¹ s ⁻¹)
DTT	45	0.086	5.2×10^5	2.3×10^5	3.7×10^5	2.0×10^5
GSH	— ^d	> 50	2.7×10^2 ^d	2.3×10^3	5.9×10^3	2.0×10^3
TSH	16	3.23	4.9×10^3	nd ^e	nd	nd
rRNase A	22	0.36	6.0×10^4	1.8×10^5	3.7×10^5	2.2×10^5
rRfBP	14	0.30	4.5×10^4	1.6×10^5	nd	nd

^aTurnover numbers are expressed in terms of thiols oxidized, not disulfides generated per second. For comparison between substrates with different numbers of thiol groups, K_m values are expressed on a per thiol basis. The oxidation of trypanothione, rRNase, and rRfBP was evaluated using discontinuous thiol titer assays with DTNB. Measurement of the turnover with DTT and GSH utilized the oxygen electrode (see Experimental Procedures). ^bRecalculated in terms of monothiols using data from avian QSOX enzymes (7). ^cData taken from the bovine milk (9) and the recombinant human QSOX1 (10). ^dRates of GSH oxidation were essentially linear up to 40 mM; the catalytic efficiency is expressed as the apparent second-order rate constant for 10 mM GSH. ^end: not done.




FIGURE 4: Comparison of the turnover numbers of wild-type *Tb*QSOX and its CxxC mutants toward DTT and rRNase. Turnover numbers were measured at 25 °C in phosphate buffer, pH 7.5, using 5 mM DTT and 200 μ M rRNase thiols (filled and open columns, respectively). Data were normalized with the wild-type protein set at 100%.

C^V–C^{VI} disulfide was nonessential *in vitro* (10). These studies, combined with earlier results from the avian QSOX (11, 50), led to a revised model for catalysis in which the thioredoxin C^I–C^{II} motif receives reducing equivalents from a protein substrate and then interacts with C^{III}–C^{IV} of the Erv/ALR domain within the same subunit of the oxidase (10). We were thus interested in comparing the behavior of the single-Trx QSOX with these prior results. In particular, we wished to examine whether the C^V–C^{VI} disulfide was catalytically important in this smaller parasite enzyme.

When C^I (C69) or C^{II} (C72) was mutated to serine, catalytic activity of *Tb*QSOX toward 5 mM DTT decreased to ~5% of wild-type protein (Figure 4). This decrease in catalytic activity was even more drastic when rRNase was used as a substrate; now only ~0.5% of wild-type activity was observed (Figure 4). These values likely represent the ability of DTT, in the absence of a functional thioredoxin domain, to directly interact with the Erv/ALR oxidase domain. Accordingly, we prepared a truncated protein (HRR-Erv) which lacked the entire Trx domain (the sequence of this construct is shown in Supporting Information, Figure S2). HRR-Erv binds FAD with essentially the same visible spectrum as that for full-length protein (Supporting Information, Figure S5). The turnover number of the HRR-Erv truncated protein using 5 mM DTT was comparable to the full-length protein containing either a C^IS or a C^{II}S mutation (150, 114, and 112 thiols oxidized per minute, respectively).

Replacement of either cysteine of the proximal disulfide (C^{III} or C^{IV}) with serine essentially abolished activity toward

both DTT and rRNase (Figure 4). The placement of this proximal disulfide within the Erv/ALR domain of *Tb*QSOX is likely to be very similar to that found in the other Erv/ALR family members (18, 19, 21). C^{III} is the interchange residue with a sulfur atom largely exposed to solvent and poised to accept an incoming mixed disulfide bond from the surface-accessible C^I cysteinyl thiolate of the *Tb*QSOX Trx domain. Following resolution of this mixed disulfide between Trx and Erv/ALR domains, the now-reduced C^{III}–C^{IV} motif can then transfer a pair of electrons to the flavin probably via the intermediacy of a C4a adduct (1, 3, 51, 52). Clearly, both C^{III} and C^{IV} residues would then be expected to be catalytically critical in *Tb*QSOX. The visible spectra of the two mutant proteins are noticeably different. The C^{III}S mutant leaves C^{IV} unpaired, and it forms a thiolate to oxidized flavin charge-transfer complex (showing a wedge-shaped long-wavelength band; Supporting Information, Figure S6). In contrast, the corresponding C^{IV}S mutant protein shows a typical oxidized flavin spectrum, but it is blue shifted by 6 nm compared to the wild-type protein (Supporting Information, Figure S6). This behavior likely reflects greater access of solvent to the flavin when the proximal disulfide is reduced and/or an increase in polarity when the hydroxyl group of serine replaces the S atom of C^{IV}.

Finally, mutations of the terminal CxxC disulfide were prepared. The visible spectrum of both C^VS and C^{VI}S mutants were comparable to that of the wild-type protein (Supporting Information, Figure S6). In terms of catalysis, neither mutant showed significant loss of activity toward DTT or rRNase (Figure 4). In aggregate, these studies show that the first two CxxC redox centers (C^I–C^{II} and C^{III}–C^{IV}) in *Tb*QSOX are crucial for the efficient oxidation of rRNase. The third conserved CxxC motif (C^V–C^{VI}) may play an important role *in vivo*, but it is not revealed in these *in vitro* experiments.

Dithionite Titrations of *Tb*QSOX. In preparation for the rapid reaction experiments to be presented later, we wished to assess how many redox centers could be revealed during dithionite titrations of the wild-type enzyme. Sodium dithionite solutions were standardized by anaerobic titrations against solutions of free FAD before being used as redox titrants of the oxidase (7, 53). The dithionite ion, S₂O₄²⁻, serves as an overall 2-electron reductant of the flavin, although reduction may proceed via the intermediacy of the SO₂⁻ radical anion (54). In addition, disulfides can be reduced indirectly by dithionite (requiring an additional 2-electron equivalent per disulfide) if these redox centers can communicate with the flavin ring. Figure 5 shows that complete reduction of the flavin in *Tb*QSOX occurs with




FIGURE 5: Dithionite titrations of wild-type, C¹S, and HRR-Erv forms of *TbQSOX*. WT *TbQSOX* in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA, was made anaerobic and titrated at 25 °C with a standardized solution of sodium dithionite yielding curves 1–7. The inset plots the reduction of the oxidase as a function of electron equivalents added per flavin (■) together with comparable titrations with C¹S (○) and HRR-Erv (□) proteins.

concomitant reduction of a redox-active disulfide. The apparent stoichiometry of 4.6 electron equivalents per QSOX monomer, slightly more than the expected 4.0 electron equivalents, might reflect kinetic sluggishness toward the end of the titration. The approximate linearity of the absorbance decrease at 456 nm with reductant added (inset Figure 5) shows that both the flavin and the redox-active disulfide are almost equipotential. However, when dithionite titrations are repeated with the C¹S mutant, removing any potential involvement of the Trx redox-active disulfide (Figure 1B), full reduction of the flavin now requires only 2-electron equivalents (open circles, inset Figure 5). These data suggest that the second redox center is C^I–C^{II} and that the proximal C^{III}–C^{IV} disulfide is considerably more reducing than either the flavin or the Trx disulfide. Removal of the entire Trx domain, as in the HRR-Erv construct, again results in a stoichiometry of 2-electron equivalents for complete reduction of the flavin prosthetic group (open squares, inset Figure 5).

Consistent with these results using *TbQSOX*, the proximal disulfide in a number of other Erv/ALR/QSOX family members is found to be significantly more reducing than the flavin prosthetic group. In Erv2p, reducing equivalents from PDI enter via a CxC shuttle (distal) disulfide and are then passed to the proximal disulfide, to the adjacent flavin prosthetic group, and finally to molecular oxygen (18, 55). The 4-electron equivalents needed to completely bleach the flavin in Erv2p are distributed, more or less evenly, between a distal disulfide and the isoalloxazine, thereby leaving the proximal disulfide largely oxidized (55). Second, the cytokine-like form of augments of liver regeneration, ALR, has both flavin and proximal disulfide redox centers. Here, 2-electron reduction of the flavin occurs before significant reduction of the proximal disulfide, again reflecting their relative redox potentials (56). Finally, partial proteolysis of avian egg white QSOX generates a fragment analogous to the HRR-Erv fragment of *TbQSOX* mentioned above (11). The subsequent removal of the redox-active thioredoxin domain in the avian QSOX causes the stoichiometry required for flavin reduction to revert from 4- to 2-electron equivalents (11). In all of these examples, the proximal (C^{III}–C^{IV}) disulfide is more reducing than either the isoalloxazine ring or any additional shuttle




FIGURE 6: Enzyme monitored turnover of DTT by *TbQSOX*. The oxidase was mixed with DTT to give final concentrations of 18.4 μ M *TbQSOX* and 2.75 mM DTT in 50 mM phosphate buffer, pH 7.5, 25 °C, containing 240 μ M dissolved oxygen. Only selected spectra between 5 ms to 3 s are shown for clarity. The inset shows absorbance at 456 and 560 nm plotted as a function of time.

disulfides that communicate reducing equivalents from distal loci to the flavin binding domain.

Monitoring Turnover with DTT and rRNase in the Stopped-Flow Spectrophotometer. Aerobic solutions of *TbQSOX* and saturating concentrations of DTT were mixed in the stopped-flow instrument, and the evolution of spectral changes was followed using a diode-array spectrophotometer (Figure 6). The inset plots the corresponding changes at 456 and 560 nm. Here, the flavin is almost completely oxidized in the steady state, showing that the reoxidation of reduced flavin species by molecular oxygen is considerably faster than the rate-determining steps in the reductive half-reaction (see later). The small (3%) decline in 456 nm absorbance in the steady state is accompanied by a correspondingly rapid formation of a weak long-wavelength band consistent with a thiolate to oxidized flavin charge-transfer complex (monitored here at 560 nm and observed, under static conditions, in the spectrum of the C^{III}S mutant; Supporting Information, Figure S6). The steady state persists for about 0.6 s before reduced flavin begins to accumulate as the dissolved oxygen is depleted. While reduction of the flavin is largely complete by 0.9 s (inset Figure 6), further smaller decreases in absorbance continue over several seconds. These minor secondary changes are too slow to be catalytically significant; they are also encountered during the reduction of *TbQSOX* by DTT under anaerobic conditions (see later).

The area under the curve for the absorbance changes at 560 nm (inset Figure 6) allows calculation of the turnover values as a function of remaining oxygen concentration (Supporting Information, Figure S7) and shows a K_m for oxygen of 10.5 μ M, with a limiting turnover of 43 s⁻¹, in good agreement with the data obtained from the oxygen electrode (Table 2). A comparable progression of changes was seen during the aerobic turnover of rRNase, with the rapid generation of the charge-transfer band and the eventual reduction of the flavin (Supporting Information, Figure S8).

Reductive Half-Reaction Using DTT: Kinetic Evidence for the Importance of a Functional Thioredoxin Domain. Figure 7 shows that anaerobic reduction of *TbQSOX* by 0.5 mM DTT rapidly generates small levels of a charge-transfer species which decays more slowly to yield reduced flavin (inset). The rate constant for the appearance of the long wavelength intermediate is shown in Figure 7B (solid squares; with a limiting rate constant of 280 s⁻¹ and an apparent K_d for DTT of 0.66 mM). This phase




FIGURE 7: Anaerobic reduction of *TbQSOX* with DTT. Anaerobic solutions of *TbQSOX* and DTT (in 50 mM phosphate buffer, pH 7.5, containing 1 mM EDTA, 5 mM glucose, 5 nM glucose oxidase, and 1 nM catalase) were mixed in a stopped-flow spectrophotometer at 25 °C to give final concentrations of 8 μ M and 0.5 mM, respectively. Panel A shows selected spectra from the diode-array detector, and the inset follows the absorbance at 560 nm using a monochromator. Panel B plots observed rate constants as a function of the concentration of DTT. Appearance of the long wavelength species at 560 nm shows a limiting rate constant of 280 s⁻¹ with an apparent K_d of 0.66 mM (♦). Disappearance of the long wavelength species (◆) and the reduction of flavin (○) show limiting rate constants of 18.8 and 17.9 s⁻¹, respectively, both with an apparent K_d of 0.31 mM.

is more than 10-fold faster than overall turnover (23 disulfides/s, 45 thiols/s; Table 2). In contrast, the concentration dependence and the limiting rate constant of the disappearance of 560 nm absorbance closely parallel the main reduction phase (Figure 7B, solid diamonds; 0.31 mM K_d and 18.8 s⁻¹). Comparable values are observed for the major phase of flavin reduction at 456 nm (accounting for ~85% of the total bleaching observed with a K_d of 0.28 mM DTT and a limiting rate of 17.9 s⁻¹; Figure 7B, open circles). The average between the two values for this slower phase (18.4 s⁻¹) is in good agreement with the 22.5 s⁻¹ expected for overall turnover (disulfides generated) under these conditions. About 15% of the total bleaching of the flavin observed at 456 nm in Figure 7A requires 50 s for completion and can be fit to two exponentials with rate constants of approximately 0.8 and 0.1 s⁻¹ (not shown). Both of these slower phases are irrelevant for aerobic turnover conditions; they likely entail eventual reduction of all three redox centers in *TbQSOX* (see later).

Anaerobic stopped-flow experiments were then repeated using the C^IS mutant of *TbQSOX* so that the CxxC motif of the Trx




FIGURE 8: Simplified model for turnover in *TbQSOX*. Reduction of the C^I-C^{II} disulfide by DTT (k_1) yields form B. A thiolate to oxidized flavin charge-transfer species (represented by the curved dotted line: involving either, or both, of forms C and D) then forms with a limiting rate of 280 s⁻¹ (k_2). The charge-transfer absorbance decays (k_3) with a limiting rate of 18 s⁻¹ yielding reduced flavin (form E). The extent to which the interdomain redox reaction between forms C and D contributes to this rate-limiting step is unknown. The anticipated C4a adduct in the reduction of the flavin is omitted from the scheme because it has not been observed experimentally. The second substrate oxygen participates in k_4 by converting form E to A. For simplicity, the scheme omits the possibility that DTT could reduce the C^I-C^{II} disulfide in forms D and E to generate 4-electron-reduced *TbQSOX* prior to the oxidative half-reaction (k_4). A contribution of 4-electron-reduced oxidase to overall turnover would be favored by high DTT concentrations and low oxygen tensions.

domain could no longer participate in reduction of the enzyme by DTT. The consequences of lack of a functional thioredoxin domain are profound. Full reduction of the flavin at 456 nm can be fit to a single exponential and occurs without the formation of a significant charge-transfer complex (Supporting Information, Figure S9). Unlike the wild-type protein, these rate constants show a linear dependence up to 20 mM DTT (Supporting Information, Figure S10). At 5 mM DTT, the resulting rate constant of 1.2 s⁻¹ is in good agreement with the overall turnover number determined in the oxygen electrode (1.25 s⁻¹ and Figure 4). Removing the Trx domain entirely (as in the HRR-Erv construct) yields a similar turnover number (0.93 s⁻¹).

Oxidative Half-Reaction. As mentioned previously, dithionite reduction of the wild-type *TbQSOX* leads to the reduction of both flavin and C^I-C^{II} redox centers (Figure 5). In contrast, the C^IS mutant can be reduced essentially exclusively at the flavin center (Figure 5), allowing its reaction with molecular oxygen to be followed without the associated spectral changes involved in

bringing the second pair of reducing equivalents from the Trx domain to the flavin. At a final concentration of 120 μM oxygen, reoxidation is very rapid and proceeds without detectable intermediates (Supporting Information, Figure S11). Assuming that the *TbQSOX* reaction with oxygen is bimolecular, as has been observed for avian QSOX (50) and for most flavin-dependent oxidases (57), the second-order rate constant would be approximately $2.5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$. For comparison, a value of $9 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ was reported for the oxygen reactivity of the reduced flavin in avian QSOX at 4 °C (50).

A Model for Turnover. Figure 8 presents a simplified depiction of catalysis with the model substrate DTT. Turnover is initiated by the reduction of $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ disulfide by DTT. Reducing equivalents from $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ are transferred to $\text{C}^{\text{III}}\text{--}\text{C}^{\text{IV}}$ via a mixed disulfide between C^{I} and C^{III} (form **C**), allowing C^{IV} to form a thiolate to oxidized flavin charge-transfer species (shown by a dotted line from C^{IV} to FAD, forms **C** and **D**). FAD is reduced by the transfer of electrons from $\text{C}^{\text{III}}\text{--}\text{C}^{\text{IV}}$ and then reoxidized by molecular oxygen to complete the catalytic cycle. In some cases, the molecular basis for the kinetic transients observed in this work is not certain. For example, while it is clear that the weak long wavelength band that appears rapidly in the steady state represents a charge-transfer complex between the thiolate form of C^{IV} and the flavin (see earlier, Figure 8), the relative contributions of forms **C** and **D** to this absorbance feature are unclear. The appearance of the **C/D** charge-transfer species occurs with a limiting rate of $\sim 280 \text{ s}^{-1}$ and is half-saturated at 0.66 mM DTT. What is the significance of this DTT concentration dependence? In an early kinetic study of avian QSOX we interpreted a comparable saturation of the apparent rate constant in terms of a binding site for DTT (50). This prior work was performed before the realization that these QSOX enzymes contain at least three redox centers and that input of reducing equivalents occurred primarily through the $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ disulfide of a thioredoxin domain. While a binding site for reducing substrates (e.g., DTT or reduced proteins) is certainly plausible, there is currently no structural or kinetic evidence for such a site in any member of the QSOX family. For simplicity, we have here assumed that the reduction of the enzyme by DTT occurs by a simple bimolecular encounter with the $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ disulfide of the Trx domain. Assuming that the conversion of **A** to **C/D** reflects two irreversible steps, k_1 and k_2 (Figure 8), and analyzing for the formation of **C/D** as in Strickland et al. (58), predicts a value for this bimolecular reduction step of $4.6 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$.

Form **D** is likely to react via a C4a thiol–flavin adduct (not shown (51, 59–61)) yielding form **E**. Currently, it is unknown whether the rate-limiting step of 18 s^{-1} reflects a slow interconversion of forms **C** and **D** (followed by a rapid reaction generating reduced flavin) or whether it actually represents the direct accumulation of form **E** from **D**. There is a further uncertainty in Figure 8: whether *TbQSOX* turnover is necessarily restricted to cycling between 0- and 2-electron reduced states. Indeed, this seems unlikely. Since DTT reacts rapidly with the $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ disulfide in the Trx domain of form **A**, it might rereact with the disulfide in form **E** (and/or form **D**) before the reoxidation event (k_4). For illustration, using 240 μM oxygen and 5 mM DTT, form **E** would be predicted to undergo a second reduction by DTT in preference to reoxidation by step k_4 (with pseudo-first-order rate constants of 1500 vs 600 s^{-1} , respectively). Partition in favor of 4-electron reduction of *TbQSOX* would be favored by low oxygen tensions. In contrast, low thiol substrate concentrations would favor the simple 0- to 2-electron

cycling scheme shown in Figure 8. Further work is aimed at a more detailed examination of the pre-steady-state behavior of *TbQSOX* and the possible roles of 4-electron-reduced forms of the oxidase in turnover.

CONCLUSIONS

This characterization of a single-Trx QSOX shows strong similarities with the double-Trx enzymes investigated to date. Lack of the second Trx domain evidently does not result in major changes in the catalytic specificity of the parasite QSOX. While *TbQSOX* is between 3- and 6-fold less efficient at the oxidation of the rRNase and rRfBP than the metazoan enzymes, reduced unfolded proteins are its best known substrates. The average $k_{\text{cat}}/K_{\text{m}}$ values for the metazoan enzymes for DTT are about one-half of those for *TbQSOX*. Both single- and double-Trx QSOXs oxidize the monothiol GSH very poorly. The dithiol trypanothione shows a $k_{\text{cat}}/K_{\text{m}}$ value per thiol that is at least 10-fold lower than that for rRNase and rRfBP.

The similarities between single- and double-Trx QSOXs extend to pre-steady-state kinetic behavior. Both enzymes rapidly form a 2-electron-reduced charge-transfer complex with dithiol substrates, and in both cases this species decays to yield reduced flavin in a process that strongly limits overall turnover. Reoxidation of both enzymes by molecular oxygen does not limit turnover, illustrated by the almost exclusively oxidized flavin component found in the steady state during enzyme-monitored aerobic turnover experiments (Figure 6 (50)). Finally, both enzymes show a low K_{m} for oxygen (10.5 and 4.6 μM for *T. brucei* and the avian QSOX (50), respectively, when the reductant is DTT).

The mutagenesis studies reported in this work complement, and extend, those performed recently with the human enzyme (10). Two particular aspects of the behavior of *TbQSOX* are of note. First, stopped-flow studies of the recombinant parasite enzyme reinforce how important the $\text{C}^{\text{I}}\text{--}\text{C}^{\text{II}}$ motif is for overall turnover. The observation that complete removal of the Trx domain by truncation has no more serious catalytic impact than a C^{I} to serine mutation in full-length *TbQSOX* suggests that the thioredoxin domain does not serve additional effector roles under the conditions of our experiments. A second issue is that one could envisage that the simpler domain structure of a protist QSOX would result in an overt catalytic role for the $\text{C}^{\text{V}}\text{--}\text{C}^{\text{VI}}$ disulfide: a motif conserved in all QSOX family members. However, we see no significant impact in *TbQSOX*: mutations at this locus behave like the $\text{C}^{\text{V}}\text{--}\text{C}^{\text{VI}}$ mutations of the human recombinant QSOX enzyme.

The pathways of oxidative protein folding in *Trypanosoma* remain to be uncovered. In addition to QSOX, the parasite has a typical Ero1 protein and several PDI-like proteins. Neither parasite flavin-dependent sulphydryl oxidase has an obvious ER retention sequence, but it is possible that one, or both, could be retained in the ER lumen by binding to a PDI homologue (as has been suggested for human Ero1- $\text{L}\alpha$ and - $\text{L}\beta$ (62)). In terms of *TbQSOX*, we show that GSH and its dithiol analogue, trypanothione, are relatively poor substrates. Thus *TbQSOX* might introduce disulfide bonds into client proteins in the secretory system of *Trypanosoma* while minimizing collateral perturbation of the small molecule thiol redox pool and the consequent added oxidative stress that this may engender (3, 49). Alternatively, *TbQSOX* may function post-ER (1, 4): in the Golgi apparatus (as has been suggested for the long form of human QSOX1 (63)), on

the plasma membrane, or following secretion from the cell. Since trypanosomes seem to contain single paralogues of QSOX and Ero1, it will be interesting to examine their relative roles in oxidative protein folding. More generally, oxidative protein folding pathways in protists might represent an additional target for antimicrobial therapies including the arsenicals (40) that have been used for over a century to combat trypanosomiasis.

ACKNOWLEDGMENT

We thank Drs. Deborah Fass and Luise Krauth-Siegel for helpful discussions and anonymous reviewers for their insight.

SUPPORTING INFORMATION AVAILABLE

Table S1 listing the sequences of primers used in this study and Figures S1–S11 showing the DNA and protein sequences of the constructs used, analysis of analytical ultracentrifugation data, UV/vis spectra of *Tb*QSOX constructs and mutants, and data from steady-state and pre-steady-state kinetic measurements. This material is available free of charge via the Internet at <http://pubs.acs.org>.

REFERENCES

- Heckler, E. J., Rancy, P. C., Kodali, V. K., and Thorpe, C. (2008) Generating disulfides with the quiescin-sulfhydryl oxidases. *Biochim. Biophys. Acta* 1783, 567–577.
- Thorpe, C., Hooper, K., Raje, S., Glynn, N., Burnside, J., Turi, G., and Coppock, D. (2002) Sulfhydryl oxidases: emerging catalysts of protein disulfide bond formation in eukaryotes. *Arch. Biochem. Biophys.* 405, 1–12.
- Thorpe, C., and Coppock, D. L. (2007) Generating disulfides in multicellular organisms: emerging roles for a new flavoprotein family. *J. Biol. Chem.* 282, 13929–13933.
- Coppock, D. L., and Thorpe, C. (2006) Multidomain flavin-dependent sulfhydryl oxidases. *Antioxid. Redox Signaling* 8, 300–311.
- Ostrowski, M. C., and Kistler, W. S. (1980) Properties of a flavo-protein sulfhydryl oxidase from rat seminal vesicle secretion. *Biochemistry* 19, 2639–2645.
- Benayoun, B., Esnard-Fève, A., Castella, S., Courty, Y., and Esnard, F. (2001) Rat seminal vesicle FAD-dependent sulfhydryl oxidase. Biochemical characterization and molecular cloning of a member of the new sulfhydryl oxidase/quiescin Q6 gene family. *J. Biol. Chem.* 276, 13830–13837.
- Hooper, K. L., Joneja, B., White, H. B., III, and Thorpe, C. (1996) A sulfhydryl oxidase from chicken egg white. *J. Biol. Chem.* 271, 30510–30516.
- Hooper, K. L., Glynn, N. M., Burnside, J., Coppock, D. L., and Thorpe, C. (1999) Homology between egg white sulfhydryl oxidase and quiescin Q6 defines a new class of flavin-linked sulfhydryl oxidases. *J. Biol. Chem.* 274, 31759–31762.
- Jaje, J., Wolcott, H. N., Fadugba, O., Cripps, D., Yang, A. J., Mather, I. H., and Thorpe, C. (2007) A flavin-dependent sulfhydryl oxidase in bovine milk. *Biochemistry* 46, 13031–13040.
- Heckler, E. J., Alon, A., Fass, D., and Thorpe, C. (2008) Human quiescin-sulfhydryl oxidase, QSOX1: probing internal redox steps by mutagenesis. *Biochemistry* 47, 4955–4963.
- Raje, S., and Thorpe, C. (2003) Inter-domain redox communication in flavoenzymes of the quiescin/sulfhydryl oxidase family: role of a thioredoxin domain in disulfide bond formation. *Biochemistry* 42, 4560–4568.
- Gruber, C. W., Cemazar, M., Heras, B., Martin, J. L., and Craik, D. J. (2006) Protein disulfide isomerase: the structure of oxidative folding. *Trends Biochem. Sci.* 31, 455–464.
- Tian, G., Xiang, S., Noiva, R., Lennarz, W. J., and Schindelin, H. (2006) The crystal structure of yeast protein disulfide isomerase suggests cooperativity between its active sites. *Cell* 124, 61–73.
- Hatahet, F., and Ruddock, L. W. (2009) Protein disulfide isomerase: a critical evaluation of its function in disulfide bond formation. *Antioxid. Redox Signaling* 11, 2807–2850.
- Atkinson, H. J., and Babbitt, P. C. (2009) An atlas of the thioredoxin fold class reveals the complexity of function-enabling adaptations. *PLoS Comput. Biol.* 5, e1000541.
- Lisowsky, T. (1994) ERV1 is involved in the cell-division cycle and the maintenance of mitochondrial genomes in *Saccharomyces cerevisiae*. *Curr. Genet.* 26, 15–20.
- Lee, J., Hofhaus, G., and Lisowsky, T. (2000) Erv1p from *Saccharomyces cerevisiae* is a FAD-linked sulfhydryl oxidase. *FEBS Lett.* 477 (1–2), 62–66.
- Gross, E., Sevier, C. S., Vala, A., Kaiser, C. A., and Fass, D. (2002) A new FAD-binding fold and intersubunit disulfide shuttle in the thiol oxidase Erv2p. *Nat. Struct. Biol.* 9, 61–67.
- Wu, C. K., Dailey, T. A., Dailey, H. A., Wang, B. C., and Rose, J. P. (2003) The crystal structure of augmentin of liver regeneration: a mammalian FAD-dependent sulfhydryl oxidase. *Protein Sci.* 12, 1109–1118.
- Vitu, E., Bentzur, M., Lisowsky, T., Kaiser, C. A., and Fass, D. (2006) Gain of function in an ERV/ALR sulfhydryl oxidase by molecular engineering of the shuttle disulfide. *J. Mol. Biol.* 362, 89–101.
- Fass, D. (2008) The Erv family of sulfhydryl oxidases. *Biochim. Biophys. Acta* 1783, 557–566.
- Hakim, M., and Fass, D. (2009) Dimer interface migration in a viral sulfhydryl oxidase. *J. Mol. Biol.* 391, 758–768.
- Barry, D., and Carrington, M. (2004) Antigenic variation, in *The Trypanosomiasis* (Maudlin, I., Holmes, P., and Miles, M., Eds.) pp 25–38, International CABI Publishing, U.K.
- Chattopadhyay, A., Jones, N. G., Nietlispach, D., Nielsen, P. R., Voorheis, H. P., Mott, H. R., and Carrington, M. (2005) Structure of the C-terminal domain from *Trypanosoma brucei* variant surface glycoprotein MITat1.2. *J. Biol. Chem.* 280, 7228–7235.
- Carrington, M., Miller, N., Blum, M., Roditi, I., Wiley, D., and Turner, M. (1991) Variant specific glycoprotein of *Trypanosoma brucei* consists of two domains each having an independently conserved pattern of cysteine residues. *J. Mol. Biol.* 221, 823–835.
- Freyman, D., Down, J., Carrington, M., Roditi, I., Turner, M., and Wiley, D. (1990) 2.9 Å resolution structure of the N-terminal domain of a variant surface glycoprotein from *Trypanosoma brucei*. *J. Mol. Biol.* 216, 141–160.
- Blum, M. L., Down, J. A., Gurnett, A. M., Carrington, M., Turner, M. J., and Wiley, D. C. (1993) A structural motif in the variant surface glycoproteins of *Trypanosoma brucei*. *Nature* 362, 603–609.
- Jones, N. G., Nietlispach, D., Sharma, R., Burke, D. F., Eyres, I., Mues, M., Mott, H. R., and Carrington, M. (2008) Structure of a glycosylphosphatidylinositol-anchored domain from a trypanosome variant surface glycoprotein. *J. Biol. Chem.* 283, 3584–3593.
- Fairlamb, A. H., Blackburn, P., Ulrich, P., Chait, B. T., and Cerami, A. (1985) Trypanothione: a novel bis(glutathionyl)spermidine co-factor for glutathione reductase in trypanosomatids. *Science* 227, 1485–1487.
- Krauth-Siegel, L. R., Comini, M. A., and Schlecker, T. (2007) The trypanothione system. *Subcell. Biochem.* 44, 231–251.
- Sevier, C. S., and Kaiser, C. A. (2006) Disulfide transfer between two conserved cysteine pairs imparts selectivity to protein oxidation by Ero1. *Mol. Biol. Cell* 17, 2256–2266.
- Mezghrani, A., Fassio, A., Benham, A., Simmen, T., Braakman, I., and Sitia, R. (2001) Manipulation of oxidative protein folding and PDI redox state in mammalian cells. *EMBO J.* 20, 6288–6296.
- Tu, B. P., Ho-Schleyer, S. C., Travers, K. J., and Weissman, J. S. (2000) Biochemical basis of oxidative protein folding in the endoplasmic reticulum. *Science* 290, 1571–1574.
- Frandsen, A. R., and Kaiser, C. A. (1999) Ero1p oxidizes protein disulfide isomerase in a pathway for disulfide bond formation in the endoplasmic reticulum. *Mol. Cell* 4, 469–477.
- Baker, K. M., Chakravarthi, S., Langton, K. P., Sheppard, A. M., Lu, H., and Bulleid, N. J. (2008) Low reduction potential of Ero1alpha regulatory disulphides ensures tight control of substrate oxidation. *EMBO J.* 27, 2988–2997.
- Daithankar, V. N., Farrell, S. R., and Thorpe, C. (2009) Augmentin of liver regeneration: substrate specificity of a flavin-dependent oxidoreductase from the mitochondrial intermembrane space. *Biochemistry* 48, 4828–4837.
- Deponte, M., and Hell, K. (2009) Disulfide bond formation in the intermembrane space of mitochondria. *J. Biochem.* 146, 599–608.
- Bihlmaier, K., Mesecke, N., Terziyska, N., Bien, M., Hell, K., and Herrmann, J. M. (2007) The disulfide relay system of mitochondria is connected to the respiratory chain. *J. Cell Biol.* 179, 389–395.
- Tokatlidis, K. (2005) A disulfide relay system in mitochondria. *Cell* 121, 965–967.
- Ramadan, D., Rancy, P. C., Nagarkar, R. P., Schneider, J. P., and Thorpe, C. (2009) Arsenic(III) species inhibit oxidative protein folding in vitro. *Biochemistry* 48, 424–432.

41. Emanuelsson, O., Brunak, S., von Heijne, G., and Nielsen, H. (2007) Locating proteins in the cell using TargetP, SignalP and related tools. *Nat. Protoc.* 2, 953–971.
42. Bryson, K., McGuffin, L. J., Marsden, R. L., Ward, J. J., Sodhi, J. S., and Jones, D. T. (2005) Protein structure prediction servers at University College London. *Nucleic Acids Res.* 33, W36–38.
43. Chenna, R., Sugawara, H., Koike, T., Lopez, R., Gibson, T. J., Higgins, D. G., and Thompson, J. D. (2003) Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res.* 31, 3497–3500.
44. Wilkinson, B., and Gilbert, H. F. (2004) Protein disulfide isomerase. *Biochim. Biophys. Acta* 1699, 35–44.
45. Wilkinson, B., Xiao, R., and Gilbert, H. F. (2005) A structural disulfide of yeast protein-disulfide isomerase destabilizes the active site disulfide of the N-terminal thioredoxin domain. *J. Biol. Chem.* 280, 11483–11487.
46. Krauth-Siegel, R. L., and Comini, M. A. (2008) Redox control in trypanosomatids, parasitic protozoa with trypanothione-based thiol metabolism. *Biochim. Biophys. Acta* 1780, 1236–1248.
47. Navon, A., Ittah, V., Landsman, P., Scheraga, H. A., and Haas, E. (2001) Distributions of intramolecular distances in the reduced and denatured states of bovine pancreatic ribonuclease A. Folding initiation structures in the C-terminal portions of the reduced protein. *Biochemistry* 40, 105–118.
48. Hooper, K. L., Sheasley, S. S., Gilbert, H. F., and Thorpe, C. (1999) Sulfhydryl oxidase from egg white: a facile catalyst for disulfide bond formation in proteins and peptides. *J. Biol. Chem.* 274, 22147–22150.
49. Rancy, P. C., and Thorpe, C. (2008) Oxidative protein folding in vitro: a study of the cooperation between quiescin-sulfhydryl oxidase and protein disulfide isomerase. *Biochemistry* 47, 12047–12056.
50. Hooper, K. L., and Thorpe, C. (1999) Egg white sulfhydryl oxidase: kinetic mechanism of the catalysis of disulfide bond formation. *Biochemistry* 38, 3211–3217.
51. Thorpe, C., and Williams, C. H. (1976) Spectral evidence for a flavin adduct in a monoalkylated derivative of pig heart lipoamide dehydrogenase. *J. Biol. Chem.* 251, 7726–7728.
52. Dmitrenko, O., and Thorpe, C. (2008) A computational analysis of the interaction between flavin and thiol(ate) groups. Implications for flavoenzyme catalysis. *J. Sulfur Chem.* 29, 415–C421.
53. Williams, C. H., Arscott, L. D., Matthews, R. G., Thorpe, C., and Wilkinson, K. D. (1979) Methodology employed for anaerobic spectrophotometric titrations and for computer-assisted data analysis. *Methods Enzymol.* 185–198.
54. Mayhew, S. G. (1978) Redox potential of dithionite and SO_2^- from equilibrium reactions with flavodoxins, methyl viologen and hydrogen plus hydrogenase. *Eur. J. Biochem.* 85, 535–547.
55. Wang, W., Winther, J. R., and Thorpe, C. (2007) Erv2p: characterization of the redox behavior of a yeast sulfhydryl oxidase. *Biochemistry* 46, 3246–3254.
56. Farrell, S. R., and Thorpe, C. (2005) Augmenter of liver regeneration: a flavin dependent sulfhydryl oxidase with cytochrome C reductase activity. *Biochemistry* 44, 1532–1541.
57. Mattevi, A. (2006) To be or not to be an oxidase: challenging the oxygen reactivity of flavoenzymes. *Trends Biochem. Sci.* 31, 276–283.
58. Strickland, S., Palmer, G., and Massey, V. (1975) Determination of dissociation constants and specific rate constants of enzyme-substrate (or protein-ligand) interactions from rapid reaction kinetic data. *J. Biol. Chem.* 250, 4048–4052.
59. Williams, C. H., Jr. (1992) Lipoamide dehydrogenase, glutathione reductase, thioredoxin reductase, and mercuric ion reductase—A family of flavoenzyme transhydrogenases, in *Chemistry and Biochemistry of Flavoenzymes* (Müller, F., Ed.) pp 121–211, CRC Press, Boca Raton, FL.
60. Miller, S. M., Massey, V., Ballou, D., Williams, C. H., Jr., Distefano, M. D., Moore, M. J., and Walsh, C. T. (1990) Use of a site-directed triple mutant to trap intermediates: demonstration that the flavin C(4a)-thiol adduct and reduced flavin are kinetically competent intermediates in mercuric ion reductase. *Biochemistry* 29, 2831–2841.
61. O'Donnell, M. E., and Williams, C. H., Jr. (1984) Reconstitution of *Escherichia coli* thioredoxin reductase with 1-deazaFAD. Evidence for 1-deazaFAD C-4a adduct formation linked to the ionization of an active site base. *J. Biol. Chem.* 259, 2243–2251.
62. Otsu, M., Bertoli, G., Fagioli, C., Guerini-Rocco, E., Nerini-Molteni, S., Ruffato, E., and Sitia, R. (2006) Dynamic retention of Ero1 α and Ero1 β in the endoplasmic reticulum by interactions with PDI and ERp44. *Antioxid. Redox Signaling* 8, 274–282.
63. Chakravarthi, S., Jessop, C. E., Willer, M., Stirling, C. J., and Bulleid, N. J. (2007) Intracellular catalysis of disulphide bond formation by the human sulphhydryl oxidase, QSOX1. *Biochem. J.* 404, 403–411.