

Molecular Basis for Enantioselectivity in the (*R*)- and (*S*)-Hydroxypropylthioethanesulfonate Dehydrogenases, a Unique Pair of Stereoselective Short-Chain Dehydrogenases/Reductases Involved in Aliphatic Epoxide Carboxylation[†]

Dariusz A. Sliwa,[‡] Arathi M. Krishnakumar,[§] John W. Peters,[§] and Scott A. Ensign^{*‡}

[‡]Department of Chemistry and Biochemistry, Utah State University, Logan, Utah 84322, and [§]Department of Chemistry and Biochemistry, Montana State University, Bozeman, Montana 59717

Received February 27, 2010; Revised Manuscript Received March 16, 2010

ABSTRACT: (*R*)- and (*S*)-2-hydroxypropyl-CoM (*R*-HPC and *S*-HPC) are produced as intermediates in bacterial propylene metabolism from the nucleophilic addition of coenzyme M to (*R*)- and (*S*)-epoxypropane, respectively. Two highly enantioselective dehydrogenases (*R*-HPCDH and *S*-HPCDH) belonging to the short-chain dehydrogenase/reductase family catalyze the conversion of *R*-HPC and *S*-HPC to 2-ketopropyl-CoM (2-KPC), which undergoes reductive cleavage and carboxylation to produce acetoacetate. In the present study, one of three copies of *S*-HPCDH enzymes present on a linear megaplasmid in *Xanthobacter autotrophicus* strain Py2 has been cloned and overexpressed, allowing the first detailed side by side characterization of the *R*-HPCDH and *S*-HPCDH enzymes. The catalytic triad of *S*-HPCDH was found to consist of Y156, K160, and S143. R211 and K214 were identified as the amino acid residues coordinating the sulfonate of CoM in *S*-HPC. R211A and K214A mutants were severely impaired in the oxidation of *S*-HPC or reduction of 2-KPC but were largely unaffected in the oxidation and reduction of aliphatic alcohols and ketones. Kinetic analyses using *R*- and *S*-HPC as substrates revealed that enantioselectivity in *R*-HPCDH (value, 944) was dictated largely by differences in k_{cat} while enantioselectivity for *S*-HPCDH (value, 1315) was dictated largely by changes in K_{m} . *S*-HPCDH had an inherent high enantioselectivity for producing (*S*)-2-butanol from 2-butanone that was unaffected by modulators that interact with the sulfonate binding site. The tertiary alcohol 2-methyl-2-hydroxypropyl-CoM (M-HPC) was a competitive inhibitor of *R*-HPCDH-catalyzed *R*-HPC oxidation, with a K_{i} similar to the K_{m} for *R*-HPC, but was not an inhibitor of *S*-HPCDH. The primary alcohol 2-hydroxyethyl-CoM was a substrate for both *R*-HPCDH and *S*-HPCDH with identical K_{m} values. The pH dependence of kinetic parameters suggests that the hydroxyl group is a larger contributor to *S*-HPC binding to *S*-HPCDH than for *R*-HPC binding to *R*-HPCDH. It is proposed that active site constraints within the *S*-HPCDH prevent proper binding of *R*-HPC and M-HPC due to steric clashes with the improperly aligned methyl group on the C2 carbon, resulting in a different mechanism for controlling substrate specificity and enantioselectivity than present in the *R*-HPCDH.

Alcohol dehydrogenases that catalyze the interconversion of secondary alcohols and ketones are often highly stereoselective with regard to the enantiomer of the alcohol oxidized or produced during the course of catalysis (1–3). However, there

are very few examples of pairs of alcohol dehydrogenases synthesized simultaneously to deal with both the *R*- and *S*-enantiomers of an alcohol produced during a metabolic pathway. One such unique pair was discovered in the pathway of propylene oxidation by the hydrocarbon-oxidizing bacteria *Xanthobacter autotrophicus* Py2 and *Rhodococcus rhodochrous* B276 (4–8). As shown in Figure 1, these bacteria initiate propylene oxidation by epoxidation to produce a mixture of the *R*- and *S*-enantiomers of epoxypropane, which are further metabolized by a coenzyme M (2-mercaptoethanesulfonate, CoM)¹ mediated sequence of reactions involving nucleophilic addition, dehydrogenation, reductive cleavage, and carboxylation. (*R*)- and (*S*)-2-hydroxypropyl-CoM (*R*- and *S*-HPC) are the substrates for the unique pair of dehydrogenases that form 2-ketopropyl-CoM (2-KPC), which is reductively cleaved and carboxylated (4).

Biochemical and molecular characterization of the *R*- and *S*-HPCDH enzymes showed that they are members of the “classical” short-chain dehydrogenase/reductase (SDR) family of alcohol dehydrogenases (4–6). Classical SDR enzymes are defined by a conserved NAD⁺-binding motif in the N-terminal portion of the protein and a catalytic triad (or tetrad) within the

[†]This work was supported by National Institutes of Health Grant GM51805 to S.A.E. and by Department of Energy Grant DE-FG02-04ER15563 to J.W.P.

^{*}To whom correspondence should be addressed: (435) 797-3969 (phone); (435) 797-3390 (fax); scott.ensign@usu.edu (e-mail).

¹Abbreviations: *R*-HPC, 2-[(*R*)-2-hydroxypropylthio]ethanesulfonate ((*R*)-2-hydroxypropyl-CoM); *S*-HPC, 2-[(*S*)-2-hydroxypropylthio]ethanesulfonate ((*S*)-2-hydroxypropyl-CoM); *S*-HPCDH, 2-[(*S*)-2-hydroxypropylthio]ethanesulfonate dehydrogenase; *rS*-HPCDH, recombinant 2-[(*S*)-2-hydroxypropylthio]ethanesulfonate dehydrogenase; *R*-HPCDH, 2-[(*R*)-2-hydroxypropylthio]ethanesulfonate dehydrogenase; *rR*-HPCDH, recombinant 2-[(*R*)-2-hydroxypropylthio]ethanesulfonate dehydrogenase; M-HPC, 2-(2-methyl-2-hydroxypropylthio)ethanesulfonate; HEC, 2-(2-hydroxyethylthio)ethanesulfonate; 2-KPC, 2-(2-ketopropylthio)ethanesulfonate (2-ketopropyl-CoM); 2-KPCC, 2-(2-ketopropylthio)ethanesulfonate carboxylase/oxidoreductase; CoM, coenzyme M (2-mercaptoethanesulfonate); SDR, short-chain dehydrogenase/reductase; Tris, tris(hydroxymethyl)aminomethane; CD, circular dichroism; ee, enantiomeric excess; *E*, enantioselectivity; SDS–PAGE, sodium dodecyl sulfate–polyacrylamide gel electrophoresis.




FIGURE 1: Pathway of propylene oxidation in *X. autotrophicus* Py2. The reactions catalyzed by *R*-HPCDH and *S*-HPCDH are shown in the boxed region.

central portion (9–13). The C-terminal domains of the SDR families diverge to confer specificity for different substrates. The *R*- and *S*-HPCDH enzymes from *X. autotrophicus* are homologous enzymes that share high sequence identity but with notable sequence differences within their C-terminal regions (5–8). These enzymes are highly specific for their respective enantiomers of HPC, exhibiting less than 0.5% activity with the opposite isomer (4).

The *R*-HPCDH from *X. autotrophicus* has been extensively characterized mechanistically and structurally, culminating in the formulation of a mechanism of catalysis and stereoselectivity shown on the left side of Scheme 1 (14–16). Stereoselectivity is conferred in large part by two positively charged arginine residues that form salt bridges with the CoM moiety of the substrate within the CoM-binding pocket of the enzyme (14, 16). This strong ionic interaction orients the hydroxyl group and hydrogen atom of the alcohol functional group in the proper orientation for general base abstraction and hydride transfer (Scheme 1) (16). By analogy, a similar mechanistic strategy is believed to operate in the *S*-HPCDH, wherein the spatial orientations of the CoM- and methyl-binding pockets are reversed relative to the hydroxyl group and hydrogen atom to be transferred (Scheme 1, right side) (14, 16, 17). To date, this model has not been tested for the *S*-HPCDH due to difficulties encountered in attempting to express the enzyme in an active form in *Escherichia coli* (14, 15).

In the present work, one of three copies of *S*-HPCDH homologues present in *X. autotrophicus* has been successfully cloned and overexpressed, providing the necessary system for testing the tenets of Scheme 1. These results show that the *R*- and *S*-HPCDH enzymes share some common mechanistic features but differ significantly in the strategies for controlling enantioselectivity

for the *S*- and *R*-HPC natural substrates, as well as nonphysiological aliphatic alcohols and ketones.



EXPERIMENTAL PROCEDURES

Materials. All commercially available chemicals were purchased from Sigma-Aldrich Chemicals, Acros Organics, or Fisher Scientific and were of analytical grade. 2-KPC, *R*-HPC, *S*-HPC, HEC, and M-HPC were synthesized as described previously (15). Chemical structures of the compounds were confirmed using ¹H NMR. The spectra of HPC enantiomers and 2-KPC were identical to those reported previously (4, 15). Purity of the synthesized chemicals as determined by reverse-phase HPLC was estimated to be ≥98%.

Cloning of the *S*-HPCDH Genes (*xecE1*, *xecE2*, and *xecE3*). Total genomic DNA was isolated from propylene-grown cells of *X. autotrophicus* strain Py2 using the Epicenter MasterPure DNA purification kit. Each of the three *xecE* genes was PCR amplified using the FailSafe PCR premix selection kit (Epicenter). Primers were designed such that the forward and reverse primers contained *Bam*HI and *Sac*I restriction site overhangs, respectively. The primers were as follows: *xecE1*, forward, GCAGGATCCAATGCTGGACGCAGAGG, and reverse, CGTGAGCTCTCATATGGCGGTCATCC; *xecE2*, forward, TAGGATCCAGTGGCGCGCGCCGCGGT, and reverse, ATGAGCTCTCATATGGCGGTCATCCCTC; *xecE3*, forward, GCAGGATCCAATGTCGAATCGCTTGAAG, and reverse, GCA-GAGCTCTCATATCGCCGTCATC. The reactions contained 200 ng of *X. autotrophicus* genomic DNA, 1.0 μM primers, FailSafe PCR enzyme mix (1.25 units), and the manufacturer's buffer H. PCR was performed using the following cycling parameters: stage 1, (95 °C × 5 min) × 1; stage 2, (94 °C × 1 min, 60 °C × 1 min, 72 °C × 50 s) × 35; stage 3, (72 °C × 10 min) × 1. The PCR products were analyzed on 1% (w/v) agarose gels, and *xecE1*, *xecE2*, and *xecE3* DNA (0.75, 0.69, and 0.77 kb, respectively) were purified using the QIAquick gel extraction kit (Qiagen). Purified DNA along with pET28-b (Novagen) and pRSFDuet-1 expression vectors were simultaneously subjected to a double digest with *Bam*HI and *Sac*I, resolved on 1% agarose gel, and purified with the QIAquick gel extraction kit. Digested *xecE* genes and pET28-b DNA were mixed and ligated with T4 DNA ligase for 2 h at 25 °C. The resulting plasmids were designated pDS41, pDS42, and pDS43 and carried the *xecE1*, *xecE2*, and *xecE3* genes, respectively. Likewise, digested *xecE* genes and pRSFDuet-1 DNA were mixed and ligated with T4 DNA ligase for 2 h at 25 °C to result in pDS51, pDS52, and pDS53 plasmids with *xecE1*, *xecE2*, and *xecE3* genes, respectively. Subsequent to DNA sequencing, all constructs were transformed into *E. coli* DH5α for plasmid maintenance and into *E. coli* BL21(DE3) CodonPlus (Stratagene) cells for protein expression.

Site-Directed Mutagenesis (SDM). All primers used to introduce point mutations were purchased from IDT. SDM of pDS53 was carried out utilizing the QuickChange site-directed mutagenesis kit (Stratagene) according to the manufacturer's protocols. All mutations were confirmed by DNA sequencing. The sequences of the mutagenic primer pairs used for each codon substitution in *xecE3* are as follows: S143A, GATCGTCAATTTTGGCGCCGTCGCTGGCCTC and GAGCCAGCGACGCGCCCAAAATTGACGATC; Y156A, GACCATGGCGCGCCGCTGCGCAGCCAAG and CTTGGCTGCGCAGGCGGCCGCGCATGTGTC; Y156F, CCATGGCGGCCTTCTGCGCAGCCAAGG and CTTGGCTGCGCAGAAGGCCGCGCATGG; K160A,

Scheme 1



CCTACTGCGCAGCCGCGGGCGCAATCGTCA and TGA-CGATTGCGCCCGCGCTGCGCAGTAGG; R211A, GAA-GTTCAGGCTCGCGCGCTGGCGAAATATCC and GGA-TATTCGCCAGCGCGGAGCCTGAACTTC; K214A, CTC-GCCGGCTGGCGGCATATCCGATCGGGC and GCCCG-ATCGGATATGCCGCCAGCCGGCGAG.

DNA Sequencing. Sequencing was performed on an AB 3730 DNA analyzer at the Utah State University Biotechnology Center DNA sequencing laboratory. The following sequencing primers were used to confirm all mutations to pDS53: ACYC-DuetUP1 primer (Novagen), GGATCTCGACGCTCTCCCT, and DuetDOWN1 primer (Novagen), GATTATGCGGCC-GTGTACAA.

Media and Growth of Bacteria. *X. autotrophicus* was grown on propylene (10% (v/v) gas phase) in a 15 L semicontinuous microferm fermenter as described previously (18). *E. coli* DH5 α was grown in standard Luria–Bertani (LB) broth supplemented with kanamycin (50 $\mu\text{g}\cdot\text{mL}^{-1}$). *E. coli* BL21(DE3) CodonPlus was grown in LB media that contained both kanamycin (50 $\mu\text{g}\cdot\text{mL}^{-1}$) and chloramphenicol (50 $\mu\text{g}\cdot\text{mL}^{-1}$). All other procedures were performed as described previously (15).

Preparation of Cell-Free Extracts. About 60 g of frozen cell paste was thawed at room temperature in 2 volumes of lysis buffer (20 mM Tris–chloride, 20% glycerol, 500 mM NaCl, 25 mM imidazole, 0.1% (w/v) Tween 20, 0.03 $\text{mg}\cdot\text{mL}^{-1}$ DNase I, at pH 8.0). Homogenized cell suspension was passed twice through a French press at 15000 psi, and the crude lysate was clarified by centrifugation at 244717g for 45 min.

Purification of S-HPCDH1, rS-HPCDH1, rS-HPCDH3, and rS-HPCDH3 Mutants. S-HPCDH was purified from propylene-grown *X. autotrophicus* as described previously (5). For the purification of rS-HPCDH1, rS-HPCDH3, and rS-HPCDH3 mutants, clarified cell extract was applied to a 2.6×4.7 cm column of Ni Sepharose 6 fast flow (Amersham) at $3.0\text{ mL}\cdot\text{min}^{-1}$ ($33.9\text{ cm}\cdot\text{h}^{-1}$). The column was rinsed at $6.0\text{ mL}\cdot\text{min}^{-1}$ ($67.8\text{ cm}\cdot\text{h}^{-1}$) with 2 column volumes of lysis buffer, followed by 80 column volumes of rinse buffer (20 mM Tris, 20% glycerol, 500 mM NaCl, 50 mM imidazole, 0.1% (w/v) Tween 20, pH 8.0). rS-HPCDH was eluted at $4.0\text{ mL}\cdot\text{min}^{-1}$ ($45.2\text{ cm}\cdot\text{h}^{-1}$) with a linear gradient of 19.2 column volumes of 0–100% elution buffer (20 mM Tris, 20% glycerol, 500 mM NaCl, 500 mM imidazole, pH 8.0). Fractions containing rS-HPCDH were concentrated over a 30000 MW cutoff membrane, diluted in a 1:20 ratio with dialysis buffer (20 mM Tris, 10% glycerol,

100 mM NaCl, pH 8.2), and concentrated again (this procedure was repeated twice). Dialyzed rS-HPCDH was flash frozen and stored in liquid nitrogen. All steps were performed at 4 $^{\circ}\text{C}$, and all buffers used had the pH adjusted at 4 $^{\circ}\text{C}$. Protein concentrations were determined on a NanoDrop spectrophotometer using theoretical extinction coefficients ($\epsilon_{280} = 10033\text{ M}^{-1}\text{ cm}^{-1}$ for S-HPCDH3, $\epsilon_{280} = 15595\text{ M}^{-1}\text{ cm}^{-1}$ for S-HPCDH1, and $\epsilon_{280} = 18512.5\text{ M}^{-1}\text{ cm}^{-1}$ for R-HPCDH1) with dialysis flow-through buffers as blanks.

SDS–PAGE and Native PAGE Analysis. SDS–PAGE (12% T) and native PAGE (4–20% T; Bio-Rad) were performed according to the Laemmli procedure (19). The apparent molecular masses of polypeptides on a SDS–PAGE gel were determined by comparison to R_f values of standard proteins. Migration of rS-HPCDH3 mutants was also compared directly to wild-type rS-HPCDH3, wild-type rR-HPCDH1, native S-HPCDH, and R-HPCDH (isolated from *X. autotrophicus*). The apparent molecular masses of polypeptides on a native PAGE gel were determined from a standard curve constructed by plotting R_f values against the log of the native molecular mass for the following standards: β -amylase (200 kDa), alcohol dehydrogenase (150 kDa), bovine serum albumin (66 kDa), and ovalbumin (43 kDa).

Gel Filtration Chromatography. The native molecular masses of rS-HPCDH3 and rS-HPCDH3 mutants were estimated by gel filtration chromatography using HPLC (Shimadzu SLC-10A) with a fluorescence detector (Shimadzu RF-10AXL) set up for excitation and emission wavelengths at 280 and 350 nm, respectively. The gel filtration column (BioSep-SEC S-2000, 300×7.8 mm; Phenomenex) was equilibrated in 50 mM Na_2HPO_4 and 200 mM NaCl, pH 7.5, containing 10% glycerol and 1 mM NAD^+ . The following molecular mass standards were used for calibration: β -amylase (200 kDa), aldolase (158 kDa), conalbumin (75 kDa), bovine serum albumin (66 kDa), ovalbumin (43 kDa), carbonic anhydrase (29 kDa), and myoglobin (18.8 kDa). All separations were performed at 4 $^{\circ}\text{C}$ at a flow rate of $0.5\text{ mL}\cdot\text{min}^{-1}$. A calibration curve was constructed by plotting the retention time against the log of the native molecular mass for standards. This plot was fit using a second-order polynomial, and the equation of this line was used to determine the log native molecular masses for rS-HPCDH3 and rS-HPCDH3 mutants using their experimentally determined retention times.

Circular Dichroism (CD). CD spectra were recorded at 25 $^{\circ}\text{C}$ on an AVIV Model 410 CD spectrophotometer using 1 nm

spectral bandwidth and a 0.1 cm path length. Stock peptide solutions were desalted on Sephadex G-25 (PD-10) columns and diluted in 10× buffer to a final concentration of 10% glycerol, 10 mM KH₂PO₄, and 100 mM KF at pH 7.0. Total enzyme concentration was approximately 0.3 mg·mL⁻¹ (10 μM). Typically, five scans were acquired over the wavelength range 190–260 nm. The residue ellipticity (θ) was calculated using *rS*-HPCDH3 molecular mass of 27085.5 Da and N_A (number of amino acid in the protein) of 268 residues.

Spectrophotometric Enzyme Assays. Assays with 2-KPC, S-HPC, R-HPC, and 2-propanol as substrates were performed in 50 mM glycine, 50 mM NaH₂PO₄, and 50 mM Tris–base (GPT buffer mix) at a pH of 7.5 (adjusted at 30 °C) as described previously (15). Assays with all other substrates (C₄ to C₆ in carbon chain length) were carried out in 50 mM GPT buffer containing 15% (v/v) glycerol. Stock solutions of synthesized substrates were standardized as described previously (15). All enzyme assays were performed in triplicate at 30 °C in a Shimadzu model UV160U spectrophotometer containing a water-jacketed cell holder for thermal control. Alcohol or ketone production was monitored by measuring the change in absorbance at 340 nm using the extinction coefficient for NADH ($\epsilon_{340} = 6.22 \text{ mM}^{-1} \text{ cm}^{-1}$). For alcohol oxidation assays, the following ranges of alcohol concentrations were used in determining kinetic constants for R-HPCDH1: R-HPC, 0.020–1.0 mM; S-HPC, 0.074–11 mM; 2-propanol, 65–3900 mM; (S)-2-butanol, 10.9–381 mM; (R)-2-butanol, 43.6–490 mM. For S-HPCDH3 the concentrations were as follows: R-HPC, 0.038–48 mM; S-HPC, 0.016–0.310 mM; 2-propanol, (S)-2-butanol, and (R)-2-butanol, the same as for R-HPCDH1. The concentration of NAD⁺ for all assays was 10 mM (26 × value of $K_{m\text{NAD}^+}$). For assays of ketone reduction, the following concentration ranges were used in determining kinetic constants: 2-KPC, 0.050–2.6 mM; 2-butanone, 10–300 mM. The concentration of NADH for these assays was 0.16 mM (4.4 × value of $K_{m\text{NADH}}$). On average, seven concentrations of substrates within the ranges indicated were chosen for the kinetic analyses. All samples were degassed/flushed with nitrogen and incubated at 30 °C water bath for 5 min prior to the enzyme addition. To determine kinetic parameters (apparent K_m and V_{\max}), initial rate values were plotted as a function of substrate concentration, and data points were fitted to the Michaelis–Menten equation using SigmaPlot 11.0.

pH Dependence of Kinetic Parameters. All assays were performed in GPT buffer mix or APT buffer mix (50 mM CH₃COONa, 50 mM NaH₂PO₄, and 50 mM Tris–base) at ten different pH values (5.0–10.0) adjusted to the desired pH at 30 °C as described previously (15). The assay results were used to construct a plot of k_{cat}/K_m vs pH, which was then fit to eq 1:

$$\log k_{\text{cat}}/K_m = \log(C/(1 + [\text{H}^+]/K_a)) \quad (1)$$

where C is the maximal $\log k_{\text{cat}}/K_m$ value and K_a is the acid dissociation constant of the catalytically important ionizing residue. Plots of $\log k_{\text{cat}}$ and $\log K_m$ vs pH were constructed in a point-to-point manner. Stability of the enzyme at pH 5.0 and 10.0 was tested according to the pH-jump method (20). The stock solution of *rS*-HPCDH3 was diluted to 0.1 mg·mL⁻¹ in 10 mM APT or GPT buffer mix containing 15% glycerol, at pH values of 5.0 and 10.0, respectively. The enzyme was equilibrated on ice at the desired pH (adjusted at 4 °C) for 5 min prior to its addition to the reaction mixture (total of 0.3 μg of *rS*-HPCDH3). Activity

assays were performed for 30 s in 100 mM GPT buffer mix, pH 7.5, at 30 °C.

Inhibition Studies. All assays were performed in GPT buffer mix at pH 7.5 with saturating concentration of NAD⁺ (5 mM). Inhibition assays for R-HPCDH1 were performed at variable concentrations of R-HPC (31, 72, 144, 287, and 615 μM) or HEC (0.8, 2.0, 4.0, 8.0, and 1.6 mM). Each assay was performed at several fixed concentrations of M-HPC (0, 0.2, 0.4, 0.8, and 1.6 mM). Initial rate data for *rR*-HPCDH1 were fit to eq 2 describing competitive inhibition of enzymatic activity:

$$v = V_{\max}[S]/(\alpha K_m + [S]) \quad (2)$$


where S is the substrate, I is the inhibitor, $\alpha = 1 + ([I]/K_{is})$, and K_{is} is the inhibition constant for the competitive component of enzyme inhibition. The type of inhibition exhibited by M-HPC was determined by graphing initial rate data in the form of double reciprocal plots ($1/v$ vs $1/[S]$ at various $[I]$). The numerical value of K_{is} was calculated from eq 2 using SigmaPlot. Inhibition assays for S-HPCDH3 were performed at varying concentrations of S-HPC (16, 31, 62, 155, and 310 μM) or HEC (0.4, 0.8, 2.0, 4.0, 8.0, and 1.6 mM) and three fixed concentrations of M-HPC (0, 1.2, and 4.9 mM). Initial rate data were fit to a rectangular hyperbola described by the standard form of the Michaelis–Menten equation.

Chiral Gas Chromatographic Assay for 2-Butanone Reduction. All assays were performed in GPT buffer mix, pH 7.5 (adjusted at 30 °C), containing 15% (v/v) glycerol. Assay components in 1 mL reaction volume were NADH (15 mM), 2-butanone (56 mM), and 0.64 mg of enzyme. Serum vials (3 mL) were crimp-sealed and degassed/flushed with nitrogen three times and allowed to equilibrate for 20 min at 30 °C in a shaking water bath (200 rpm). The assay was initiated by adding enzyme. After 1 h incubation 250 μL of headspace gas was removed and injected into a Shimadzu GC-8A gas chromatograph outfitted with a Supelco β-Dex 225 (30 m × 0.53 mm) column. GC parameters used were as previously described (14).

RESULTS AND DISCUSSION

Genetic and Bioinformatic Analysis of the Enzymes of Epoxide Metabolism. A previous study demonstrated that the genes of alkene and epoxide metabolism are present on a 320 kb linear megaplasmid in *X. autotrophicus* Py2 (21). Furthermore, the genes encoding the enzymes of epoxide carboxylation were found to be clustered in an operon where *xecA*, *xecC*, *xecD*, and *xecE* encode the key enzymes shown in Figure 1 (5, 22). The *xecD* gene (encoding R-HPCDH) was successfully cloned, overexpressed, and purified in a fully active state, allowing for detailed mechanistic and structural studies to be performed (14–16). Unfortunately, all attempts to clone and express the counterpart S-HPCDH in a soluble and active form were unsuccessful despite numerous attempts (15), and the mechanistic and structural characterization of this enzyme has thus lagged behind that of the R-HPCDH.

Recently, the genome of *X. autotrophicus* Py2 was sequenced, assembled, and annotated (<http://genome.jgi-psf.org/xanau/xanau.home.html>). Unexpectedly, an analysis of the linear megaplasmid of *X. autotrophicus* shows the presence of multiple copies of the *xec* genes, as well as for the putative enzymes of coenzyme M biosynthesis. The *xec* paralogues are distributed among three apparent operons and share high identities. The *xecD* and *xecE* genes whose protein sequences were reported previously (5, 22)



Cloning, Expression, and Biochemical Characterization of *xecE* Homologues. Each of the three S-HPCDH homologues identified on the linear megaplasmid was expressed using an optimized expression system with the inclusion of N-terminal six-histidine tags. Only small amounts (< 0.1 mg/g of cell paste) of the XecE1 (S-HPCDH1) protein were present in the soluble fraction, with most of the protein being in inclusion bodies. The small amount of soluble protein, when purified, had a specific activity nearly identical to the native enzyme, demonstrating that the enzyme can be expressed in an active form, albeit at levels too low for detailed biochemical and structural characterization. The second copy, XecE2 (S-HPCDH2), contained apparent mutations in the N-terminal region, most notably the lack of the classic GXXGXG cofactor sequence motif that is found within the Rossman fold (10). This protein was expressed with more protein

Kinetic Parameters for rS-HPCDH1, rS-HPCDH3, and rR-HPCDH1 with Physiological Substrates. In the present work, the kinetic analysis of rR-HPCDH1 performed previously (14) was repeated in side by side experiments with the two active XecE homologues so that the three enzymes can be compared directly. Although expressed only at very low levels,

Table 1: Kinetic Parameters for *R*- and *S*-HPCDH with Physiological Substrates in the Forward and Reverse Directions^a

substrate	<i>rR</i> -HPCDH1			<i>rS</i> -HPCDH1			<i>rS</i> -HPCDH3		
	K_m (μ M)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1} s^{-1}$)	K_m (μ M)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1} s^{-1}$)	K_m (μ M)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1} s^{-1}$)
<i>R</i> -HPC	96 \pm 7	48 \pm 1	5.0 $\times 10^5$	5200 \pm 300	2.4 \pm 0.06	4.6 $\times 10^2$	9100 \pm 1100	5.5 \pm 0.2	6.0 $\times 10^2$
<i>S</i> -HPC	220 \pm 70	0.12 \pm 0.01	5.3 $\times 10^2$	220 \pm 20	28 \pm 0.8	1.3 $\times 10^5$	31 \pm 0.9	25 \pm 0.2	7.9 $\times 10^5$
2-KPC	68 \pm 10	29 \pm 2	4.2 $\times 10^5$	240 \pm 10	20 \pm 0.2	8.3 $\times 10^4$	270 \pm 13	11 \pm 0.2	3.9 $\times 10^4$

^aAssay for *R*-HPC and *S*-HPC oxidation by *rR*-HPCDH1 contained 1.0 and 46 μ g of enzyme, respectively. Assay for *R*-HPC and *S*-HPC oxidation by *rS*-HPCDH1 and *rS*-HPCDH3 contained 5.0 and 1.0 μ g of enzyme, respectively. Assay for 2-KPC reduction contained 1 μ g of enzyme. Apparent k_{cat} and K_m values are reported as means \pm standard deviations. All assays were performed in triplicate at 30 °C with fixed concentrations of NAD⁺ (10 mM) or NADH (0.17 mM). Apparent kinetic constants were determined by fitting experimental data to the standard form of the Michaelis–Menten equation.

rS-HPCDH1 was included in these analyses in order to see if the two XecE homologues had any significant differences in catalytic properties. These results are presented in Table 1.

The apparent k_{cat} and K_m values determined for *R*-HPC and *S*-HPC oxidation, and 2-KPC reduction, by *rR*-HPCDH1 are similar to those reported previously (14). With regard to *rS*-HPCDH1 and *rS*-HPCDH3, the apparent k_{cat} values for *S*-HPC oxidation are nearly identical to each other, while the apparent K_m is 7-fold lower for *rS*-HPCDH3. For 2-KPC reduction, *rS*-HPCDH1 exhibits a 2-fold higher k_{cat} but nearly identical K_m relative to *rS*-HPCDH3. As noted above, *S*-HPCDH1 and *S*-HPCDH3 share high identity (74%) but are not identical enzymes, so it is not surprising that some differences in kinetic parameters are observed.

The most surprising results of the kinetic analyses are the large relative differences in k_{cat} and K_m for the *opposite enantiomers* when characterized as substrates for the *R*- and *S*-dehydrogenases. As shown in Table 1, *rR*-HPCDH1 catalyzed the oxidation of *S*-HPC with a k_{cat} that is 402 times less than that for *R*-HPC. In contrast, *rS*-HPCDH1 and *rS*-HPCDH3 catalyzed the oxidation of *R*-HPC with k_{cat} values only 11 and 4.5 times less than that for *S*-HPC but with substantially higher K_m values (24 and 290 times higher) (Table 1).

A comparison of the catalytic efficiencies (k_{cat}/K_m) of the three enzymes for the *R*- and *S*-enantiomers provides additional insights. The catalytic efficiencies for the “natural” enantiomers for each enzyme are in the range of $(1–8) \times 10^5$. By comparison, the catalytic efficiencies for the opposite enantiomers are about 3 orders of magnitude lower. When enantioselectivity (*E*) is defined as the ratio of k_{cat}/K_m for the natural enantiomer to k_{cat}/K_m for the opposite enantiomer, the following values are obtained: $E_{R-HPCDH1} = 944$, $E_{S-HPCDH1} = 283$, and $E_{S-HPCDH3} = 1315$. Thus, the HPCDH enzymes are highly efficient at discriminating between the HPC enantiomers, with enantioselectivity controlled largely by differences in k_{cat} for *R*-HPCDH and predominantly by differences in K_m for the two *S*-HPCDH enzymes.

rS-HPCDH1 and *rS*-HPCDH3 are similar but not identical in terms of their catalytic properties, verifying that the homologues are redundant *S*-HPC-specific enzymes. The subsequent studies of *S*-HPCDH are focused on *rS*-HPCDH3, since only this protein could be expressed in sufficient amounts for the detailed analyses described below.

The apparent K_m values for NAD⁺ and NADH with *R*-HPC and 2-KPC as substrates were determined for *rS*-HPCDH3 and gave the following values: K_m for NAD⁺, 191 \pm 21 μ M; K_m for NADH, 8.42 \pm 1.58 μ M. By comparison, the K_m values for *rR*-HPCDH1 using a complete bisubstrate kinetic analysis were previously reported to be 457 and 36.6 μ M for NAD⁺ and NADH, respectively (15).




FIGURE 3: Changes of kinetic parameters with pH for *rS*-HPCDH3-catalyzed oxidation of *S*-HPC: (A) k_{cat} vs pH, (B) k_{cat}/K_m vs pH, and (C) K_m vs pH, represented in log scale. The plots in (A) and (C) are shown as simple line plots. The line in (B) was generated by a fit of the experimental data to eq 1.

pH Dependence of the Kinetic Parameters for *rS*-HPCDH3. Kinetic parameters were determined for the oxidation of *S*-HPC by *rS*-HPCDH3 at a range of pH values and plotted as $\log k_{cat}$ vs pH, $\log k_{cat}/K_m$ vs pH, and $\log K_m$ vs pH as was done previously for oxidation of *R*-HPC by *rR*-HPCDH1 (15). These results are presented in Figure 3. The overall trends are similar to those reported for *rR*-HPCDH1 but with some important differences (15) suggesting a similar fundamental chemical mechanism for substrate oxidation. The differences in the pH analyses for the two enzymes are highlighted below.

Figure 3A shows a fairly steady increase in $\log k_{cat}$ as the pH increases (slope 0.08), suggesting that isomerization of the enzyme–NAD⁺ complex could be a major rate-determining step, as described for other dehydrogenases (23). The same effect was seen for *rR*-HPCDH1 (15), although the change in k_{cat} was more pronounced (over 6-fold increase vs 3-fold increase for *rS*-HPCDH3).

Linear regression of the plot of $\log k_{cat}/K_m$ vs pH from pH 5.0 to pH 8.0 (Figure 3B) gave a slope of 0.998 with an R^2 value of

Table 2: Summary of Amino Acid Substitutions Made to the Putative *rS*-HPCDH3 Catalytic Residues Using Site-Directed Mutagenesis^a

amino acid	postulated catalytic role(s) ^b	substitution	K_m (μ M <i>S</i> -HPC)	k_{cat} (s^{-1})	k_{cat}/K_m ($M^{-1} s^{-1}$)
S143	H-bond donor to substrate/charge stabilization of the transition state	none S143A	31 ± 0.9 9500 ± 2500	25 ± 0.2 0.013 ± 0.001	7.9×10^5 1.4×10^0
Y156	general acid/base	Y156A Y156F	1800 ± 200 no activity ^c	0.65 ± 0.02	3.5×10^2
K160	lowers pK_a of Y156/coenzyme binding	K160A	no activity ^c		

^aAssays contained 66 μ g of S143A or 20 μ g of Y156A, 5 mM NAD⁺, and variable concentrations of *S*-HPC. ^bPostulated catalytic roles are based on a general trend found in most SDR enzymes (including *R*-HPCDH). ^cNo activity was defined as no measurable change in the absorbance at 340 nm (for NADH) in assays containing 100 μ g of protein, 2.5 mM *S*-HPC, and 20 mM NAD⁺. Apparent k_{cat} and K_m values are reported as means \pm standard errors. All other values are reported as means only. All assays were performed in triplicate at 30 °C with fixed concentrations of NAD⁺. Apparent kinetic constants were determined by fitting experimental data to the standard form of the Michaelis–Menten equation.

0.980, suggesting the importance of a single ionizable residue that must be deprotonated for catalysis. A fit of the data in Figure 3B to eq 1 provides a pK_a value for this residue of 7.9. Based on pH studies of other SDR enzymes, the ionizable residue most likely represents the tyrosine of the catalytic triad, which serves as the general acid/base for catalysis. For comparison, the pK_a value for the corresponding ionizable group of *rR*-HPCDH1 was reported to be a lower value of 6.9 (15), while the prototype alcohol dehydrogenases from *Drosophila melanogaster* and *Drosophila lebanonensis* were reported to be 7.1 and 7.6, respectively (24, 25).

The data presented in Figure 3C show a dramatic decrease in K_m (321-fold decrease from pH 5.0 to pH 9.0) as the pH is increased. Taken together with the above results, this suggests that the deprotonated tyrosine general base plays an important role in both alcohol binding (K_m effect) and catalysis (k_{cat} and k_{cat}/K_m effects). The increase in K_m from pH 9 to pH 10 could reflect deprotonation of another functional group important in coordinating the sulfonate of CoM (K214, as described below). The decrease in K_m for *rS*-HPCDH3 was much greater than that observed for *rR*-HPCDH1, where only a 14-fold decrease in K_m was observed (15). This suggests that the general acid/base of *S*-HPCDH3 is more important for binding of the alcohol substrate in the (*S*)-dehydrogenase than the (*R*)-dehydrogenase, an idea that is supported by the further studies described below.

Site-Directed Mutagenesis of the Catalytic Triad. As shown in Figure 2, the positions of the residues in the primary sequences believed to constitute the catalytic triad are conserved between the *R*- and *S*-HPCDH enzymes. To confirm the importance of the key residues, site-directed mutants were constructed. As shown in Table 2, the Y156F and K160A mutants were completely inactive. The S143A mutant exhibited a small amount of activity but with a catalytic efficiency that was reduced by more than 5 orders of magnitude relative to wild type. Interestingly, the Y156A mutant exhibited a k_{cat} that was 2.6% of the activity of the wild-type enzyme, with a K_m value 60-fold higher than wild type. The observation that the Y156A but not Y156F mutant retains some activity suggests that the removal of the bulky phenyl group allows the enzyme to facilitate hydrogen atom abstraction from the hydroxyl group by another mechanism. An analysis of the primary sequences of the HPCDH enzymes shows that a cysteine residue is next to the tyrosine base in the *S*-enzymes but not the *R*-enzyme (Figure 2). A cysteine is also adjacent to the catalytic triad in the SDR 17 β -hydroxysteroid dehydrogenase, and the Y to A substitution in that enzyme also exhibits a small amount of activity (26). It is conceivable that C157 substitutes as the general base in the Y156A mutant, albeit with a reduced catalytic efficiency,

although this possibility has not been pursued further in this study.

Identification and Site-Directed Mutagenesis of the Sulfonate Binding Residues. For *rR*-HPCDH1, two arginine residues, R152 and R196, were shown to interact with the sulfonate of CoM via ionic interactions (Scheme 1, left side) (14, 16). Site-directed mutagenesis of either of these residues to alanine resulted in substantially reduced k_{cat} values and increased K_m values for 2-KPC reduction. Importantly, the same amino acid substitutions did not significantly alter k_{cat} or K_m when 2-butanone, a nonphysiological substrate lacking the sulfonate, was used as the substrate (14). Thus, the sulfonate-coordinating arginine residues of *R*-HPCDH1 are only required for effective catalysis with the natural substrate.

By analogy, positively charged residues within a pocket are believed to interact with the sulfonate moiety of 2-KPC and *S*-HPC in *S*-HPCDH, but with differential placement relative to the methyl binding pocket such that the hydroxyl group and hydrogen atom of the substrate are oriented properly for catalysis (14, 16). To facilitate the identification of these residues, a homology model was constructed for *S*-HPCDH3 with substrates modeled in using the *R*-HPCDH1 structure with 2-KPC bound as a guide (16).

As shown in Figure 4A, the positioning of the catalytic triad residues, verified for *S*-HPCDH3 by the mutational analyses in Table 2, is conserved between the two *S*-HPCDH enzymes. As predicted, the active sites of the enzymes differ in the spatial orientations of the CoM and methyl groups, with the positions of the hydrogen atom and hydroxyl group being fixed relative to NAD⁺ and the tyrosine general base. Figure 4B shows a different view of the active sites, highlighting the interactions known (for *rR*-HPCDH1) and proposed (for *rS*-HPCDH3) to be important in binding the sulfonate of CoM. While two arginines coordinate the sulfonate within *R*-HPCDH1, the model suggests coordination by an arginine and a lysine for *S*-HPCDH3 (as well as *S*-HPCDH1 as seen in the multiple sequence alignment of Figure 2).

As shown in Table 3, substitution of either R211 or K214 by alanine resulted in substantially reduced catalytic efficiencies for *S*-HPCDH3 with the natural substrates in both the forward (oxidative, *S*-HPC as substrate) and reverse (reductive, 2-KPC as substrate) directions. Interestingly, the most dramatic effects of the mutations were on K_m values, which were 50–74 times higher in the oxidative direction and 41–43 times higher in the reductive direction. There was a significant (40–80%) decrease in k_{cat} due to the mutations for the forward reaction but very little change for the reductive direction. By comparison, 2-KPC reduction by




FIGURE 4: Superimposed active sites of *R*-HPCDH1 and *S*-HPCDH3 based on the crystal structure of *R*-HPCDH1 and a homology model of *S*-HPCDH3. The cartoon structures and carbon atoms of amino acid residues of *R*-HPCDH1 (PDB ID 2cfc) and the homology model for *S*-HPCDH3 are colored gray and green, respectively. NAD^+ is shown in magenta. *R*-HPC and *S*-HPC were modeled using the crystal structure for *S*-HPC bound at the active site of *R*-HPCDH1 as described previously (16). In both views, *R*-HPC (gray carbon atoms) and *S*-HPC (green carbon atoms) are modeled at the active sites such that the positions of the hydroxyl group and hydrogen atom occupy the same positions. The methyl groups and the methylene groups linking the hydroxypropyl groups to CoM are overlaid on top of each other to highlight the different spatial orientations of these groups in *R*- and *S*-HPC. Panel A: Superimposed structures highlighting the interactions of substrates with the catalytic triads. Panel B: Superimposed structures highlighting the interactions of substrates with the amino acids that coordinate the sulfonate of CoM.

Table 3: Kinetic Parameters for Wild-Type *rS*-HPCDH3 and R211A and K214A with Various Substrates ^a

enzyme	K_m (mM)	change in K_m (x-fold)	k_{cat} (s^{-1})	change in k_{cat} (x-fold)	k_{cat}/K_m ($M^{-1} s^{-1}$)
Substrate: <i>S</i> -HPC					
wild type	0.031 ± 0.001	1.0	25 ± 0.2	1.0	7.9×10^5
R211A	1.6 ± 0.1	49	16 ± 0.4	0.64	1.0×10^4
K214A	2.3 ± 0.1	73	5.8 ± 0.1	0.23	2.5×10^3
Substrate: 2-KPC					
wild type	0.28 ± 0.01	1.0	11 ± 0.2	1.0	3.9×10^4
R211A	11 ± 0.4	41	8.6 ± 0.2	0.80	7.5×10^2
K214A	12 ± 0.9	43	9.6 ± 0.3	0.89	8.1×10^2
Substrate: 2-Butanone					
wild type	120 ± 5	1.0	0.044 ± 0.001	1.0	3.7×10^2
R211A	180 ± 30	1.5	0.072 ± 0.05	1.6	4.0×10^2
K214A	72 ± 6	0.61	0.039 ± 0.001	0.90	5.4×10^2
Substrate: 2-Propanol					
wild type	1400 ± 60	1.0	2.0 ± 0.5	1.0	1.4×10^3
R211A	720 ± 40	0.51	1.7 ± 0.03	0.85	2.4×10^3
K214A	950 ± 20	0.67	1.7 ± 0.2	0.86	1.8×10^3

^aAssay for *S*-HPC oxidation contained 0.20 μg of wild-type *rS*-HPCDH3 or 4.0 μg of R211A and K214A while assay for 2-KPC reduction contained 0.20 μg of wild-type *rS*-HPCDH3, 5.0 μg of R211A, and 2.0 μg of K214A. Assay for 2-butanone reduction contained 197 μg of enzyme while assay for 2-propanol oxidation contained 8.0 μg of enzyme. Apparent k_{cat} and K_m values are reported as means \pm standard deviations. All other values are reported as means only. All assays were performed in triplicate at 30 °C with fixed concentrations of NAD^+ (10 mM) or NADH (0.17 mM). Apparent kinetic constants were determined by fitting experimental data to the standard form of the Michaelis–Menten equation.

the R152A and R196A mutants of *R*-HPCDH1 was impaired sizably in both K_m and k_{cat} values (14).

The k_{cat} and K_m values measured for the reduction of the aliphatic ketone 2-butanone and the oxidation of the aliphatic alcohol 2-propanol were largely unaffected by the amino acid changes in *S*-HPCDH3 (Table 3). These aliphatic substrates had dramatically higher K_m values and lower k_{cat} values relative to the natural substrates. The fact that the kinetic parameters did not change substantially in the site-directed mutants demonstrates that the sulfonate-binding residues are not important for aliphatic substrates where no favorable interaction with the positively charged residues would occur.

Similar results were obtained when 2-butanone was analyzed as a substrate for wild-type *R*-HPCDH1 and the R152A and R196A mutants (14).

S-HPCDH3 Has an Inherent Stereoselectivity for 2-Butanone Reduction Not Present in *R*-HPCDH1. As shown in Table 4, *S*-HPCDH3 has a very high inherent stereoselectivity, producing (*S*)-2-butanol with a 98.4% enantiomeric excess (ee). These results are dramatically different than those observed for *R*-HPCDH1, where (*S*)-2-butanol is also produced in higher amounts than (*R*)-2-butanol, but in only a 44% enantiomeric excess (Table 4 and ref 14). Thus, even though the *R*- and *S*-HPCDH enzymes are highly specific for production

Table 4: Enantioselectivity of 2-Butanone Reduction by Wild-Type *rS*-HPCDH3, *rS*-HPCDH3 Mutants, and *rR*-HPCDH1^a

enzyme	no additions			+1 mM ethanesulfonate		
	% (<i>S</i>)-2-butanol	% (<i>R</i>)-2-butanol	ee (<i>S</i>)-2-butanol	% (<i>S</i>)-2-butanol	% (<i>R</i>)-2-butanol	ee (<i>S</i>)-2-butanol
<i>rS</i> -HPCDH3	99.2 ± 0.2	0.82 ± 0.15	98.36	98.0 ± 0.08	1.97 ± 0.08	96.06
<i>rS</i> -HPCDH3 R211A	99.4 ± 0.2	0.58 ± 0.22	98.84	99.4 ± 0.2	0.64 ± 0.15	98.72
<i>rS</i> -HPCDH3 K214A	91.6 ± 0.3	8.4 ± 0.31	83.18	90.9 ± 0.2	9.12 ± 0.20	81.76
<i>rR</i> -HPCDH1	71.9 ± 1.3	28.0 ± 1.3	43.88	92.7 ± 0.1	7.33 ± 0.05	85.34

^aAll assays were performed in triplicate at 30 °C using 0.64 mg of enzyme, 15 mM NADH, and 56 mM 2-butanone. Percent yields are reported as means ± standard deviations.

Table 5: Kinetic Parameters for *rS*-HPCDH3- and *rR*-HPCDH1-Catalyzed Oxidation of 2-Butanol^a

substrate	K_m (mM)	V_{max} (units/mg)	k_{cat} (s ⁻¹)	k_{cat}/K_m (M ⁻¹ s ⁻¹)	change in k_{cat} (x-fold)	change in K_m (x-fold)	enantioselectivity (<i>E</i>)
<i>rS</i> -HPCDH3							
(<i>R</i>)-2-butanol	68 ± 3	2.2 ± 0.03	1.0	14.8	1.0	1.0	0.16
(<i>S</i>)-2-butanol	28 ± 2	5.8 ± 0.07	2.6	92.8	2.6	0.4	6.3
<i>rR</i> -HPCDH1							
(<i>R</i>)-2-butanol	220 ± 10	3.77 ± 0.06	1.9	8.69	1.0	1.0	0.33
(<i>S</i>)-2-butanol	350 ± 50	18.8 ± 0.9	9.3	26.4	5.0	1.6	3.0

^aAll assays were performed in triplicate at 30 °C with fixed concentrations of NAD⁺ (10 mM). Assays of 2-butanone oxidation catalyzed by *rS*-HPCDH3 and *rR*-HPCDH1 contained 25 and 21 μg of enzyme, respectively. Apparent kinetic constants were determined by fitting experimental data to the standard form of the Michaelis–Menten equation. Apparent V_{max} and K_m values are reported as means ± standard deviations. All other values are reported as means only. Enantioselectivity was defined as $(k_{cat}/K_m)_{R\text{-enantiomer}}/(k_{cat}/K_m)_{S\text{-enantiomer}}$ for (*R*)-2-butanone oxidation and as $(k_{cat}/K_m)_{S\text{-enantiomer}}/(k_{cat}/K_m)_{R\text{-enantiomer}}$ for (*S*)-2-butanone oxidation.

of *R*-HPC and *S*-HPC from 2-KPC reduction, they both form excesses of (*S*)-2-butanol from 2-butanone.


An interesting feature of the *R*-HPCDH1 is that the enantioselectivity of 2-butanone reduction can be “modulated” to produce higher amounts of (*S*)-2-butanol by including CoM or alkylsulfonates in the assay (14). These “enantioselective modulators” modulated stereospecificity in a saturable fashion, with a theoretical yield of 100% (*S*)-2-butanol at the saturation points (14). The effects of these enantioselective modulators were abolished in mutants in the sulfonate-coordinating arginines (R152A and R196A) (14). These results were interpreted as follows: the alkylsulfonates bind in the CoM binding pocket, constraining 2-butanone to bind in the active site with the methyl group rather than the ethyl group oriented toward the bound alkylsulfonate to prevent steric clashes. In this orientation, the sp² hybridized carbonyl of 2-butanone is necessarily constrained for hydride transfer to the plane that produces (*S*)-2-butanol (14). As shown in Table 4, the addition of ethanesulfonate to wild-type *rS*-HPCDH3 and the R211A and K214A mutants had very little effect on the stereochemical outcome of 2-butanone reduction relative to the effect observed with *rR*-HPCDH1, demonstrating that the modulating effect is a phenomenon only seen with *R*-HPCDH1.

Dehydrogenation of 2-Butanol Enantiomers by *rR*- and *rS*-HPCDH Enzymes. The studies of 2-butanone reduction were expanded to examine the kinetic parameters for the reverse reaction, i.e., (*R*)- and (*S*)-2-butanone oxidation to 2-butanone (Table 5). Both enzymes exhibited a preference for (*S*)-2-butanol as the substrate, with enantioselectivity values of 6.23 and 3.04 for *rS*-HPCDH3 and *rR*-HPCDH1, respectively. The K_m values for 2-butanone reduction were about 1000 times higher than for 2-KPC.

*2-(2-Methyl-2-hydroxypropylthio)ethanesulfonate Is a Competitive Inhibitor of Natural Substrate Oxidation by *rR*-HPCDH1 but Not *rS*-HPCDH3.* As shown in Table 1, *rR*-HPCDH1 exhibits a K_m for *S*-HPC in the same range as *R*-HPC, while *rS*-HPCDH3 exhibits a K_m for *R*-HPC that is nearly 300 times higher than for *S*-HPC. These results indicate that *rR*-HPCDH1 is able to bind either enantiomer with high affinity, although for the improper enantiomer, the hydroxyl and hydrogen of the substrate are misaligned, resulting in lower turnover (Table 1). In contrast, the high K_m value for *R*-HPC with *rS*-HPCDH3 suggests that the other enantiomer does not bind well in the first place, possibly due to steric constraints when the methyl group is misaligned. As shown in Chart 1, 2-(2-methyl-2-hydroxypropylthio)ethanesulfonate (M-HPC) is an achiral analogue of both *R*-HPC and *S*-HPC where the hydrogen atoms of each alcohol are replaced by a methyl group, resulting in a tertiary alcohol that cannot undergo oxidation. M-HPC is thus an equivalent mimic of both HPC enantiomers and can be studied as a possible inhibitor of both enzymes to shed light on the K_m differences for the opposite enantiomers discussed above.

As shown in Figure 5A, M-HPC was a competitive inhibitor of *R*-HPC oxidation by *R*-HPCDH1, with a K_{is} of 290 ± 14 μM. This value is in the range of the K_m for the natural substrate *R*-HPC (96 μM) and the opposite enantiomer *S*-HPC (224 μM). In marked contrast to these results, no detectable inhibition was observed for M-HPC vs the natural substrate *S*-HPC with *S*-HPCDH3 (Figure 5B). This result correlates with the observation that *R*-HPC has a dramatically higher K_m for *S*-HPCDH3 relative to the natural substrate (9100 vs 31 μM). This experiment was repeated with *rS*-HPCDH1, and the same result was obtained: no detectable inhibition was seen with M-HPC (Figure 5C). Thus, the addition of the methyl group in the improper position relative to the CoM and hydroxyl groups has

Chart 1



very different effects on binding affinity for the *R*- and *S*-specific dehydrogenases.

2-(2-Hydroxyethylthio)ethanesulfonate (HEC) Is a Substrate for both the *R*- and *S*-HPCDH Enzymes with Identical K_m Values. HEC is an achiral mimic of both *R*-HPC and *S*-HPC in which the methyl group is replaced by a hydrogen (see Chart 1). HEC was found to be a substrate for both *rR*-HPCDH1 ($k_{\text{cat}} = 0.55 \pm 0.02 \mu\text{M}$, $K_m = 960 \pm 100 \mu\text{M}$) and *rS*-HPCDH3 ($k_{\text{cat}} = 3.8 \pm 0.3 \mu\text{M}$, $K_m = 970 \pm 250 \mu\text{M}$). Thus, the loss of the methyl groups imparting chirality resulted in an enzyme substrate with identical (within experimental error) K_m values for both enzymes, suggesting that for this substrate the CoM moiety and hydroxyl group have become equal determinants in binding affinity. As a final experiment, M-HPC was investigated as an inhibitor of HEC oxidation to see if the result of Figure 5 would hold with the achiral substrate. M-HPC was a competitive inhibitor of HEC oxidation by *R*-HPCDH1 with a K_{is} of $110 \pm 10 \mu\text{M}$ but was not an inhibitor of HEC oxidation by *S*-HPCDH3.

Different Strategies Control Enantioselectivity in the R- and S-HPCDH Enzymes. The results presented above are interpreted as follows: *R*-HPCDH1 can bind either enantiomer of HPC with the CoM moiety oriented properly in the sulfonate-binding pocket consisting of R152 and R196 (Figure 4), but *S*-HPCDH3 is unable to bind *R*-HPC in this fashion, presumably due to steric clashes imposed by the presence of the misaligned methyl group on the C2 atom. Both enzymes bind HEC with identical affinities since no methyl group is present. Thus, a high-affinity ternary complex of *S*-HPC, NAD^+ , and *R*-HPCDH forms, but the misalignment of the hydrogen and hydroxyl groups




FIGURE 5: Effects of 2-(2-methyl-2-hydroxypropylthio)ethanesulfonate (M-HPC) on *R*- and *S*-HPC oxidation by *R*-HPCDH1, *S*-HPCDH3, and *S*-HPCDH1. Panel A: Competitive inhibition of *R*-HPCDH1-catalyzed *R*-HPC oxidation by M-HPC. The double reciprocal plots for assays performed in the presence of different concentrations of M-HPC are shown in the main diagram. Data points represent the average of triplicate experiments. The solid lines were generated by nonlinear least-squares fits of the v vs S data, shown in the inset, to the equation for a rectangular hyperbola using SigmaPlot. M-HPC concentrations: (●) 0 mM, (○) 0.2 mM, (▼) 0.4 mM, (Δ) 0.8 mM, and (■) 1.6 mM. Panels B and C: v vs S plots for *S*-HPC oxidation by *S*-HPCDH3 and *S*-HPCDH1, respectively, in the presence of different concentrations of M-HPC. The lines were generated by fitting the data to the standard form of the Michaelis–Menten equation. M-HPC concentrations: (●) 0 mM, (○) 1.2 mM, and (▼) 4.9 mM.

on C2 relative to NAD^+ and the tyrosine general base results in a 403-fold lower turnover rate.

In contrast to *R*-HPCDH1, *S*-HPCDH3 apparently cannot bind *S*-HPC analogues (*R*-HPC and M-HPC) with CoM oriented properly in the sulfonate-binding pocket that consists of R211 and K214. The most logical explanation for this is that the misaligned methyl groups on C2 required for this high-affinity binding are not accommodated due to steric clashes with amino acid side chain(s). Thus, *R*-HPC binds to *S*-HPCDH3 with a 290-fold lower affinity but in an orientation where the hydroxyl and hydrogen on C2 can be more properly aligned with tyrosine 156 and NAD^+ , such that k_{cat} only decreases by 4.5-fold relative to the natural substrate *S*-HPC.

The results of the pH studies further indicate that the hydroxyl group at C2 is a larger determinant in the binding of *S*-HPC to *S*-HPCDH3 than for binding of *R*-HPC to *R*-HPCDH1. Note that the K_m for the natural substrate is 3-fold lower for *S*-HPCDH3 than for *R*-HPCDH1, which could be due to higher affinity binding of the hydroxyl group. However, the K_m values for HEC, lacking the methyl group, are identical for both enzymes. Thus, the methyl group of *S*-HPC may facilitate

the alignment of the hydroxyl group in this higher affinity position.

Physiological Implications of These Studies. The differential control of enantioselectivity in the *R*- and *S*-HPCDH enzymes may have evolved to reflect the roles these enzymes play in propylene metabolism. As shown in Figure 1, alkene monooxygenase from *X. autotrophicus* produces a mixture of (*R*)- and (*S*)-epoxypropene, although (*R*)-epoxypropene is produced in a 90% enantiomeric excess (4). Both (*R*)- and (*S*)-epoxypropene are substrates for epoxyalkane:CoM transferase, which results in the production of *R*- and *S*-HPC, respectively. Although (*S*)-epoxypropene (and by extrapolation, *S*-HPC) are the minor products of the metabolic pathway, it is still essential for these compounds to be further metabolized by the bacterium, since epoxides are highly toxic. While epoxyalkane:CoM transferase detoxifies (*S*)-epoxypropene by conversion to *S*-HPC, *S*-HPC must proceed through the pathway of epoxide carboxylation to regenerate free CoM; if it did not, the CoM pool would be wastefully sequestered. *R*-HPCDH1 will still operate efficiently even with a similar K_m value for the opposite enantiomer, since *S*-HPC is predicted to be produced at about 20-fold lower concentrations than *R*-HPC. In contrast, in order for *S*-HPCDH to be efficient at the lower concentrations of *S*-HPC encountered in the cell, it needs to have a lower K_m for the natural substrate and substantially higher K_m for the opposite enantiomer to work efficiently. To summarize, the different amounts of *R*- and *S*-HPC that accumulate in the cells due to the inherent stereoselectivity of the alkene monooxygenase are proposed to have led to the evolution of different strategies for controlling efficient substrate flux through the pathway. *R*-HPC oxidation is controlled by k_{cat} , since *R*-HPCDH does not need to discriminate substrate binding at the much lower concentrations of the opposite enantiomer present, while control of *S*-HPC oxidation is controlled by K_m , since discrimination of substrate binding is crucial with the higher concentrations of the inhibitory enantiomer present. In this context it should be noted that *R*- and *S*-HPCDH were found to have comparable specific activities in cell extracts of *X. autotrophicus* when grown on propylene, with each predicted to account for about 1% of soluble cell protein (4–6). This observation reiterates that substrate flux is controlled by differences in mechanisms of these enzymes and not by differences in levels of expression.

The fact that *X. autotrophicus* has redundant copies of the dehydrogenases (as well as the other enzymes of epoxide metabolism) further highlights the importance of these enzymes to the bacterium. The results of Table 1 suggest some differences in the kinetic properties for two of the redundant *S*-enzymes. The copies we purified from *X. autotrophicus* in our previous studies (4–6, 18, 27) consisted primarily (or wholly) of the enzymes in the first operon based on yields from the purification schemes and their biochemical properties. Now that we have a greater understanding of the organization of the epoxide carboxylation genes, we can apply molecular genetics to determine how important the individual copies are to bacterial growth and survival under different conditions.

ACKNOWLEDGMENT

We thank Dr. Michael Kay and Matthew Weinstock, University of Utah School of Medicine, for technical assistance with the CD spectrometer.

REFERENCES

1. Kroutil, W., Mang, H., Edegger, K., and Faber, K. (2004) Recent advances in the biocatalytic reduction of ketones and oxidation of sec-alcohols. *Curr. Opin. Chem. Biol.* 8, 120–126.
2. Nakamura, K., Yamanaka, R., Matsuda, T., and Harada, T. (2003) Recent developments in asymmetric reduction of ketones with biocatalysts. *Tetrahedron: Asymmetry* 14, 2659–2681.
3. Goldberg, K., Schroer, K., Lütz, S., and Liese, A. (2007) Biocatalytic ketone reduction—a powerful tool for the production of chiral alcohols—part I: processes with isolated enzymes. *Appl. Microbiol. Biotechnol.* 76, 237–248.
4. Allen, J. R., Clark, D. D., Krum, J. G., and Ensign, S. A. (1999) A role for coenzyme M (2-mercaptoethanesulfonic acid) in a bacterial pathway of aliphatic epoxide carboxylation. *Proc. Natl. Acad. Sci. U.S.A.* 96, 8432–8437.
5. Allen, J. R., and Ensign, S. A. (1997) Purification to homogeneity and reconstitution of the individual components of the epoxide carboxylase multiprotein enzyme complex from *Xanthobacter* strain Py2. *J. Biol. Chem.* 272, 32121–32128.
6. Allen, J. R., and Ensign, S. A. (1999) Two short-chain dehydrogenases confer stereoselectivity for enantiomers of epoxypropene in the multiprotein epoxide carboxylating systems of *Xanthobacter* strain Py2 and *Nocardia corallina* B276. *Biochemistry* 38, 247–256.
7. Ensign, S. A. (2001) Microbial metabolism of aliphatic alkenes. *Biochemistry* 40, 5845–5853.
8. Ensign, S. A., and Allen, J. R. (2003) Aliphatic epoxide carboxylation. *Annu. Rev. Biochem.* 72, 55–76.
9. Jörnvall, H., Persson, B., Krook, M., Atrian, S., González-Duarte, R., Jeffery, J., and Ghosh, D. (1995) Short-chain dehydrogenases/reductases. *Biochemistry* 34, 6003–6013.
10. Kavanagh, K. L., Jörnvall, H., Persson, B., and Oppermann, U. (2008) The SDR superfamily: functional and structural diversity within a family of metabolic and regulatory enzymes. *Cell. Mol. Life Sci.* 65, 3895–3906.
11. Oppermann, U., Filling, C., Hult, M., Shafqat, N., Wu, X., Lindh, M., Shafqat, J., Nordling, E., Kallberg, Y., Persson, B., and Jörnvall, H. (2003) Short-chain dehydrogenases/reductases (SDR): the 2002 update. *Chem.-Biol. Interact.* 143–144, 247–253.
12. Persson, B., Kallberg, Y., Bray, J. E., Bruford, E., Dellaporta, S. L., Favia, A. D., Duarte, R. G., Jörnvall, H., Kavanagh, K. L., Kedishvili, N., Kisiela, M., Maserk, E., Mindnich, R., Orchard, S., Penning, T. M., Thornton, J. M., Adamski, J., and Oppermann, U. (2009) The SDR (short-chain dehydrogenase/reductase and related enzymes) nomenclature initiative. *Chem.-Biol. Interact.* 178, 94–98.
13. Persson, B., Krook, M., and Jörnvall, H. (1991) Characteristics of short-chain alcohol dehydrogenases and related enzymes. *Eur. J. Biochem.* 200, 537–543.
14. Clark, D. D., Boyd, J. M., and Ensign, S. A. (2004) The stereoselectivity and catalytic properties of *Xanthobacter autotrophicus* 2-[(*R*)-2-hydroxypropylthio]ethanesulfonate dehydrogenase are controlled by interactions between C-terminal arginine residues and the sulfonate of coenzyme M. *Biochemistry* 43, 6763–6771.
15. Clark, D. D., and Ensign, S. A. (2002) Characterization of the 2-[(*R*)-2-hydroxypropylthio]ethane sulfonate dehydrogenase from *Xanthobacter* strain Py2: product inhibition, pH dependence of kinetic parameters, site-directed mutagenesis, rapid equilibrium inhibition, and chemical modification. *Biochemistry* 41, 2727–2740.
16. Krishnakumar, A. M., Nocek, B. P., Clark, D. D., Ensign, S. A., and Peters, J. W. (2006) Structural basis for stereoselectivity in the (*R*)- and (*S*)-hydroxypropylthioethanesulfonate dehydrogenases. *Biochemistry* 45, 8831–8840.
17. Krishnakumar, A. M., Sliwa, D., Endrizzi, J. A., Boyd, E. S., Ensign, S. A., and Peters, J. W. (2008) Getting a handle on the role of coenzyme M in alkene metabolism. *Microbiol. Mol. Biol. Rev.* 72, 445.
18. Allen, J. R., and Ensign, S. A. (1997) Characterization of three protein components required for functional reconstitution of the epoxide carboxylase multienzyme complex from *Xanthobacter* strain Py2. *J. Bacteriol.* 179, 3110–3115.
19. Laemmli, U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
20. Cook, P. F., and Cleland, W. W. (2007) Enzyme kinetics and mechanism, Garland Science, New York.
21. Krum, J. G., and Ensign, S. A. (2001) Evidence that a linear megaplasmid encodes enzymes of aliphatic alkene and epoxide metabolism and coenzyme M (2-mercaptoethanesulfonate) biosynthesis in *Xanthobacter* strain Py2. *J. Bacteriol.* 183, 2172–2177.
22. Swaving, J., Weijers, C. A., van Ooyen, A. J., and de Bont, J. A. M. (1995) Complementation of *Xanthobacter* Py2 mutants defective

- in epoxyalkane degradation, and expression and nucleotide sequence of the complementing DNA fragment. *Microbiology* 141, 477–484.
23. Grimshaw, C. E., Cook, P. F., and Cleland, W. W. (1981) Use of isotope effects and pH studies to determine the chemical mechanism of *Bacillus subtilis* L-alanine dehydrogenase. *Biochemistry* 20, 5655–5661.
24. Winberg, J. O., Brendskag, M. K., Sylte, I., Lindstad, R. I., and McKinley-McKee, J. S. (1999) The catalytic triad in *Drosophila* alcohol dehydrogenase: pH, temperature and molecular modelling studies. *J. Mol. Biol.* 294, 601–616.
25. Brendskag, M. K., McKinley-McKee, J. S., and Winberg, J. O. (1999) *Drosophila lebanonensis* alcohol dehydrogenase: pH dependence of the kinetic coefficients. *Biochim. Biophys. Acta* 1431, 74–86.
26. Puranen, T. J., Poutanen, M. H., Peltoketo, H. E., Vihko, P. T., and Vihko, R. K. (1994) Site-directed mutagenesis of the putative active site of human 17 β -hydroxysteroid dehydrogenase type. *Biochem. J.* 304, 289–293.
27. Clark, D. D., Allen, J. R., and Ensign, S. A. (2000) Characterization of five catalytic activities associated with the NADPH:2-ketopropyl-coenzyme M [2-(2-ketopropylthio)ethanesulfonate] oxidoreductase/carboxylase of the *Xanthobacter* strain Py2 epoxide carboxylase. *Biochemistry* 39, 1294–1304.