


Correlation of Thermal Stability and Structural Distortion of DNA Interstrand Cross-Links Produced from Oxidized Abasic Sites with Their Selective Formation and Repair

Souradyuti Ghosh and Marc M. Greenberg*

Department of Chemistry, Johns Hopkins University, 3400 North Charles Street, Baltimore, Maryland 21218, United States

S Supporting Information

ABSTRACT: C4'-oxidized (C4-AP) and C5'-oxidized abasic sites (DOB) that are produced following abstraction of a hydrogen atom from the DNA backbone reversibly form cross-links selectively with dA opposite a 3'-adjacent nucleotide, despite the comparable proximity of an opposing dA. A previous report on UvrABC incision of DNA substrates containing stabilized analogues of the ICLs derived from C4-AP and DOB also indicated that the latter is repaired more readily by nucleotide excision repair [Ghosh, S., and Greenberg, M. M. (2014) *Biochemistry* 53, 5958–5965]. The source for selective cross-link formation was probed by comparing the reactivity of ICL analogues of C4-AP and DOB that mimic the preferred and disfavored cross-links with that of reagents that indirectly detect distortion by reacting with the nucleobases. The disfavored C4-AP and DOB analogues were each more reactive than the corresponding preferred cross-link substrates, suggesting that the latter are more stable, which is consistent with selective ICL formation. In addition, the preferred DOB analogue is more reactive than the respective C4-AP ICL, which is consistent with its more efficient incision by UvrABC. The conclusions drawn from the chemical probing experiments are corroborated by UV melting studies. The preferred ICLs exhibit melting temperatures higher than those of the corresponding disfavored isomers. These studies suggest that oxidized abasic sites form reversible interstrand cross-links with dA opposite the 3'-adjacent thymidine because these products are more stable and the thermodynamic preference is reflected in the transition states for their formation.



DNA–DNA interstrand cross-links comprise an important family of DNA damage whose members are absolute blocks of replication and transcription.^{1,2} ICLs are typically produced by exogenous bis-electrophiles that react in the minor or major grooves with staggered nucleobases on opposite strands.^{3–6} Recently, ICLs that result from reaction of various abasic lesions (Figure 1) with dA, dC, and/or dG on the





Figure 1. Abasic site structures.

complementary strand have been characterized.^{7–10} Such cross-links emanate from AP, as well as the oxidized abasic lesions, C4-AP and DOB, that are in equilibrium with ring-opened, electrophilic aldehyde forms (Figure 1 and Scheme 1). C4-AP yields two types of ICLs, one in which both strands are intact (Scheme 1) and one in which the oxidized abasic site has undergone strand scission.^{11–14} The former C4-AP ICL and

Scheme 1. ICLs and Stabilized ICLs from C4-AP and DOB




that produced from DOB are formed reversibly and selectively with the dA opposite the 3'-adjacent thymidine (Scheme 2).¹⁵ Chemically stable analogues [X–Y and W–V (Scheme 1)] that facilitate the characterization of these reversibly formed ICLs have been prepared.¹⁶ ICLs (and analogues) arising from

Received: July 30, 2015

Revised: September 16, 2015

Published: October 1, 2015

Scheme 2. ICL Positioning from C4-AP and DOB



oxidized abasic sites (C4-AP and DOB) are substrates for bacterial nucleotide excision repair, which transforms some of these molecules into double-strand breaks.^{17,18} The chemically stabilized C4-AP ICL analogue is a poorer substrate than that resulting from DOB, but the structural basis for this difference is unknown.¹⁸ Experiments designed to probe the source for the differences in NER efficiency and selective formation of ICLs between C4-AP and DOB and the dA opposite the 3'-adjacent thymidine are described herein.

The right-handed helical nature of duplex DNA provides a rationalization for reaction with the N6-amino group of a dA opposite the 3'-adjacent thymidine compared to the same nucleotide opposite a 5'-thymidine because of the proximity of the C1-aldehyde to the major groove. However, molecular models do not explain why DOB and C4-AP do not readily form ICLs with an opposing nucleotide. Models suggest that the distances between the C1-aldehyde of DOB (see the Supporting Information) or C4-AP (Figure 2) and the N6-amino group of an opposing dA are comparable (within 0.4 Å) to those of the same functionality on a dA opposite the 3'-adjacent thymidine. We considered the possibility that an ICL between C4-AP or DOB and an opposing dA was less stable




Figure 2. Molecular model of C4-AP in a 5'-d(TXT)/(AAA) sequence (X = C4-AP).

than one resulting from reaction of an oxidized abasic site with a dA opposite the 3'-adjacent thymidine and that this would be reflected in the kinetics and equilibria of the reversibly formed ICLs. We hypothesized that the structures of the chemically stabilized ICL analogues would correlate with the actual products produced from DOB and C4-AP (Scheme 1), and that the unobserved, putatively less stable molecules consisting of cross-links with opposing dA's would experience greater base pairing disruption. Relative perturbations of duplex structure may also correlate with ICL susceptibility to NER. Although the mechanism for NER recognition of DNA damage is still under investigation, the proteins are believed to recognize macroscopic duplex distortion.^{6,19–23} Greater distortion of DOB ICL analogues than of those derived from C4-AP would correlate with comparisons of NER.¹⁸ Base pairing disruptions in the ICLs should translate into greater reactivity with chemical agents (e.g., dimethyl sulfate, diethyl pyrocarbonate, potassium permanganate, and hydroxyl radical) that probe nucleobase accessibility, as well as lower UV melting temperatures.^{24–35}

MATERIALS AND METHODS

Materials and General Methods. Oligonucleotides were prepared on an Applied Biosystems Inc. 394 DNA synthesizer. Commercially available DNA synthesis reagents were obtained from Glen Research Inc. DNA substrates used in this study are presented in Figure 3. T4 polynucleotide kinase, Klenow exo[−], terminal deoxynucleotide transferase, and T4 DNA ligase were obtained from New England Biolabs. [α -³²P]dCTP, [α -³²P]-dTTP, [γ -³²P]ATP, and [α -³²P]cordycepin 5'-triphosphate were purchased from PerkinElmer. Analyses of radiolabeled oligonucleotides were conducted using a Storm 840 Phosphor-imager and ImageQuant TL software. C18-Sep-Pak cartridges were obtained from Waters.

General Procedure for the Hydroxyl Radical Cleavage Reaction of 1–3 and 8–10. The appropriately ³²P-labeled substrate (5000–15000 cpm) was suspended in 8 μ L of water and mixed with 10 μ L of 2 \times oxidation buffer [200 mM NaCl, 20 mM sodium phosphate (pH 7.2), 2 mM sodium ascorbate, and 0.5 mM hydrogen peroxide]. An Fe-EDTA solution [2 μ L, 1 mM EDTA and 0.5 mM Fe(NH₄)₂(SO₄)₂·6H₂O] was added, and the reaction was allowed to proceed for 3–5 min. The reaction was quenched by addition of 100 mM thiourea (10 μ M) and the sample precipitated with 3 M NaOAc (4 μ L), 1 μ g/ μ L calf thymus DNA (5 μ L), and cold ethanol (115 μ L). After centrifugation, removal of the supernatant, and drying under vacuum, the DNA was suspended in formamide loading buffer [10 μ L, 50% formamide, 5 mM EDTA (pH 8.0), 0.1% xylene cyanol, and 0.1% bromophenol blue], heated (90 °C, 2 min), cooled (0 °C, 5 min), and analyzed by polyacrylamide gel electrophoresis (PAGE) (20% denaturing) using a Storm 840 phosphorimager and Imagequant TL software.

General Procedure for DEPC Treatment of 1–3 and 8–12. The appropriately ³²P-labeled substrate (1–3) (5000–15000 cpm) was reacted (30 μ L) with 1.04 M DEPC in PBS buffer [10 mM sodium phosphate (pH 7.2) and 100 mM NaCl] for 10 min at room temperature. The reaction tubes were briefly vortexed every 2 min during the reaction. Following incubation with DEPC, the samples were precipitated with 5 M NH₄OAc (5.2 μ L), 1 μ g/ μ L calf thymus DNA (5 μ L), water (10 μ L), and cold ethanol (155 μ L). The precipitated samples were treated with 1 M piperidine (50 μ L) for 20 min at 90 °C and dried in vacuo. The samples were




Figure 3. Cross-linked DNA substrates used in this study.

resuspended in water (50 μ L) and again dried in vacuo. The DNA was resuspended in formamide loading buffer [10 μ L, 50% formamide, 5 mM EDTA (pH 8.0), 0.1% xylene cyanol, and 0.1% bromophenol blue], heated (90 $^{\circ}$ C, 2 min), cooled (0 $^{\circ}$ C, 5 min), and analyzed by PAGE (20% denaturing) using a Storm 840 phosphorimager and ImageQuant TL software. Reactions with substrates 8–12 were conducted as described above except the concentration of DEPC was 1.75 M and the total volume of the reaction was 35.5 μ L.

General Procedure for KMnO_4 Treatment of 1–3 and 8–12. The appropriately ^{32}P -labeled substrate (1–3) (5000–15000 cpm) was reacted (20 μ L) with 0.3 mM KMnO_4 in PBS buffer [10 mM sodium phosphate (pH 7.2) and 100 mM NaCl] for 10 min at room temperature. The reaction tubes were briefly vortexed every 2 min during the reaction. Following incubation with KMnO_4 , the reactions were quenched by the addition of allyl alcohol (1 μ L) and vortexing. The samples were precipitated with 5 M NH_4OAc (5.2 μ L), 1 $\mu\text{g}/\mu\text{L}$ calf thymus DNA (5 μ L), water (20.5 μ L), and cold ethanol (155 μ L). The precipitated samples were treated with 1 M piperidine (50 μ L) for 20 min at 90 $^{\circ}$ C and dried in vacuo. The samples were resuspended in water (50 μ L) and again dried in vacuo. The samples were resuspended in formamide loading buffer [10 μ L, 50% formamide, 5 mM EDTA (pH 8.0),

0.1% xylene cyanol, and 0.1% bromophenol blue], heated (90 $^{\circ}$ C, 2 min), cooled (0 $^{\circ}$ C, 5 min), and analyzed by PAGE (20% denaturing) using a Storm 840 phosphorimager and ImageQuant TL software. Reactions with substrates 8–12 were conducted as described above except the concentration of KMnO_4 was 0.6 mM.

General Procedure for DMS Treatment of 1–3. The appropriately ^{32}P -labeled substrate (1–3) (5000–15000 cpm) was reacted (20 μ L) with 25 mM DMS (diluted from a 250 mM DMS solution in ethanol) in PBS buffer [10 mM sodium phosphate (pH 7.2) and 100 mM NaCl] for 12 min at room temperature. The reaction tubes were briefly vortexed every 2 min during the reaction. Following incubation, the samples were precipitated with 5 M NH_4OAc (5.2 μ L), 1 $\mu\text{g}/\mu\text{L}$ calf thymus DNA (5 μ L), water (20.5 μ L), and cold ethanol (155 μ L). The precipitated samples were treated with 1 M piperidine (50 μ L) for 20 min at 90 $^{\circ}$ C and dried in vacuo. The samples were resuspended in water (50 μ L) and again dried in vacuo. The samples were resuspended in formamide loading buffer [10 μ L, 50% formamide, 5 mM EDTA (pH 8.0), 0.1% xylene cyanol, and 0.1% bromophenol blue], heated (90 $^{\circ}$ C, 2 min), cooled (0 $^{\circ}$ C, 5 min), and analyzed by PAGE (20% denaturing) using a Storm 840 phosphorimager and ImageQuant TL software.




Figure 4. Chemical reactivity of DOB ICL analogues in 5'- ^{32}P -b-1–3 with (A) DEPC at A_{11} relative to A_8 , (B) DMS at G_{13} relative to G_6 , and (C) DEPC at X_{14} relative to A_8 ($X = A$ in 2 and 3, and $X = \text{cross-linked } A$ in 1). P values were calculated using the Student's t test (four replicates).

General Procedure for Hydroxylamine Treatment of 1–3. A 2 M stock solution (10 mL) of hydroxylamine (pH 6.0) was prepared by dissolving hydroxylamine hydrochloride (1.34 g) in autoclaved water (5 mL). Diethylamine (approximately 1.2 mL) was added to adjust the pH to 6.0. Autoclaved water was added to this solution to achieve a final volume of 10 mL. The stock solution was diluted with autoclaved water to make a 1.04 M hydroxylamine solution. The 2 M stock solution can be stored in a -20°C freezer and used for up to 1 week. The appropriately ^{32}P -labeled substrate (1–3) (5000–15000 cpm) was reacted (26 μL) with 0.80 M hydroxylamine (pH 6.0) (diluted from a 1.04 M stock solution) in PBS buffer [10 mM sodium phosphate (pH 7.2) and 100 mM NaCl] for 10 min at room temperature. The reaction tubes were briefly vortexed every 2 min during the reaction. Following incubation, the samples were precipitated with 5 M NH_4OAc (5.2 μL), 1 $\mu\text{g}/\mu\text{L}$ calf thymus DNA (5 μL), water (20.5 μL), and cold ethanol (155 μL). The precipitated samples were treated with 1 M piperidine (50 μL) for 20 min at 90°C and dried in vacuo. The samples were resuspended in water (50 μL) and again dried in vacuo. The samples were resuspended in formamide loading buffer [10 μL , 50% formamide, 5 mM EDTA (pH 8.0), 0.1% xylene cyanol, and 0.1% bromophenol blue], heated (90°C , 2 min), cooled (0°C , 5 min), and analyzed by PAGE (20% denaturing) using a Storm 840 phosphorimager and ImageQuant TL software.

UV Melting Temperature Experiments. UV melting temperature experiments were conducted in quartz cells with a path length of 1 cm. The samples were prepared by mixing oligonucleotides (1 μM), PIPES (pH 7.0, 10 mM), NaCl (100 mM), and MgCl_2 (10 mM). For substrates 4–7, the temperature was held at 13°C for 2 min and then increased linearly to 85°C at a rate of $1.0^\circ\text{C}/\text{min}$ while the absorbance was monitored at 260 nm. The cuvette rack was constantly flushed with N_2 flow to avoid moisture condensation in the lower temperature range ($<17^\circ\text{C}$). For substrates 13 and 14, the temperature was varied from 20 to 85°C . For each experiment, the absorbance (260 nm) of a solution containing the same buffer but without the oligonucleotides was recorded as a function of temperature and subtracted from the absorbance (260 nm) of the oligonucleotide solutions. The data were processed using Origin 6.1 by plotting the absorbance versus temperature and fitting the curve to the equation $y = A_2 + (A_1 - A_2)/\{1 + \exp[(x - x_0)/dx]\}$, where y is the absorbance, A_1 is the initial absorbance (left horizontal asymptote), A_2 is the final absorbance (right horizontal asymptote), x_0 is the center (point of inflection or T_m), and dx is the change in the X value corresponding to the most significant change in the Y value.

RESULTS

Substrate Design and Preparation. Substrates containing ICL analogues of C4-AP and DOB (Figure 3) were synthesized by a combination of solid-phase oligonucleotide synthesis and enzymatic ligation, as previously described.^{16,18} Substrates containing the cross-link with the abasic sites opposite the 3'-adjacent dT ("observed") and with the cross-linked dA opposite the abasic site analogues ("unobserved") were prepared. The "observed" ICLs correspond to those that are (preferentially) formed by C4-AP and DOB (Scheme 2). Specific ligation protocols for previously unreported substrates are described in the Supporting Information. The chemo-enzymatic approach provided a convenient method for controlling which strand and terminus in the cross-linked substrate was labeled with ^{32}P . Labeling each terminus was desirable because analysis of the reactivity of nucleotides flanking the ICLs proceeded with the highest resolution when the nucleotides were between the radiolabel and the cross-link. [Please note that the strand labeled in each substrate is termed "t" for the strand containing the abasic site analogue (top strand in Figure 3) or "b" for the strand containing the dA that is involved in the cross-link (bottom strand in Figure 3).] This approach worked as planned for preparing the desired ^{32}P -labeled cross-linked substrates containing the stabilized DOB ICL [X–Y (Scheme 1 and Figure 3)]. However, slight impurities were detected when 5'- ^{32}P -b-8 and 5'- ^{32}P -b-9 were prepared. Consequently, substrates 11 and 12 were employed for examining the reactivity of the 5'-bottom portion of ICLs containing the C4-AP analogues. Although the overall lengths of substrates 11 and 12 are greater than the lengths of the others (8 and 9) containing the C4-AP ICL analogue (W–V), the flanking sequences over which reactivity was examined (four or five nucleotides) were identical. In addition, studies conducted on DOB ICLs revealed that the reaction of 2'-deoxycytidines with hydroxylamine was significantly less precise in our hands than that of other chemical probing reagents. Consequently, C4-AP cross-link substrates were designed to be devoid of G–C base pairs near the cross-link.

General Aspects of Chemical Reactivity Studies.

Reactivities at all sites were calibrated in two ways. Reactions were conducted to low conversion (single-hit conditions), and ratios of cleavage at a nucleotide of interest proximal to the ICL relative to a distal position were recorded. In addition, reactivity at a given position within an ICL was calibrated by comparison to that of the other ICL and a control substrate lacking a cross-link. Reactions at dA, dG, and T were examined using diethyl pyrocarbonate (DEPC), dimethyl sulfate (DMS), and potassium permanganate (KMnO_4), respectively. Reactivity at dC using hydroxylamine was used sparingly because of unsatisfactory precision in many instances. (Please note that




Figure 5. Chemical reactivity of DOB ICL analogues in 3'-³²P-b-1-3 with DEPC at (A) X_{15} relative to A_{23} ($X = A$ in 1 and 3, and $X =$ cross-linked A in 2), (B) A_{16} relative to A_{23} , and (C) A_{17} relative to A_{23} . P values were calculated using the Student's t test (four replicates).

for the sake of convenience the deoxy descriptor is dropped in the discussion below.) Similarly, hydroxyl radical cleavage was not very useful for detecting differences in structure, and results using this reagent are reported only for substrates that provide information about the reactivity of the abasic site analogues themselves.

Chemical Reactivity of the DOB ICL Analogue.

Reactivity of 5'-³²P-b-1-3 was examined using DEPC and DMS (Figure 4). Reaction at A_{14} of the unobserved ICL (2) with DEPC (Figure 4C) was significantly greater than in the duplex control, indicating a high degree of distortion at this position. In contrast, reaction at A_{14} in the observed ICL (1), which is the cross-linked purine, was negligible. This suggests that the abasic site analogue bonded to the N6 position hinders DEPC reaction at the N7 position. Per below, this was a general observation for the cross-linked positions in all of the substrates. Reaction at G_{13} in the unobserved ICL (2) was still greater than that at the corresponding nucleotide in 1, despite the greater distance from the cross-linking position (Figure 4B). However, there was no enhanced reactivity in either cross-linked substrate compared to that of the control duplex (3) at A_{11} , indicating that any structural distortion induced by the ICLs dissipated at that position (Figure 4A).

As mentioned above, reaction of 3'-³²P-b-1-3 (Figure 5) with DEPC indicated that the 2'-deoxyadenosine involved in the unobserved ICL (A_{15} in 2) was inaccessible. However, the 2'-deoxyadenosine adjacent to the cross-linked purine nucleotide (A_{15}) in observed ICL 1 was ~2-fold more reactive than the comparable nucleotide in duplex DNA (3). At the same time, the 2'-deoxyadenosine adjacent to the cross-linked purine nucleotide (A_{16}) in the unobserved cross-linked substrate (2) was more reactive than the comparable position (A_{15}) in the observed ICL (1) or A_{16} in the duplex (3). The greater reactivity at A_{16} in unobserved ICL 2 compared to that of A_{15} of 1, both of which are adjacent to the cross-linked dA, is even more striking when one considers that the latter is opposite the abasic site, which provides no opportunity for Watson-Crick base pairing. The difference in reactivity between ICLs at A_{17} was negligible, but both were more susceptible to DEPC than the control duplex (3). Enhanced reactivity at these positions could be due to fraying of the base paired complement in 1 and 2. Fraying of the noncovalently bound strand may also contribute to the enhanced reactivity at T_{44} and T_{45} with $KMnO_4$ in cross-linked substrates 1 and 2 (Figure 6). This is particularly apparent at the 3'-terminus of the strand (T_{45}) where reactivity in either ICL is >6-fold greater than in control duplex 3. However, the reactivity in the unobserved ICL (2) remains >6 times higher when the nucleotide (T_{44}) is one nucleotide further removed from the ICL. In comparison,




Figure 6. Chemical reactivity of DOB ICL analogues in 5'-³²P-t-1-3 with $KMnO_4$ at (A) T_{44} relative to T_{38} and (B) T_{45} relative to T_{38} . P values were calculated using the Student's t test (four replicates).

reactivity at T_{44} in 5'-³²P-t-1 is reduced to ~3-fold greater than in duplex 3.

In contrast, potassium permanganate reacts much more readily at T_{47} in the observed ICL (3'-³²P-t-1) than it does in the duplex control or the unobserved ICL (3'-³²P-t-2). However, T_{47} is still >6-fold reactive in 2 than it is in duplex 3 (Figure 7B). Evaluation of reactivity at T_{47} was unusual in that it was calibrated against cleavage at a different nucleotide (G_{54}). This was because cleavage at other thymidines distal from the cross-link within the 3'-arm of the top strand in 1-3 was too low to reliably measure. The abasic site (X_{46}) also reacts more readily with hydroxyl radical in the unobserved cross-link (3'-³²P-t-2) than it does in the analogue of the observed molecule (Figure 7A). However, both abasic sites react more readily with hydroxyl radical than does the control duplex. Although reaction at C_{48} is indistinguishable in 3'-³²P-t-1-3 due to the larger error (Figure 7C), the unobserved ICL (2) is more susceptible to DMS treatment at G_{49} (Figure 7D) than either of the other substrates. Reaction of T_{50} is indistinguishable in the three substrates (data not shown), indicating that any structural alteration in the ICLs has dissipated by that position.

Chemical Reactivity of the C4-AP ICL Analogue.

As discussed above, reactivity on the 5'-arm in the bottom strand of the cross-linked C4-AP analogue was examined (Figure 8) using longer substrates (11 and 12) but compared against a shorter duplex (10) containing the same sequence in the relevant region. The reactivity of the thymidines ($KMnO_4$) and




Figure 7. Chemical reactivity of DOB ICL analogues in 3'-³²P-t-1–3 at (A) X_{46} relative to G_{54} ($X = T$ in 3, and $X =$ cross-linked abasic site in 1 and 2) with hydroxyl radical, (B) T_{47} relative to G_{54} with $KMnO_4$, (C) C_{48} relative to C_{55} with hydroxylamine, and (D) G_{49} relative to G_{54} with DMS. P values were calculated using the Student's t test (four replicates).




Figure 8. Chemical reactivity of C4-AP ICL analogues in 5'-³²P-b-10–12 upon (A) reaction with DEPC, (B) reaction with $KMnO_4$, and (C) reaction with DEPC. P values were calculated using the Student's t test (four replicates).




Figure 9. Chemical reactivity of C4-AP ICL analogues in 3'-³²P-b-8–10 at (A) X_{18} relative to A_{28} ($X = A$ in 8 and 10, and $X =$ cross-linked A in 9) with DEPC, (B) T_{19} relative to T_{29} with $KMnO_4$, and (C) A_{20} relative to A_{28} with DEPC. P values were calculated using the Student's t test (four replicates).

A's (DEPC) bonded via the 5'-phosphate of the cross-linked A was quantified. The cross-linked A (A_{29}) in the analogue of the observed C4-AP ICL (11) is protected against DEPC, whereas the respective nucleotide in the analogue of the unobserved cross-link (12) is adjacent to the cross-linked position and is very reactive (Figure 8C). The difference in reactivity between the observed (11) and unobserved (12) ICL analogues rapidly dissipates. The unobserved ICL is only slightly more reactive at T_{28} (Figure 8B), and the difference is negligible just two nucleotides further removed from the cross-linked position [A_{27} (Figure 8A)].

The nucleotides bonded to the 3'-phosphate of the cross-linked dA [A_{18} in 8 (Figure 9A) and T_{19} in 9 (Figure 9B)] were significantly more reactive than the control duplex (10). In each instance, these are the positions adjacent to the cross-linked dA. However, the reactivity at T_{19} in the observed C4-AP ICL analogue (8) was anomalously high (Figure 9B), and far greater than that of the 5'-adjacent A_{18} , which is opposite the cross-linked abasic site. Reaction at T_{19} of the observed ICL is also significantly greater than at T_{19} in the unobserved C4-AP cross-link product. The reason for the hyperreactivity of T_{19} in 3'-³²P-b-8 is unknown. More enhanced reactivity in the unobserved ICL (9) relative to the analogue of the observed ICL (8) is




Figure 10. Chemical reactivity of C4-AP ICL analogues in 5'-³²P-t-8-10 at (A) A₅₂ relative to G₄₈ with DEPC, (B) T₅₃ relative to T₄₅ with KMnO₄, and (C) A₅₄ relative to G₄₈ with DEPC. *P* values were calculated using the Student's *t* test (four replicates).




Figure 11. Chemical reactivity of C4-AP ICL analogues in 3'-³²P-t-8-10 at (A) V₅₅ relative to A₆₇ (X = T in 10, and X = cross-linked abasic site in 8 and 9) with hydroxyl radical, (B) T₅₆ relative to T₆₈ with KMnO₄, (C) A₅₇ relative to A₆₇ with DEPC, and (D) T₅₈ relative to T₆₈ with KMnO₄. *P* values were calculated using the Student's *t* test (four replicates).

restored upon moving one nucleotide farther from the ICL [A₂₀ (Figure 9C)], but the reactivities of the cross-linked substrates are indistinguishable at T₂₁ (data are not shown).

The nucleotides in the 5'-arm of the top strands in the C4-AP ICL analogues are consistently more reactive in the unobserved (5'-³²P-t-9) cross-link substrate than in 8, which is designed to model the observed cross-link (Figure 10). The differential reactivity was greatest at the 5'-adjacent nucleotide to the abasic site [A₅₄ (Figure 10C)], where the extent of DEPC-induced cleavage was more than 6 times greater in 9 than in the control duplex. With the exception of reaction with hydroxyl radical (Figure 11A), the top strand in the unobserved analogue (9) was also consistently more reactive than 8 on the 3'-side of the cross-link (Figure 11). However, the differences in cleavage in 3'-³²P-t-8 and 3'-³²P-t-9 are considerably smaller, and reactions of the abasic site analogues (Figure 11A) are indistinguishable from one another, as well as with respect to the control duplex (3'-³²P-t-10). Overall, the C4-AP ICL analogues are significantly less reactive than the respective DOB compounds (Figure 7A) with hydroxyl radical.

UV Melting Comparison of ICL Analogues. The UV melting temperatures of the observed and unobserved cross-links were measured at 1 μM at pH 7.0 in PIPES (10 mM), MgCl₂ (10 mM), and NaCl (100 mM). Two transitions were observed for the DOB ICL analogue substrates [4 and 5 (Figure 12A)]. The lower-temperature transition (*T_m*, 31.0 ± 0.3 °C for 4 and 28.1 ± 0.6 °C for 5) corresponds to melting of




Figure 12. UV melt of (A) DOB ICL analogues (4 and 5) and (B) C4-AP ICL analogues (13 and 14).

the noncovalently bound strand. As expected, the presence of this strand does not affect the melting of the cross-linked strands [4 and 5 (Figure 12A) and 6 and 7 (Supporting

Information and Table 1)]. Importantly, the melting of the cross-link required a temperature ($\sim 4^\circ\text{C}$) for the observed

Table 1. UV Melting Temperatures of ICL Analogues

ICL analogue	ICL T_M ($^\circ\text{C}$) ^a
DOB observed (4)	69.8 ± 0.8
DOB unobserved (5)	65.3 ± 0.1
C4-AP observed (13)	59.1 ± 0.2
C4-AP unobserved (14)	57.4 ± 0.4

^aMeasurements are the average \pm the standard deviation of three independent measurements.

ICL (4) considerably higher than that of the cross-link in which the dA and abasic site are opposite one another (unobserved, 5).

A similar trend was observed when the C4-AP ICL analogues (13 and 14) were examined (Figure 12B and Table 1). The substrate containing the cross-linked components opposite one another (14), which represents the orientation that is not observed experimentally, melted at a temperature $\sim 1.7^\circ\text{C}$ lower than that for the orientation in which the cross-linked dA is opposite the thymidine that is bonded to the 3'-phosphate of the abasic site (13). Although the absolute temperatures for melting of the C4-AP analogue ICLs are lower than those containing the DOB cross-link, the difference in T_M between the observed (13) and unobserved (14) is smaller in the former. This suggests that the C4-AP ICL analogue imparts a smaller perturbation on the structure and stability of the DNA than the DOB cross-link analogue.

DISCUSSION

Spontaneous interstrand cross-link formation by abasic sites, which require NER, increases the potential biochemical impact of these DNA lesions.³⁶ Reversible cross-linking to a dA opposite a 3'-adjacent thymidine by C4-AP and DOB oxidized abasic sites occurs with very high selectivity over other sequences (Scheme 2).^{12–15} Inspection of molecular models (Figure 2) does not provide an obvious rationalization for this selectivity. Consequently, UV melting experiments and chemical reactivity studies were conducted with stabilized analogues of the ICLs to discern a difference in the stabilities of the isomeric cross-linked molecules.

In general, the cross-linked 2'-deoxyadenosine was unreactive with DEPC, and certainly less reactive than a comparable nucleotide in duplex DNA. One possible explanation for this is that the cross-linked abasic site model hinders approach to the N7 position of the cross-linked dA and/or reduces its nucleophilicity. Another general observation is that the "top" strands in the cross-linked substrates, which contain the abasic site analogues, tend to be more reactive than the "bottom" strands containing the cross-linked dAs. This suggests significant solvent exposure and structural disruption in the cross-linked strands containing the abasic site analogues.

Although there are some exceptions, a comparison of the reactivity of the "observed" (e.g., 1, 8, and 11) and "unobserved" (e.g., 2, 8, and 12) stabilized cross-link substrates (see also Scheme 2) at individual nucleotides showed that the latter were consistently more reactive. This is consistent with greater distortion of the unobserved cross-links than of the analogues of observed ICLs. These inferential observations were corroborated by UV melting measurements (Table 1) of stabilized analogues of C4-AP (13 and 14) and DOB (4 and 5)

ICLs. Substrates containing cross-links between opposing nucleotides melted at temperatures $1.7\text{--}4.5^\circ\text{C}$ lower than those of the analogues of their "observed" counterparts. Overall, UV melting temperatures and chemical reactivity of stabilized analogues of these ICLs (Scheme 1) revealed that the preferred cross-links are more stable and their structures less perturbed than those of comparable unobserved cross-linked substrates, providing a rationale for their preferential formation.

A comparison of the reactivity in the respective "observed" ICL substrates relative to the corresponding duplexes lacking the cross-link indicates that the DOB analogue induces a greater overall enhancement (Figure 13A) than does the C4-




Figure 13. Nucleotide by nucleotide comparison of ICL reactivity relative to analogous duplex: (A) DOB-containing ICL and (B) C4-AP-containing ICL.

AP-like lesion (Figure 13B). The anomalous reactivity at T_{19} in 3'-³²P-b-8 is the one outlier. This is especially true upon comparison of the reactivity of the strands containing the abasic site analogues ("top" strands). In particular, the abasic site analogue in the DOB substrate (Figure 7A) is considerably more reactive than the respective C4-AP-type abasic sites (Figure 11A) with hydroxyl radical. In addition, the thymidine opposite the cross-linked dA [X_{14} (Figure 13A)] and the first two nucleotides in the flanking strand (T_{45} and T_{44}) are far more reactive than in the duplex. The greater reactivity at T_{44} and T_{45} in the DOB ICL may be attributed to their higher level of breathing because they are in effect at the 3'-terminus of a duplex. Overall, the greater reactivity suggests that cross-links derived from DOB perturb the structure of DNA more than does C4-AP. This is consistent with the repair incision of DNA containing the DOB ICL being more facile than that involving C4-AP by UvrABC, which recognizes macroscopic distortions in the nucleic acid substrate.^{18–22}

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.biochem.5b00860.

Procedures for preparing ICL substrates, UV melting plots of analogues of 6 and 7 lacking the flanking strand, ESI-MS of cross-linked oligonucleotides, and sample autoradiograms of chemical probing reactions (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*Department of Chemistry, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218. Telephone: 410-516-8095. Fax: 410-516-7044. E-mail: mgreenberg@jhu.edu.

Funding

We are grateful for support of this research by the National Institute of General Medical Sciences (GM-063028).

Notes

The authors declare no competing financial interest.

■ ABBREVIATIONS

AP, apurinic/apyrimidinic site; C4-AP, C4'-oxidized abasic site; BER, base excision repair; DOB, 5'-(2-phosphoryl-1,4-dioxobutane); dsb, double-strand break; ICL, interstrand cross-link; NER, nucleotide excision repair; DMS, dimethyl sulfate; DEPC, diethyl pyrocarbonate.

■ REFERENCES

- (1) Räsche, M., Knipscheer, P., Enoiu, M., Angelov, T., Sun, J., Griffith, J. D., Ellenberger, T. E., Schärer, O. D., and Walter, J. C. (2008) Mechanism of replication-coupled DNA interstrand crosslink repair. *Cell* 134, 969–980.
- (2) Noll, D. M., Mason, T. M., and Miller, P. S. (2006) Formation and repair of interstrand cross-links in DNA. *Chem. Rev.* 106, 277–301.
- (3) Kozekov, I. D., Turesky, R. J., Alas, G. R., Harris, C. M., Harris, T. M., and Rizzo, C. J. (2010) Formation of deoxyguanosine cross-links from calf thymus DNA treated with acrolein and 4-hydroxy-2-nonenal. *Chem. Res. Toxicol.* 23, 1701–1713.
- (4) Stone, M. P., Cho, Y.-J., Huang, H., Kim, H.-Y., Kozekov, I. D., Kozekova, A., Wang, H., Minko, I. G., Lloyd, R. S., Harris, T. M., and Rizzo, C. J. (2008) Interstrand DNA cross-links induced by α,β -unsaturated aldehydes derived from lipid peroxidation and environmental sources. *Acc. Chem. Res.* 41, 793–804.
- (5) Schärer, O. D. (2005) DNA interstrand crosslinks: Natural and drug-induced DNA adducts that induce unique cellular responses. *ChemBioChem* 6, 27–32.
- (6) Clauson, C., Schärer, O. D., and Niedernhofer, L. (2013) Advances in understanding the complex mechanisms of DNA interstrand cross-link repair. *Cold Spring Harbor Perspect. Biol.* 5, a012732.
- (7) Price, N. E., Catalano, M. J., Liu, S., Wang, Y., and Gates, K. S. (2015) Chemical and structural characterization of interstrand cross-links formed between abasic sites and adenine residues in duplex DNA. *Nucleic Acids Res.* 43, 3434–3441.
- (8) Catalano, M. J., Liu, S., Andersen, N., Yang, Z., Johnson, K. M., Price, N. E., Wang, Y., and Gates, K. S. (2015) Chemical structure and properties of interstrand cross-links formed by reaction of guanine residues with abasic sites in duplex DNA. *J. Am. Chem. Soc.* 137, 3933–3945.
- (9) Price, N. E., Johnson, K. M., Wang, J., Fekry, M. I., Wang, Y., and Gates, K. S. (2014) Interstrand DNA-DNA crosslink formation between adenine residues and abasic sites in duplex DNA. *J. Am. Chem. Soc.* 136, 3483–3490.

- (10) Johnson, K. M., Price, N. E., Wang, J., Fekry, M. I., Dutta, S., Seiner, D. R., Wang, Y., and Gates, K. S. (2013) On the formation and properties of interstrand DNA-DNA cross-links forged by reaction of an abasic site with the opposing guanine residue of 5'-Cap sequences in duplex DNA. *J. Am. Chem. Soc.* 135, 1015–1025.
- (11) Regulus, P., Duroux, B., Bayle, P.-A., Favier, A., Cadet, J., and Ravanat, J.-L. (2007) Oxidation of the sugar moiety of DNA by ionizing radiation or bleomycin could induce the formation of a cluster DNA lesion. *Proc. Natl. Acad. Sci. U. S. A.* 104, 14032–14037.
- (12) Szczepanski, J. T., Jacobs, A. C., and Greenberg, M. M. (2008) Self-promoted DNA interstrand cross-link formation by an abasic site. *J. Am. Chem. Soc.* 130, 9646–9647.
- (13) Szczepanski, J. T., Jacobs, A. C., Majumdar, A., and Greenberg, M. M. (2009) Scope and mechanism of interstrand cross-link formation by the C4'-oxidized abasic site. *J. Am. Chem. Soc.* 131, 11132–11139.
- (14) Szczepanski, J. T., Hiemstra, C. N., and Greenberg, M. M. (2011) Probing DNA interstrand cross-link formation by an oxidized abasic site using nonnative nucleotides. *Bioorg. Med. Chem.* 19, 5788–5793.
- (15) Guan, L., and Greenberg, M. M. (2009) DNA interstrand cross-link formation by the 1,4-dioxobutane abasic lesion. *J. Am. Chem. Soc.* 131, 15225–15231.
- (16) Ghosh, S., and Greenberg, M. M. (2014) Synthesis of cross-linked DNA containing oxidized abasic site analogues. *J. Org. Chem.* 79, 5948–5957.
- (17) Szczepanski, J. T., Jacobs, A. C., Van Houten, B., and Greenberg, M. M. (2009) Double strand break formation during nucleotide excision repair of a DNA interstrand cross-link. *Biochemistry* 48, 7565–7567.
- (18) Ghosh, S., and Greenberg, M. M. (2014) Nucleotide excision repair of chemically stabilized analogues of DNA interstrand cross-links produced from oxidized abasic sites. *Biochemistry* 53, 5958–5965.
- (19) Schärer, O. D. (2013) Nucleotide excision repair in eukaryotes. *Cold Spring Harbor Perspect. Biol.* 5, a012609.
- (20) Pakotiprapha, D., Samuels, M., Shen, K., Hu, J. H., and Jeruzalmi, D. (2012) Structure and mechanism of the UvrA,UvrB DNA damage sensor. *Nat. Struct. Mol. Biol.* 19, 291–298.
- (21) Pakotiprapha, D., Liu, Y., Verdine, G. L., and Jeruzalmi, D. (2009) A structural model for the damage-sensing complex in bacterial nucleotide excision repair. *J. Biol. Chem.* 284, 12837–12844.
- (22) Pakotiprapha, D., Inuzuka, Y., Bowman, B. R., Moolenaar, G. F., Goosen, N., Jeruzalmi, D., and Verdine, G. L. (2008) Crystal structure of bacillus stearothermophilus UvrA provides insight into ATP-modulated dimerization, UvrB interaction, and DNA binding. *Mol. Cell* 29, 122–133.
- (23) Knipscheer, P., Raschle, M., Smogorzewska, A., Enoiu, M., Ho, T. V., Schärer, O. D., Elledge, S. J., and Walter, J. C. (2009) The fanconi anemia pathway promotes replication-dependent DNA interstrand cross-link repair. *Science* 326, 1698–1701.
- (24) Yakubovskaya, M. G., Belyakova, A. A., Gasanova, V. K., Belitsky, G. A., and Dolinnaya, N. G. (2010) Comparative reactivity of mismatched and unpaired bases in relation to their type and surroundings. Chemical cleavage of DNA mismatches in mutation detection. *Biochimie* 92, 762–771.
- (25) Jarem, D. A., Huckaby, L. V., and Delaney, S. (2010) AGG interruptions in (CGG)(n) DNA repeat tracts modulate the structure and thermodynamics of non-B conformations in vitro. *Biochemistry* 49, 6826–6837.
- (26) Sun, D.-A., and Hurley, L. H. (2010) Biochemical techniques for the characterization of G-quadruplex structures: EMSA, DNA footprinting, and DNA polymerase stop assay. *Methods Mol. Biol.* 608, 65–79.
- (27) Maxam, A. M., and Gilbert, W. (1980) Sequencing end-labeled DNA with base-specific chemical cleavages. *Methods Enzymol.* 65, 499–513.
- (28) Ambrose, B. J. B., and Pless, R. C. (1987) DNA sequencing: Chemical methods. *Methods Enzymol.* 152, 522–539.
- (29) Fox, K. R., and Grigg, G. W. (1988) Diethylpyrocarbonate and permanganate provide evidence for an unusual DNA conformation

induced by binding of the antitumor antibiotics bleomycin and phleomycin. *Nucleic Acids Res.* 16, 2063–2075.

(30) Shcherbakova, I., and Mitra, S. (2009) Hydroxyl-radical footprinting to probe equilibrium changes in RNA tertiary structure. *Methods Enzymol.* 468, 31–46.

(31) Price, M. A., and Tullius, T. D. (1992) Using hydroxyl radical to probe DNA structure. *Methods Enzymol.* 212, 194–219.

(32) Gelfand, C. A., Plum, G. E., Grollman, A. P., Johnson, F., and Breslauer, K. J. (1998) Thermodynamic consequences of an abasic lesion in duplex DNA are strongly dependent on base sequence. *Biochemistry* 37, 7321–7327.

(33) McCarthy, J. G., Williams, L. D., and Rich, A. (1990) Chemical reactivity of potassium permanganate and diethyl pyrocarbonate with B DNA: Specific reactivity with short A-tracts. *Biochemistry* 29, 6071–6081.

(34) Bishop, E. P., Rohs, R., Parker, S. C. J., West, S. M., Liu, P., Mann, R. S., Honig, B., and Tullius, T. D. (2011) A map of minor groove shape and electrostatic potential from hydroxyl radical cleavage patterns of DNA. *ACS Chem. Biol.* 6, 1314–1320.

(35) Tullius, T. D., and Greenbaum, J. A. (2005) Mapping nucleic acid structure by hydroxyl radical cleavage. *Curr. Opin. Chem. Biol.* 9, 127–134.

(36) Greenberg, M. M. (2014) Abasic and oxidized abasic site reactivity in DNA: Enzyme inhibition, crosslinking, and nucleosome catalyzed reactions. *Acc. Chem. Res.* 47, 646–655.