

2P.7 F₁F₀ ATP synthase mutants in *Chlamydomonas*: Stability and oligomycin resistance mediated by atypical Asa7 protein; interaction between chloroplastic and mitochondrial bioenergetics

Marie Lapaille¹, Adelma Escobar-Ramírez², Hervé Degand³, Denis Baurain⁴, Marc Thiry⁵, Emilie Perez¹, Marc Boutry³, Diego Gonzalez-Halphen², Claire Remacle¹, Pierre Cardol¹

¹Université de Liège, Botany Institute, Belgium

²Universidad Nacional Autónoma de México,

Instituto de Fisiología Celular, Mexico

³Université catholique de Louvain, Institut des Sciences de la vie, Belgium

⁴Université de Liège, Faculty of Veterinary Medicine, Belgium

⁵University of Liège, Laboratory of Cell and Tissue Biology, Belgium

E-mail: pierre.cardol@ulg.ac.be

In yeast, mammals, and land plants, F₁F₀ ATP synthase (complex V) comprises about 15 conserved subunits. In this work, we show that complex V from chlorophycean green algae has an atypical subunit composition of its peripheral stator and dimerization module, with 9 subunits of unknown evolutionary origin (Asa subunits), while complex V has a canonical subunit composition in other classes of green algae. In addition, growth, respiration and ATP levels in Chlorophyceae are also barely affected by oligomycin concentrations that affect representatives of the other classes of Chlorophytes. We then isolated *Chlamydomonas* mutants lacking either beta subunit (Atp2) or an atypical subunit (Asa7). The Atp2 mutant is an obligate phototroph lacking complex V assembly and ATP synthesis coupled to the respiration. In addition, Atp2-deficient mitochondria are deprived of cristae, and rearrangements of the photosynthetic apparatus and thylakoid organization are observed. In contrast, the loss of Asa7 subunit has no impact on cell bioenergetics or organelles structures, but it destabilizes the enzyme dimeric form *in vitro* and renders growth, respiration and ATP level sensitive to oligomycin. Altogether, our results suggest that the loss of canonical components of the stator happened at the root of chlorophycean lineage and was accompanied by the recruitment of novel polypeptides. Such a massive modification of stator features might have conferred novel properties, including the stabilization of the enzyme dimeric form and the shielding of the proton channel. Our study also contributes to the understanding of the yet poorly-studied bioenergetic interactions between organelles in photosynthetic organisms.

doi:10.1016/j.bbabbio.2010.04.104

2P.8 Purification and characterisation of F-ATPase from three species of fungi

Thomas J. Charlesworth, Ian M. Fearnley, John V. Bason, Scott A. Ferguson, Michael J. Runswick, John E. Walker
Medical Research Council, Mitochondrial Biology Unit,
Wellcome Trust/MRC Building, Hills Road, Cambridge, CB2 0XY, UK
E-mail: walker@mrc-mbu.cam.ac.uk

F-ATPase has been purified from mitochondria from *Pichia angusta*, *Pichia pastoris* and *Yarrowia lipolytica* by affinity chromatography with a monomeric pH insensitive form of the bovine yeast inhibitor protein, IF₁. The subunit compositions of the complexes have been characterized by SDS-PAGE, by mass spectrometric analysis of tryptic peptides, and by measurement of the masses of the subunits by LC-MS [1]. The impact of phospholipids on specific activities and the sensitivity to oligomycin of the complexes has been studied. The enzymes have been reconstituted in phospholipid vesicles and proton pumping was measured with the pH sensitive probe amino-6-chloro-2-methoxy-acridine (ACMA). ACMA quenching was observed after addition of ATP, halted by addition of oligomycin and reversed by addition of the ionophore gramicidin. The thermostability of the

purified complexes was investigated by following the ATP hydrolysis activity from 25 °C to 70 °C. As expected, the *P. angusta* enzyme was the most thermostable. It has been selected for crystallisation trials.

Reference

[1] Carroll J, Fearnley IM, Wang Q, Walker JE, (2009) *Anal. Biochem.* **395**: 249–255.

doi:10.1016/j.bbabbio.2010.04.105

2P.9 3-Iodothyronamine favours IF₁ release from F₀F₁ ATP synthase

Sebastiano Cumero¹, Stefania Contessi¹,
Riccardo Zucchi², Irene Mavelli¹

¹University of Udine, Department of Biomedical Sciences and Technologies, Italy

²University of Pisa, Department of Human and Environment Sciences, Italy
E-mail: stefania.contessi@uniud.it

F₀F₁ ATP synthase has been demonstrated as a good molecular target for drugs in the treatment of various diseases and regulation of energy metabolism so that the development of new F₀F₁ ATP synthase-directed agents is encouraged nowadays. On the basis of a structural similarity with well characterised enzyme inhibitors, such as resveratrol, the inhibitory capacity of 3-iodothyronamine (T₁AM), a naturally occurring derivative of thyroid hormone, has been investigated [1]. First, T₁AM effect on the activity of the mitochondrial respiratory chain was excluded. Then, T₁AM direct inhibitory effect on ATP synthase/ase activity has been investigated on F₀F₁ whole enzyme using preparations of submitochondrial particles (SMP) as model. When ΔΨ is sustained by succinate, IC₅₀ values indicate that T₁AM has an inhibitory potency on ATP synthase activity similar to that of resveratrol. On the contrary, when ΔΨ is inhibited by rotenone and the enzyme can only hydrolyze ATP, T₁AM shows an activating effect at a low concentration (10–50 μM) followed by a limited inhibitory effect at a higher concentration (50–150 μM). The activating action results more marked on SMP preparations having a higher content of the natural inhibitor protein (IF₁) [2], while it disappears on IF₁-free AS particles. Such effect is consistent with IF₁ release from the enzyme as a result of T₁AM binding. To support this evidence, immunoblotting analysis was performed on high-IF₁ containing SMP after treatment with T₁AM. The dose-response curve shows that IF₁ release occurs starting from the nanomolar T₁AM range suggesting that this mechanism could be of physiologic relevance and distinct from the inhibitory one. To help into the indirect localization of T₁AM inhibitory binding site, purified IF₁-free F₁ was used as a model. Kinetic analysis shows that T₁AM exerts a mixed type inhibition of ATPase activity as resveratrol does with a similar inhibitory potency. Nevertheless, competition experiments show that the inhibitory effects of T₁AM and resveratrol are additive indicating that the binding of the two inhibitors to F₁ is not mutually exclusive. This suggests distinct binding sites for the two drugs in agreement with peculiar T₁AM action. The binding site/s of T₁AM is/are now under investigation by further kinetic analyses and by molecular modelling with reference to available F₁ native and inhibited structures.

References

[1] Chiellini G *et al.* (2007) *FASEB J.* **21**: 1597–1608.
[2] Gledhill JR *et al.* (2007) *Proc. Natl. Acad. Sci.* **104**: 15671–15676.

doi:10.1016/j.bbabbio.2010.04.106