

Marine nitrogen: Phosphorus stoichiometry and the global N:P cycle

JOHN A. DOWNING

*Department of Animal Ecology, Iowa State University, 124 Science II, Ames, Iowa
50011-3221, USA*

Key words: freshwater, limitation, marine, nitrogen, phosphorus, ratio, stoichiometry, trace elements

Abstract. Nitrogen supply is often assumed to limit marine primary production. A global analysis of total nitrogen (N) to phosphorus (P) molar ratios shows that total N:P is low ($<16:1$) in some estuarine and coastal ecosystems, but up to $100:1$ in open oceans. This implies that elements other than N may limit marine production, except in human impacted, estuarine or coastal ecosystems. This pattern may reconcile conflicting enrichment studies, because N addition frequently increases phytoplankton growth where total N:P is expected to be low, but P, Fe, or Si augment phytoplankton growth in waters where total N:P is high. Comparison of total N:P stoichiometry between marine and freshwaters yields a model of the form of the aquatic N:P cycle.

Introduction

Marine primary production yields >90 billion kg of food to the world economy each year (FAO 1993). Marine ecosystems store 50-times more inorganic carbon than the earth's atmosphere, so marine primary production may play an important role in establishing global climate (Mackenzie et al. 1993; Ritschard 1992). Factors regulating marine production are therefore of broad societal interest.

There is currently disagreement about the nutrient elements limiting marine primary production (Howarth 1988; Smith 1984). On one hand, some have concluded (e.g. Boynton et al. 1982) that because ratios of dissolved inorganic nitrogen (N) and phosphorus (P) are often lower than the average intracellular N:P ratio of marine organisms (Redfield 1934; Redfield et al. 1963), nitrogen is the element in shortest supply and therefore must limit marine primary production. On the other hand, geochemical budgets suggest that inorganic phosphorus should be in shortest supply (Meybeck 1982) in part because atmospheric N_2 can be fixed (Redfield 1958; Vitousek & Howarth 1991). Experimental additions of inorganic nutrients to seawater samples from various marine ecosystems suggest that several elements may play limiting

roles, including N, P, Si and Fe (Boynton et al. 1982; Martin & Fitzwater 1988; Martin et al. 1994).

Although N and P are present in inorganic, organic and particulate forms, much of marine nutrient stoichiometry has been based only upon the relative amounts of dissolved inorganic N and P found in marine ecosystems. Frequently measured NO_2^- , NO_3^- , NH_4^+ and PO_4^{3-} may represent only a fraction of the N and P that can be used by the biotic community, however. Many aquatic organisms, from bacteria to fish, consume and cycle dissolved organic matter (e.g. Amon & Benner 1994; Bentzen et al. 1992; Glibert et al. 1991; Hoegh-Goldberg 1994), although dissolved organic N and P may be cycled at different rates (e.g. Smith et al. 1986). Particulate N and P can also be ingested and rapidly converted to dissolved inorganic and organic forms by zooplankton (e.g. Ikeda et al. 1982). Thus, many analyses of nutrient stoichiometry in marine ecosystems are based on analyses of fragments of the complete N and P pools.

Analyses of total N and P pools are used to distinguish N from P limitation in freshwaters. Phosphorus, usually measured as PO_4^{3-} was initially recognized as the principal limiting nutrient in freshwaters, but limitation by nitrogen has since been reported in many freshwater ecosystems (Elser et al. 1990). Dissolved N and P are cycled so rapidly by living organisms that they are absorbed as fast as they are produced and thus are often difficult to detect in the soluble inorganic state. Because both organic and inorganic N and P can be used by aquatic organisms, analyses of the total amounts of nutrient elements (i.e. the sum of all dissolved, particulate, organic and inorganic nutrients) yield better measurements of freshwater nutrient supply than analyses of soluble, inorganic N and P alone (Sakamoto 1966).

This approach has been successfully applied in the management of freshwater ecosystems. A recent study shows that comparison of the total N:P ratios of whole water samples with the average N and P needs of algal cells is useful in distinguishing freshwater ecosystems limited by N from those limited by P (Downing & McCauley 1992). Ecosystems with N:P molar ratios less than the average required cellular ratio of 16:1 (Hecky & Kilham 1988) are generally N-limited and those with ratios >16:1 are P-limited. Unfortunately, no analysis of marine data on total N and P stoichiometry has been performed due to the scarcity of total N and P data in marine environments. The purpose of this analysis is to summarize existing marine data to find how total inventories of N and P vary in the world's marine ecosystems.

Methods

Marine data on total N and total P are rare because marine studies seldom analyze all fractions of N and P. Some estimates of total N and P concentrations are scattered throughout the world literature, including data from a polluted European estuary, Chesapeake estuary, a North Carolina inlet, several harbors, the Pacific Ocean near Peru, California and New Zealand, the Gulf of Bothnia, the Baltic, the Gulf of Finland, the Strait of Gibraltar, the Indian Ocean, the Adriatic, and the Mediterranean. Several sources of total N and total P data are analyzed here: Ben-Taleb et al. (1987), Black et al. (1981), Coste et al. (1988), Dame et al. (1986), Degobbis & Gilmartin (1990), Jordan et al. (1991), Lahdes & Leppanen (1988) Le Rouzic & Bertru (1992), Perttila et al. (1980), Pietikainen et al. (1978), Pitkanen & Malin (1980), Ramadan et al. (1984), Rydberg & Sundberg (1986), Updegraff et al. (1977), Valderrama (1981), and Williams (1967). Data were restricted to averages for depth strata at sampling stations where analyses of total N and P or all components (inorganic, organic and particulate) were made using accepted standard methods. These methods usually consist of a digestion of samples for TP analysis followed by a colorimetric detection of PO_4^{3-} . Total nitrogen was analyzed by several methods, frequently determining concentrations of N fractions separately then summing, or by digestion to a soluble form, followed by inorganic N determination. Although some of these analyses may underestimate the abundance of dissolved, organic N (Suzuki et al. 1985), such underestimates, if present, would tend toward underestimation of TN:TP ratios (Karl et al. 1993). Standard TP and TN methods employed were those described in: Dal Pont et al. (1974), D'Elia et al. (1977), Grasshoff (1976), Grasshoff (1983), Menzel & Corwin (1965), Strickland & Parsons (1968), Strickland & Parsons (1972), Treguer & Le Corre (1975), and Valderrama (1981).

Results and discussion

We collected published data on total N and P inventories (dissolved inorganic and organic plus particulate forms) for 191 sample series taken at 88 marine sampling stations throughout the world. The total nitrogen concentration in marine ecosystems, ranging from estuaries to oligotrophic seas, only varied by about 30-fold (6–200 μM), whereas total phosphorus varied 650-fold (0.03–20 μM) (Figure 1). Therefore, total N:P ratios varied between about 5 and 310 (in moles), averaging N:P = 37, over all.

The majority of marine data analyzed here show N:P greater than the ratio required by planktonic marine algae (Figure 2). The N:P in living marine organisms varies greatly but averages about 16 (Hecky & Kilham 1988;




Figure 1. Relationship between total nitrogen and total phosphorus concentrations of marine ecosystems. Total N and P data were usually averages calculated from sets of samples taken at the same site on at least 3 different dates within the same year. Stations in the upper photic zone were defined conservatively as those <50 m in depth (Tett 1990) while nearshore sites were determined from information presented in published studies. The dashed line is the approximate average ratio of N and P in living marine plankton.

Meybeck 1982; *cf* Elser & Hassett 1994), therefore ambient N:P < 16 would indicate N limitation, while N:P > 16 would suggest P limitation of production. N:P is especially high in the upper photic zone where most of the ocean's planktonic production occurs and where nutrient limitation of production is most acute (Tett 1990). Only 12% of the open ocean photic zone sites had total N:P < 16:1. Only 3% had N:P indicating strong nitrogen deficiency (N:P < 10), 19% had N:P near to biological requirements (10 < N:P < 20), while 78% of open, photic zone sites had N:P indicating a deficiency of phosphorus or some element other than nitrogen (Figure 2). Both parametric and nonparametric statistical analyses show that the average total N:P ratio ($\overline{N:P} = 43$) for open ocean photic zone sites (conservatively estimated as <50 m depth) was significantly ($p < 0.0001$) greater than 16:1. Oligotrophic sites have highest total N:P illustrated by the negative correlation between the logarithm of N:P and the logarithm of total P (Figure 3; $p < 0.00001$, $r^2 = 0.81$). The significant tendency for oligotrophic, open seas to have N:P greater than 16, indicates that P or some nutrient other than N limits production in open seas. Ratios of N:P are frequently >50:1 in the nutrient poor sites examined (Figure 3), often as great as 100:1. Total N:P > 16 occurred at 61




Figure 2. Nitrogen deficiency in marine waters relative to the usual N:P of living organisms. Shown are total N:P calculated from inventories of dissolved and particulate inorganic and organic N and P at several marine sampling stations. Environments include a broad variety of conditions and represent station-depth averages. Stations in the upper photic zone were defined conservatively as those <50 m in depth (Tett 1990) while nearshore sites were determined from information presented in published studies.

sites in 34 ecosystems, analyzed independently in 13 different studies, using several different standard methods. Provided that there is not a great divergence in the fraction of total N and P that are biologically labile, total N and P stoichiometry suggests that phosphorus or other factors limit production in the open oceans.

The data analyzed here suggest that N:P is very high in oligotrophic, open seas but often very low in estuaries and coastal ecosystems (Table 1). Estuarine, coastal, or enclosed parts of the sea had N:P ratios less than the required cellular ratio of 16:1 significantly more frequently (χ^2 -test, $p = 0.005$) than samples taken from the surface waters of the open ocean (Figures




Figure 3. Total N:P ratios of data plotted in Figure 1 distinguishing types of marine environments. Deep and shallow sites were determined as in Figure 2. Data obtained from estuaries, harbors and bays were differentiated as such by the authors of studies. The dashed line is the approximate average ratio of N and P in living marine plankton.

Table 1. Frequency of total N:P molar ratios (Figure 3) that are greater than the approximate average ratio of N and P in living marine plankton (16:1). Ratios >16 imply that nitrogen is probably not the principal limiting nutrient.

Environment	Frequency of total N:P molar ratios:	
	<16	≥ 16
Open oceans, $<50\text{m}$ depth	14	104
Open oceans, surface films	3	6
Estuaries, harbors, bays	11	17
Deep, open oceans, $>50\text{m}$ depth	23	13
Total	51	140

2–3; Table 1). Nitrogen is therefore probably the primary limiting nutrient in estuarine and coastal ecosystems, especially those heavily loaded by human influence (Howarth et al. 1995).

The unexpectedly great frequency of high N:P in open oceans is surprising because, except for a few cases (e.g. Kron et al. 1991), most analyses have concluded that N:P is usually <16 in marine ecosystems. This pattern has probably been missed because measurements of dissolved organic, particulate

or whole N and P fractions have lagged behind measurements of NO_2^- , NO_3^- , NH_4^+ and PO_4^{3-} .

Some modern studies measuring several N and P pools at oligotrophic, open ocean sites also indicate total N:P frequently > 16 in the photic zone. For example, the fractionation of N and P at an oligotrophic station in the open Pacific can be approximated from data collected under the Hawaii Ocean Time-series project of the Joint Global Ocean Flux Study (Winn et al. 1993). Water samples, collected in February, March, April, May, June, July, August, September, October, and December of 1991, were analyzed for several N and P fractions. N and P in different fractions and depths can be estimated from averaged N and P profiles in different sample series. Averaging over all cruises during the year, the approximate average total nitrogen concentration in the upper 200 m was $9.7 \mu\text{M}$ and total phosphorus was $0.4 \mu\text{M}$. The annual average N:P of the photic zone was therefore about 24, but average N:P of the photic zone during individual cruises was as high as 120 and was > 16 on 8 out of 10 dates. Karl et al. (1995) have also discussed the lack of evidence for N limitation shown in these samples. Although Karl et al. (1993) suggest that the average N:P (considering dissolved forms alone) does not depart from the Redfield ratio, more recent work shows that photic zone TDN:TDP ratios in the photic zone are strongly skewed toward high, non-Redfield ratios (David M. Karl, pers. comm.).

In spite of the high total N:P found at this site, traditional analyses of dissolved inorganic N:P would have suggested N limitation. The average dissolved inorganic N:P in the upper 200 m was usually < 5 . The particulate and dissolved organic fractions, however, had N:P as high as 200:1 in many samples. The fact that N:P was very high in particulate matter further counter-indicates N-limitation at this open-ocean site since much of the particulate matter may be living plankton.

Other studies also show that particulate and dissolved organic N:P can be very high at a variety of oligotrophic sites in the open sea (Copin-Montégut & Copin-Montégut 1983; Jackson & Williams 1985). The organic N and P pools are therefore essential to our understanding of N and P cycling in the seas but measurements of the size of these pools and their dynamics have been beyond the reach of standard marine chemical methodology. We therefore need much more extensive knowledge of the total stoichiometry of marine ecosystems.

Differences in the total N:P stoichiometry among marine ecosystems may reconcile the conflicting results obtained in nutrient bioassays. Many such analyses have experimentally enriched nutrients in natural seawater samples to see which nutrients lead to increased phytoplankton growth (Table 2). In polluted estuaries and bays, where total N:P ratios are frequently low (Figure 3), such bioassays usually find that N addition leads to greatest increases in

algal activity (Granéli 1987; Le Rouzic & Bertru 1992; Ryther & Dunstan 1971), although some coastal systems may vary seasonally from N to P limitation (D'Elia et al. 1986; D'Elia et al. 1992; Fisher et al. 1992; Rinne & Tarkiainen 1975) depending upon the N:P of inflowing waters. Less polluted coastal waters also show N limitation (Edmondson 1956; Smayda 1974; Tarkiainen et al. 1974) but addition of both N and P together often results in greater phytoplankton growth than addition of N alone (Granéli 1984; Rudek et al. 1991; Vince & Valiela 1973). The N:P supply ratio may therefore be very close to 16:1 in some less polluted coastal systems. In the open ocean, however, where I found total N:P generally >16 (Figure 3), nitrogen is rarely the principal nutrient limiting phytoplankton abundance or activity (Table 2). Instead, nutrient bioassays usually find that production is limited by phosphorus (Berland et al. 1980; Berland et al. 1987; Bonin et al. 1989; Lapointe 1986), iron (DiTullio et al. 1993; Martin & Fitzwater 1988; Menzel & Ryther 1961; Menzel et al. 1963; Ryther & Guillard 1959; Tranter & Newell 1963), or silica (Smayda 1971). Nitrogen is probably rarely the principal element limiting phytoplankton in the open oceans because its supply appears to be ample relative to phosphorus.

Global N:P model

The combined patterns of variation of total N:P ratios in marine and fresh-water environments provides a model of the global N:P cycle. In upstream freshwaters, where input is mostly derived from high N:P precipitation and high N:P run-off from undisturbed soils (Downing & McCauley 1992), P concentrations are low, N:P is high (Figure 4) and production is strongly P-limited. As water moves downstream, it is enriched by high P, low N:P run-off from terrestrial systems, P increases and N:P declines, resulting in frequent N-limitation of primary production (Elser et al. 1990) and blooms of N-fixing cyanobacteria (Smith 1983). This enrichment may proceed to a varying degree, depending upon the size, land use, and human inhabitation in the drainage system (Peierls et al. 1991).

When nutrient laden rivers flow into coastal marshes and estuaries, anoxic sediments and organic matter are usually abundant, leading to rapid denitrification (Seitzinger 1988), which may explain the drop in N:P as freshwater discharge passes through estuarine environments (Figure 4). As coastal waters move out into open seas, P may be sedimented more rapidly than N, as evidenced by low N:P in deep waters (Figures 2–3), marine sediments and sedimentary rocks (Downing & McCauley 1992). Phosphorus concentrations decline more rapidly than N concentrations in open seas, perhaps because atmospheric N_2 can be fixed in surface waters (Capone & Carpenter 1982;

Table 2. Examples of nutrient enrichment studies arranged in approximately increasing order of expected total N:P, based on the pattern observed in Figure 3. Elements that increased algal growth, production or biomass are listed in decreasing order of their importance (most important first), as perceived by the authors of the studies. "N+P" indicates that adding N and P together increased biomass over addition of N or P alone.

Place	Reference	Limiting elements
Polluted, enclosed estuary: Gulf of Morbihan, France	Le Rouzic & Bertru (1992)	N, N+P
Shallow, polluted Long Island Bays	Ryther & Dunstan (1971)	N
Brackish Laholm Bay	Granéli (1987)	N
Chesapeake Bay	Fisher et al. (1992)	N, P*
Chesapeake Bay	D'Elia et al. (1986; 1992)	N, P†
Neuse River Estuary, North Carolina.	Rudek et al. (1991)	N, N+P
New York Harbor, polluted Long Island Sound	Ryther & Dunstan (1971)	N
Lower Narragansett Bay	Smayda (1974)	N
Polluted Baltic near Helsinki	Rinne & Tarkiainen (1975)	N, P*
Baltic near Helsinki	Tarkiainen et al. (1974)	N, P‡
Woods Hole Harbor	Edmondson (1956)	N, P
Woods Hole Harbor	Vince & Valiela (1973)	N, N+P
Oresund narrows near Copenhagen	Granéli (1984)	N, N+P
Ocean between Bermuda and Puerto Rico	Smayda (1971)	N, P, Si
Open Ocean between Montauk Pt. and Bermuda	Ryther & Guillard (1959)	Fe, N+P
Indian Ocean	Tranter & Newell (1963)	Fe, N+P
Sargasso Sea	Menzel & Ryther (1961)	Fe, N+P
Sargasso Sea	Ryther & Guillard (1959)	Fe, N+P
Sargasso Sea	Menzel et al. (1963)	Fe, N+P
Sargasso Sea, Sargassum	Lapointe (1986)	P
Subarctic Pacific	Martin & Fitzwater (1988)	Fe
Mediterranean Sea	Bonin et al. (1989)	P
Mediterranean Sea	Berland et al. (1977)	P
Mediterranean Sea	Berland et al. (1987)	P, N+P

* changes seasonally. † changes according to N:P of inflowing waters. ‡ usually N limited.

Smith & Veeh 1989) and aeolian N deposition can be substantial (Fanning 1989). Surface N:P ratios can increase dramatically in offshore marine environments, leading to frequent limitation of production by P or nutrients other than N.

Taken together, the results of this study and those of Downing & McCauley (1992) suggest the form of general longitudinal N:P profiles of temperate zone rivers in developed countries (Figure 5A). Upland, headwater streams should have high N:P that largely reflect nutrient ratios in precipitation. Thus, in pristine areas, N:P ratios should be very high because N in rain is enriched by atmospheric fixation and there should be little P in aeolian deposition. N:P in precipitation is sometimes lower in urbanized or agricultural areas, and may alter headwater N:P (Loehr 1974). Downstream freshwaters will usually be




Figure 4. Observed global aquatic N:P cycle. Average trends shown are locally weighted sequentially smoothed fits to the actual observations. Trends were determined using LOWESS, a model-free, unbiased method for finding average trends in data (Cleveland & McGill 1985). LOWESS parameters were $\Delta = 0$, $n\text{-steps} = 2$, $f = 0.5$. The freshwater line is from Downing & McCauley (1992) and the marine line is fitted to the data in Figure 1.

richer in both N and P but have lowered N:P unless P is selectively removed or N is enriched.

A conservative mixing of two solutions that have different relative concentrations of two elements results in a linear relationship between the concentrations of the two elements in all possible admixtures of the two solutions. Therefore, if the variations in observed N:P are simply due to conservative enrichment of rainwater by polluted waters or the dilution of river water by dilute oceanic waters, relationships between N and P should be approximately linear across marine and freshwater systems. In freshwater systems, there is a clear change in mechanism of relative N and P enrichment as one moves from headwaters to downstream systems (Figure 5B). Since the slope of the relationship is directly related to the N:P of the enriching solution, this pattern is suggestive of a progressive decrease in the N:P of effluents. In marine systems however, the decline in N concentration with decreasing P in open oceans is nearly linear, suggesting that variations in marine N:P may be related to the conservative dilution of freshwater effluent by oceanic waters.

This study adds to the mounting evidence that productivity is limited by elements other than N in many of the world's open oceans (Martin & Fitzwater 1988; Martin et al. 1994). These data imply that $N:P > 16$ is found




Figure 5. Predicted gradient in total N and P along the axis of rivers in temperate zone developed countries. Trends are actual LOWESS relationships (see Figure 4) in data from upstream and downstream lakes (Downing & McCauley 1992) and inshore vs. offshore marine ecosystems. (A) shows trends in N:P while (B) shows trends in N concentration. Data are arranged in order of increasing enrichment (0–20 μM P) in freshwaters and in order of decreasing enrichment (20–0 μM P) in marine systems.

most frequently below total P concentrations of 1 μM . Broad-scale surveys of total P in marine environments suggest that concentrations >1 μM may be rare away from shore or upwellings (Gibbs et al. 1986; Rochford 1958). The concept of nitrogen limitation of marine productivity and carbon cycling may therefore have its principal relevance to the shallow and coastal areas of the world's oceans where waters are polluted by low N:P freshwaters.

One might correctly suggest that <200 sets of observations of total N and total P can scarcely be expected to represent all of the world's oceans. However, it is significant that these data are all of the total N and P inventories that I could discover in an extensive literature review and letter writing campaign. This is the most extensive collection of data on total N and P stoichiometry in marine waters. It strongly suggests that open ocean nutrient ratios do not support the marine nitrogen limitation model.

Acknowledgements

I thank W.L. Downing, E. Gorham, R.W. Howarth, D.M. Karl, E. McCauley, M. Pace, R. Peters, L. Pommeroy, E.L. Schmidt, S.V. Smith, V.H. Smith, D. Tilman and an anonymous reviewer for comments on drafts of the manuscript, G. Copin-Montégut and P.M. Williams for encouragement on this project, T.E. Jordan for sharing raw data, and D. Tilman for providing space, research facilities and discussions during my sabbatical leave at University of Minnesota. Supported by an operating grant from the Natural Sciences and Engineering Research Council of Canada.

References

- Amon RMW & Benner R (1994) Rapid cycling of high-molecular-weight dissolved organic matter in the ocean. *Nature* (London) 369: 549–552
- Ben-Taleb KS, El-Khattal AO, Ramadan ZM & El-Belaazi AM (1987) Some physico-chemical characteristics of Tajura sea-shore water: East of Tripoli (S.P.L.A.J.) *Bulletin of the Marine Biological Research Center of Tajura* 8: 89–124
- Bentzen E, Taylor WD & Millard ES (1992) The importance of dissolved organic phosphorus to phosphorus uptake by limnetic plankton. *Limnol. and Oceanography* 37: 217–231
- Berland BR, Bonin DJ & Maestrini SY (1977) Role du phosphore dans la limitation de la production primaire dans les eaux Méditerranéennes. *Actualités de Biochimie Marine* 2: 243–246
- Berland BR, Burlakova ZP, Georgieva LV, Izmetstieva M, Khologov VI, Krupatkina DK, Maestrini SY & Zaika VE (1987) Phytoplankton estival de la mer du Levant, biomasse et facteurs limitants. *IFREMER Acta Colloquia* 5: 61–83
- Black RE, Lukatelich RJ, McComb AJ & Rosher JE (1981) Exchange of water, salt, nutrients and phytoplankton between Peel Inlet, Western Australia, and the Indian Ocean. *Australian J. Marine and Freshwater Res.* 32: 709–720
- Bonin DJ, Bonin MC & Berman T (1989) Mise en évidence expérimentale des facteurs nutritifs limitants de la production du micro-nanoplancton et de l'ultraplancton dans une eau cotière de la Méditerranée orientale (Haïfa, Israël). *Aquatic Sciences* 51(2): 129–152
- Boynton WR, Kemp WM & Keefe CW (1982) A comparative analysis of nutrients and other factors influencing estuarine phytoplankton production. In: Kennedy VS (Ed) *Estuarine Comparisons* (pp 69–70). Academic, NY
- Capone D & Carpenter EJ (1982) Nitrogen fixation in the marine environment. *Science* 217: 1140–1142
- Cleveland W & McGill SR (1985) Graphical perception and graphical methods for analyzing scientific data. *Science* 229: 828–833
- Copin-Montégut C & Copin-Montégut G (1983) Stoichiometry of carbon, nitrogen, and phosphorus in marine particulate matter. *Deep-Sea Research* 30: 31–46
- Coste B, Le Corre P & Minas HJ (1988) Re-evaluation of the nutrient exchanges in the Strait of Gibraltar. *Deep-Sea Research* 35: 767–775
- D'Elia CF, Steudler PA & Corwin N (1977) Determination of total nitrogen in aqueous samples using persulfate digestion. *Limnology and Oceanography* 22: 760–764
- D'Elia CF, Sanders JG & Boynton WR (1986) Nutrient enrichment studies in a coastal plain estuary: Phytoplankton growth in large-scale, continuous cultures. *Can. J. Fisheries and Aquatic Sci.* 43: 397–406
- D'Elia CF, Harding LW Jr., Leffler M & Mackiernan GB (1992) The role and control of nutrients in Chesapeake Bay. *Water Science and Technology* 26: 2635–2644

- Dal Pont GK, Hogan M & Newell B (1974) Laboratory techniques in marine chemistry II – a manual. CSIRO Australia Division of Fisheries and Oceanography, Report 55
- Dame R, Chrzanowski T, Bildstein K, Kjerfve B, McKellar H, Nelson D, Spurrier J, Stancyk S, Stevenson H, Vernberg J & Zingmark R (1986) The outwelling hypothesis and North Inlet, South Carolina. *Marine Ecology Progress Series* 33: 217–229
- Degobbi D & Gilmartin M (1990) Nitrogen, phosphorus, and biogenic silicon budgets for the northern Adriatic Sea. *Oceanologica Acta* 13: 31–45
- DiTullio GR, Hutchins DA & Bruland KW (1993) Interaction of iron and major nutrients controls phytoplankton growth and species composition in the tropical North Pacific Ocean. *Limnol. and Oceanography* 38: 495–508
- Downing JA & McCauley E (1992) The nitrogen:phosphorus relationship in lakes. *Limnol. and Oceanography* 37: 936–945
- Edmondson WT (1956) Factors affecting productivity in fertilized salt water. *Deep-Sea Research Supplement* 3: 451–464
- Elser JJ & Hassett RP (1994) A stoichiometric analysis of the zooplankton-phytoplankton interaction in marine and freshwater ecosystems. *Nature* 370: 211–213
- Elser JJ, Marzolf ER & Goldman CR (1990) Phosphorus and nitrogen limitation of phytoplankton in freshwaters of North America: A review and critique of experimental enrichments. *Can. J. Fisheries and Aquatic Sci.* 47: 1468–1477
- Fanning KA (1989) Influence of atmospheric pollution on nutrient limitation in the ocean. *Nature* 339: 460–463
- FAO (1993) *Fishery Statistics Yearbook*, vol. 72 Rome.
- Fisher TR, Peele ER, Ammerman JW & Harding LW Jr (1992) Metabolic adaptations of deep-sea benthic foraminifera to seasonally varying food input. *Marine Ecology Progress Series* 82: 51–63
- Gibbs CF, Tomczak M & Longmore AR (1986) The nutrient regime of Bass Strait. *Australian J. Marine and Freshwater Res.* 37: 451–466
- Glibert PM, Garside C, Fuhrman JA & Roman MR (1991) Time-dependent coupling of inorganic and organic nitrogen uptake and regeneration in the plume of the Chesapeake Bay estuary and its regulation by large heterotrophs. *Limnol. and Oceanography* 36: 895–909
- Granéli E (1984) Algal growth potential and limiting nutrients for phytoplankton production in Öresund water of Baltic and Kattegat origin. *Limnologica* 15: 563–569
- Granéli E (1987) Nutrient limitation of phytoplankton biomass in a brackish water bay highly influenced by river discharge. *Estuarine, Coastal and Shelf Science* 25: 555–565
- Grasshoff K (1976) *Methods of Seawater Analysis*. 1st edn. Verlag Chemie Weinheim
- Grasshoff K (1983) *Methods of Seawater Analysis*. 2nd edn. Verlag Chemie Weinheim
- Hecky RE & Kilham P (1988) Nutrient limitation of phytoplankton in freshwater and marine environments: A review of recent evidence on the effects of enrichment. *Limnol. and Oceanography* 33: 796–822
- Hoegh-Goldberg O (1994) Uptake of dissolved organic matter by larval stage of the crown-of-thorns starfish *Acanthaster planci*. *Marine Biology* 120: 55–63
- Howarth RW (1988) Nutrient limitation of net primary production in marine ecosystems. *An. Rev. of Ecology* 19: 89–110
- Howarth RW, Jensen H, Marino R & Postma H (1995) Transport to and processing of phosphorus in near-shore and oceanic waters (pp 323–345). In: Tiessen H (Ed) *Phosphorus in the Global Environment: Transfers, Cycles and Management*. SCOPE 54, Wiley & Sons, Chichester
- Ikeda T, Carleton JH, Mitchell AW & Dixon P (1982) Ammonia and phosphate excretion by zooplankton from the inshore waters of the Great Barrier Reef. II. Their *in situ* contributions to nutrient regeneration. *Australian J. Marine and Freshwater Res.* 33: 683–698
- Jackson GA & Williams PM (1985) Importance of dissolved organic nitrogen and phosphorus to biological nutrient cycling. *Deep-Sea Research* 32: 223–235

- Jordan TE, Correll DL, Miklas J & Weller DE (1991) Nutrients and chlorophyll at the interface of a watershed and an estuary. *Limnol. and Oceanography* 36: 251–267
- Karl DM, Leteller R, Hebel D, Tupas L, Dore J, Christian J & Winn C (1995) Ecosystem changes in the North Pacific subtropical gyre attributed to the 1991–92 El Niño. *Nature* (London) 373: 230–234
- Karl DM, Tien G, Dore J & Winn CD (1993) Total dissolved nitrogen and phosphorus concentrations at US-JGOFS station ALOHA: Redfield reconciliation. *Marine Chemistry* 41: 203–208
- Krom MD, Kress N, Brenner S & Gordon LI (1991) Phosphorus limitation of primary productivity in the eastern Mediterranean Sea. *Limnol. and Oceanography* 36: 424–432
- Lahdes E & Leppanen JM (1988) Cycling of organic matter during the vernal growth period in the open Northern Baltic proper. II. Nutrient development and chemical composition of particulate matter. *Finnish Marine Research* 255: 19–35
- Lapointe BE (1986) Phosphorus-limited photosynthesis and growth of *Sargassum natans* and *Sargassum fluitans* (Phaeophyceae) in the western North Atlantic. *Deep-Sea Research* 33: 391–399
- Le Rouzic & Bertru BG (1992) Détermination par bioessai de la biodisponibilité des ressources azote et phosphore, dans les eaux du Golfe du Morbihan. *Revue des Sciences de l'Eau* 5: 97–111
- Mackenzie FT, Ver LM, Sabine C, Lane M & Lerman A (1993) C, N, P, S global biogeochemical cycles and modeling of global change. In: Wollast R, Mackenzie FT & Chou L (Eds) *Interactions of C, N, P and S, Biogeochemical Cycles and Global Change* (pp 1–61). Springer-Verlag, Berlin
- Martin JH & Fitzwater SE (1988) Iron deficiency limits phytoplankton growth in the north-east pacific subarctic. *Nature* 331: 341–343
- Martin JH, Coale KH, Johnson KS, Fitzwater SE, Gordon RM, Tanner SJ, Hunter CN, Elrod VA, Nowicki JL, Coley TL, Barber RT, Lindley S, Watson AJ, Van Scoy K, Law CS, Liddicoat MI, Ling R, Stanton T, Stockel J, Collins C, Anderson A, Bidigare R, Ondrusek M, Latasa M, Millero FJ, Lee K, Yao W, Zhang JZ, Friederich G, Sakamoto C, Chavez F, Buck K, Kolber Z, Greene R, Falkowski P, Chisholm SW, Hoge F, Swift R, Yungel J, Turner S, Nightingale P, Hatton A, Liss P & Tindale NW (1994) Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean. *Nature* 371: 123–129
- Menzel DW & Corwin N (1965) The measurement of total phosphorus in seawater based on the liberation of organically bound fractions by persulfate oxidation. *Limnol. and Oceanography* 10: 280–282
- Menzel DW & Ryther JH (1961) Nutrients limiting the production of phytoplankton in the Sargasso Sea, with special reference to iron. *Deep-Sea Research* 7: 276–281
- Menzel DW, Hulburt EM & Ryther JH (1963) The effects of enriching Sargasso Sea water on the production and species composition of the phytoplankton. *Deep-Sea Research* 10: 209–219
- Meybeck M (1982) Carbon, nitrogen, and phosphorus transport by world rivers. *Am. J. Sci.* 282: 401–450
- Peierls B, Caracao N, Pace M & Cole J (1991) Human influence on river nitrogen. *Nature* 350: 386–387
- Perttilä MP, Tulkki P & Pietikainen S (1980) Mean values and trends of hydrographical and chemical properties in the Gulf of Finland 1962–1978. *Finnish Marine Research* 247: 38–50
- Pietikainen S, Niemi A, Tulkki P & Aurimaa K (1978) Mean values and trends of physical and chemical properties in the Gulf of Bothnia 1962–1975. *Finnish Marine Research* 244: 64–75
- Pitkänen H & Malin V (1980) The mean values and trends of some water quality variables in winter in the Gulf of Finland 1966–1978. *Finnish Marine Research* 247: 51–60

- Ramadan ZM, Ben-Taleb KS & Trozosinska A (1984) Ecological conditions in the Mediterranean coastal zone. A case study: Fishery Harbour of Tripoli, Libya. *Bulletin of the Marine Research Center of Tripoli* 5: 69–106
- Redfield AC (1934) On the proportions of organic derivatives in sea water and their relation to the composition of plankton. In: James Johnstone Memorial Volume (pp 176–192). Univ. Press, Liverpool
- Redfield AC (1958) The biological control of chemical factors in the environment. *Am. Sci.* 46: 205–221
- Redfield AC, Ketchum BH & Richards FA (1963) The influence of organisms on the composition of sea-water. In: Hill MN (Ed) *The Sea*, vol. 2 (pp 26–77) Wiley, NY
- Rinne IE & Tarkiaainen E (1975) Chemical factors affecting algal growth off Helsinki. *Merentutkimuslait. Julk. Havsforskningsinst. Skr.* 239: 91–99
- Ritschard RL (1992) Marine algae as a CO₂ sink. *Water Air Soil Poll.* 64: 289–303
- Rochford D (1958) Total phosphorus as a means of identifying East Australian water masses. *Deep-Sea Research* 5: 89–110
- Rudek J, Paerl HW, Mallin MA & Bates PW (1991) Seasonal and hydrological control of phytoplankton nutrient limitation in the lower Neuse River Estuary, North Carolina. *Marine Ecology Progress Series* 75: 133–142
- Rydberg L & Sundberg J (1986) Seasonal nutrient supply to coastal waters. In: *Marine Interfaces Ecohydrodynamics*, 17th Annual Liege Colloquium on Ocean Hydrodynamics (pp 467–485). Elsevier Oceanography Series 42
- Ryther JH & Dunstan WM (1971) Nitrogen, phosphorus, and eutrophication in the coastal marine environment. *Science* 171: 1008–1013
- Ryther JH & Guillard RRL (1959) Enrichment experiments as a means of studying nutrients limiting to phytoplankton production. *Deep-Sea Research* 6: 65–69
- Sakamoto M (1966) Primary production by phytoplankton community in some Japanese lakes and its dependence on lake depth. *Archiv für Hydrobiologie* 62: 1–28
- Seitzinger SP (1988) Denitrification in freshwater and coastal marine ecosystems: Ecological and geochemical significance. *Limnol. and Oceanography* 33: 702–724
- Smayda TJ (1974) Bioassay of the growth potential of the surface water of lower Narragansett Bay over an annual cycle using the diatom *Thalassiosira pseudonana* (oceanic clone, 13-1). *Limnol. and Oceanography* 19: 889–901
- Smith SV (1984) Phosphorus vs. nitrogen limitation in the marine environment. *Limnol. and Oceanography* 29: 1149–1160
- Smith SV & Veeh HH (1989) Mass balance of biogeochemically active materials (C, N, P) in a hypersaline gulf. *Estuarine, Coastal and Shelf Science* 29: 195–215
- Smith SV, Kimmerer WJ & Walsh TW (1986) Vertical flux and biogeochemical turnover regulate nutrient limitation of net organic production in the North Pacific Gyre. *Limnol. and Oceanography* 31: 161–167
- Smith VH (1983) Low nitrogen to phosphorus ratios favor dominance by blue-green algae in lake phytoplankton. *Science* 221: 669–671
- Strickland JH & Parsons TR (1968) *A Practical Handbook of Seawater Analysis*. Fisheries Res. Board Canada, Ottawa
- Strickland JDH & Parsons TR (1972) *A Practical Handbook of Seawater Analysis*. Fisheries Res. Board Canada 167
- Suzuki Y, Sugimura Y & Ito T (1985) A catalytic oxidation method for the determination of total nitrogen dissolved in seawater. *Marine Chemistry* 16: 83–97
- Smayda TJ (1971) Further enrichment experiments using the marine centric diatom *Cyclotella nana* (clone 13-1) as an assay organism. In: Costlow JD (Ed) *Fertility of the Sea*, vol. 2 (pp 493–511). Gordon and Breach, NY
- Tarkiaainen E, Rinne I & Niemisto L (1974) On the chemical factors regulating the primary production of phytoplankton in the Baltic proper. *Merentutkimuslait. Julk. Havsforskningsinst. Skr.* 238: 39–52

- Tett P (1990) The photic zone. In: Herring PJ, Campbell AK, Whitfield M & Maddock L (Eds) *Light and Life in the Sea* (pp 59–87). Cambridge
- Tranter DJ & Newell BS (1963) Enrichment experiments in the Indian Ocean. *Deep-Sea Research* 10: 1–9
- Treguer P & Le Corre P (1975) *Manuel d'analyse des sels nutritifs dans l'eau de mer*. Lab. Océanogr. Chimique, Univ. Bretagne Occident. Brest
- Updegraff DM, Stanton DJ & Spencer MJ (1977) Surface waters of Waimea Inlet and Nelson Haven: A preliminary assessment of quality. *New Zealand Journal of Marine and Freshwater Research* 11: 559–575
- Valderrama JC (1981) The simultaneous analysis of total nitrogen and total phosphorus in natural waters. *Marine Chemistry* 10: 109–122
- Vince S & Valiela I (1973) The effects of ammonium and phosphate enrichments on chlorophyll a, pigment ratio, and species composition of phytoplankton of Vineyard Sound. *Marine Biology* 19: 69–73
- Vitousek PM & Howarth RW (1991) Nitrogen limitation on land and in the sea: How can it occur? *Biogeochem.* 13: 87–115
- Williams PM (1967) Sea surface chemistry: Organic carbon and organic and inorganic nitrogen and phosphorus in surface films and subsurface waters. *Deep-Sea Research* 14: 791–800
- Winn C, Lukas R, Karl D & Firing E (1993) *Hawaii Ocean Time-series Data Report 3, 1991*. University of Hawaii, Honolulu