
Social common mole-rats enhance outbreeding via extra-pair mating

Jacqueline M Bishop, Colleen O'Ryan and Jennifer U.M Jarvis

Biol. Lett. 2007 **3**, 176-179
doi: 10.1098/rsbl.2006.0607

References

This article cites 20 articles, 1 of which can be accessed free
<http://rsbl.royalsocietypublishing.org/content/3/2/176.full.html#ref-list-1>

Article cited in:
<http://rsbl.royalsocietypublishing.org/content/3/2/176.full.html#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Social common mole-rats enhance outbreeding via extra-pair mating

Jacqueline M. Bishop^{1,*}, Colleen O’Ryan² and Jennifer U. M. Jarvis¹

¹Department of Zoology, and ²Department of Molecular and Cell Biology, University of Cape Town, Private Bag X3, Rondebosch 8001, South Africa

*Author for correspondence (jbishop@botzoo.uct.ac.za).

Females in many species engage in matings with males that are not their social mates. These matings are predicted to increase offspring heterozygosity and fitness, and thereby prevent the deleterious effects of inbreeding. We tested this hypothesis in a cooperative breeding mammal, the common mole-rat *Cryptomys hottentotus hottentotus*. Laboratory-based studies suggested a system of strict social monogamy, while recent molecular studies indicate extensive extra-pair paternity despite colonies being founded by an outbred pair. Our data show that extra-pair and within-colony breeding males differed significantly in relatedness to breeding females, suggesting that females may gain genetic benefits from breeding with non-resident males. Extra-colony male mating success was not based on heterozygosity criteria at microsatellite loci; however, litters sired by extra-colony males exhibited increased heterozygosity. While we do not have the data that refute a relationship between individual levels of inbreeding (H_s) and fitness, we propose that a combination of both male and female factors most likely explain the adaptive significance of extra-pair mating whereby common mole-rats maximize offspring fitness by detecting genetic compatibility with extra-pair mates at other key loci, but it is not known which sex controls these matings.

Keywords: relatedness; genetic compatibility; microsatellite; extra-pair paternity; mate choice; *Cryptomys*

1. INTRODUCTION

While evidence abounds demonstrating that females of many species produce young via extra-pair males, the adaptive significance for this behaviour still remains unclear. A number of studies suggest that further to the acquisition of food resources, parental care and ‘good genes’, selection for extra-pair mates may primarily be driven by genetic compatibility and the costs of inbreeding (Tregenza & Weddell 2000; Griffith *et al.* 2002; Wolff & Macdonald 2004). Negative fitness consequences associated with inbreeding are well documented (Thornhill 1993; Coltman *et al.* 1998; Keller & Waller 2002; Bean *et al.* 2004) and evidence for individual levels of

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2006.0607> or via <http://www.journals.royalsoc.ac.uk>.

inbreeding as fitness predictors is accumulating (Hansson *et al.* 2001; Hansson & Westerberg 2002; Foerster *et al.* 2003). By mating with males that have dissimilar alleles, females increase levels of heterozygosity in their offspring; this may preclude the effects of deleterious alleles and probably increases heterozygosity at loci that enhance fitness, e.g. the major histocompatibility (MHC) gene complex (Petrie & Kempenaers 1998; Tregenza & Weddell 2000).

The common mole-rat (*Cryptomys hottentotus hottentotus*) is widely distributed in South Africa, where colonies comprising two to fourteen individuals permanently occupy a discrete network of burrows, locating food as they blindly extend foraging tunnels (Spinks *et al.* 2000). Laboratory-based studies suggested that a system of strict social monogamy characterized the mating system of this species, and mark-recapture field studies confirmed that colonies were founded by unrelated pairs (Bennett 1989, 1992). Nevertheless, microsatellite data collected from our field study populations revealed unexpectedly high levels of paternity assigned to males that were not colony residents (extra-colony males, ECMs; Bishop *et al.* 2004). The common mole-rat is a cooperative breeder that lives in colonies characterized by a generally linear social hierarchy, where the breeding male is the most dominant individual (Bennett 1989). Therefore, why should females mate with males that do not contribute to the maintenance and provisioning of her resident colony? In many social species, constraints on mate choice may result in females paired with genetically similar males (Cohas *et al.* 2006). Subordinate common mole-rat females do not breed within their home colony. Instead, they appear to delay reproduction until such time that ecological conditions favour dispersal and independent reproduction (Spinks *et al.* 2000); this may result in social pairings between genetically suboptimal mates, which in turn may favour extra-pair matings.

In this study, we were interested in whether relatedness is an important factor influencing extra-pair mating patterns in the common mole-rat. If female mole-rats engage in extra-pair matings to enhance the fitness of their offspring, we can predict that (i) females would choose ECM that were less genetically similar than within-pair males, (ii) ECM would have higher levels of heterozygosity than within-pair males, and (iii) offspring sired by extra-pair males would have higher levels of heterozygosity than those sired by within-pair males. Because size is an important determinant of male social status, we also investigated whether any difference exists in the size of colony breeding males and those ECM that succeeded in gaining paternity. Resident breeding males within a colony are most probably the largest male or one of the largest males (Bishop *et al.* 2004); subordinate males, however, are fully reproductively functional and merely behaviourally quiescent (Spinks *et al.* 1997).

2. MATERIAL AND METHODS

We studied 13 colonies of the common mole-rat at two sites representing the broad ecological range of the species. Individuals were live-trapped using modified Hickman traps from September 1992 to November 1996; all individuals were sexed, weighed and marked as part of long-term mark-recapture studies (detailed in Spinks *et al.* 2000). Seven microsatellite loci were used to

investigate paternity for a total of 73 offspring collected from the two study sites (detailed in Bishop *et al.* 2004); the loci had a combined exclusion probability of more than 0.95 and the mean probability of identity (P_{ID}) for potential sires in the two populations was 2.1×10^{-4} (see electronic supplementary material). In contrast to the suggested mating system of strict monogamy, ECM were assigned paternity with high levels of confidence to approximately 30% of offspring at both sites (Bishop *et al.* 2004). In this study, we investigate measures of genetic similarity and heterozygosity between females and within-colony breeding males (WCMs), as well as ECMs to which paternity was assigned. WCMs were those males known to have sired young within their resident colony. As a measure of individual inbreeding, we calculated standardized heterozygosity (H_s ; proportion of heterozygous loci/mean heterozygosity of typed loci; Coltman *et al.* 1999). Measures of relatedness (R) within and among colonies and between females and males were calculated using genotypic data from seven microsatellite loci in RELATEDNESS v. 5.0.8 (Queller & Goodnight 1989). Average relatedness among adult males and females in our study populations (expected $R=0$) was 0.03 ± 0.0009 s.e.m. calculated by jackknifing; in a randomized sample of 100 litters, the average relatedness among full siblings (expected $R=0.5$) was 0.45 ± 0.01 . All test statistics were performed in STATISTICA v. 7.0 (StatSoft Inc.).

3. RESULTS AND DISCUSSION

Our results suggest that common mole-rats enhance outbreeding by engaging in extra-colony matings. Females were generally less related to ECMs than to resident WCMs with whom they sired young (figure 1; WCMs, mean R to females \pm s.e.m. = 0.17 ± 0.06 ; ECMs, mean $R = -0.05 \pm 0.06$; one-tailed paired t -test, $t = 2.19$, d.f. = 17, $p < 0.025$). Furthermore, the mean difference in R between females and ECMs was found to be significantly greater than the observed differences in R within and among colonies of mole-rats (see electronic supplementary material; two-sample randomization test, $p = 0.04$), suggesting that extra-pair mating occurs non-randomly with respect to levels of non-kin relatedness. Overall however, females who mated with non-resident males were not more related to their colony breeding male compared with females whose litters were only sired by the colony male (mean R of females to colony males with extra-colony paternity \pm s.e.m. = 0.22 ± 0.08 ; mean R of females to colony males without extra-colony paternity \pm s.e.m. = 0.15 ± 0.07 ; t -test, $t = 0.51$, d.f. = 19, $p = 0.61$). This might be because a sample of individuals at any point in time will represent both established and recently founded colonies in which variation in turnover of reproductive females will influence opportunities for mating with ECMs. Reproductive tenure within colonies will provide some females greater opportunity to mate with ECMs, while females that had recently founded a colony would not necessarily engage in extra-pair matings immediately; unfortunately, it is not clear from our data whether older females with long-term reproductive tenure were more likely to engage in extra-colony paternity than young, recently dispersed and mated females.

Do females mate with ECM that are more heterozygous than their colony mates? Our results suggest that ECM mating success is not based on heterozygosity criteria at neutral loci, such as microsatellites. Overall, males that lost paternity to ECMs did not differ in heterozygosity from those that sired all their colony offspring (mean $H_s \pm$ s.e.m. of cuckolded males = 0.61 ± 0.05 , mean H_s of non-cuckolded males = $0.56 \pm$




Figure 1. Distribution of pairwise estimates of relatedness between females and their colony mates versus extra-colony males. R was calculated according to Queller & Goodnight (1989).

0.05; Mann-Whitney U -test, $n = 12, 19$; $U = 103$, $Z = 0.446$, $p = 0.65$), and colony males did not differ in microsatellite heterozygosity compared with their cuckolders (mean $H_s \pm$ s.e.m. of WCMs = 0.62 ± 0.04 , mean H_s of ECMs = 0.61 ± 0.03 ; Wilcoxon's matched pairs test, $n = 17$, $Z = 0.21$, $p = 0.83$; figure 2a). Nevertheless, litters from extra-pair matings displayed increased levels of heterozygosity; mean H_s for litters sired by ECMs was significantly greater than for those sired by WCMs (mean $H_s \pm$ s.e.m. of ECM litters = 0.65 ± 0.03 , mean H_s of WCM litters = 0.55 ± 0.02 ; one-tailed t -test, $t = -2.03$, d.f. = 35, $p = 0.025$). As a result, female mole-rats appear to maximize offspring heterozygosity by mating with genetically unrelated males rather than heterozygous males. At present, we do not have data that associate microsatellite heterozygosity in mole-rats with measures of individual fitness, but we cannot exclude the possibility that females seek increased variation and/or compatibility at additional paternal loci known to influence fitness, and measures of R may reflect differences between males at various important loci under moderate or strong selection (Reed & Frankham 2001).

Finally, dominant males are generally the largest individuals within colonies and are thought more likely to gain access to oestrous females than smaller subordinate males (Bennett 1992). However, our data suggest that WCMs appear to be cuckolded time and again by generally smaller, foreign males (mean WCM weight \pm s.e.m. = $87.7 \text{ g} \pm 2.7$, mean ECM weight = $72.2 \text{ g} \pm 3.3$; Wilcoxon's matched pairs test, $n = 18$, $Z = 2.59$, $p = 0.009$; figure 2b). We believe that this is unlikely to be associated with female choice for size; mole-rats have highly reduced eyes and spend much of their lives within their subterranean burrow systems. They do not use visual cues to determine whether individuals are foreign or conspecifics; instead, they rely on colony odour to identify colony members (Spinks *et al.* 1998). Thus, a more probable explanation is that breeding females simply have a greater chance of encountering subordinate males from foreign colonies; subordinate males work harder, foraging and extending the tunnel system, and are more likely to disperse than established breeding males or indeed breeding females, and these factors may in turn




Figure 2. Pairwise comparisons of colony male and extra-colony male mate choice ($n=18$ dyads). (a) Standardized heterozygosity (H_s) and (b) size of male (g). Dotted line represents values that are identical.

lead to increased encounters with foreign oestrous females. Undoubtedly though, both dominant and subordinate ECMs increase their reproductive fitness by successfully siring offspring in colonies to whose maintenance they do not contribute.

In summary, our results suggest that female common mole-rats enhance outbreeding by mating with non-resident males that are less related to themselves than their social mate. While this behaviour leads to increased offspring heterozygosity, male reproductive success does not appear to be dependent on individual heterozygosity at microsatellite loci. We propose that both males and females maximize offspring fitness by detecting genetic compatibility with extra-pair mates at additional key loci. Given the life history, ecology and reproductive physiology of the species, we can reject a number of alternative hypotheses explaining the adaptive value of extra-pair mating behaviour (reviewed in Wolff & Macdonald 2004). Detecting genetic compatibility requires a mechanism to determine the degree to which extra-pair mates differ from their colony mates. In the absence of visual cues, mole-rats appear to be an excellent model to test whether odour signatures

indicate compatibility between males and females at loci, such as the MHC.

The authors thank J. Emmerson who kindly allowed us to study and trap animals on his property. We are grateful to N. C. Bennett, M. J. O'Riain and A. C. Spinks for their insightful discussions and data collection; F. P. D. Cotterill and three anonymous reviewers provided constructive comments on the manuscript. This work was supported by research grants from the South African National Research Foundation and the Cape Town University Research Council.

Bean, K., Amos, W., Pomeroy, P. P., Twiss, S. D., Coulson, T. N. & Boyd, I. L. 2004 Patterns of parental relatedness and pup survival in the grey seal (*Halichoerus grypus*). *Mol. Ecol.* **13**, 2365–2370. (doi:10.1111/j.1365-294X.2004.02199.x)

Bennett, N. C. 1989 The social structure and reproductive biology of the common mole-rat, *Cryptomys hottentotus hottentotus* and remarks on the trends in reproduction and sociality in the family Bathyergidae. *J. Zool. Lond.* **219**, 45–59.

Bennett, N. C. 1992 Aspects of social behaviour in a captive colony of the common mole-rat, *Cryptomys hottentotus hottentotus*. *Z. Säugertier.* **57**, 294–309.

Bishop, J. M., Jarvis, J. U. M., Spinks, A. C., Bennett, N. C. & O'Riain, C. 2004 Molecular insight into patterns of colony composition and paternity in the common mole-rat *Cryptomys hottentotus hottentotus*. *Mol. Ecol.* **13**, 1217–1229. (doi:10.1111/j.1365-294X.2004.02131.x)

Cohas, A., Yoccoz, N. G., Da Silva, A., Goosens, B. & Allainé, D. 2006 Extra-pair paternity in the monogamous alpine marmot (*Marmota marmota*): the roles of social setting and female mate choice. *Behav. Ecol. Sociobiol.* **59**, 597–605. (doi:10.1007/s00265-005-0086-8)

Coltman, D. W., Bowen, W. D. & Wright, J. M. 1998 Birth weight and neonatal survival of harbour seal pups are positively correlated with genetic variation measured by microsatellites. *Proc. R. Soc. B* **265**, 803–809. (doi:10.1098/rspb.1998.0363)

Coltman, D. W., Pilkington, J. G., Smith, J. A. & Pemberton, J. M. 1999 Parasite-mediated selection against inbred soay sheep in a free-living island population. *Evolution* **53**, 1259–1267. (doi:10.2307/2640828)

Foerster, K., Delhey, K., Johnson, A., Lijfield, J. T. & Kempenaers, B. 2003 Females increase offspring heterozygosity and fitness through extra-pair matings. *Nature* **425**, 714–717. (doi:10.1038/nature01969)

Griffith, S. C., Owens, I. P. F. & Thuman, K. A. 2002 Extra pair paternity in birds: a review of interspecific variation and adaptive function. *Mol. Ecol.* **11**, 2195–2212. (doi:10.1046/j.1365-294x.2002.01613.x)

Hansson, B., Bensch, S., Hasselquist, D. & Åkesson, M. 2001 Microsatellite diversity predicts recruitment of sibling great reed warblers. *Proc. R. Soc. B* **268**, 1287–1291. (doi:10.1098/rspb.2001.1640)

Hansson, B. & Westerberg, L. 2002 On the correlation between heterozygosity and fitness in natural populations. *Mol. Ecol.* **11**, 2467–2474. (doi:10.1046/j.1365-294X.2002.01644.x)

Keller, L. F. & Waller, D. M. 2002 Inbreeding effects in wild populations. *Trends Ecol. Evol.* **17**, 230–241. (doi:10.1016/S0169-5347(02)02489-8)

Petrie, M. & Kempenaers, B. 1998 Extra-pair paternity in birds: explaining variation between species and populations. *Trends Ecol. Evol.* **13**, 52–58. (doi:10.1016/s0169-5347(97)01232-9)

- Queller, D. C. & Goodnight, K. 1989 Estimating relatedness using genetic markers. *Evolution* **43**, 258–275. (doi:10.2307/2409206)
- Reed, D. H. & Frankham, R. 2001 How closely correlated are molecular and quantitative measures of genetic variation? A meta-analysis. *Evolution* **43**, 258–275.
- Spinks, A. C., van der Horst, G. & Bennett, N. C. 1997 Influence of breeding season and reproductive status on male reproductive characteristics in the common mole-rat, *Cryptomys hottentotus hottentotus*. *J. Reprod. Fert.* **109**, 79–86.
- Spinks, A. C., O’Riain, M. J. & Polokow, D. A. 1998 Intercolonial encounters and xenophobia in the common mole-rat, *Cryptomys hottentotus hottentotus* (Bathyergidae): the effects of aridity, sex and reproductive status. *Behav. Ecol.* **69**, 224–234.
- Spinks, A. C., Jarvis, J. U. M. & Bennett, N. C. 2000 Comparative patterns of philopatry and dispersal in two common mole-rat populations: implications for the evolution of mole-rat sociality. *J. Anim. Ecol.* **69**, 224–234. (doi:10.1046/j.1365-2656.2000.00388.x)
- Thornhill, N. W. 1993 *The natural history of inbreeding and outbreeding: theoretical and empirical perspectives*. Chicago, IL: University of Chicago Press.
- Tregenza, T. & Weddell, N. 2000 Genetic compatibility, mate choice and patterns of parentage: invited review. *Mol. Ecol.* **9**, 1013–1027. (doi:10.1046/j.1365-294X.2000.00964.x)
- Wolff, J. O. & Macdonald, D. W. 2004 Promiscuous females protect their offspring. *Trends Ecol. Evol.* **19**, 127–134. (doi:10.1016/j.tree.2003.12.009)