
Cytokinin-mediated leaf manipulation by a leafminer caterpillar

David Giron, Wilfried Kaiser, Nadine Imbault and Jérôme Casas

Biol. Lett. 2007 **3**, 340-343
doi: 10.1098/rsbl.2007.0051

References

This article cites 20 articles, 7 of which can be accessed free
<http://rsbl.royalsocietypublishing.org/content/3/3/340.full.html#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Biol. Lett.* go to: <http://rsbl.royalsocietypublishing.org/subscriptions>

Cytokinin-mediated leaf manipulation by a leafminer caterpillar

David Giron^{1,*}, Wilfried Kaiser¹, Nadine Imbault² and Jérôme Casas¹¹Institut de Recherche sur la Biologie de l'Insecte, UMR-CNRS-6035, and ²EA2106 Biomolécules et Biotechnologies Végétales, UFR Sciences et Techniques, Université François Rabelais, 37200 Tours, France

*Author for correspondence (david.giron@univ-tours.fr).

A large number of hypotheses have been proposed to explain the adaptive significance and evolution of the endophagous-feeding mode, nutritional benefits being considered to be one of the main advantages. Leaf-mining insects should feed on most nutritional tissues and avoid tissues with high structural and/or biochemical plant defences. This selective feeding behaviour could furthermore be reinforced by manipulating the plant physiology, as suggested by the autumnal formation of 'green islands' around mining caterpillars in yellow leaves. The question we address here is how such metabolic manipulation occurs and what the nutritional consequences for the insect are. We report a large accumulation of cytokinins in the mined tissues which is responsible for the preservation of functional nutrient-rich green tissues at a time when leaves are otherwise turning yellow. The analogy with other plant manipulating organisms, in particular galling insects, is striking.

Keywords: leafminer; cytokinins; green island; nutrient acquisition; endophytic organisms

1. INTRODUCTION

Leaf mining is a means by which some plant-eating insects consume live foliage while simultaneously dwelling inside it. A large number of hypotheses have been proposed to explain the adaptive significance and the evolution of this endophagous-feeding mode. The mine presumably functions as a shelter from the detrimental effects of the physical environment, as a protection from attack by natural enemies, enables an increased development rate due to higher temperatures and is potentially a means of making food choices at a micro-scale (Connor & Taverner 1997; Pincebourde & Casas 2006; Pincebourde *et al.* 2006). Following this 'selective feeding hypothesis', many authors have shown that leafminers select those plant tissues with a high nutrient content and avoid those tissues with a high structural and/or chemical defence (Hering 1951; Feeny 1970; Connor & Taverner 1997; Scheirs *et al.* 2001).

Leafminers develop in a small confined space with limited availability of nutritive tissue. They might also strongly suffer from alteration of leaf physiology caused, for example, by premature leaf abscission or

leaf senescence. Indeed, unlike gall-forming insects—another main type of endophytophagous insects—leaf mining does not lead to the proliferation of undifferentiated nutrient-rich plant tissues.

The observation that larval attacks may be responsible for the formation of autumnal 'green islands', green leaf tissues around larvae, is very old and well known (e.g. Tubeuf 1916). Furthermore, empirical evidence also suggests a primary role of cytokinins in 'green island' formation by leafminers (Englebrecht *et al.* 1969). These phytohormones play a crucial role in controlling leaf senescence and have been clearly shown to be involved in plant–bacteria, plant–fungi and plant–galling insects interactions (Barry *et al.* 1984; Smart *et al.* 1991; Leicht 1994; Gan & Amasino 1995).

A markedly visible sign of leaf senescence is the gradual loss of chlorophyll in autumn and concomitant yellowing, but senescence is also a metabolic turning point biochemically characterized by the transition from nutrient assimilation to nutrient remobilization (Masclaux *et al.* 2000; Soudry *et al.* 2005). Leafminers are therefore potentially deprived from the required nutrients for completing their development before winter.

Here, we report that leafminers manipulate their environment using three types of cytokinins. This system allows leafminers to generate a microenvironment with all the nutrient supply needed for their survival in an otherwise senescent environment.

2. MATERIAL AND METHODS

Phyllonorycter blancardella F. (Lepidoptera: Gracillariidae) is a polyvoltine leaf-mining microlepidopteran. The first three larval instars are sap feeders and set the outline of the mine and, therefore, the total surface available to later stages. The two following instars are selective tissue feeders.

Samples were collected early in autumn 2005 on 12-year-old apple (*Malus domestica* Borkh.) cv. 'Allstars' trees grown in a biologically managed orchard. Collected leaves were directly dissected on site, frozen on dry ice and stored at -80°C until analysis. Samples were controlled for leaf age and natural exposition in the orchard (Sun versus shade—position in the rows). Mined age (only one mine per leaf) was also controlled and corresponds to L4 tissue-feeding larvae.

In order to study the spatial variation of cytokinins and nutrient concentrations, mined tissues, ipsilateral tissues (leaf tissues situated on the same side of the main vein as the mine) and contralateral tissues (leaf tissues on the opposite side of the main vein as the mine) were dissected according to figure 1. Leaf-miner insects and frass were also removed from the mine. Non-infected green and yellow apple leaves were dissected as previously and used as a control.

(a) Cytokinin quantification in mined and non-mined tissues according to leaf senescence

Leaf samples were lyophilized and pulverized. Similar amounts of control, mined, ipsilateral and contralateral tissues were extracted overnight in aqueous methanol containing butylated hydroxytoluene as an antioxidant.

(i) Purification

Purification was performed using a nitrocellulose prefilter (4.5 μm , Sartorius, Germany) connected to an Oasis cartridge (Waters, USA) with a Teflon filter (0.2 μm , Sartorius, Germany) at the outlet.

(ii) Separation

Eluates were reduced by rotary evaporation, taken up with acidified water and injected into a reverse-phase HPLC column (Waters, USA). After sample injection, the column was eluted at 1 ml min^{-1} with a convex gradient of acidified water and pure acetonitrile (HPLC grade) until the eluant was 100% acetonitrile. Retention time of cytokinins was determined by separate and pooled injections of pure zeatin (Z), ribosyl zeatin (ZR), isopentenyladenine (iP) and isopentenyl adenosine (iPA) (Sigma, USA) as standards. Fractions centred on the retention time of each cytokinin were collected and evaporated to dryness in a speed-vac concentrator (Savant, USA).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2007.0051> or via <http://www.journals.royalsoc.ac.uk>.



Figure 1. An apple tree leaf infected with the tentiform leafminer *Phyllonorycter blancardella* larva in autumn. The 'Green island' (feeding area) exhibits intact green chlorophyll-containing tissues, while the remaining leaf tissues undergo leaf senescence. The white spots on the mine are feeding windows, where all but the epidermis has been consumed by the caterpillar. Letters show areas used for cytokinin and nutrient content analysis: 'a' mined, 'b' ipsilateral and 'c' contralateral plant tissues.

(iii) *Quantification*

Aliquots of each fraction corresponding to the retention time of each hormone standard were subjected to the retention time of each hormone standard were subjected to enzyme-linked immunosorbent assay (ELISA) as previously published (Jourdain *et al.* 1997). ELISA quantification could be done owing to a strict separation of each compound by HPLC fractionation. Optical densities were measured at 405 nm. See electronic supplementary materials for details.

(b) *Nutrient quantification in mined and non-mined tissues according to leaf senescence*

Quantification of the amount of nutrients in leaf samples was carried out using colorimetric techniques successfully used for plant material (Yemm & Willis 1954; Jones *et al.* 1989). Briefly, samples were lyophilized, weighed and ground. Nutrients were measured using a spectrophotometer after reaction with Anthrone reagent for sugars and Bradford reagent for proteins. See electronic supplementary materials for details.

(c) *Statistical analysis*

Statistical analyses were performed using R v. 2.3.0 software. Data were analysed by non-parametric Kruskal–Wallis tests followed by Behrens–Fisher *post hoc* tests.

3. RESULTS

(a) *Cytokinin content of mined area exceeds that of non-infected or control areas*

To evaluate the role of phytohormones on 'green island' formation, we compared cytokinin contents in mined and non-mined tissues (figure 2). As expected, cytokinin content decreases with leaf senescence, some compounds such as isopentenyladenine, falling below the detection level (green versus yellow controls). However, the global cytokinin content of mined areas far exceeded that of uninfected areas (ipsi- and contralateral) and controls (green and yellow uninfected leaves). This phenomenon is particularly true for isopentenyl adenosine and isopentenyladenine, both compounds being at the beginning of the synthesis pathway of cytokinins. High levels of cytokinins are also observed in the near vicinity of the mine, ipsilateral tissues showing an amount of iP higher than contralateral tissues.




Figure 2. Spatial changes in the levels of cytokinins of senescing apple leaves. (a) Isopentenyladenosine, (b) isopentenyladenine and (c) zeatin concentrations in mined, ipsilateral and contralateral plant tissues are expressed as means \pm s.e. Non-infected green or yellow apple leaves were used as a control. Homologous groups are indicated by identical characters (Kruskal–Wallis test: iP, $p < 0.01$; iPA, $p < 0.05$; Z, $p < 0.01$).

(b) *Nutrient content of mined tissues does not decrease in senescing leaves*

We next conducted experiments investigating the nutritional benefits of the hormonal manipulation. Both food quality parameters decreased in senescing leaves falling from 135.56 ± 25.35 to $53.25 \pm 4.27 \mu\text{g mg}^{-1}$ for sugars ($n=15$) and from 29.10 ± 0.91 to $16.22 \pm 1.67 \mu\text{g mg}^{-1}$ for proteins ($n=15$; figure 3). Mined areas showed similar content of carbohydrates and proteins as uninfected green tissues. Carbohydrate and protein amounts in distant




Figure 3. Spatial changes in the levels of (a) sugars and (b) proteins of senescing apple leaves. Results for mined, ipsilateral and contralateral plant tissues are expressed as means \pm s.e. Non-infected green or yellow apple leaves were used as a control. Homologous groups are indicated by identical characters (Kruskal–Wallis test: *sugars*, $p < 0.01$; *proteins*, $p < 0.05$).

regions of infected leaves were similar or slightly higher than uninfected yellow controls.

4. DISCUSSION

'Green islands' are characterized by similar or even greater levels of zeatin, isopentyladenine and isopentenyl adenosine than non-senescing green leaves. This explains the stay-green phenotype of mined areas, while other regions of the same leaf are turning yellow. The origin of these cytokinins is however currently unknown and several hypotheses can be proposed. First, high levels of cytokinins in mined areas could result from the overexpression of the plant genes which produce cytokinins in the infected zone. Oral secretions of feeding insects contain many specific compounds able to interact with plant gene expression as shown in the upregulation of plant defences (e.g. Arimura *et al.* 2005). Second, cytokinins in mined tissues could result from an accumulation process rather than a synthesis *per se*. Feeding activity of leafminers is known to alter biophysical properties of leaves and consequently respiration and transpiration rates (Pincebourde *et al.* 2006). Such processes strongly impact the flow of nutrients within a leaf and could potentially result in the accumulation of cytokinins in mined areas. Third,

insects themselves could potentially synthesize cytokinins. Large quantities of cytokinins have already been measured in labial glands of leafminers (Engelbrecht *et al.* 1969) and the ability to produce cytokinins has been shown in several plant-associated organisms like fungi and bacteria (Barry *et al.* 1984; Stevens & Berry 1988; Morris 1995).

During the senescence of apple tree leaves, the levels of proteins and carbohydrates decrease, but mined tissues displayed a similar level of both proteins and sugars as non-senescing uninfected green leaves. Owing to the maintenance of the photosynthetic apparatus within the mine—as a direct consequence of high levels of phytohormones—natural production of sugars and proteins 'on site' cannot be excluded. However, modifications of the water flow and accumulation of nutrient-attracting cytokinins in the mined area could favour the translocation of nutrients from the leaf to the mine as well. In any case, both processes result in a net accumulation of nutrients in the mined tissues with potential competition between the plant and the mine for nutrients originating from senescing cells (change in the source–sink relationships—Balibrea *et al.* 2004; Wingler *et al.* 2006).

The maintenance of functional green tissues is, of course, of considerable ecological value to the development of the larvae as it allows the insect to maintain a favourable nutritional environment in an otherwise degenerating context. The 'green island', enriched in cytokinins, enables the larvae to retain nutrients within the mine and to undergo a long development later in the season. Interestingly, remote parts of infected leaves have higher concentration of cytokinins than uninfested yellow leaves (except for iP). Yellow leaves also totally lack iP, while this class of cytokinin is present in all tissues of infected leaves. Both results are a clear proof of manipulation of the plant physiology by the herbivore and over a much larger scale than the mine itself. The observed metabolic changes taking place in the mine and the manipulation of plant physiology induced by the insect strengthens the hypothesis that mines behave independently from the hosting leaf, operating a metabolic machinery of their own. Similar conclusions have been drawn with galling insects also known to produce galling tissues with high levels of cytokinins (e.g. Leicht 1994; Carol & Davies 2001). Despite great morphological differences between galls and mines, our results suggest a probable generalized process associated with the colonization of plants by endophagous organisms. Such intimate associations probably facilitated biochemical and hormonal cross-talk between insects and plants, setting the ground for host-plant manipulation by insects.

This work was supported by the ANR grant no. ANR-05-JCJC-0203-01. We also thank two anonymous reviewers for their comments on the manuscript and L. H. for full access to his orchard and helpful discussions.

Arimura, G., Kost, C. & Boland, W. 2005 Herbivore-induced volatiles induce the emission of ethylene in neighboring lima bean plants. *Plant J.* **29**, 87–98. (doi:10.1016/j.bbailp.2005.03.001)

- Balibrea, M. E., Gonzales Garcia, M. C., Fatima, T., Ehneß, R., Lee, T. K., Proels, R., Tanner, W. & Roitsh, T. 2004 Extracellular invertase is an essential component of cytokinin-mediated delay of senescence. *The Plant Cell* **16**, 1276–1287. (doi:10.1105/tpc.018929)
- Barry, G. F., Rogers, S. G., Fraley, R. T. & Brand, L. 1984 Identification of a cloned cytokinin biosynthetic gene. *Proc. Natl Acad. Sci. USA* **81**, 4776–4780. (doi:10.1073/pnas.81.15.4776)
- Carol, C. M. & Davies, J. P. 2001 Cytokinins in the ball gall of *Solidago altissima* and in the gall forming larvae of *Eurosta solidiginis*. *New Phytol.* **151**, 203–212. (doi:10.1046/j.1469-8137.2001.00158.x)
- Connor, E. F. & Taverner, M. P. 1997 The evolution and adaptive significance of the leaf-mining habit. *Oikos* **79**, 6–25. (doi:10.2307/3546085)
- Engelbrecht, L., Orban, U. & Heese, W. 1969 Leaf-miner caterpillars and cytokinins in the green islands of autumn leaves. *Nature* **223**, 319–321. (doi:10.1038/223319a0)
- Feeny, P. 1970 Seasonal changes in oak leaf tannins and nutrients as a cause of spring feeding by wintermoth caterpillars. *Ecology* **51**, 565–581. (doi:10.2307/1934037)
- Gan, S. & Amasino, R. M. 1995 Inhibition of leaf senescence by autoregulated production of cytokinin. *Science* **270**, 1986–1988. (doi:10.1126/science.270.5244.1986)
- Hering, E. M. 1951 *Biology of leaf miners*. Berlin, Germany: Junk-Gravenhage.
- Jones, C. G., Hare, J. D. & Compton, S. J. 1989 Measuring plant protein with the Bradford assay. *J. Chem. Ecol.* **15**, 979–992. (doi:10.1007/BF01015193)
- Jourdain, I., Lelu, M.-A. & Label, P. 1997 Hormonal change during growth of somatic embryogenic masses in hybrid larch. *Plant Physiol. Biochem.* **35**, 741–749.
- Leicht, I. J. 1994 Induction and development of the bean gall caused by *Pontania proxima*. In *Plant galls: organisms, interactions, population* (ed. M. A. J. Williams), pp. 283–300. New York, NY: Oxford University Press.
- Masclaux, C., Valadier, M. H., Brugiere, N., Morot-Gaudry, J. F. & Hirel, B. 2000 Characterization of the sink/source transition in tobacco (*Nicotiana tabacum* L.) shoots in relation to nitrogen management and leaf senescence. *Planta* **211**, 510–518. (doi:10.1007/s004250000310)
- Morris, R. O. 1995 Genes specifying auxin and cytokinin biosynthesis in prokaryotes. In *Plant hormones: physiology, biochemistry, and molecular biology* (ed. P. J. Davies), pp. 98–118. Dordrecht, The Netherlands: Kluwer Academic Publishers.
- Pincebourde, S. & Casas, J. 2006 Multitrophic biophysical budgets: thermal ecology of an intimate herbivore insect–plant interaction. *Ecol. Monogr.* **76**, 175–194.
- Pincebourde, S., Frak, E., Sinoquet, H., Regnard, J.-L. & Casas, J. 2006 Herbivory mitigation through increased water-use efficiency in a leaf-mining moth–apple tree relationship. *Plant Cell Environ.* **29**, 2238–2247. (doi:10.1111/j.1365-3040.2006.01598.x)
- Scheirs, J., De Bruyn, L. & Verhagen, R. 2001 Optimization of adult performance determines host choice in a grass miner. *Proc. R. Soc. B* **267**, 2065–2069. (doi:10.1098/rspb.2000.1250)
- Smart, C. M., Scofield, S. R., Bevan, M. W. & Dyer, T. A. 1991 Delayed leaf senescence in tobacco plants transformed with *tmr*, a gene for cytokinin production in *Agrobacterium*. *Plant Cell* **3**, 647–656. (doi:10.1105/tpc.3.7.647)
- Soudry, E., Ulitzur, S. & Gepstein, S. 2005 Accumulation and remobilization of amino acids during senescence of detached and attached leaves: *in planta* analysis of tryptophan levels by recombinant luminescent bacteria. *J. Exp. Bot.* **56**, 695–702. (doi:10.1093/jxb/eri054)
- Stevens, G. A. & Berry, A. M. 1988 Cytokinin secretion by *Frankia* sp. HFPArI3 in defined medium. *Plant Physiol.* **87**, 15–16. (doi:0032-0889/88/87/0015/02/\$01.00/0)
- Tubeuf, C. V. 1916 Die von Parasiten bewohnten grünen Inseln verfilbender Blätter. *Naturw. Z. Forst- u. Landw.* **14**, 42–46.
- Yemm, E. W. & Willis, A. J. 1954 The estimation of carbohydrates in plant extract by anthrone. *Biochem. J.* **57**, 508–514.
- Wingler, A., Purdy, S., MacLean, J. A. & Pourteau, N. 2006 The role of sugars in integrating environmental signals during the regulation of leaf senescence. *J. Exp. Bot.* **57**, 391–399. (doi:10.1093/jxb/eri279)