

## ERRATUM

**Cytogenetic analysis of buccal cells from shoeworkers and pathology and anatomy laboratory workers exposed to *n*-hexane, toluene, methyl ethyl ketone and formaldehyde**SEMA BURGAZ, ONUR ERDEM, GONCA ÇAKMAK,  
NURTEN ERDEM, ASUMAN KARAKAYA, & ALI ESAT KARAKAYA*Biomarkers* 2002;7:151–161

The authors would like to note that MN frequencies in the abstract and in table 4 of the above mentioned article were incorrectly printed. These should have appeared as ‰ where necessary.

The corrected table is presented below.

Table 4. Results of analysis of MN (‰) frequencies in buccal mucosa cells of control, group I and II workers according to smoking habits.

	Control (n)	Shoe-workers (n) Group I*	FA workers (n) Group II
All	0.33 ± 0.30 (18)	0.62 ± 0.45 (21)	0.71 ± 0.56 (28)
Range	0.00–1.00	0.00–1.67	0.00–2.00
Smokers	0.33 ± 0.19 (7)	0.69 ± 0.52 (13)	0.89 ± 0.52 (12)
Range	0.00–0.67	0.00–1.67	0.33–2.00
Nonsmokers	0.33 ± 0.37 (11)	0.50 ± 0.31 (8)	0.58 ± 0.57 (16)
Range	0.00–1.00	0.00–1.00	0.00–1.67
Number of cells analysed	54 × 10 <sup>3</sup>	63 × 10 <sup>3</sup>	84 × 10 <sup>3</sup>

\*Buccal cells from one individual were not available.