



Corrigendum


Corrigendum to “DNA gyrase (GyrB)/topoisomerase IV (ParE) inhibitors: Synthesis and antibacterial activity” [Bioorg. Med. Chem. Lett. 19 (2009) 894]

Stephen P. East^{a,*}, Clara Bantry White^a, Oliver Barker^a, Stephanie Barker^b, James Bennett^b, David Brown^b, E. Andrew Boyd^a, Christopher Brennan^a, Chandana Chowdhury^a, Ian Collins^b, Emmanuelle Convers-Reignier^a, Brian W. Dymock^a, Rowena Fletcher^b, David J. Haydon^b, Mihaly Gardiner^a, Stuart Hatcher^a, Peter Ingram^a, Paul Lancett^b, Paul Mortenson^a, Konstantinos Papadopoulos^a, Carol Smee^b, Helena B. Thomaidis-Brears^b, Heather Tye^a, James Workman^a, Lloyd G. Czaplewski^{b,*}

^a Evotec (UK) Ltd, Discovery Services, 114 Milton Park, Abingdon, Oxfordshire OX14 4SA, UK

^b Prolysis Ltd, Oxford University Begbroke Science Park, Sandy Lane, Yarnton, Oxfordshire OX5 1PF, UK

The authors regret that there was an error in [Scheme 3](#). The correct scheme is shown here.



Scheme 3. Reagents and conditions: (i) *n*-BuLi (3 equiv), THF, -78°C , then CO_2 ; (ii) SOCl_2 , MeOH, rt to 75°C , 18% (over 2 steps); (iii) 3-pyridyl boronic acid, $\text{Pd}(\text{dppf})\text{Cl}_2$, K_3PO_4 , MeOH: 1,4-dioxane (1:4), 100°C , 39%; (iv) LiOH, 1,4-dioxane, H_2O , rt, 88%; (v) HNR^1R^2 , AlMe_3 , 1,4-dioxane, toluene, 45°C , ~90%.