



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl

Corrigendum

Corrigendum to “Discovery of novel taspine derivatives as antiangiogenic agents” [Bioorg. Med. Chem. Lett. 20 (2010) 718]

Jie Zhang, Yanmin Zhang, Sanqi Zhang, Sicen Wang, Langchong He *

School of Medicine, Xi'an Jiaotong University, No. 76, Yanta Weststreet, Xi'an, Shaanxi Province 710061, PR China

The authors regret that the concentration unit of IC₅₀ values in the full paper including Table 2 appeared incorrectly. The correct concentration unit of IC₅₀ values is μM instead of nM. The correct format of Table 2 appears below.

Table 2
Antiproliferative activities of target compounds on human cancer and normal cells

Compound	Cell lines (IC ₅₀ , μM)		
	CACO-2 ^a	A549 ^b	ECV304 ^c
Taspine	153	4.18	22.3
2	110,03	453	ND
3	3100	111	ND
4	52.5	158	2.67
5	ND	180	61

ND is not determined.

^a CACO-2, human epithelial colorectal adenocarcinoma cells.

^b A549, carcinomic human alveolar basal epithelial cells.

^c ECV304, human umbilical endothelial cell line.

DOI of original article: [10.1016/j.bmcl.2009.11.073](https://doi.org/10.1016/j.bmcl.2009.11.073)

* Corresponding author. Tel./fax: +86 29 82655451.

E-mail address: helc@mail.xjtu.edu.cn