



Contents lists available at ScienceDirect

Bioorganic & Medicinal Chemistry Letters

journal homepage: www.elsevier.com/locate/bmcl



Synthesis of ‘clickable’ acylhomoserine lactone quorum sensing probes: Unanticipated effects on mammalian cell activation

Amanda L. Garner^{a,b,c}, Jing Yu^{a,b}, Anjali Kumari Struss^{a,b}, Colin A. Lowery^{a,b,c}, Jie Zhu^{a,b}, Sook Kyung Kim^{a,b}, Junguk Park^{a,b,c}, Alexander V. Mayorov^{a,b,c}, Gunnar F. Kaufmann^{a,b,c}, Vladimir V. Kravchenko^b, Kim D. Janda^{a,b,c,d,*}

^a Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

^b Department of Immunology and Microbial Sciences, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

^c The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA, USA

^d Worm Institute of Medical Research (WIRM), The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, CA 92037, USA

ARTICLE INFO

Article history:

Received 11 November 2010

Accepted 29 November 2010

Available online 4 December 2010

Keywords:

Acylhomoserine lactones

Click chemistry

Pseudomonas aeruginosa

Quorum sensing

ABSTRACT

Alkynyl- and azido-tagged 3-oxo-C₁₂-acylhomoserine lactone probes have been synthesized to examine their potential utility as probes for discovering the mammalian protein target of the *Pseudomonas aeruginosa* autoinducer, 3-oxo-C₁₂-acylhomoserine lactone. Although such substitutions are commonly believed to be quite conservative, from these studies, we have uncovered a drastic difference in activity between the alkynyl- and azido-modified compounds, and provide an example where such structural modification has proved to be much less than conservative.

© 2010 Elsevier Ltd. All rights reserved.

Quorum sensing is a diffusion-based process whereby bacteria secrete various signaling molecules, or autoinducers, as a population sensor and a means to control gene expression in response to changes in cell number; thus, allowing the bacteria to behave as a multicellular organism.¹ Importantly, this process has been correlated with the expression of virulence factors and biofilm formation,¹ and a ‘deafening’ of bacteria to these signaling molecules should provide a mechanism by which to target the bacteria without exerting a selective pressure thereby decreasing the possibility of resistance development.^{2,3} As a result, much effort has been invested in developing small molecule probes,^{4,5} autoinducer analogs^{4,5} and protein biologics^{6–8} for studying quorum sensing.

In particular, quorum sensing in the gram-negative bacteria *Pseudomonas aeruginosa* has received significant attention due to the species’ role as an opportunistic pathogen in humans.⁹ Most prominent is the role of *P. aeruginosa* in patients suffering from cystic fibrosis.¹⁰ Affected individuals suffer from frequent bacterial lung infections due to impaired lung defense function, and these infections are often a cause of morbidity.¹⁰ With respect to quorum sensing, it has now been established that this signaling process is required for *P. aeruginosa* to cause disease, including lung infections in cystic fibrosis patients.^{11–14} More specifically, the ability

of the bacteria to grow into sessile, drug-resistant biofilms at high cell densities makes such infections difficult to treat, and resistance is beginning to be observed with currently available antibiotics.¹⁵

To conduct its quorum sensing-controlled activities, including biofilm formation, *P. aeruginosa* uses *N*-acyl *L*-homoserine lactones (AHLs) as its primary autoinducers (Fig. 1).¹ AHL 1 activates LasR¹⁶ and AHL 2 activates RhIR,¹⁷ and activation of both receptor proteins is required for quorum sensing.¹ Since LasR activation by AHL 1 is required to initiate the quorum sensing cascade in this species,¹ many studies have been conducted toward deciphering the role of this chemical signal in both quorum sensing and its resulting human pathologies. Recently, our group^{18,19} and others^{20,21} have explored the impact of AHL 1 in mammalian cells. For example, AHL 1 inhibits innate immune responses through modulation of NF-κB signaling induced by pro-inflammatory stimuli such as tumor necrosis factor and the Toll-like receptor (TLR) agonists, thereby potentially promoting persistent infection.¹⁹ Notably, this molecule directly activates mammalian cells through a mechanism distinct from the TLR and other canonical




Figure 1. Autoinducers used by *Pseudomonas aeruginosa*.

* Corresponding author. Tel.: +1 858 784 2515; fax: +1 858 784 2595.

E-mail address: kdjanda@scripps.edu (K.D. Janda).

pathogen-associated receptor pathways.¹⁸ Because of the potential significance of these findings, our group and others²² have been actively involved in identifying the biochemical target(s) of this important quorum sensing signaling molecule.

Click chemistry, more specifically Cu^I-catalyzed azide-alkyne 1,3-dipolar cycloaddition chemistry,^{23–25} has emerged as a powerful method for bioconjugation.^{26,27} Coupled with activity-based protein profiling (ABPP), this method grants the ability to probe the molecular interactions of a molecule of choice by modifying the compound with a bioorthogonal azide or alkyne tag.^{28,29} While this substitution may seem conservative, in fact, this modification can have drastic impact on the activity of the small molecule, although such findings are often not reported. Moreover, as high affinity is a requirement for use of a probe in target identification studies, particularly in cases when the unknown target may be of low abundance,³⁰ choice of this modification may prove critical for such future studies. Herein, we describe our efforts to synthesize clickable AHL probes for target identification studies and demonstrate the critical nature that the choice of an alkyne or azide modification has on AHL activity in mammalian cells.

Although much is known about the structural requirements for AHL-mediated activation in *P. aeruginosa*,^{4,5} very little has been concluded about the requirements for biological activity in mammalian cells. Previously, our group demonstrated that the L-homoserine lactone, 3-oxo substituent and an alkyl chain of 10–14 carbons were necessary to activate mammalian cells.¹⁸ To begin our studies, we first synthesized alkynyl analogs **5a** and **5b**, as an alkyne seemed to be the most conservative modification since it is composed of only carbons similar to the natural tail of AHL **1**. Additionally, previous studies have showed that detection with rhodamine-azide compounds yields lower background signals than detection with the corresponding rhodamine-alkyne tags in click chemistry-mediated ABPP.²⁹ Although a diazirine moiety was used in a previously reported AHL-based probe,²² we hypothesized that the acylhomoserine lactone could act as a covalent tag on its own, particularly since lactone natural products are well known to covalently modify protein targets in vivo.³¹ The synthesis of alkynyl analogs **5a** and **5b** is shown in Scheme 1, and proceeded via a straightforward route starting from the commercially available alkynyl fatty acids **3a** and **3b**. Briefly, alkynyl acids **3** were first activated with DCC followed by nucleophilic displacement of the resultant activated esters with Meldrum's acid and coupling with L-homoserine lactone **4** to yield alkynyl AHL analogs **5**. For carbon chain lengths, we chose to synthesize both a shorter 9 carbon tail and a longer 13 carbon tail for comparison.


To assess a potential utility of these alkynyl AHL analogs as probes for target identification studies, we tested their biological activities in primary macrophages. In brief, bone marrow-derived macrophages (BMDM) were treated with an alkynyl AHL derivative (50 μM) over a period of 90 min, and cellular extracts were examined by Western blot analysis for the biochemical markers indicative of AHL **1**-mediated signaling events including phosphorylation of protein kinase p38, phosphorylation of eukaryotic translation initiation factor 2α (eIF2α) and the cleavage of poly(ADP)-ribose polymerase (PARP), as previously described.¹⁸ AHL **1** (3-oxo-C₁₂-

AHL) and C₁₂-AHL (no 3-oxo group, structure not shown) were used as positive and negative controls, respectively. As Figure 2 shows, only alkynyl analog **5a** exhibited activity; however, its activity was greatly reduced compared to that of the parent compound **1**.

Knowing that the diminished activity of alkynyl analog **5a** and apparent lack of activity in analog **5b** would dampen any efforts toward identifying the mammalian target(s) of AHL **1**, we next synthesized an azide derivative in hope of identifying a more viable AHL analog for our desired future studies. Since the shorter alkyne probe showed no activity, we chose to examine only a long alkyl chain. Azido analog **8a**, which contains 12 carbons in its tail, was synthesized in a similar manner as the alkynyl derivatives (Scheme 2). In this case, however, azido-carboxylic acid **7** was used as a starting material and was obtained via nucleophilic substitution of bromo-acid **6** with sodium azide (Scheme 2). We then tested the activity of this azido derivative in BMDM. Interestingly, unlike the alkynyl derivatives, the azido AHL analog exhibited activity very similar to that of the parent AHL (Fig. 3). Thus, substitution at the tail terminus clearly has a drastic impact on AHL activity. This finding is unanticipated since an azido group, although overall neutral, is polar and carries charged nitrogen atoms, while an alkynyl group is fully neutral and nonpolar similar to the natural AHL tail. Although purely speculative, it is possible that the azido moiety makes additional hydrogen bond or van der Waal contacts³² that aids in its binding of the putative AHL target(s). Importantly, this potential probe exhibits much higher activity in mammalian cells than a previously reported diazirine-modified alkynyl AHL that was used to label and isolate *P. aeruginosa* LasR.²² As an additional control compound, we also synthesized the corresponding D-homoserine lactone derivative (**8b**, Scheme 2), and similar to previous experiments with this enantiomer,¹⁸ no activity was observed (Fig. 3). Thus, **8b** could prove useful as a structurally similar negative control for potential target identification strategies.

In addition to testing these AHL derivatives in mammalian cells, we also examined their impact as potential autoinducers in *P. aeruginosa*. Autoinducer assays were conducted using *P. aeruginosa* luminescence reporter strain PAO-JP2, a PAO1 *lasI/rhlI* double mutant (PAO-JP2-*luxABCDE*).^{33,34} As Figure 4 shows, only alkynyl AHL analog **5a** showed autoinducer activity similar to that of the parent AHL **1**. These data indicate that the structural requirements for autoinducer activity and activity in mammalian cells are distinct, and nonpolar, all carbon alkynyl probes may be more useful for identifying bacterial AHL receptors.

In conclusion, we have synthesized alkynyl- and azido-tagged 3-oxo-C₁₂-AHL analogs as potential ABPP probes for discovering the mammalian target of this important *P. aeruginosa* quorum sensing signaling molecule. Of particular significance, we have found a drastic difference in activity between the alkynyl- and azi-



Scheme 1. Synthesis of alkynyl AHL analogs.







Figure 2. Biological activities of AHL **1**, C₁₂-HSL (A-C12; negative control) and its alkynyl analogs **5a** and **5b** in mammalian cells. BMDM were treated as indicated, and cellular extracts were analyzed by Western blot for the cleavage of PARP, phosphorylated forms of eIF2α (p-eIF2α) or p38 (p-p38) and actin as a loading control.



Scheme 2. Synthesis of azido AHL analogs.

Figure 3. Biological activities of azido analogs **8a** and **8b** in mammalian cells.Figure 4. Autoinducer activities of alkynyl- and azido-AHL analogs in *P. aeruginosa*. Relative luminescence units were normalized with respect to cell viability (OD_{600}).

do-tagged AHL derivatives in that only the azide-containing analog possessed activity similar to that of the natural compound. Thus, these data demonstrate that click chemistry-amenable substitutions may not be as conservative as previously thought, and that no simple guidelines can exist for such click chemistry-mediated ABPP endeavors. Both alkynyl- and azido-modified compounds should be initially examined for activity before examination as potential probes. Additionally, the discovery that polar, yet neutral substituents at the tail are accepted for mammalian cell activity may open up the possibility for the synthesis of new, more potent AHL derivatives. Target identification and additional synthetic studies are currently ongoing and will be reported in due course.

Acknowledgments

We thank M. M. Meijler (Ben Gurion University) for kindly providing *P. aeruginosa* strain PAO-JP2. We gratefully acknowledge the

financial support of the NIH (Grants AI079503 and DE018452 to K.D.J., and Grant AI085324 to G.F.K.), the Skaggs Institute for Chemical Biology, and a NIH postdoctoral fellowship (F32-DK083179, A.L.G.).

Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bmcl.2010.11.122.

References

- Waters, C. M.; Bassler, B. L. *Annu. Rev. Cell. Dev. Biol.* **2005**, *21*, 319.
- Rasmussen, T. B.; Givskov, M. *Int. J. Med. Microbiol.* **2006**, *296*, 149.
- Suga, H.; Smith, K. M. *Curr. Opin. Chem. Biol.* **2003**, *7*, 586.
- Lowery, C. A.; Salzameda, N. T.; Sawada, D.; Kaufmann, G. F.; Janda, K. D. *J. Med. Chem.* doi:10.1021/jm901742e.
- Mattmann, M. E.; Blackwell, H. E. *J. Org. Chem.* **2010**, *75*, 6737.
- Kaufmann, G. F.; Sartorio, R.; Lee, S. H.; Mee, J. M.; Altobelli, L. J., III; Kujawa, D. P.; Jeffries, E.; Clapham, B.; Meijler, M. M.; Janda, K. D. *J. Am. Chem. Soc.* **2006**, *128*, 2802.
- De Lamo Marin, S.; Xu, Y.; Meijler, M. M.; Janda, K. D. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 1549.
- Kaufmann, G. F.; Park, J.; Mee, J. M.; Ulevitch, R. J.; Janda, K. D. *Mol. Immunol.* **2008**, *45*, 2710.
- Lyczak, J. B.; Cannon, C. L.; Pier, G. B. *Microbes Infect.* **2000**, *2*, 1051.
- Lyczak, J. B.; Cannon, C. L.; Pier, G. B. *Clin. Microbiol. Rev.* **2002**, *15*, 194.
- Pearson, J. P.; Feldman, M.; Iglewski, B. H.; Prince, A. *Infect. Immun.* **2000**, *68*, 4331.
- Tang, H. B.; DiMango, E.; Bryan, R.; Gambello, M.; Iglewski, B. H.; Goldberg, J. B.; Prince, A. *Infect. Immun.* **1996**, *64*, 37.
- Davies, D. G.; Parsek, M. R.; Pearson, J. P.; Iglewski, B. H.; Costerton, J. W.; Greenberg, E. P. *Science* **1998**, *280*, 295.
- Calfee, M. W.; Coleman, J. P.; Pesci, E. C. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 11633.
- Drenkard, E. *Microbes Infect.* **2003**, *5*, 1213.
- Pearson, J. P.; Gray, K. M.; Passador, L.; Tucker, K. D.; Eberhard, A.; Iglewski, B. H.; Greenberg, E. P. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91*, 197.
- Pearson, J. P.; Passador, L.; Iglewski, B. H.; Greenberg, E. P. *Proc. Natl. Acad. Sci. U.S.A.* **1995**, *92*, 1490.
- Kravchenko, V. V.; Kaufmann, G. F.; Mathison, J. C.; Scott, D. A.; Katz, A. Z.; Wood, M. R.; Brogan, A. P.; Lehmann, M.; Mee, J. M.; Iwata, K.; Pan, Q.; Fearn, C.; Knaus, U. G.; Meijler, M. M.; Janda, K. D.; Ulevitch, R. J. *J. Biol. Chem.* **2006**, *281*, 28822.
- Kravchenko, V. V.; Kaufmann, G. F.; Mathison, J. C.; Scott, D. A.; Katz, A. Z.; Grauer, D. C.; Lehmann, M.; Meijler, M. M.; Janda, K. D.; Ulevitch, R. J. *Science* **2008**, *321*, 259.
- Thomas, G. L.; Bohner, C. M.; Williams, H. E.; Walsh, C. M.; Ladlow, M.; Welch, M.; Bryant, C. E.; Spring, D. R. *Mol. Biosyst.* **2006**, *2*, 132.
- Li, L.; Hooi, D.; Chhabra, S. R.; Pritchard, D.; Shaw, P. E. *Oncogene* **2004**, *23*, 4894.
- Dubinsky, L.; Jarosz, L. M.; Amara, N.; Krief, P.; Kravchenko, V. V.; Krom, B. P.; Meijler, M. M. *Chem. Commun.* **2009**, 7378.
- Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596.
- Tornøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057.
- Meldal, M.; Tornøe, C. W. *Chem. Rev.* **2008**, *108*, 2952.
- Wang, Q.; Chan, T. R.; Hilgraf, R.; Fokin, V. V.; Sharpless, K. B.; Finn, M. G. *J. Am. Chem. Soc.* **2003**, *125*, 3192.
- Best, M. D. *Biochemistry* **2009**, *48*, 6571.
- Speers, A. E.; Adam, G. C.; Cravatt, B. F. *J. Am. Chem. Soc.* **2003**, *125*, 4686.

29. Speers, A. E.; Cravatt, B. F. *Chem. Biol.* **2004**, *11*, 535.
30. Sato, S.; Murata, A.; Shirakawa, T.; Uesugi, M. *Chem. Biol.* **2010**, *17*, 616.
31. Konaklieva, M. I.; Plotkin, B. J. *Mini-Rev. Med. Chem.* **2005**, *5*, 73.
32. Petsalakis, E. I.; Chrysina, E. D.; Tiraidis, C.; Hadjiloi, T.; Leonidas, D. D.; Oikonomakos, N. G.; Aich, U.; Varghese, B.; Loganathan, D. *Bioorg. Med. Chem.* **2006**, *14*, 5316.
33. Duan, K.; Surette, M. G. *J. Bacteriol.* **2007**, *189*, 4827.
34. Amara, N.; Mashiach, R.; Amar, D.; Krief, P.; Spieser, S. A. H.; Bottomley, M. J.; Aharoni, A.; Meijler, M. M. *J. Am. Chem. Soc.* **2009**, *131*, 10610.