



New prodrugs of Adefovir and Cidofovir

Tomáš Tichý^{a,*}, Graciela Andrei^b, Martin Dračínský^a, Antonín Holý^a, Jan Balzarini^b, Robert Snoeck^b, Marcela Krečmerová^a

^a Gilead Sciences & IOCB Research Centre, Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, v.v.i., Flemingovo nám. 2, Prague 166 10, Czech Republic

^b Rega Institute for Medical Research, Katholieke Universiteit Leuven, Minderbroedersstraat 10, Leuven B-3000, Belgium

ARTICLE INFO

Article history:

Received 14 January 2011

Revised 25 March 2011

Accepted 9 April 2011

Available online 22 April 2011

Keywords:

Adefovir

Cidofovir

Antivirals

Prodrugs

Acyclic nucleoside phosphonate

Phosphonate ester

In vitro evaluation

ABSTRACT

New Adefovir (PMEA) prodrugs with a pro-moiety consisting of decyl or decyloxyethyl chain bearing hydroxyl function(s), hexaethyleneglycol or a (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl unit were prepared starting from the tetrabutylammonium salt of the phosphonate drug and an appropriate alkyl bromide or tosylate. Analogously, two esters of Cidofovir [(S)-HPMPC] bearing a hexaethyleneglycol moiety were prepared. The activity of the prodrugs was evaluated in vitro against different virus families. A loss in the antiviral activities of the hydroxylated decyl or decyloxyethyl esters and hexaethyleneglycol esters of PMEA against human immunodeficiency virus (HIV) and herpesviruses [including herpes simplex virus (HSV), varicella-zoster virus (VZV), and human cytomegalovirus (CMV)] occurred in comparison with the parent compound. On the other hand, the (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of PMEA showed significant activities against HIV and herpesviruses. (S)-HPMPC prodrugs exhibited anti-cytomegalovirus activities in the same range as the parent drug, whereas the anti-HSV and anti-VZV activities were one- to seven-fold lower than that of Cidofovir.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Acyclic nucleoside phosphonates (ANPs) are a group of compounds with remarkable antiviral activities. Their development has resulted in three approved drugs, and research on ANPs continues to provide new active compounds.^{1,2} The presence of the phosphonate group is responsible for their ionic character when subjected to physiological pH. The ionized molecule is not easily permeable through the gastrointestinal wall and biological membranes. In order to achieve better oral bioavailability, the phosphonate group of the drug can be transformed to a phosphonic ester or amidate, which is enzymatically cleaved to the parent drug after passing the intestinal barrier, or later inside the cells. A considerable number of ANP prodrugs have been evaluated, but only a few of them passed the preclinical studies. This likely arises from the complexity in achieving the right balance between the suitable chemical, physical, and pharmacological prodrug properties.

The basic aim of this work is to apply several less common pro-moieties to 9-[(2-phosphonylmethoxy)ethyl]adenine (**1**) (PMEA, Adefovir). PMEA has demonstrated a broad spectrum of antiviral activity¹ against human immunodeficiency virus (HIV) and other retroviruses and is active against various DNA viruses, including the hepatitis B virus and seventeen herpesviruses. The bis(pivaoyl-

oxymethyl)ester of PMEA [Adefovir Dipivoxyl, bis(POM)-PMEA] (Fig. 1) has been approved as an oral prodrug for the treatment of hepatitis B. The orally administered prodrug is absorbed and then cleaved by carboxyesterases in the serum generating pivalate and formaldehyde as its byproducts. Formaldehyde is the significant contributor to the overall cytotoxic effects of the prodrug.³

Another masking group of the pharmacochimistry of antibiotics—the dioxolenone unit—has been previously utilized to mask the carboxylic function analogously to the pivaoyloxymethyl variety. However, rare examples of dioxolenone esters on phosphate or phosphonate have been published.⁴ The bioavailability and stability of (5-substituted 2-oxo-1,3-dioxolen-4-yl)methyl esters have been explored. Methyl as a substituent in position 5 proved to be optimal with respect to oral absorbability as well as stability of the prodrug.⁵ We have aimed at the preparation of a conjugate of this particular dioxolenone unit and **1**.

Lipophilic esters constitute another important class of phosphate and phosphonate prodrugs. Some alkyl⁶ and alkoxyalkyl⁷ esters of nucleotides or acyclic nucleoside phosphonates have already been advanced into clinical studies. Their antiviral activities are superior when compared to the parent drugs. On the other hand, their physical properties are far from being optimal. They suffer from low solubility in water. Additionally, a simple alkyl chain is subjected to a ω -oxidation process resulting in inactivation of the prodrug. Therefore, we have focused on those pro-moieties consisting of the aliphatic chain modified by means of addition(s)

* Corresponding author. Tel.: +420 220 183 475; fax: +420 220 183 578.

E-mail addresses: tichy78@uochb.cas.cz, tomastichy78@email.cz (T. Tichý).

of hydroxyl group(s) or insertions of oxygen atoms. A hydroxylated alkyl or alkoxyalkyl unit or a hexaethyleneglycol unit was attached by ester linkage to the phosphonate group of Adefovir as a model drug. The influence of the above-mentioned modifications on the *in vitro* antiviral activities is reported here. We also studied the effects of the addition of the hexaethylene glycol unit in Cidofovir [(S)-HPMPC, **2**] as prodrugs closely related to Cidofovir hexadecyloxypropyl and octadecyloxyethyl esters, that is, HDP (S)-HPMPC and ODE (S)-HPMPC, respectively⁷ (Fig. 1). HDP (S)-HPMPC (CMX001) is currently being developed for use in the prophylactic and preemptive therapy of dsDNA viral infections.

Dioxolenone prodrugs are expected to be cleaved by serum or tissue esterases yielding the parent phosphonate, carbon dioxide and relatively nontoxic acetoin⁵, according to Scheme 1. The lipid phosphonate diesters are cleaved into appropriate monoesters after permeation through the wall of the small intestine.⁸ Unlike the dioxolenone ester of phosphonate, lipid monoesters are stable in plasma due to their hydrolytic stability and the absence of phospholipases. The remaining ester linkage is cleaved by phosphoesterases C inside the cells.^{7b} It was believed that the only role of the lipid pro-moiety is the facilitation of the passive transport through the phospholipid membranes. Recently, it was shown that the attachment of the masking alkoxyalkyl unit to the phosphonate pharmacophore affects pharmacokinetics more complexly.^{7b}

2. Results

We have prepared several structural types of PMEA prodrugs which are outlined in Section 1 (see Fig. 2). The prodrugs of **1** were prepared in doublets of corresponding mono and diesters with the exception of compound **15**. The presented pro-moieties are 10-hydroxydecyl unit (compounds **3** and **5**), 9,10-dihydroxydecyl unit (compounds **4** and **6**), 2-[(10-hydroxydecyl)oxy]ethyl unit (compounds **7** and **9**), 2-[(9,10-dihydroxydecyl)oxy]ethyl unit (compounds **8** and **10**), hexaethyleneglycol units (compounds **11–14**) or (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl unit (compound **15**). The length of the hydroxylated alkyl chain in compounds **3–10** was limited by the accessibility of the precursors (9-decen-1-ol, 10-undecen-1-ol). Hexaethyleneglycol pro-moiety was coupled also to Cidofovir (**2**), and thus prodrugs **16** and **17** were prepared.

2.1. Chemistry

2.1.1. Alkylation of the phosphonate function

The synthesis of the prodrugs is based on nucleophilic substitution reactions. The acidic P–OH group can be deprotonated and the negatively charged oxygen atom acts as a nucleophile in the reac-

tion with a functionalized alkyl bromide or tosylate. Their preparation is depicted in Section 2.1.2 below. Tetrabutylammonium hydroxide was used as a base for the deprotonation of alkyl phosphonic acid **1**. The resulting tetrabutylammonium salt is soluble in organic solvents and reacts with 2 equiv of an appropriately protected alkyl bromide or tosylate in moderate yields. The acid labile protecting groups of the prepared phosphonate diesters **18–22** were cleaved in a subsequent step by conventional methods (AcOH, Dowex 50). The monoesters **3, 4, 7, 8, 11, 12** were prepared from appropriate diesters by means of the published procedure⁹ involving a treatment with excess of lithium azide. Resulting monoesters are sufficiently stable towards the additional nucleophilic attack of the azide ion due to the presence of charged oxygen atom of the phosphonate function. Thus, the appropriate monoester was isolated as a sole product.

Diisopropylethylamine was utilized first as a base for the preparation of the (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of PMEA. Monitoring of the reaction mixture by TLC showed the formation of a relatively nonpolar product which was thought to be the appropriate diester. However, it disappeared during the working up the reaction mixture and monoester **15** was isolated as the sole product (see Scheme 3). Although the preparation procedure involving one equivalent of tetrabutylammonium hydroxide as a base seemed to be more suitable, the yield was still low (16%).

Cidofovir was alkylated starting from its cyclic form (**23**).^{7a} The tetrabutylammonium salt of **23** was treated with tosylate **28** or **29** (see Section 2.1.2). The resulting cyclic phosphonate esters yielded the monoesters (**16** or **17**) after the final working up (see Scheme 4).

2.1.2. The preparation of functionalized alkyl bromides and tosylates

The (5-Methyl-2-oxo-1,3-dioxolen-4-yl)methyl bromide for the preparation of **15** was synthesized according to the published procedure.⁵ Bromides for the preparation of **18–21** were prepared from commercially available 9-decen-1-ol or 10-undecen-1-ol. Their hydroxyl groups were replaced with bromine and the double bond was oxidized to 1,2-diol by the treatment with osmium tetroxide and *N*-methylmorpholine-*N*-oxide (Scheme 5). The diol system in **22** was protected with an isopropylidene group, and the prepared bromide **23** was reacted in a subsequent step with ethylene glycol. The arising alcohol was converted to bromide **24**. The diol system in **25** was oxidized by the action of sodium periodate followed by reductive treatment and MOM-protection procedure. The resulting bromide **26** was treated in the same two-step-way as compound **23** yielding bromide **27**. Bromides **23, 24, 26, 27** were used for the alkylation of PMEA according to Scheme 2. The acid labile protecting groups (isopropyliden, MOM) of the prepared phosphonate esters **18–21** were cleaved





Figure 1. The structures of Adefovir (**1**), Cidofovir (**2**) and some of their prodrugs.



Scheme 1. The activation of the considered prodrug types





Figure 2. The structures of the synthesized prodrugs.

in a subsequent step by conventional methods (AcOH, Dowex 50) (see Scheme 2).


The hexaethyleneglycol moiety was coupled to phosphonates **1** and **2** via its monotosylate. Hexaethyleneglycol was tosylated with one equivalent of tosylchloride with silver oxide as a base.¹⁰ The remaining hydroxyl group was protected as THP ether. The reaction of the prepared **28** with sodium ethoxide was followed by THP deprotection and tosylation. The three-step procedure yielded compound **29** (see Scheme 6). Both tosylates **28** and **29** were utilized for PMEA (**1**) or cHPMPC (**23**) alkylation (see Scheme 2). The THP protecting group of synthesized phosphonate esters **22** and **24** was cleaved in the presence of Dowex 50.

3. Biological activities

The antiviral activity of the different compounds was evaluated against various DNA viruses, including poxviruses [i.e., vaccinia virus (VACV)], herpesviruses [i.e., herpes simplex virus type 1 (HSV-1) and type 2 (HSV-2), thymidine kinase-deficient HSV-1 (acyclovir-resistant, ACV^r), varicella-zoster virus (VZV) and human cytomegalovirus (HCMV)] (Table 1). The compounds were also evaluated against retroviruses [(i.e., human immunodeficiency virus type 1 (HIV-1) and type 2 (HIV-2)] (Table 2). All of the compounds were also examined against several RNA viruses, including vesicular stomatitis virus (VSV), Coxsackie B4, respiratory syncytial



Scheme 2. The preparation of prodrugs **3–14**. Reagents and conditions: (a) Bu₄NOH, MeOH; (b) **23**, **24**, **26–29**, DMF, 100 °C; (c) LiN₃, DMF, 100 °C; (d) HCl, MeOH, 65 °C; (e) Dowex 50WX8-400 [H⁺], MeOH, H₂O, 60 °C; (f) AcOH, H₂O, 50 °C.




Scheme 3. The preparation of prodrug **15**. Reagents and conditions: (a) Bu₄NOH, MeOH; (b) (5-Methyl-2-oxo-1,3-dioxolen-4-yl)methyl bromide, DMF.


virus (RSV), parainfluenza virus type 3, reovirus-1, Sindbis virus and Punta Toro virus (Table 2). None of the prodrugs showed

activity against the different RNA viruses, except for HIV-1 and HIV-2. However, a loss in the antiviral activities of the hydroxylated decyl or decyloxyethyl and hexaethyleneglycol esters of PMEAs (compounds **3–14**) against HIV was observed since compound **7** {2-[(10-Hydroxydecyl)oxy]ethyl ester of PMEAs} was the only compound among this type of prodrugs showing some activity against HIV at nontoxic concentrations. Thus, compound **7** was able to inhibit HIV replication with 50% effective concentrations (EC₅₀) which were ~10-fold higher than those observed for PMEAs. In contrast, the (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of PMEAs (compound **15**) had anti-HIV activities (EC₅₀ values of 10.6 μM and 9.3 μM for HIV-1 and HIV-2, respectively) equivalent to that of PMEAs (EC₅₀ values of 10.4 and 8.6 μM for HIV-1 and HIV-2, respectively).


The 2-[(10-Hydroxydecyl)oxy]ethyl ester of PMEAs (**7**) exhibited also activities against HSV-1, HSV-2 and VZV. Similarly to HIV, the



Scheme 4. The preparation of prodrugs **16** and **17**. Reagents and conditions: (a) DCC, N,N'-dicyclohexyl-4-morpholinecarboxamidate, DMF, 100 °C; (b) tetrabutylammonium hydroxide, MeOH; (c) **28** or **29**, DMF, 100 °C; (d) AcOH, H₂O; (e) NH₃, H₂O.



Scheme 5. The preparation of protected bromides for the alkylation of **1** or **23**. Reagents and conditions: (a) CBr_4 , PPh_3 , CH_2Cl_2 , $0^\circ\text{C} \rightarrow \text{rt}$; (b) *N*-methylmorpholine-*N*-oxide, OsO_4 , acetone, H_2O , rt ; (c) 2,2-dimethoxypropane, TsOH , rt ; (d) NaH , ethyleneglycol, DMF , rt ; (e) NaIO_4 , H_2O , THF , rt ; (f) NaBH_4 , H_2O , THF , rt ; (g) $\text{CH}_3\text{OCH}_2\text{Br}$, CH_2Cl_2 , $i\text{Pr}_2\text{EtN}$, 0°C .



Scheme 6. The preparation of protected hexaethylene glycol monotosylates for the alkylation of **1** and **23**. Reagents and conditions: (a) TsCl , Ag_2O , KI , CH_2Cl_2 , 0°C ; (b) 3,4-dihydro-2*H*-pyran, PPTS , CH_2Cl_2 , 40°C ; (c) EtOH , NaH , THF , r.t. ; (d) Dowex 50WX8-400 $[\text{H}^+]$, MeOH , rt ; (e) TsCl , Et_3N , CH_2Cl_2 , rt .

anti-herpesvirus activities of the prodrugs **3–14** decreased when compared to the parent compound, with compound **7** being the most active one among these prodrugs. In the case of HSV and VZV, the drop in antiviral activities was inferior to that observed for HIV since the EC_{50} values for compound **7** against HSV and VZV were three- to four-fold lower than those for PMEA. However, compound **7** was totally inactive against CMV. The (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester **15** had anti-herpesvirus activities. The activity of compound **15** against herpesviruses was equivalent (HSV-1 and VZV) or two- to four-fold (HSV-2) lower than that observed for the parent compound PMEA. Other esters of PMEA, that is, compounds **6**, **8**, **11** and **12**, showed weak activities only against VZV, with EC_{50} values in the range of 18–114 μM , which is 3- to 15-fold higher than the EC_{50} values for the parent compound. None of the PMEA prodrugs showed activity against vaccinia virus or feline herpesvirus, like the parent compound PMEA.

The two prodrugs of (S)-HPMPC, that is, compounds **16** {14-hydroxy-3,6,9,12-tetraoxaheptadecyl ester of (S)-1-[3-hydroxy-2-(phosphonomethoxy)propyl]cytosine} and **17** {3,6,9,12,15-pentaoxa-aeicosyl ester of (S)-1-[3-hydroxy-2-(phosphonomethoxy)propyl]cytosine} were able to inhibit the replication of the different herpesviruses tested and the poxvirus vaccinia virus. Both prodrugs had activities equivalent to that of (S)-HPMPC against CMV, with EC_{50} values in the range of 0.6–0.95 μM as compared to 0.9 μM for (S)-HPMPC. However, compounds **16** and **17** had six- to seven-fold lower activities than (S)-HPMPC against VZV. Compound **17** had activities against HSV, feline herpesvirus, and vaccinia virus similar to those found for (S)-HPMPC while the activities of compound **16** against these viruses were 5- to 15-fold lower.

As already mentioned, prodrugs **16** and **17** represent the penta oxy-analogs of alkoxyalkyl prodrugs developed by Hostetler.^{7b} Table 3 compares some antiviral activities of **12** (HIV), **16** and **17** (CMV) with analogous simple alkoxyalkyl ester.^{11,12} The comparison demonstrates that insertion of oxygen atoms into the chain in **16** and **17** caused 400- to 1000-fold drop of activity against CMV. The loss of anti-HIV activity of **12** is even more significant ($>10^7$ -fold compared to HDP-PMEA). Comparison of activities of

compounds **3–11** with HDP-PMEA is not possible because different chain length plays a role. We can only speculate based on the relationship between activity and chain length, as observed with Cidofovir prodrugs.¹¹ The adequate chain shortening from 20 atoms (HDP) to 12 atoms (octyloxypropyl) caused circa 1000-fold drop in anti-CMV activity. However, the decrease in antiviral activity of compounds **3–8** is more significant ($\sim 10^7$ -fold compared to HDP-PMEA) and is hardly accountable solely to the chain shortening.

The superior antiviral activity of HDP esters is explained by the interplay of increased permeation through the cell membrane and the by-pass of the phosphorylation step leading to the rapid formation of bioactive Cidofovir diphosphate.^{7b} Since the prodrugs **3–14**, **16** and **17** are less active than the parent drug in some cases, decreased cellular uptake or slow cleavage of the promoiety by intracellular phospholipase C may be the reason for their poor antiviral activities.

4. Conclusions

A set of new amphiphilic prodrugs of PMEA (Adefovir) and (S)-HPMPC (Cidofovir) has been prepared and evaluated in vitro. The phosphonate group of PMEA was masked with a hexaethyleneglycol unit or hydroxylated decyl or decyloxyethyl chain. The first prodrug type was applied also to (S)-HPMPC (Cidofovir). As a less toxic analog of Adefovir Dipivoxyl, the (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of PMEA was prepared. However, complete masking of the phosphonate group with the dioxolenone unit was not accomplished. A loss in the antiviral activities of hexaethyleneglycol esters and hydroxylated decyl or decyloxyethyl esters of PMEA was noted while the (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of PMEA showed significant activities against HIV and herpesviruses. (S)-HPMPC prodrugs bearing hexaethyleneglycol pro-moiety exhibited significant activities against cytomegalovirus, which were in the same range as (S)-HPMPC. Activities against herpesviruses and poxviruses were one- to fifteen-fold lower than that of (S)-HPMPC. Considering these results, we assume that the prodrugs **3–14** and **16–17** are taken up less efficiently or are not suitable substrates for phospholipase C.

Table 1
Antiviral and cytotoxic activity of the compounds against herpes- and poxviruses in cell culture

Compound	Antiviral activity: EC ₅₀ (μM) ^a								Cytotoxicity (μM)			
	HSV-1		HSV-2	VZV		HCMV		Feline	Vaccinia virus	Cell morphology	Cell growth	
	KOS	KOS ACV ^r	G strain	OKA	07/1.	AD-169	Davis	herpesvirus		(MCC) ^b	(CC ₅₀) ^c	
	(HEL)	(HEL)	(HEL)	(HEL)	(HEL)	(HEL)	(HEL)	(CRFK)	(HEL)	(HEL)	(HEL)	CRFK
3	>233	>233	>233	114	64.3	>233	>233	>233	>233	>233	>233	>233
4	>224	>224	>224	>224	>224	>224	>224	>9	>224	>224	>224	9.4
5	>171	>171	>171	>34	>34	>34	>34	>171	>171	171	>171	>171
6	>162	>162	>162	99	60.7	>162	>162	>162	>162	>162	>162	>162
7	106	122	122	26	18.4	>211	>211	>211	>211	>211	>211	>211
8	>204	>204	>204	61.4 ± 57.4	≥ 72.7 ± 41.5	>204	>204	>204	>204	>204	>204	>204
9	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	148	62.5	>148
10	>142	>142	>142	>142	>142	>28.3	>142	>142	>142	≥ 142	>142	>142
11	>186	>186	>186	18.6 ± 5.2	40.9 ± 18.4	>186	>186	>186	>186	>186	>186	>186
12	>177	>177	>177	74.3	69	>177	>177	>177	>177	>177	146	>177
13	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125
14	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117
15	51.9	51.9	98.6	1.92 ± 0.8	7.6 ± 2.4	259	259	>259	>259	>259	117.2 ± 71.8	>259
16	16.5	12.9	7.4	1.91 ± 0.6	0.60 ± 0.01	0.80 ± 0.20	0.70 ± 0.05	14.5	107	>184	>184	>184
17	1.75	3.5	1.4	1.78 ± 1.09	0.64 ± 0.04	0.95 ± 0.18	0.61 ± 0.13	2.8	15.7	>175	>175	>175
Acyclovir	0.27 ± 0.15	27.1	0.23 ± 0.15	3.0 ± 2.2	69.3 ± 23.5	N.D. ^d	N.D.	N.D.	>250	>222	≥ 1421 ± 422	N.D.
Brivudin	0.028 ± 0.021	274.3	45.7 ± 41.7	0.013 ± 0.01	≥ 105.4 ± 36.9	N.D.	N.D.	N.D.	4.0 ± 2.3	>150	470 ± 165	N.D.
Ganciclovir	0.03 ± 0.02	15.7	0.027 ± 0.0058	N.D.	N.D.	6.0 ± 3.4	6.54 ± 4.0	8.1	>250	>392	435 ± 447	>394
(S)-HPMPC	1.6 ± 1.0	2.0 ± 1.4	1.43 ± 0.60	0.29 ± 0.25	0.10 ± 0.06	0.92 ± 0.82	0.94 ± 0.67	N.D.	7.0 ± 3.8	>317	234 ± 182	N.D.
PMEA	42.5 ± 13.2	41.4 ± 6.6	24.9 ± 14.6	6.7 ± 4.1	5.0 ± 0.8	117.6 ± 48.0	114.6 ± 28.2	N.D.	>183	>183		

^a Effective concentration required to reduce virus-induced cytopathicity by 50%.

^b Minimum cytotoxic concentration that causes a microscopically detectable alteration of the cell morphology.

^c Cytotoxic concentration required to reduce cell growth by 50%.

^d Not determined.

Table 2
Antiviral and cytotoxic activity of the compounds against RNA and retroviruses in cell culture

Compound	Antiviral activity: EC ₅₀ (μM) ^a														Cytotoxicity (μM)				
	HIV-1	HIV-2	VSV		RSV	Coxsackie B4		Para influenza-3	Reovirus-1	Sindbis	Punta Toro	Feline coronavirus (CRFK)	Influenza A H1N1 (MDCK)	Influenza A H3N2 (MDCK)	Influenza B (MDCK)	Cell morphology (MCC) ^b	Cell growth (CC ₅₀) ^c		
	(CEM)	(CEM)	(HEL)	(HeLa)	(HeLa)	(HeLa)	(Vero)	(Vero)	(Vero)	(Vero)	(Vero)					Vero	MDCK	HeLa	(CEM)
3	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233	>233
4	>224	>224	>224	>224	>224	>224	>224	>224	>224	>224	>224	>9	>224	>224	>224	>224	>224	>224	>224
5	>171	>171	>171	>171	99 ± 0	>171	>34	>34	>34	>34	>34	>171	>171	>171	>171	171	>171	>171	>171
6	>162	>162	>162	>162	>162	>162	>162	>162	>162	>162	>162	>162	>32.4	>32.4	>32.4	>162	>162	>162	>162
7	106/122	122/95	>211	>211	>211	>211	>211	>211	>211	>211	>211	>211	>42	>42	>42	>211	>211	>211	>211
8	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204	>204
9	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>29.7	>5.9	>5.9	>5.9	148	29.7	148	13.5
10	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142	>142
11	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186	>186
12	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177	>177
13	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125	>125
14	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117	>117
15	10.6 ± 8.3	9.3 ± 1.7	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	>259	88 ± 9.1
16	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184	>184
17	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175	>175
Oseltamivir	N.D. ^d	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	12 ± 6.5	90 ± 21	23.5 ± 15.2	N.D.	>100	N.D.	N.D.
Ribavirin	N.D.	N.D.	N.D.	14.8 ± 10.7	3.0 ± 10.7	69 ± 50	>250	121 ± 99	≥ 174 ± 96	>250	125 ± 98	N.D.	9.8 ± 1.5	12.5 ± 5.2	7.3 ± 2.4	N.D.	>100	>250	N.D.
PMEA	10.4 ± 4.9	8.6 ± 3.4	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	>250	164 ± 37

^a Effective concentration required to reduce virus-induced cytopathicity by 50%.

^b Minimum cytotoxic concentration that causes a microscopically detectable alteration of cell morphology.

^c Cytotoxic concentration required to reduce cell growth by 50%

^d Not determined.

Table 3Antiviral activity of compounds **16** and **17** compared to TDP-(S)-HPMPC and HDP-(S)-HPMPC and of compound **12** compared to HDP-PMEA

Compound	Number of atoms in alkyloxyalkyl chain	Type of alkyloxyalkyl chain	CMV (AD-169)/EC ₅₀ (μM)
16	18	Hydroxytetraoxaheptadecyl-	0.80 ± 0.20
TDP (S)-HPMPC	18	Tetradecyloxypropyl-	0.002 ± 0.002 ^a
17	20	Pentaoxaecicosyl-	0.95 ± 0.18
HDP (S)-HPMPC	20	Hexadecyloxypropyl-	0.0009 ± 0.0001 ^a
(S)-HPMPC			0.92 ± 0.82
(S)-HPMPC			1.20 ± 0.43 ^a
			HIV-1/EC ₅₀ (μM)
12	20	Pentaoxaecicosyl-	>177
HDP-PMEA	20	Hexadecyloxypropyl-	0.000015 ± 0.00003 ^b
PMEA			10.4 ± 4.9
PMEA			1.1 ± 0.6 ^b

^a Data from Ref. 11, where anti-CMV assays were performed in human foreskin fibroblasts (HFF)^b Data from Ref. 12, where anti-HIV assays were performed in MT-2 cells with the HIV-1 LAI strain.

5. Experimental

Unless stated otherwise, solvents were evaporated at 40 °C and compounds were dried at 100 Pa. The purification of the products by reverse phase HPLC technique was performed on a Waters Delta 600 instrument with a Waters 2487 Dual λ Absorbance Detector using Luna Phenomenex® C-18 preparative columns (10 μM, 21 × 250 mm, flow 12 ml/min); the elution conditions are given in the text. The column chromatography was performed on 60 μM silica gel (Fluka). The ¹H and ¹³C NMR spectra were measured in CDCl₃, D₂O or DMSO-*d*₆ on a Bruker Avance II 600 spectrometer (¹H at 600 MHz, ¹³C at 151 MHz) or Bruker Avance II 500 spectrometer (¹H at 500 MHz, ¹³C at 125.7 MHz). The spectra were referenced to TMS or dioxane (δ 3.75 and 67.19) as internal standards or to the residual solvent signal (δ 2.50 and 39.7 for DMSO). The ³¹P NMR spectra were measured on Bruker Avance II 500 spectrometer (202.3 MHz) in CDCl₃ using H₃PO₄ as the external standard. The general numbering for the assignment of the NMR signals is depicted in Figure 3. Mass spectra were measured

by the ESI technique using LCQ Fleet or LTQ Orbitrap XL (Thermo Fisher Scientific). Most of the chemicals and ion-exchange resins were purchased from Sigma-Aldrich (Czech Republic).

5.1. General procedure for the preparation of PMEA diesters


PMEA (**1**) (273 mg, 1.0 mmol) was suspended in MeOH (5 ml) and dissolved by a solution of tetrabutylammonium hydroxide (2.0 ml, 1 M solution in MeOH). The solution was then evaporated and co-distilled with isopropanol (5 ml) and toluene (3 × 5 ml). The syrupy residue was dissolved in DMF (4 ml) and appropriate bromide (**23**, **24**, **26** or **27**) or tosylate (**28** or **29**) (2.1 mmol) was added. The mixture was stirred at 100 °C (8–12 h), evaporated, after which the residue was chromatographed on a silica gel column (80 g) in 10% MeOH/CHCl₃ (for compounds **12** and **13**) or 6% MeOH/CHCl₃ for all of the others.

5.1.1. Bis(10-Hydroxydecyl) ester of 9-[2-(phosphonomethoxy)ethyl]adenine (**5**)

Compound **18**, prepared by a general procedure from **1** and **26**, was refluxed in methanolic hydrochloric acid (5 ml, pH ~2) for 6 h. The mixture was evaporated, co-distilled with ethanol and crystallized from EtOH–Et₂O. Yield: 270 mg (46%) of crystals; mp 101–104 °C. ¹H NMR (DMSO-*d*₆, ppm) δ: 1.20–1.28 (m, 24H, H-3''–H-8''), 1.39 (m, 4H, H-9''), 1.48 (m, 4H, H-2''), 3.36 (t, 4H, J_{10'-9''} = 6.6, H-10''), 3.80–3.88 (m, 6H, H-1'', H-3'), 3.91 (t, 2H, J_{2'-1'} = 5.0, H-2'), 4.43 (t, 2H, J_{1'-2'} = 5.0, H-1'), 8.44 (s, 1H, H-8), 8.51 (s, 1H, H-2), 8.80 (b s, NH), 9.46 (b s, NH). ¹³C NMR (DMSO-*d*₆, ppm) δ: 25.10 (C-3''), 25.71 (C-8''), 28.71, 29.13, 29.15, 29.24 (C-4'', C-5'', C-6'', C-7''), 30.11 (d, J_{2'-P} = 5.5, C-2''), 32.73 (C-9''), 43.42 (C-1'), 60.89 (C-10''), 63.83 (d, J_{3'-P} = 162.7, C-3'), 65.76 (d, J_{1'-P} = 6.5, C-1''), 70.19 (d, J_{2'-P} = 11.2, C-2'), 118.03 (C-5), 144.24 (C-8), 145.39 (C-2), 148.76 (C-4), 150.72 (C-6). ³¹P NMR (CDCl₃, ppm) δ: 21.38 (m). ESI-MS, *m/z*: 584.2 (100) [M–H][–], 585.2 (32) [M–H][–], 620.0 (18), 652.2 (18). ESI-HRMS calcd for C₂₈H₅₁N₅O₆P 584.3582, found: 584.3578 [M–H][–].

5.1.2. Bis(9,10-Dihydroxydecyl) ester of 9-[2-(phosphonomethoxy)ethyl]adenine (**6**)

Compound **19**, prepared by a general procedure from **1** and **23**, was treated with Dowex 50WX8-400 [H⁺] in methanol/water (1:1, 20 ml) at 60 °C for 1 h. The resin was filtered off, washed with conc. ammonia/methanol (1:4). Yield: 225 mg (46%) of crystallizing syrup. Crystallization from EtOH–Et₂O; mp 72–73 °C. ¹H NMR (DMSO-*d*₆, ppm) δ: 1.16–1.27 (m, 20H, H-3''–H-6'', H-7''a, H-8''a), 1.35–1.41 (m, 4H, H-7''b, H-8''b), 1.47 (m, 4H, H-2''), 3.19–3.27 (m, 4H, H-10''), 3.37 (m, 2H, H-9''), 3.80–3.89 (m, 8H, H-1'', H-3', H-2'), 4.32 (t, 2H, J_{1'-2'} = 5.1, H-1'), 4.35 (d, 2H, J_{OH-9''} = 5.0, OH), 4.43 (t,

**Figure 3.** The general numbering for the assignment of the NMR signals.

2H, $J_{\text{OH}-10''} = 5.7$, OH), 7.19 (b s, 2H, NH₂), 8.06 (s, 1H, H-8), 8.13 (s, 1H, H-2). ¹³C NMR (DMSO-*d*₆, ppm) δ : 25.13 (C-3''), 25.38 (C-7''), 28.74, 29.23, 29.44 (C-4'', C-5'', C-6''), 30.13 (d, $J_{2''-P} = 5.5$, C-2''), 33.62 (C-8''), 42.58 (C-1'), 63.86 (d, $J_{3'-P} = 162.7$, C-3'), 65.79 (d, $J_{1''-P} = 6.5$, C-1''), 66.21 (C-10''), 70.48 (d, $J_{2'-P} = 11.5$, C-2'), 71.29 (C-9''), 118.77 (C-5), 141.18 (C-8), 149.71 (C-4), 152.55 (C-2), 156.14 (C-6). ³¹P NMR (CDCl₃, ppm) δ : 21.37 (m). ESI-MS, *m/z*: 616.2 (100) [M-H][−], 617.2 (30) [M-H][−], 662.0 (27), 684.2 (10), 900.1 (12). ESI-HRMS calcd for C₂₈H₅₁N₅O₈P 616.3481, found: 616.3455 [M-H][−].

5.1.3. Bis[2-[(10-Hydroxydecyl)oxy]ethyl] ester of 9-[2-(phosphonomethoxy)ethyl]adenine (9)

Compound **20**, prepared by a general procedure from **1** and **27**, was treated with Dowex 50WX8-400 [H⁺] in methanol/water (1:1, 20 ml) at 60 °C for 10 h. The resin was filtered off, washed with conc. ammonia/methanol (1:4). Yield: 276 mg (41%) of solid; mp 52 °C. ¹H NMR (DMSO-*d*₆, ppm) δ : 1.20–1.27 (m, 24H, H-5''–H-10''), 1.38 (m, 4H, H-11''), 1.44 (m, 4H, H-4''), 3.30–3.38 (m, 8H, H-3'', H-12''), 3.45 (t, 4H, H-2'', $J_{2''-1''} = 4.7$, H-2''), 3.87–3.89 (m, 4H, H-3', H-2'), 3.99 (m, 4H, H-1''), 4.30–4.33 (m, 2H, H-1'', OH), 7.18 (b s, 2H, NH₂), 8.07 (s, 1H, H-8), 8.12 (s, 1H, H-2). ¹³C NMR (DMSO-*d*₆, ppm) δ : 25.70 (C-10''), 25.80 (C-5''), 29.10, 29.16, 29.22, 29.28, 29.34 (C-4'', C-6'', C-7'', C-8'', C-9''), 32.74 (C-11''), 42.58 (C-1'), 60.91 (C-12''), 63.99 (d, $J_{3'-P} = 163.2$, C-3'), 65.03 (d, $J_{1''-P} = 6.5$, C-1''), 69.29 (d, $J_{2'-P} = 5.6$, C-2''), 70.44 (C-3''), 70.49 (d, $J_{2'-P} = 11.6$, C-2'), 118.75 (C-5), 141.19 (C-8), 149.68 (C-4), 152.53 (C-2), 156.13 (C-6). ³¹P NMR (CDCl₃, ppm) δ : 22.17 (m). ESI-MS, *m/z*: 255.3 (11), 283.3 (21), 672.2 (100) [M-H][−], 673.3 (32) [M-H][−], 694.2 (24), 740.3 (13), 956.1 (11). ESI-HRMS calcd for C₃₂H₅₉N₅O₈P 672.4107, found: 672.4085 [M-H][−].

5.1.4. Bis[2-[(9,10-Dihydroxydecyl)oxy]ethyl] ester of 9-[2-(phosphonomethoxy)ethyl]adenine (10)

Compound **21**, prepared by a general procedure from **1** and **24**, was treated with Dowex 50WX8-400 [H⁺] in methanol/water (1:1, 20 ml) at 60 °C for 1 h. The resin was filtered off, washed with conc. ammonia/methanol (1:4). Yield: 279 mg (40%) of thick syrup. ¹H NMR (DMSO-*d*₆, ppm) δ : 1.17–1.28 (m, 20H, H-5''–H-8'', H-9''a, H-10''a), 1.34–1.41 (m, 4H, H-9''b, H-10''b), 1.45 (m, 4H, H-4''), 3.23 (m, 4H, H-12''), 3.31–3.38 (m, 8H, H-3'', H-11''), 3.45 (t, 4H, $J_{2''-1''} = 4.6$, H-2''), 3.87–3.89 (m, 4H, H-2', H-3'), 3.99 (m, 4H, H-1''), 4.31–4.33 (m, 4H, C-1', OH), 4.41 (t, 2H, $J_{\text{OH}-12''} = 5.7$, OH), 7.18 (b s, 2H, NH₂), 8.08 (s, 1H, H-8), 8.12 (s, 1H, H-2). ¹³C NMR (DMSO-*d*₆, ppm) δ : 25.37 (C-9''), 25.81 (C-5''), 29.09, 29.29, 29.35, 29.46 (C-4'', C-6'', C-7'', C-8''), 33.60 (C-10''), 42.60 (C-1'), 63.99 (d, $J_{3'-P} = 163.3$, C-3'), 65.04 (d, $J_{1''-P} = 6.4$, C-1''), 66.19 (C-12'), 69.29 (d, $J_{2'-P} = 5.7$, C-2''), 70.45 (C-3''), 70.50 (d, $J_{2'-P} = 11.2$, C-2'), 71.28 (C-11''), 118.75 (C-5), 141.21 (C-8), 149.69 (C-4), 152.54 (C-2), 156.13 (C-6). ³¹P NMR (CDCl₃, ppm) δ : 22.84 (m). ESI-MS, *m/z*: 704.2 (100) [M-H][−], 705.5 (33) [M-H][−], 772.2 (11). ESI-HRMS calcd for C₃₂H₅₉N₅O₁₀P 704.4005, found: 704.3981 [M-H][−].

5.1.5. Bis(14-Hydroxy-3,6,9,12-tetraoxooctadecyl) ester of 9-[2-(phosphonomethoxy)ethyl]adenine (13)

The ester **22**, prepared by a general procedure from **1** and **28**, was contaminated with tetrabutylammonium salts. The mixture was treated with Dowex 50WX8-400 [H⁺] in methanol at 40 °C. The suspension was put on a Dowex 50WX8-400 [H⁺] column. Rinsing with water was followed by 1% ammonia. The UV-absorbing eluate was evaporated. Yield: 375 mg (47%) of syrup. ¹H NMR (DMSO-*d*₆, ppm) δ : 3.40 (m, 4H, H-11''), 3.46–3.51 (m, 40H, H-2''–H-10'', H-12''), 3.87–3.90 (m, 4H, H-2', H-3'), 4.00 (m, 4H, H-1''), 4.32 (t, 2H, $J_{1'-2'} = 5.1$, H-1'), 4.59 (t, $J_{\text{OH}-12''} = 5.5$, OH), 7.20 (b

s, 2H, NH₂), 8.08 (s, H-8), 8.13 (s, H-2). ¹³C NMR (DMSO-*d*₆, ppm) δ : 42.64 (C-1'), 60.40 (C-12''), 63.95 (d, $J_{3'-P} = 162.7$, C-3'), 65.04 (d, $J_{1''-P} = 6.5$, C-1''), 69.66–70.00 (m, C-2''–C-10''), 70.46 (d, $J_{2'-P} = 11.2$, C-2'), 72.54 (C-11''), 118.76 (C-5), 141.29 (C-8), 149.70 (C-4), 152.57 (C-2), 156.14 (C-6). ³¹P NMR (CDCl₃, ppm) δ : 22.84 (m). ESI-MS, *m/z*: 413.0 (18), 423.9 (31), 802.3 (14) [M+H]⁺, 824.4 (100) [M+Na]⁺, 825.4 (37) [M+Na]⁺, 826.4 (10) [M+Na]⁺. ESI-HRMS calcd for C₃₂H₆₁N₅O₁₆P 802.3845, found: 802.3844 [M+H]⁺.

5.1.6. Bis(3,6,9,12,15,18-Hexaoxaicosyl) ester of 9-[2-(phosphonomethoxy)ethyl]adenine (14)

The product obtained according to the general procedure using tosylate **29** was contaminated with tetrabutylammonium bromide. It was further purified on a Dowex 50WX8-400 [H⁺] column (5 ml). Rinsing with water was followed by an elution of 1% ammonia. The UV-absorbing fractions were evaporated and co-distilled with ethanol and toluene. Yield: 430 mg (50%) of syrup. ¹H NMR (DMSO-*d*₆, ppm) δ : 1.08 (t, 6H, $J_{13''-14''} = 7.0$, H-14''), 3.41 (q, 4H, $J_{14''-13''} = 7.0$, H-13''), 3.43–3.52 (m, 44H, H-2''–H-12''), 3.87–3.90 (m, 4H, H-2', H-3'), 4.00 (m, 4H, H-1''), 4.32 (t, 2H, $J_{1'-2'} = 5.2$, H-1'), 7.18 (b s, 2H, NH₂), 8.08 (s, 1H, H-8), 8.13 (s, 1H, H-2). ¹³C NMR (DMSO-*d*₆, ppm) δ : 15.32 (C-14''), 42.63 (C-1'), 63.96 (d, $J_{3'-P} = 162.8$, C-3'), 65.04 (d, $J_{1''-P} = 6.5$, C-1''), 65.74 (C-13''), 69.42–70.03 (m, C-2''–C-12''), 70.46 (d, $J_{2'-P} = 11.0$, C-2'), 118.77 (C-5), 141.28 (C-8), 149.70 (C-4), 152.56 (C-2), 156.15 (C-6). ³¹P NMR (CDCl₃, ppm) δ : 22.84 (m). ESI-MS, *m/z*: 333.3 (15), 440.9 (22), 451.9 (42), 858.3 (17) [M+H]⁺, 880.4 (100) [M+Na]⁺. ESI-HRMS calcd for C₃₆H₆₉N₅O₁₆P 858.4471, found: 858.4468 [M+H]⁺.

5.2. General procedure for the preparation of PMEA monoesters

The appropriate diester (0.16 mmol) and LiN₃ (78 mg, 1.6 mmol) in DMF (1 ml) was heated to 100 °C. The conversion was monitored by means of TLC (CHCl₃/MeOH, 9:1). The conversion was complete in 10–16 h.

5.2.1. 10-Hydroxydecyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (3)

The product was crystallized overnight from the reaction mixture. Recrystallization from H₂O–EtOH afforded 50 mg (73%); mp 181–183 °C. ¹H NMR (D₂O, ppm) δ : 0.85–0.95 (m, 6H), 1.03 (m, 2H), 1.11 (m, 2H, H-3''–H-7''), 1.18–1.24 (m, 4H, H-2'', H-8''), 1.48 (m, 2H, H-9''), 3.55–3.62 (m, 6H, H-3', H-1'', H-10''), 3.86 (t, 2H, $J_{2'-1'} = 4.9$, H-2'), 4.40 (t, 2H, $J_{1'-2'} = 4.9$, H-1'), 8.15 (s, 1H, H-2), 8.18 (s, 1H, H-8). ¹³C NMR (D₂O, ppm) δ : 25.27 (C-3''), 25.64 (C-8''), 28.75, 29.10, 29.15, 29.17 (C-4''–C-7''), 30.33 (d, $J_{2''-P} = 6.1$, C-2''), 31.95 (C-9''), 44.34 (C-1'), 62.49 (C-10''), 66.04 (d, $J_{1''-P} = 5.8$, C-1''), 66.54 (d, $J_{3'-P} = 157.9$, C-3'), 71.12 (d, $J_{2'-P} = 14.1$, C-2'), 118.79 (C-5), 143.58 (C-8), 149.25 (C-4), 152.80 (C-2), 155.94 (C-6). ³¹P NMR (D₂O, ppm) δ : 18.03 (m). ESI-MS, *m/z*: 428.2 (100) [M-H][−], 429.3 (21) [M-H][−], 856.8 (15) [2M-H][−], 879.3 (11); ESI-HRMS calcd for C₁₈H₃₁N₅O₅P 428.2068, found: 428.2072 [M-H][−].

5.2.2. 9,10-Dihydroxydecyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (4)

The product was crystallized overnight from the reaction mixture. The recrystallization from H₂O–EtOH afforded 52 mg (73%); mp 257–258 °C. ¹H NMR (D₂O, ppm) δ : 0.85–1.44 (m, 14H, H-2''–H-8''), 3.44 (dd, $J_{\text{gem.}} = 11.7$, $J_{10a''-9''} = 7.0$, H-10''a), 3.54–3.58 (m, 3H, H-1'', H-10''b), 3.61 (d, 2H, $J_{3'-P} = 9.2$), 3.65 (m, 1H, H-9''), 3.87 (m, 2H, H-2'), 4.42 (m, 2H, H-1'), 8.18 (s, 1H, H-2), 8.20 (s, H-8). ¹³C NMR (D₂O, ppm) δ : 25.28, 25.30 (C-3'', C-7''), 28.74, 29.09, 29.21 (C-4'', C-5'', C-6''), 30.34 (d, $J_{2''-P} = 5.7$, C-2''), 32.85

(C-8''), 44.36 (C-1'), 66.05 (d, $J_{1'-p} = 6.1$, C-1''), 66.09 (C-10''), 66.53 (d, $J_{3'-p} = 157.8$, C-3'), 71.09 (d, $J_{2'-p} = 14.0$, C-2'), 72.49 (C-9''), 118.88 (C-5), 143.68 (C-8), 149.38 (C-4), 152.87 (C-2), 156.03 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.02 (m). ESI-MS, m/z : 444.3 (100) $[\text{M}-\text{H}]^-$, 445.3 (19) $[\text{M}-\text{H}]^-$. ESI-HRMS calcd for $\text{C}_{18}\text{H}_{31}\text{N}_5\text{O}_6\text{P}$ 444.2026, found: 444.2017 $[\text{M}-\text{H}]^-$.

5.2.3. 2-[(10-Hydroxydecyl)oxy]ethyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (7)

The reaction mixture was evaporated, the residue was deionized on Dowex 50WX8-400 column ($[\text{H}^+]$, 4 ml, gradient water \rightarrow 2% aqueous ammonia) and then on a Dowex 1 column ($[\text{AcO}^-]$, 4 ml, gradient water \rightarrow 1% aqueous acetic acid). Yield: 67 mg (88%), mp 169–171 °C. ^1H NMR (D_2O , ppm) δ : 1.02–1.21 (m, 10H, H-5''–H-9''), 1.25 (m, 2H, H-10''), 1.30 (m, 2H, H-4''), 1.49 (m, 2H, H-11''), 3.21 (t, 2H, $J_{3''-4''} = 6.6$, H-3''), 3.35 (m, 2H, H-2''), 3.57 (t, 2H, $J_{12'-11'} = 6.7$, H-12''), 3.64 (d, 2H, $J_{3'-p} = 9.1$, H-3'), 3.78 (m, 2H, H-1''), 3.89 (m, 2H, H-2''), 4.42 (m, 2H, H-1'), 8.17 (s, 1H, H-2), 8.21 (s, 1H, H-8). ^{13}C NMR (D_2O , ppm) δ : 25.65 (C-10''), 25.74 (C-5''), 29.02, 29.06, 29.12, 29.23 (C-4'', C-6''–C-9''), 31.94 (C-11''), 44.37 (C-1'), 62.50 (C-12''), 64.34 (d, $J_{1'-p} = 5.7$, C-1''), 66.82 (d, $J_{3'-p} = 158.8$, C-3'), 70.23 (d, $J_{2'-p} = 6.9$, C-2''), 71.17 (d, $J_{2'-p} = 13.7$, C-2'), 71.30 (C-3''), 118.84 (C-5), 143.57 (C-8), 149.33 (C-4), 152.89 (C-2), 156.01 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.30 (m). ESI-MS, m/z : 472.3 (100) $[\text{M}-\text{H}]^-$, 473.3 (22) $[\text{M}-\text{H}]^-$, 944.8 (44), 945.9 (16) $[2\text{M}-\text{H}]^-$. ESI-HRMS calcd for $\text{C}_{20}\text{H}_{35}\text{N}_5\text{O}_6\text{P}$ 472.2330, found: 472.2333 $[\text{M}-\text{H}]^-$.

5.2.4. 2-[(9,10-Dihydroxydecyl)oxy]ethyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (8)

The reaction mixture was evaporated, the residue was deionized on a Dowex 50WX8-400 column ($[\text{H}^+]$, 4 ml, gradient water \rightarrow 2% aqueous ammonia) and then on Dowex 1 column ($[\text{AcO}^-]$, 4 ml, gradient water \rightarrow 1% aqueous acetic acid). Yield: 72 mg (92%), mp 156 °C. ^1H NMR (D_2O , ppm) δ : 1.02–1.46 (m, 14H, H-4''–H-10''), 3.20 (t, 2H, $J_{3''-4''} = 6.7$, H-3''), 3.33 (m, 2H, H-2''), 3.44 (dd, 1H, $J_{\text{gem}} = 11.7$, $J_{12b''-11''} = 7.0$, H-12b''), 3.57 (dd, 1H, $J_{\text{gem}} = 11.7$, $J_{12a''-11''} = 3.8$, H-12a''), 3.62–3.68 (m, 4H, H-3', H-11''), 3.76 (m, 2H, H-1''), 3.89 (m, 2H, H-2''), 4.43 (m, 2H, H-1'), 8.19 (s, 1H, H-2), 8.22 (s, 1H, H-8). ^{13}C NMR (D_2O , ppm) δ : 25.30 (C-9''), 25.72 (C-5''), 28.98, 29.05, 29.15, 29.24 (C-4'', C-6''–C-8''), 32.86 (C-10''), 44.37 (C-1'), 64.33 (d, $J_{1'-p} = 5.7$, C-1''), 66.09 (C-12''), 66.82 (d, $J_{3'-p} = 158.4$, C-3'), 70.22 (d, $J_{2'-p} = 6.6$, C-2''), 71.18 (d, $J_{2'-p} = 13.6$, C-2'), 71.30 (C-3''), 72.50 (C-11''), 118.87 (C-5), 143.61 (C-8), 149.38 (C-4), 152.92 (C-2), 156.04 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.28 (m). ESI MS: 488.2 (100) $[\text{M}-\text{H}]^-$, 489.3 (25) $[\text{M}-\text{H}]^-$. ESI-HRMS calcd for $\text{C}_{20}\text{H}_{35}\text{N}_5\text{O}_7\text{P}$ 488.2280, found: 488.2279 $[\text{M}-\text{H}]^-$.

5.2.5. 17-Hydroxy-3,6,9,12,15-pentaoxaheptadecyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (11)

The reaction mixture was evaporated, the residue was deionized on a Dowex 50WX8-400 column ($[\text{H}^+]$, 4 ml, water) and then on Dowex 1 column ($[\text{AcO}^-]$, 4 ml, gradient water \rightarrow 0.5% aqueous acetic acid). Yield: 70 mg (81%) of thick syrup. ^1H NMR (DMSO , ppm) δ : 3.40 (m, 2H, H-11''), 3.45–3.49 (m, 20H, H-2''–H-10'', H-12''), 3.70 (d, 2H, $J_{3'-p} = 8.4$, H-3'), 3.86–3.90 (m, 4H, H-2', H-1''), 4.32 (t, 2H, $J_{1'-2'} = 5.2$, H-1'), 7.48 (b s, 2H, NH_2), 8.15, 8.16 (s, s, H-2, H-8). ^{13}C NMR (DMSO , ppm) δ : 42.82 (C-1'), 60.44 (C-12''), 64.32 (d, $J_{1'-p} = 5.7$, C-1''), 65.30 (d, $J_{3'-p} = 160.3$, C-3'), 69.82–70.13 (m, C-2''–C-10''), 70.36 (d, $J_{2'-p} = 10.6$, C-2'), 72.56 (C-11''), 118.68 (C-5), 141.75 (C-8), 149.52 (C-4), 151.69 (C-2), 155.53 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.38 (m). ESI-MS, m/z : 255.5 (14), 283.4 (19), 536.3 (100) $[\text{M}-\text{H}]^-$, 618.0 (27). ESI-HRMS calcd for $\text{C}_{20}\text{H}_{35}\text{N}_5\text{O}_{10}\text{P}$ 536.2127, found: 536.2129 $[\text{M}-\text{H}]^-$.

5.2.6. 3,6,9,12,15,18-Hexaoxaicosyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (12)

The reaction mixture was evaporated, the residue was deionized on a Dowex 50WX8-400 column ($[\text{H}^+]$, 4 ml, water) and then on a Dowex 1 column ($[\text{AcO}^-]$, 4 ml, gradient water \rightarrow 0.5% aqueous acetic acid). Yield: 72 mg (80%) of thick syrup. ^1H NMR (DMSO , ppm) δ : 1.08 (t, 3H, $J_{14''-13''} = 7.0$, H-14''), 3.40 (q, 2H, $J_{13''-14''} = 7.0$, H-13''), 3.43–3.49 (m, 22H, H-2''–H-12''), 3.70 (d, 2H, $J_{3'-p} = 8.4$, H-3'), 3.86–3.90 (m, 4H, H-2', H-1''), 4.32 (t, 2H, $J_{1'-2'} = 5.2$, H-1'), 7.51 (b s, 2H, NH_2), 8.15, 8.16 (s, s, H-2, H-8). ^{13}C NMR (DMSO , ppm) δ : 15.40 (C-14''), 42.90 (C-1'), 64.36 (d, $J_{1'-p} = 5.6$, C-1''), 65.35 (d, $J_{3'-p} = 160.1$, C-3'), 65.83 (C-13''), 69.49–70.12 (m, C-2''–C-12''), 70.39 (d, $J_{2'-p} = 11.0$, C-2'), 118.68 (C-5), 141.91 (C-8), 149.52 (C-4), 151.51 (C-2), 155.39 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.39 (m). ESI-MS, m/z : 564.2 (100) $[\text{M}-\text{H}]^-$, 565.2 (25) $[\text{M}-\text{H}]^-$, 586.2 (13), 645.9 (11), 1150.6 (23), 1151.6 (10). ESI-HRMS calcd for $\text{C}_{22}\text{H}_{39}\text{N}_5\text{O}_{10}\text{P}$ 564.2440, found: 564.2440 $[\text{M}-\text{H}]^-$.

5.2.7. (5-Methyl-2-oxo-1,3-dioxolen-4-yl)methyl ester of 9-[2-(phosphonomethoxy)ethyl]adenine (15)

A suspension of **1** (450 mg, 1.65 mmol) in methanol (5 ml) was neutralized with a 1 M solution of methanolic tetrabutylammonium bromide (1.65 ml, 1.65 mmol). PMEA (**1**) was dissolved, the solution was evaporated and co-distilled with ethanol and toluene. The residue was dissolved in anhydrous DMF (10 ml) and (5-methyl-2-oxo-1,3-dioxolen-4-yl)methylbromide⁵ (350 mg, 1.81 mmol) was added. The mixture was stirred for four days at room temperature. The resulting precipitate (400 mg) was purified on a Dowex 1 column ($[\text{AcO}^-]$, 50 ml, gradient water \rightarrow 1% aqueous acetic acid). Unreacted **1** was eluted at first, then **15**. The fractions containing the product were lyophilized. Yield: 100 mg (16%); mp 196 °C. ^1H NMR ($\text{DMSO}-d_6$, ppm) δ : 1.99 (s, 3H, H-4''), 3.67 (d, 2H, $J_{3'-p} = 9.2$, H-3'), 3.88 (m, 2H, H-2'), 4.29 (bd, 2H, $J_{1'-p} = 7.0$, H-1''), 4.41 (m, 2H, H-1'), 8.13 (s, H-2), 8.14 (s, H-8). ^{13}C NMR ($\text{DMSO}-d_6$, ppm) δ : 8.74 (C-4''), 44.29 (C-1'), 55.26 (d, $J_{1'-p} = 4.4$, C-1''), 66.94 (d, $J_{3'-p} = 157.5$, C-3'), 71.18 (d, $J_{2'-p} = 14.3$, C-2'), 118.53 (C-5), 135.25 (d, $J_{2'-p} = 6.8$, C-2''), 140.35 (C-3''), 143.52 (C-8), 149.23 (C-4), 152.79 (C-2), 154.63 (CO), 155.75 (C-6). ^{31}P NMR (D_2O , ppm) δ : 18.20 (m). ESI-MS, m/z : 384.1 (100) $[\text{M}-\text{H}]^-$, 385.1 (13) $[\text{M}-\text{H}]^-$. ESI-HRMS: calcd for $\text{C}_{13}\text{H}_{15}\text{N}_5\text{O}_7\text{P}$ 384.0715, found: 384.0717 $[\text{M}-\text{H}]^-$.

5.3. General procedure for the preparation of (S)-HPMPC esters

Cyclic (S)-HPMPC (**23**) (261 mg, 1.0 mmol, prepared according to lit.^{7a}) was suspended in MeOH (5 ml) and dissolved by a solution of tetrabutylammonium hydroxide (1.0 ml, 1 M solution in MeOH). The solution was evaporated and co-distilled with isopropanol (5 ml) and toluene (3 \times 5 ml). The syrupy residue was dissolved in DMF (4 ml) and the appropriate tosylate (**28** or **29**) (1.05 mmol) was added. The mixture was stirred at 100 °C (8 h), evaporated and the residue was chromatographed on a silica gel column (80 g) in 16% MeOH/ CHCl_3 . The obtained cyclic ester of HPMPC was contaminated with tetrabutylammonium salts.

5.3.1. 14-Hydroxy-3,6,9,12-tetraoxaheptadecyl ester of (S)-1-[3-hydroxy-2-(phosphonomethoxy)propyl]cytosine (16)

The crude ester **24**, prepared by general procedure from **23** and **28**, was heated in 50% aqueous acetic acid (10 ml) at 50 °C for 2 h. The mixture was then evaporated, co-distilled with H_2O . The residue was deionized on a Dowex 50WX8-400 $[\text{H}^+]$ column (12 ml). Rinsing with water was followed with 0.5% ammonia. The UV-adsorbing eluate was concentrated and put on a Dowex 1 $[\text{AcO}^-]$ column (12 ml). The column was washed with water and the product was eluted with 0.25% aqueous AcOH. The UV-adsorbing fractions were evaporated, co-distilled with water, ethanol and

toluene. Yield: 205 mg (38%) of syrup. ^1H NMR (DMSO, ppm) δ : 3.40–3.65 (m, 27H, H-1'b, H-3', H-4', H-2''–H-12''), 3.77 (m, 1H, H-2'), 3.86 (m, 2H, H-1''), 4.02 (dd, 1H, $J_{\text{gem}} = 13.9$, $J_{1'a-2'} = 2.9$, H-1'a), 5.82 (d, 1H, $J_{5-6} = 7.5$, H-5), 7.70 (d, 1H, $J_{6-5} = 7.5$, H-6), 8.73, 9.09 (b s, b s, 2H, NH_2). ^{13}C NMR (DMSO, ppm) δ : 50.17 (C-1'), 60.39 (C-3', C-12''), 63.43 (d, $J_{1''-p} = 5.4$, C-1''), 64.77 (d, $J_{4'-p} = 168.0$, C-4'), 69.89–70.0 (m, C-3''–C-10''), 70.46 (d, $J_{2''-p} = 5.8$, C-2''), 72.53 (C-11''), 79.05 (C-2'), 93.09 (C-5), 149.42 (C-6), 150.89 (C-2), 162.04 (C-4). ^{31}P NMR (D_2O , ppm) δ : 18.49 (m). ESI-MS, m/z : 255.5 (14), 283.4 (19), 536.3 (100) $[\text{M}-\text{H}]^-$, 618.0 (27). ESI-HRMS calcd for $\text{C}_{20}\text{H}_{35}\text{N}_5\text{O}_{10}\text{P}$ 536.2127, found: 536.2129 $[\text{M}-\text{H}]^-$.

5.3.2. 3,6,9,12,15-Pentaoxaecicosyl ester of (S)-1-[3-hydroxy-2-(phosphonomethoxy)propyl]cytosine (17)

The crude ester **26**, prepared by a general procedure from **23** and **29**, was deionized on a Dowex 50WX8-400 $[\text{H}^+]$ column (12 ml), gradient water \rightarrow 0.5% ammonia, after which concentrated UV-adsorbing eluate was put on a Dowex 1 $[\text{AcO}^-]$ (12 ml), gradient water \rightarrow 0.25% acetic acid. Yield: 255 mg (46%) of syrup. ^1H NMR (DMSO, ppm) δ : 1.09 (t, 3H, $J_{14''-13''} = 7.0$, H-14''), 3.41 (q, 2H, $J_{13''-14''} = 7.0$, H-13''), 3.43–3.64 (m, 27H, H-1'b, H-3', H-4', H-2''–H-12''), 3.76 (m, 1H, H-2'), 3.86 (m, 2H, H-1''), 4.01 (dm, 1H, $J_{\text{gem}} = 13.6$, H-1'a), 5.79 (m, 1H, H-5), 7.67 (m, 1H, H-6), 8.51, 8.91 (b s, b s, 2H, NH_2). ^{13}C NMR (DMSO, ppm) δ : 15.33 (C-14''), 50.11 (C-1'), 60.42 (C-3'), 63.50 (C-1''), 64.73 (d, $J_{4'-p} = 158.8$, C-4'), 65.74 (C-13''), 69.43–70.02 (m, C-3''–C-12''), 70.45 (d, $J_{2''-p} = 6.0$, C-2''), 79.14 (C-2'), 93.09 (C-5), 149.16 (C-6), 151.2 (C-2), 162.3 (C-4). ^{31}P NMR (D_2O , ppm) δ : 18.48 (m). ESI-MS, m/z : 570.2 (100) $[\text{M}-\text{H}]^-$, 571.2 (25) $[\text{M}-\text{H}]^-$, 592.1 (13), 1162.9 (22), 1163.9 (10). ESI-HRMS calcd for $\text{C}_{22}\text{H}_{41}\text{N}_3\text{O}_{12}\text{P}$ 570.2433, found: 570.2435 $[\text{M}-\text{H}]^-$.

5.4. Synthesis of functionalized alkyl bromides and tosylates

5.4.1. General procedure for the preparation of diols **22** and **25**

Triphenylphosphine (9.23 g, 35.2 mmol) was added in several portions to a cooled (0 °C) solution of 9-decen-1-ol or 10-undecen-1-ol (32.0 mmol) and CBr_4 (33.6 mmol, 11.1 g) in dichloromethane (50 ml). The reaction mixture was stirred for 1 h at 0 °C and 1 h at room temperature and subsequently evaporated. The residue was filtered through a short silica gel column (5 \times 10 cm) in hexane as a mobile phase. The eluate was evaporated and the crude bromide was dissolved in acetone (150 ml) and *N*-methyl-morpholine *N*-oxide (10 ml of 50% aqueous solution, 48.0 mmol) and osmium tetroxide (2.0 g of 4% aqueous solution) were added. The reaction mixture was stirred for 3 h at room temperature, then concentrated (\sim 50 ml) and partitioned between ether (100 ml) and an aqueous solution of sodium thiosulfate (100 ml). The aqueous layer was washed with ether again (50 ml). The combined ether portions were dried over MgSO_4 , filtered and evaporated. The residue was purified on a short silica gel column (5 \times 10 cm) in ethyl acetate/hexane (3:2). The fractions were evaporated and the product was crystallized from toluene-hexane. Yield: 6.85 g of 10-bromo-1,2-decandiol or 7.31 g of 11-bromo-1,2-dodecanediol (85%).

5.4.1.1. 10-Bromodecane-1,2-diol (22). Yield 6.85 g (85%). Anal. Calcd for $\text{C}_{10}\text{H}_{21}\text{BrO}_2$: C, 47.44; H, 8.36; Br, 31.56. Found: C, 47.59; H, 8.52; Br, 31.43. ESI MS, m/z : 275.1 (100) $[\text{M}+\text{Na}]^+$, 277.1 (94) $[\text{M}+\text{Na}]^+$, 526.1 (19), 528.7 (24). ESI-HRMS calcd for $\text{C}_{10}\text{H}_{21}\text{O}_2\text{BrNa}$ 275.06171, found: 275.06182 and 277.05968 $[\text{M}+\text{Na}]^+$.

5.4.1.2. 11-Bromododecane-1,2-diol (25). Yield 7.31 g (85%). Anal. Calcd for $\text{C}_{11}\text{H}_{23}\text{BrO}_2$: C, 49.45; H, 8.68; Br, 29.90. Found: C,

49.65; H, 8.96; Br, 29.57. ESI MS, m/z : 289.1 (100) $[\text{M}+\text{Na}]^+$, 291.1 (94) $[\text{M}+\text{Na}]^+$, 554.7 (27), 556.6 (37). ESI-HRMS calcd for $\text{C}_{11}\text{H}_{23}\text{O}_2\text{BrNa}$ 289.07736, found: 289.07755 and 291.07537 $[\text{M}+\text{Na}]^+$.

5.4.2. 4-(8-Bromooctyl)-2,2-Dimethyl-1,3-dioxolane (23)

Diol **22** (6.80 g, 26.9 mmol) in 2,2-dimethoxypropane (50 ml) was treated with toluenesulfonic acid (100 mg, 0.53 mmol) at room temperature for 30 min. The solution was then extracted with an aqueous solution of NaHCO_3 (50 ml). The aqueous solution was washed with ether (50 ml) again and the combined organic fractions were dried (MgSO_4) and evaporated. The crude product (7.79 g, 99%) was used in the next step without further purification. ^1H NMR (CDCl_3 , ppm) δ : 1.23–1.45 (m, 10H, H-3–H-7), 1.36 (s, 3H, CH_3), 1.41 (s, 3H, CH_3), 1.50 (m, 1H, H-8), 1.64 (m, 1H, H-8), 1.85 (m, 2H, H-2), 3.41 (t, 2H, $J_{2-1} = 6.9$), 3.50 (t, 1H, $J_{10b-9} = 7.4$, H-10b), 4.02–4.10 (m, 2H, H-10a, H-9). ^{13}C NMR (CDCl_3 , ppm) δ : 25.70 (C-7), 25.74 (CH_3), 26.95 (CH_3), 28.10 (C-3), 28.63, 29.27, 29.51 (C-4, C-5, C-6), 32.77 (C-2), 33.56 (C-8), 33.99 (C-1), 69.50 (C-10), 76.11 (C-9), 108.56 (O–C–O). ESI MS, APCI MS: no mol. peak.

5.4.3. 4-[8-(2-Bromoethoxy)octyl]-2,2-dimethyl-1,3-dioxolane (24)

Ethylene glycol (900 l, 16 mmol) was slowly added to a cooled (0 °C) suspension of NaH (400 mg of 60% suspension in mineral oil, 10 mmol) in dry DMF (3 ml). The suspension was stirred overnight at room temperature, after which bromide **23** (2.35 g, 8.0 mmol) in DMF (3 ml) was added. The mixture was stirred for 24 h. The excess of NaH was decomposed with several drops of MeOH. The solvents were evaporated and the residue was partitioned between chloroform (40 ml) and water (40 ml). The aqueous layer was treated with water again. The combined organic extracts were dried (MgSO_4) and evaporated. The residue was chromatographed on a silica gel column (60 g) in 10% \rightarrow 20% acetone/hexane. The fractions containing the main product were evaporated. The syrupy residue (1.14 g) was dissolved in dichloromethane (10 ml) and CBr_4 (1.46 g, 4.4 mmol) was added. The solution was cooled to 0 °C and triphenylphosphine (1.21 g, 4.6 mmol) was added. The reaction mixture was stirred for 1 h at 0 °C, 1 h at room temperature and subsequently evaporated. The residue was chromatographed on a silica gel column in 5 \rightarrow 10% acetone/hexane. Yield: 1.35 g (50%) of liquid. ^1H NMR (CDCl_3 , ppm) δ : 1.23–1.42 (m, 10H, H-5–H-9), 1.35 (q, 3H, $J_{\text{Me-Me}} = 0.7$, CH_3), 1.41 (q, 3H, $J_{\text{Me-Me}} = 0.7$, CH_3), 1.50 (m, 1H, H-10), 1.58 (m, 2H, H-4), 1.63 (m, 1H, H-10), 3.45–3.51 (m, 5H, H-12a, H-3, H-1), 3.74 (t, 2H, $J_{2-1} = 6.3$, H-2), 4.03 (m, 1H, H-12b), 4.07 (m, 1H, H-11). ^{13}C NMR (CDCl_3 , ppm) δ : 25.71 (C-9), 25.74 (CH_3), 25.99 (C-5), 26.93 (CH_3), 29.28, 29.39, 29.55 (C-6, C-7, C-8), 30.48 (C-1), 33.56 (C-10), 69.50 (C-12), 70.57 (C-2), 71.27 (C-3), 76.12 (C-11), 108.54 (O–C–O). ESI-MS: no mol. peak.

5.4.4. 1-Bromo-10-(methoxymethoxy)decane (26)

A solution of diol **25** (4.13 g, 16.3 mmol) in THF/ H_2O (1:1, 200 ml) was cooled to 0 °C and NaIO_4 (5.3 g, 24.8 mmol) was added while stirring. NaBH_4 (1.1 g, 29.1 mmol) was added 1 h later at 0 °C. The reaction mixture was stirred overnight at room temperature and then concentrated to \sim 100 ml. The aqueous residue was extracted with EtOAc (3 \times 50 ml), combined organic layers were dried (MgSO_4) and evaporated. The residue was dissolved in CH_2Cl_2 (50 ml), the solution was cooled (0 °C) and treated with $i\text{Pr}_2\text{EtN}$ (3.8 ml, 21.8 mmol) and methoxymethyl bromide (1.7 ml, 20.8 mmol). The solution was allowed to heat to room temperature and stirred overnight. The mixture was then washed with an aqueous solution of NaHCO_3 (50 ml), the dichloromethane portion was dried (MgSO_4) and evaporated. The residue was chromatographed

on a silica gel column (200 g) in acetone/hexane (6:100). Yield: 3.55 g (77%) of liquid. ^1H NMR (CDCl_3 , ppm) δ : 1.27–1.88 (m, 16H, H-2–H-9), 3.36 (s, 3H, CH_3), 3.41 (t, 2H, $J_{1-2} = 6.9$, H-1), 3.52 (m, 2H, H-10), 4.62 (s, 2H, O- CH_2 -O). ^{13}C NMR (CDCl_3 , ppm) δ : 26.17–32.80 (m, C-2–C-9), 34.02 (C-1), 55.07 (CH_3), 67.81 (C-10), 96.37 (O-C-O). ESI MS, m/z : 288.3 (47), 289.3 (18), 301.2 (50), 303.2 (26) $[\text{M}+\text{Na}]^+$, 305.2 (25) $[\text{M}+\text{Na}]^+$, 310.3 (95), 335.1 (21), 413.3 (25), 492.4 (44), 509.2 (40), 597.0 (64), 663.2 (100), 685.5 (47), 722.1 (34). ESI-HRMS calcd for $\text{C}_{12}\text{H}_{25}\text{O}_2\text{BrNa}$ 303.09301, found: 303.09293 and 305.09089 $[\text{M}+\text{Na}]^+$.

5.4.5. 1-(2-Bromoethoxy)-10-(methoxymethoxy)decane (27)

Ethyleneglycol (900 l, 16 mmol) was slowly added to a cooled (0°C) suspension of NaH (400 mg of 60% suspension in mineral oil, 10 mmol) in dry DMF (3 ml). The suspension was stirred for 1 h at room temperature, then bromide **26** (2.28 g, 8.1 mmol) in DMF (3 ml) was added. The mixture was stirred for 24 h. The excess of NaH was decomposed with several drops of MeOH. The solvents were evaporated and the residue was partitioned between chloroform and water. The aqueous layer was treated with water again. The combined organic extracts were dried (MgSO_4) and evaporated. The residue was chromatographed on a silica gel column (60 g) in 10%–20% acetone/hexane. Fractions containing the main product were evaporated. The syrupy residue (1.07 g, 4.01 mmol) was dissolved in dichloromethane (10 ml) and CBr_4 (1.46 g, 4.4 mmol) was added. The solution was cooled to 0°C and triphenylphosphine (1.21 g, 4.6 mmol) was added. The reaction mixture was stirred for 1 h at 0°C , 1 h at room temperature and then evaporated. The residue was chromatographed on a short silica gel column in 10% acetone/hexane. Yield: 1.24 g (47%) of liquid. ^1H NMR (CDCl_3 , ppm) δ : 1.26–1.61 (m, 16H, H-4–H-11), 3.46 (t, 2H, $J_{1-2} = 6.3$, H-1), 3.48 (t, 2H, $J_{3-4} = 6.7$, H-3), 3.52 (t, 2H, $J_{12-11} = 6.7$, H-12), 3.74 (t, 2H, $J_{2-1} = 6.3$, H-2), 4.62 (s, 2H, O- CH_2 -O); ^{13}C NMR (CDCl_3 , ppm) δ : 26.02–29.73 (m, C-4–C-11), 30.49 (C-1), 55.07 (CH_3), 67.86 (C-12), 70.59 (C-2), 71.32 (C-3), 96.37 (O-C-O). ESI MS, m/z : 342.1 (100), 344.1 (87), 347.2 (70) $[\text{M}+\text{Na}]^+$, 349.2 (65) $[\text{M}+\text{Na}]^+$, 590.1 (54), 592.1 (90), 594.1 (52), 597.2 (49), 599.2 (28). ESI-HRMS calcd for $\text{C}_{14}\text{H}_{29}\text{O}_3\text{BrNa}$ 347.11923, found: 347.11938 and 349.11714 $[\text{M}+\text{Na}]^+$.

5.4.6. 17-[(Tetrahydro-2H-pyran-2-yl)oxy]-1-tosyloxy-3,6,9,12,15-pentaoxaheptadecan (28)

A mixture of hexaethylene glycol (7.13 g, 25.3 mmol), Ag_2O (8.78 g, 37.9 mmol) and KI (0.84 g, 5.05 mmol) in CH_2Cl_2 (80 ml) was sonicated (15 min). The suspension was cooled down to -30°C and a solution of tosylchloride (4.91 g, 25.8 mmol) in dichloromethane (100 ml) was added dropwise while stirring. The mixture was then gradually heated up to 0°C and kept for 15 min at this temperature. The mixture was dried (MgSO_4) and salts were filtered off. The filtrate was evaporated and the syrupy residue chromatographed on a silica gel column (300 g). The appropriate ditosylate was eluted with EtOAc (0.75 g), the desired monotosylate was eluted with 10% MeOH/EtOAc (8.70 g). The fractions containing monotosylate were evaporated, then co-distilled with toluene (2×50 ml). The residue was dissolved in dichloromethane (80 ml), 2,3-dihydro-2H-pyran (3.0 ml, 28.9 mmol) and pyridinium *p*-toluenesulphonate (200 mg) were added and the mixture was refluxed (6 h). The reaction mixture was cooled down and washed with water (50 ml). The aqueous layer was extracted with EtOAc (4×40 ml). The combined organic portions were dried (MgSO_4) and chromatographed on a silica gel column 300 g in gradient EtOAc \rightarrow 5% MeOH/EtOAc. Yield: 7.75 g (59%). ^1H NMR (CDCl_3 , ppm) δ : 1.48–1.63 (m, 4H, H-3'a, H-4'a, H-5', THP), 1.72 (m, 1H, H-3'b, THP), 1.83 (m, 1H, H-4'b, THP), 2.45 (s, 3H, CH_3), 3.48–4.17 (m, 22H, H-6', THP, H-1–H-10), 4.63 (m, 1H, H-2', THP), 7.34 (m, 2H, H-3'', Ts), 7.80 (m, 2H, H-2'', Ts); ^{13}C NMR (CDCl_3 , ppm) δ : 19.46

(C-4', THP), 21.62 (CH_3), 25.40 (C-5', THP), 30.53 (C-3', THP), 61.67–72.58 (m, C-6', THP, C-1–C-10), 98.93 (C-2', THP), 127.96 (C-2'', Ts), 129.79 (C-3'', Ts), 132.99 (C-1'', Ts), 144.76 (C-4'', Ts). ESI MS, m/z : 538.1 (100), 539.1 (36), 540.1 (10), 543.2 (35) $[\text{M}+\text{Na}]^+$.

5.4.7. 1-Tosyloxy-3,6,9,12,15,18-hexaoxaicosan (29)

Absolute EtOH (0.5 ml, 8.5 mmol) was added dropwise to a stirred suspension of 60% NaH (0.36 g, 9.0 mmol) in anhydrous THF (10 ml). The mixture was stirred for 24 h at room temperature. Subsequently, a solution of tosylate **28** (2.63 g, 5.05 mmol) in THF (20 ml) was added. The mixture was stirred for three additional days at r.t. The excess of NaH was carefully decomposed with MeOH and the solution was concentrated *in vacuo*. The residue was partitioned between EtOAc (20 ml) and H_2O (20 ml), and the aqueous layer was extracted with EtOAc (3×20 ml) again. The combined organic portions were dried (MgSO_4), evaporated and purified on a silica gel column (50 g) in EtOAc. The fractions containing the main product (1.69 g) were evaporated and treated with Dowex 50WX8-400 $[\text{H}^+]$ in methanol (50 ml) overnight. The resin was then filtered off, washed with methanol, and combined filtrates were evaporated and co-distilled with toluene (3×10 ml). The residue (1.25 g) was tosylated with tosylchloride (1.54 g, 8.08 mmol) in a mixture of CH_2Cl_2 (20 ml) and Et_3N (1.3 ml, 9.3 mmol) at room temperature for 24 h. The mixture was washed with water (20 ml), and the aqueous layer was extracted with EtOAc (3×20 ml) again. Combined organic fractions were dried (MgSO_4) and evaporated. The syrupy residue was purified on a silica gel column (70 g) in 20% hexane/EtOAc \rightarrow EtOAc. Yield: 1.70 g (72%). ^1H NMR (CDCl_3 , ppm) δ : 1.21 (t, 3H, $J_{12-11} = 7.0$, H-12), 2.54 (s, 3H, CH_3 , Ts), 3.52 (q, 2H, $J_{11-12} = 7.0$, H-11), 3.57–3.66, 3.69, 4.16 (m, m, m, 16H, 2H, H-1–H-10), 7.34 (m, 2H, H-3', Ts), 7.80 (m, 2H, H-2', Ts); ^{13}C NMR (CDCl_3 , ppm) δ : 15.12 (C-12), 21.62 (CH_3 , Ts), 66.60 (C-11), 68.65, 69.21, 69.78, 70.48, 70.52, 70.53, 70.55, 70.58, 70.62, 70.72 (C-1–C-10), 127.96 (C-2', Ts), 129.79 (C-3', Ts), 132.98 (C-1', Ts), 144.76 (C-4', Ts). ESI MS, m/z : 482.1 (100), 483.1 (24), 487.2 (42), 488.2 (10) $[\text{M}+\text{Na}]^+$. ESI-HRMS calcd for $\text{C}_{21}\text{H}_{36}\text{O}_9\text{SNa}$ 487.19722, found: 487.19737 $[\text{M}+\text{Na}]^+$.

5.5. Antiviral activity assays

The compounds were evaluated against the following viruses: herpes simplex virus type 1 (HSV-1) strain KOS, thymidine kinase-deficient (TK-) HSV-1 KOS strain resistant to ACV (ACV^r), herpes simplex virus type 2 (HSV-2) strains Lyons and G, varicella-zoster virus (VZV) strain Oka, TK- VZV strain 07-1, human cytomegalovirus (HCMV) strains AD-169 and Davis, feline herpesvirus, vaccinia virus Lederle strain, human immunodeficiency virus (HIV) type 1 (IIIB) and type 2 (ROD), respiratory syncytial virus (RSV) strain Long, vesicular stomatitis virus (VSV), Coxsackie B4, Parainfluenza 3, Reovirus-1, Sindbis, Punta Toro, feline coronavirus, influenza A virus subtypes H1N1 and H3N2, and influenza B virus. The antiviral assays, other than HIV, were based on the inhibition of virus-induced cytopathicity or plaque formation in human embryonic lung (HEL) fibroblasts, African green monkey cells (Vero), human epithelial cells (HeLa), Crandell-Rees feline kidney cells (CRFK), or Madin Darby canine kidney cells (MDCK). Confluent cell cultures in microtiter 96-well plates were inoculated with 100 CCID₅₀ of virus (1 CCID₅₀ being the virus dose to infect 50% of the cell cultures) or with 20 plaque forming units (PFU). After 1–2 h adsorption period, the residual virus was removed, and the cell cultures were incubated in the presence of varying concentrations of the test compounds. Viral cytopathicity or plaque formation (VZV) was recorded as soon as it reached completion in the control virus-infected cell cultures that were not treated with the test

compounds. Antiviral activity was expressed as the EC₅₀ or concentration required reducing virus-induced cytopathogenicity or viral plaque formation by 50%. The methodology of the anti-HIV assays was as follows: human CEM ($\sim 3 \times 10^5$ cells/cm³) were infected with 100 CCID₅₀ of HIV-1 (IIIB) or HIV-2 (ROD)/ml and seeded in 200-μL wells of a microtiter plate containing appropriate dilutions of the test compounds. After four days of incubation at 37 °C, HIV-induced CEM giant cell formation was examined microscopically.

5.6. Cytotoxicity assays

Cytotoxicity measurements were based on the inhibition of cell growth. HEL cells were seeded at a rate of 5×10^3 cells/well into 96-well microtiter plates and allowed to proliferate for 24 h. Then, medium containing different concentrations of the test compounds was added. After three days of incubation at 37 °C, the cell number was determined with a Coulter counter. The cytostatic concentration was calculated as the CC₅₀, or the compound concentration required reducing cell proliferation by 50% relative to the number of cells in the untreated controls. CC₅₀ values were estimated from graphic plots of the number of cells (percentage of control) as a function of the concentration of the test compounds. Alternatively, the cytotoxicity of the test compounds was expressed as the minimum cytotoxic concentration (MCC) or the compound concentration that caused a microscopically detectable alteration of cell morphology.

Acknowledgments

This work is a part of the Research Project of the Institute AV0Z40550506 and Centre of New Antivirals and Antineoplastics

1M0508 supported by the Ministry of Education, Youth and Sports of the Czech Republic. It was also supported by Gilead Sciences Inc. (Foster City, CA, USA) and the ‘Geconcerteerde Onderzoeksacties (GOA), Krediet nr. 10/014’ of the K.U. Leuven. The authors would like to thank to the staff of the Mass Spectrometry (Dr. J. Cvačka, Head), the Analytical Department of the institute (Dr. S. Matějková, Head) and Mrs. Julie Křelinová for technical assistance. We also thank Anita Camps, Steven Carmans, Frieda De Meyer, Leentje Persoons, Leen Ingels, Wim Van Dam and Lies Van den Heurck for their excellent technical assistance with the antiviral assays.

References and notes

- De Clercq, E. *Antiviral Res.* **2007**, 75, 1.
- Lee, W. A.; Martin, J. S. *Antiviral Res.* **2006**, 71, 254.
- Zídek, Z.; Kmoníčková, E.; Holý, A. *Biomed. Pap. Med. Fac. Univ. Palacky Olomouc Czech. Repub.* **2005**, 149, 315.
- Sun, C. Q.; Cheng, P. T. W.; Stevenson, J.; Dejneka, T.; Brown, B.; Wang, T. C.; Robl, J. A.; Poss, M. A. *Tetrahedron Lett.* **2002**, 43, 1161.
- Sakamoto, F.; Ikeda, S.; Tsukamoto, G. *Chem. Pharm. Bull.* **1984**, 32, 2241.
- Saneyoshi, M.; Morozumi, M.; Kodama, K.; Machida, H.; Kuninaka, A.; Yoshino, H. *Chem. Pharm. Bull.* **1980**, 28, 2915.
- (a) Beadle, J. R.; Hartline, C.; Aldern, K. A.; Rodriguez, N.; Harden, E.; Kern, E. R.; Hostetler, K. Y. *Antimicrob. Agents Chemother.* **2002**, 46, 2381; (b) Hostetler, K. Y. *Antiviral Res.* **2009**, 82, A84.
- (a) Serafinowska, H. T.; Ashton, R. J.; Bailey, S.; Harnden, M. R.; Jackson, S. M.; Sutton, D. J. *Med. Chem.* **1995**, 38, 1372; (b) Starrett, J. E.; Tortolani, D. R.; Russell, J.; Hitchcock, M. J. M.; Whiterock, V.; Martin, J. C.; Mansuri, M. M. *J. Med. Chem.* **1994**, 37, 1857.
- Holý, A. *Synthesis* **1998**, 29, 381.
- Loiseau, F. A.; Hii, K. K.; Hill, A. M. *J. Org. Chem.* **2004**, 69, 639.
- Wan, W. B.; Beadle, J. R.; Hartline, C.; Kern, E. R.; Ciesla, S. L.; Valiaeva, N.; Hostetler, K. Y. *Antimicrob. Agents Chemother.* **2005**, 49, 656.
- Valiaeva, N.; Beadle, J. R.; Aldern, K. A.; Trahan, J.; Hostetler, K. Y. *Antiviral Res.* **2006**, 72, 10.