

SUBJECT INDEX TO VOLUME 5

- Active enzyme centrifugation, 93
- Active enzyme gel chromatography,
 - computer simulations for, 351
 - of hydrogenases, 339
- Albumin, bovine serum, 27, 34–37, 47–49
 - instability of dilute solutions, 49
- Alpha-bungarotoxin, 27–39
- Archibald method, 161–164
- Associating systems
 - isodesmic self-associations, 171–183
 - mixed associations, 185–196
 - specific self-associations, 161–164, 165–170, 171–183, 185–196
- Base sequence of RNA, from degradation kinetics, 369
- Binding, of proflavine to *M. Lysodeikticus* DNA, 319
- Bouyant densities, proteins, 143
- Bouyant titrations, proteins, 143–145
 - proteins, chemically modified, 150–153
 - synthetic polypeptides, 146–150
- Calorimetry, of DNA–poly-L-lysine interaction, 363
- Calorimetry, of poly G, 359
- Carboxyhemoglobine, transport parameters from
 - scanning gel chromatography, 327
- Computer simulations, for active enzyme gel chromatography, 351
- Creep recovery, of fibrin clots, 377
- Density gradient centrifugation
 - sedimentation equilibrium of proteins, 137–157
 - simulation, 107–135
 - tumor cells, 99–106
- Density markers, 139–140
- Diffusion coefficient, single file, 389
- Diffusion coefficients
 - and sedimentation velocity, 248–250
 - from UV scanner, 51–52
- DNA
 - anomalous sedimentation, 265–270
 - binding of proflavine to *M. Lysodeikticus* DNA, 319
 - calorimetric study of interaction with poly-L-lysine, 363
 - E. Coli*, unsheared, sedimentation, 271–293
 - isolation of fractions from sonicated calf thymus DNA, 285
 - light scattering, sedimentation and viscosity, 301
 - persistence length, 301
- Einstein relation, deviation from, 389
- Fibrin clots, shear creep and creep recovery of, 377
- Fluorescence monitoring, 27–39
- Gel chromatography
 - of active enzyme-substrate complexes, 339
 - scanning, 327
- Gradient forming solute properties, 126–127
- Hemoglobin, 6, 24
- Hemocyanin, 9
 - lobster, 161–164
 - shrimp, 165–170
- Histones, 186–193
- Homoserine dehydrogenase, transport properties from
 - active enzyme gel chromatography, 339
- Hydration, proteins, 143
- Isotherm, of proflavine binding to DNA, 319
- Johnston–Ogston effect, 255–264
- Kinetics, continuous degradation, 369
- Laser, ultra-violet, 27–39
- L-Glutamate dehydrogenase, transport properties
 - from active enzyme gel chromatography, 339
- Light scattering, of DNA, 301
- Lipoproteins, floatation, 127–131
- Methaemoprotein, NMR relaxation rates, 395
- M. Lysodeikticus* DNA, binding of proflavine to, 319

- Molecular weight distribution, 199–205, 207–241
 correction for non-ideality, 220–231
Myoglobin, sperm whale, 72–73
- NMR, relaxation rates in methaemoprotein solutions, 395
- Optical multichannel analyzer, 55–75
- Overspeeding, 193–195
- Persistence length, of double stranded DNA, 301
- Photon counting, 90
- Polyguanylic acid, calorimetry at neutral pH, 359
- Poly-L-lysine, calorimetric study of interaction with DNA, 363
- Polyphenyl-quinoxaline, 202–204
- Precession, 22, 75
- Pressure, osmotic, 209–211
- Pressure effects, 140–141, 161–164
- Proflavine, binding to *M. Lysodeikticus* DNA, 319
- Proteins, plasma, 128–131
- Pycnometer, magnetic float, 203–204
- Rheology, of fibrin clots, 377
- RNA, base sequence determination from continuous degradation kinetics, 369
- Scanner, with on-line computer, 19–26, 27–39, 43–53, 77–96
 television, 55–75
- Scanning gel chromatography, 327
- Schlieren optics, scanning, 50
- Sedimentation equilibrium
- bovine serum albumin, 47–50
- dextran, 213–229
- hemocyanin, 166–169
- hemoglobin, 24
- histones, 191–195
- myoglobin, 73
- polyphenyl-quinoxaline, 202
- Sedimentation, of DNA, 301
- Sedimentation, particle, 133–134
- Sedimentation velocity
- anomalous, 265–270
- bovine serum albumin, 35–37, 51, 92
- dextran, 211–213, 229
- diffusion effects, 248–250
- DNA, unsheared, 271–283
- electric charge effects, 243–251
- Johnston–Ogston effect, 255–264
- Shear creep, of fibrin clots, 377
- Simulation, 107–135, 171–183, 185–196, 201–202, 255–264
- Temperature enhancement of NMR relaxation rates, in methaemoprotein solutions, 395
- Tikhonov regularization, 200–201
- Tumor cells, 99–106
- Viscosity, of DNA, 301
- Volume, partial specific
- dextran, 209
- polyphenyl-quinoxaline, 204
- Volume, reaction, 161–164
- Zone, centrifugation of macromolecules, 132–133