

single channel recordings consistently contain many very rapid transient closings. These features are believed to be important for a detailed understanding of the kinetics of muscarinic channel activation.

In summary, we have identified a K^+ current in single atrial myocytes that is induced by activation of a muscarinic receptor. It exhibits prominent inward-going rectification, a property which in heart is of considerable functional significance activation of muscarinic receptors can hyperpolarize the atrium without completely abolishing the plateau or the upstroke phases of the action potential. More experimental and analytical work is needed to identify the detailed mechanism of this current change, and to compare it with the actions of acetylcholine on the Ca^{++} current.

This work was supported by National Institutes of Health grants HL27454 and HL24820; and by grants 81-243, 81-835 from the American Heart Association and its Texas Affiliate.

Received for publication 5 May 1983.

REFERENCES

1. Giles, W., and S. J. Noble. 1976. Changes in membrane currents in bullfrog atrium produced by acetylcholine. *J. Physiol. (Lond.)* 261:103-123.
2. Garnier, D., J. Narjeot, C. Ojeda, and O. Rougier. 1978. The action of acetylcholine on background conductance in frog atrial trabeculae. *J. Physiol. (Lond.)* 274:381-396.
3. Hume, J. R., and W. Giles. 1981. Active and passive electrical properties of single bullfrog atrial cells. *J. Gen. Physiol.* 78:18-43.
4. Hamill, O. P., A. Marty, E. Neher, B. Sakmann, and F. J. Sigworth. 1981. Improved patch clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch. Eur. J. Physiol.* 391:85-100.
5. Hume, J. R., and W. Giles. 1983. Ionic currents in single isolated bullfrog atrial cells. *J. Gen. Physiol.* 81:153-194.
6. Hutter, O. F. 1961. Ion movements during vagus inhibition of the heart. In *Nervous Inhibition*. E. Florey, editor. Pergamon Press, Oxford. 114-124.

FUNCTIONAL RECONSTITUTION OF RAT STRIATAL DOPAMINE AGONIST RECEPTORS INTO ARTIFICIAL LIPID BIMOLECULAR MEMBRANES

RANDALL B. MURPHY AND VITALY VODYANOV

Department of Chemistry and Laboratory of Radiation and Solid State Physics, New York University, New York, New York 10003

Functional reconstitution of ionic-channel-linked receptor systems into lipid bimolecular membranes (BLM) allows, in principle, the examination of single channels not only from highly purified preparations but also from essentially unenriched membrane fragments. Appropriate pharmacological criteria must be used to define the receptor operationally. The present study reports the reconstitution of an ionic-channel-linked dopamine agonist binding site from rat corpus striatum into artificial, essentially solvent-free lipid bimolecular membranes.

METHODS

Rat striata, obtained from an outbred Sprague-Dawley strain, were homogenized using a Teflon-in-glass homogenizer in buffer (15 mM K^+ -MOPS, pH 7.40, 50 mM sucrose). The homogenate was sedimented at 1,000 g for 5 min (4°C), and the pellet discarded. The supernatant was sedimented at 100,000 g for 15 min. That pellet was resuspended in the above buffer, processed in a sonicator for 2 min, and sedimented (2 min, 12,000 g); the resulting supernatant was "the homogenate" in all subsequent procedures.

The homogenate can be shown to bind dopaminergic ligands such as [3H]-spiroperidol (in the presence of 200 nM ketanserin) with a picomolar K_d .¹ We prepared essentially solvent-free lipid bimolecular

membranes, 2 mm², added homogenate, and measured changes in current under voltage-clamp conditions using previously described methods (2-4).

RESULTS

A striking increase in DC conductance occurs when the dopaminergic agonist apomorphine (7.5 nM final concentration) is added to the *cis* side of homogenate-treated striatal membrane. This change is noticeable ~30 s after agonist addition, and reaches a maximum within 2-4 min. Addition of a higher (25 nM) concentration of apomorphine resulted in a still greater increase in conductivity. Pretreating the membrane with the potent dopaminergic antagonist (+)-butaclamol (5) 5 min prior to adding apomorphine (final butaclamol concentration, *cis* side, 500 nM), we observed no statistically significant activation of the DC conductance by subsequent application of apomorphine up to 300 nM. However, the addition of (-)-butaclamol (final concentration 500 nM), which is ~10,000-fold less potent in its binding to dopamine receptors than its enantiomer (6), had no significant effect upon the dose-dependence of the apomorphine-induced conductance increase. These data are illustrated in Fig. 1.

¹Murphy, R. B. Unpublished observations.




FIGURE 1 Effect of (+)- and (-)-butaclamol on apomorphine-induced conductance increase in lipid bimolecular membranes containing membrane material from rat striatum. Data are illustrated from 11 different membranes all treated with homogenate at a Lowry protein concentration of 50 $\mu\text{g}/\text{ml}$. \circ , no added butaclamol; \blacktriangle , 500 nM (-)-butaclamol added 5 min before apomorphine treatment; \blacksquare , 500 nM (+)-butaclamol added 5 min before apomorphine treatment. A straight line within the limits of experimental precision at $G/G_0 = 0.85$ was observed if pretreatment was carried out with (+)-butaclamol at a concentration of 100 nM. Membrane: egg PC, 20 mM KCl, 20 mM NaCl, 2 mM CaCl_2 , pH 7.4. Clamp voltage: +20 mV.

We also examined the effect of the selective dopaminergic agonist ADTN (7), and observed a qualitatively similar activation to that produced by apomorphine. Adding apomorphine and ADTN up to 1,000 nM concentration to lipid BLM that was untreated with striatal homogenates produced no statistically significant changes in DC conductivity or membrane capacitance.

It is possible to express the concentration dependence of the observed current changes as a linearized Scatchard isotherm. For the high-affinity region the apomorphine K_d is ~ 1.5 nM; the ADTN is ~ 1.2 nM.

Fig. 2 is a record of single-channel activity elicited by apomorphine addition to a membrane treated with striatal homogenate. Analysis of these records yields a conductance of 12.1 ± 0.3 pS (SEM) ($N = 52$) for the apomorphine-elicited channels. The mean open time of these channels is 12 ± 2.6 s (SEM).

Although activation remains unchanged in a variety of bathing media containing chloride, no activation is observed when 40 mM potassium sodium tartarate is used. With choline chloride (40 mM) we observed single channel fluctuations excited by apomorphine (10 nM).

These data are consistent with the hypothesis that a dopamine receptor site can be functionally reconstituted into an artificial, planar, and essentially solvent-free lipid bimolecular membrane. The question remains as to which dopamine receptor subtype we observe. The agonist dissociation constant values that we see are inconsistent with the




FIGURE 2 Single-channel fluctuations from apomorphine treatment of planar lipid bimolecular membranes containing material from membrane fraction of rat striatal homogenates. Bathing electrolyte as in Fig. 1: Membrane: essentially solvent-free bacterial phosphatidylethanolamine (Avanti Biochemicals, Inc., Birmingham, AL); initial capacitance 0.75 $\mu\text{F}/\text{cm}^2$. One vertical mm represents 1.43×10^{-13} A, one horizontal cm represents 10.0 s. A, no homogenate incorporation (bare membrane). B, spontaneous fluctuations induced by the addition of striatal homogenate (final protein concentration 50 $\mu\text{g}/\text{mL}$). C, the steady current fluctuations after the addition of 5 nM apomorphine. All records were obtained under a clamp voltage of +60 mV, which was reduced to 0 mV at the end of each record. Only a portion of each record is illustrated.

association of the agonists with a putative D1 site, and suggest association with either D2 or D3 sites.

This research was supported in part by grant NADA0007 from the U. S. National Bureau of Standards, by grant MH-32755 from the National Institute of Mental Health, and by grant BNS-8118761 from the National Science Foundation.

Received for publication 23 May 1983.

REFERENCES

1. List, S., and P. Seeman. 1981. Resolution of dopamine and serotonin receptor components of [^3H] -spiperone binding to rat brain regions. *Proc. Natl. Acad. Sci. USA*. 78:2620-2624.
2. Vodyanoy, V., and R. B. Murphy. 1983. Single-channel fluctuations in bimolecular lipid membranes induced by rat olfactory epithelial homogenates. *Science (Wash., DC)*. 220:717-719.
3. Vodyanoy, V., and R. B. Murphy. 1982. Solvent-free lipid bimolecular membranes of large surface area. *Biochim. Biophys. Acta*. 687:189-194.
4. Vodyanoy, V., P. Halverson, and R. B. Murphy. 1982. Hydrostatic stabilization of solvent-free lipid bimolecular membranes. *J. Colloid Interface Sci.* 88:247-257.
5. Seeman, P. 1980. Brain dopamine receptors. *Pharmacol. Rev.* 33:229-313.
6. Creese, I. 1983. Receptor interactions of neuroleptics. In *Neuroleptics: Neurochemical, Behavioral, and Clinical Perspectives*. J. T. Coyle and S. J. Enna, editors. Raven Press, New York. 183-222.
7. Seeman, P., G. N. Woodruff, and J. A. Poat. 1979. Similar binding of [^3H]-ADTN and [^3H]-apomorphine to calf brain dopamine receptors. *Eur. J. Pharmacol.* 55:137-142.