

Investigation on the Mechanism of Crystallization of Soluble Protein in the Presence of Nonionic Surfactant

Yanwei Jia, Janaky Narayanan, Xiang-Yang Liu, and Yu Liu
Department of Physics, National University of Singapore, Singapore 117542

ABSTRACT The mechanism of crystallization of soluble, globular protein (lysozyme) in the presence of nonionic surfactant C₈E₄ (tetraoxyethylene glycol mono-octyl ether) was examined using both static and dynamic light scattering. The interprotein interaction was found to be attractive in solution conditions that yielded crystals and repulsive in the noncrystallizing solution conditions. The validity of the second virial coefficient as a criterion for predicting protein crystallization could be established even in the presence of nonionic surfactants. Our experiments indicate that the origin of the change in interactions can be attributed to the adsorption of nonionic surfactant monomers on soluble proteins, which is generally assumed to be the case with only membrane proteins. This adsorption screens the hydrophobic attractive force and enhances the hydration and electrostatic repulsive forces between protein molecules. Thus at low surfactant concentration, the effective protein-protein interaction remains repulsive. Large surfactant concentrations promote protein crystallization, possibly due to the attractive depletion force caused by the intervening free surfactant micelles.

INTRODUCTION

The difficulty of growing protein crystals of suitable size for x-ray crystallography is an impediment to molecular structure determination of proteins. In aqueous solutions, membrane proteins tend to form amorphous aggregates. In the presence of nonionic surfactant, membrane proteins maintain their physiological properties and functions and hence remain soluble. Most importantly, membrane proteins can still be crystallized by more or less conventional means within such an environment (1,2). Therefore, nonionic surfactants are widely used to crystallize membrane proteins since two membrane proteins, bacteriorhodopsin and porin, were successfully crystallized for the first time in 1980 (1,2). As soluble proteins share, to some extent, the aggregation problems of the membrane proteins, the mild nonionic surfactant may also be of use in crystallizing them. A few groups have successfully crystallized soluble proteins with nonionic surfactants (3,4). However, the mechanism of the soluble protein crystallization in the presence of nonionic surfactant is not well understood.

George and Wilson illustrated the importance of intermolecular interactions in the crystal growth of proteins (5). The protein-protein interaction can be characterized by the osmotic second virial coefficient, B_{22} . A positive value of B_{22} refers to predominantly repulsive interactions and a negative value to attractive interactions. It has been suggested that an empirical criterion for soluble protein crystallization is $-8 \times 10^{-4} < B_{22} < -2 \times 10^{-4} \text{ ml mol/g}^2$. The applicability of this criterion to the crystallization of membrane proteins in the presence of nonionic surfactants has been investigated recently (6). However, it is also desirable to examine the mechanism of soluble protein crystallization in the presence

of nonionic surfactants or other amphiphilic molecules and elucidate the importance of B_{22} .

In this work, we use lysozyme as the model protein to examine the protein interactions in nonionic surfactant solutions using static light scattering (SLS) and dynamic light scattering (DLS). The validity of the criterion for B_{22} is discussed by comparing the experimental results with predicted values. At the end, the role played by nonionic surfactant in the modification of protein interactions is investigated in an effort to unravel the mechanism of crystallization.

MATERIALS AND METHODS

Materials

Hen egg white lysozyme (six times purified by crystallization) purchased from Seikagaku (East Falmouth, MA) was used as the model protein. Tetraoxyethylene glycol mono-octyl ether (C₈E₄) purchased from Sigma (St. Louis, MO) was used as the model nonionic surfactant in protein crystallization. High purity deionized water ($\sim 18.2 \text{ M}\Omega$) from Millipore (Billerica, MA) Milli-Q system (0.22 μm) was used for preparing 0.1 M sodium acetate buffer at pH 4.5. The protein was dissolved in the buffer and filtered (0.2 μm) to make a stock solution and was kept in the refrigerator at 4°C for further use. Protein stock solution, surfactant, and buffer were mixed together in required stoichiometric ratios to obtain the desired concentration of the protein and surfactant for the following experiments. Both protein and surfactant were used as received without further purification. All measurements were carried out at 20°C ($\pm 0.1^\circ\text{C}$).

Protein crystallization

Protein crystallization was carried out by batch method using a sample volume of 40 μl . The mixture of C₈E₄/lysozyme solutions were kept in the Eppendorf tubes with the cap sealed by parafilm. The samples were kept in water bath at 20°C. Observation was made under microscope periodically.

Refractive index increment (dn/dc)

Abbe refractometer, T4 type (ATAGO, Tokyo, Japan) was used to measure the refractive index, n , of the mixed solution of lysozyme and C₈E₄ with an

accuracy of 0.001. The light source was a sodium lamp (wavelength = 589 nm). The measurements were carried out at 20°C for a series of lysozyme concentrations, c , in the range 0–50 mg/ml, in C_8E_4 solutions of different concentrations. The refractive index increment, dn/dc , was obtained from the regression analysis of the plot of n versus c .

Static light scattering

All light scattering measurements were performed at 20°C using a Brookhaven light scattering instrument (Holtsville, NY) with BI 9000AT correlator. The light source was a 5 mW He-Ne laser with wavelength, λ , of 633 nm and vertical polarization. Since the molecular size of each of the samples used was $<\lambda/20$, no angular dependence for the excess scattered intensity was expected and all light scattering data were recorded at an angle of 90°.

For dilute protein solutions, the SLS equation is given by Brown (7),

$$\frac{Kc}{R_{90}} = \frac{1}{M_w} + 2B_{22}c, \quad (1)$$

where

$$K = \frac{4\pi^2 n_0^2 (dn/dc)^2}{N_A \lambda^4}, \quad (2)$$

n_0 is a refractive index of the solvent, dn/dc the refractive index increment for the protein/solvent pair, N_A the Avogadro number, λ the wavelength of the incident light in vacuum, c the concentration of the protein (g/ml), and R_{90} the excess Rayleigh ratio at an angle of 90°. The protein concentration was <10 mg/ml.

The absolute R_{90} values were determined from the excess scattered intensities (intensities above that due to solvent, which in our case is the 0.1 M NaAc with/without added surfactant, and background) by calibration of the photometer using toluene as standard. Equation 1 indicates that from a plot of Kc/R_{90} versus c , the molecular weight of the protein M_w can be obtained from the intercept and the second virial coefficient B_{22} can be obtained from the slope of the curve.

Dynamic light scattering

DLS measurements utilize the temporal correlations of the scattering intensity fluctuations, which are related to the Brownian motion of the solute (8,9). For dilute solutions,

$$D_m(c) = D_0(1 + k_d c), \quad (3)$$

where $D_m(c)$ is the mutual diffusion coefficient which is concentration dependant. The parameter D_0 is related to R_H , the hydrodynamic radius of scatterers, and η , the solvent viscosity, through the Stokes-Einstein equation,

$$D_0 = \frac{k_B T}{6\pi\eta R_H}. \quad (4)$$

The parameter k_d can be expressed as

$$k_d = 2M_w B_{22} - k_f - 2\nu, \quad (5)$$

where k_f is the coefficient of the linear term in the development of the friction coefficient factor and ν is the partial specific volume of the protein molecule. k_f can be determined from self-diffusion measurements using pulsed gradient spin-echo (PGSE) NMR. Combining Eqs. 3 and 5, the self-diffusion coefficient D_0 can be obtained from the intercept of linear regression plot of D_m versus c . Moreover, because both k_f and ν are always positive, a positive k_d indicates a positive B_{22} , and a negative B_{22} gives negative values for k_d .

Surface tension

Surface tension was determined using the Wilhelmy plate method with a K14 Krüss tensiometer (accuracy 0.01 mN/m) (Hamburg, Germany). The

measurements were performed at 20°C by gradual increase of surfactant concentration. An appropriate volume of protein stock solution was added to keep the protein concentration constant at 0.2 mg/ml or 2 mg/ml while changing the surfactant concentration. The solution was stirred and allowed to rest for 10 min before each run of measurement. The equilibrium surface tension was calculated by averaging 10 data points collected at the end of each run until the standard deviation became <0.01 mN/m.

RESULTS AND DISCUSSION

Crystallization

A series of lysozyme concentrations from 40 mg/ml to 100 mg/ml and C_8E_4 concentrations from 0.5% to 30% were screened for crystallization. The results are shown in Table 1. Irrespective of the protein concentration, no crystallization occurred in the 0.5% C_8E_4 solution, and the solution remained clear even after 3 months. In 30% C_8E_4 , lysozyme crystals appeared in all the trials (with lysozyme concentration >40 mg/ml) in ~ 1 month. For the intermediate C_8E_4 concentration, amorphous aggregates and/or crystals were obtained at some protein concentrations. In some cases, crystals grew up from the amorphous aggregates. However, these cases were less reproducible.

The crystallization of soluble proteins such as lysozyme, horse heart cytochrome c, bovine red blood cell ubiquitin etc., in the presence of nonionic surfactant, C_8E_4 , has been investigated by Mustafa et al. (3). They used the vapor diffusion method (sitting drop) and batch method (layering 5 μ l of surfactant solution on top of 5 μ l protein solution and allowing slow evaporation of water by keeping one end unsealed). Our results show the same optimal crystallization condition as obtained by them in the batch method. Furthermore, in our study, the concentration of both protein and surfactant were fixed in crystallization trials by sealing the crystallization tube. This would benefit the study of protein molecular interactions by the light scattering experiment as the solvent condition becomes a known parameter.

Refractive index increment (dn/dc)

The refractive index increment, dn/dc , is a prerequisite for the data analysis of SLS (vide Eqs. 1 and 2). As shown in Table 2, the refractive index of the solvent (C_8E_4 in buffer),

TABLE 1 Lysozyme crystallization results in the presence of C_8E_4

Concentration of C_8E_4 (v/v %)	Concentration of lysozyme (mg/ml)	Crystallization results
0.5%	40–100	CS
10%	50–100	CS/AA
20%	40–80	CS/AA
	90	crystal
30%	40–100	crystal

CS, clear solution; AA, amorphous aggregation, with crystal growth from aggregates in a few cases.

TABLE 2 Refractive index and refractive index increment, dn/dc , of lysozyme in C_8E_4 solution

Solvent	n_s	dn/dc (ml/g)
Buffer only	1.335	0.177
0.5% C_8E_4	1.335	0.167
5% C_8E_4	1.341	0.185
10% C_8E_4	1.347	0.142
20% C_8E_4	1.360	0.118
30% C_8E_4	1.371	0.114


n_s , the refractive index of solvent (0.1 M NaAc buffer at pH 4.5 with/without added surfactant C_8E_4).

n_s , increased with C_8E_4 concentration gradually. In the case of buffer alone, n_s was 1.335, whereas for 30% v/v C_8E_4 , n_s was 1.371. In the absence of C_8E_4 , the dn/dc of lysozyme in buffer was 0.177 ml/g, consistent with that from the literatures (10,11). In the presence of C_8E_4 , the variation of dn/dc of lysozyme with C_8E_4 concentration was not very large for low surfactant concentrations, but at high C_8E_4 concentration (30% v/v), the dn/dc decreased to 0.114 ml/g, ~60% of that without C_8E_4 .

We have to mention here that generally dialysis needs to be performed, and the dialysate should be used for dn/dc and SLS measurements. This is to keep the chemical potential of all other solutes except protein a constant during these measurements. The dn/dc of protein in the presence of SDS (sodium dodecyl sulfate) had been studied by other groups (12–15). Dialysis time varied from 1 day to several months (12,14,15). However, results show that the dialysis equilibrium is difficult to achieve even after several weeks and is a function of the cutoff size of the membrane used (12). The dn/dc keeps increasing with time because the solution is not in equilibrium. The reason for the long equilibration time is the slow diffusion of surfactant. The micelles cannot pass through the membrane. The higher the micelle concentration, the longer the time needed to reach equilibrium. Very high surfactant concentrations were used in our crystallization trials. Hence we expected the same tendency of dn/dc in dialysis while using C_8E_4 solutions. Also a huge amount of C_8E_4 is needed to perform dialysis. The dn/dc values reported here are at constant solute concentration rather than at constant chemical potential. Thus we estimate apparent rather than true molecular weights and second virial coefficients.

Static light scattering

Fig. 1 shows the plot of Kc/R_{90} against the lysozyme concentration, c , in the presence of different concentrations of C_8E_4 . As per Eq. 1, the intercept corresponds to the reciprocal molecular weight of protein, and the slope gives the value of the second virial coefficient. The data do not coincide at a single point on the ordinate axis. The molecular weight for lysozyme in buffer is estimated to be 13.7 kDa (Table 3), close to the value given by the manufacturer and the literature (14.3 kDa). In the presence of C_8E_4 , the esti-

**FIGURE 1** Plot of Kc/R versus lysozyme concentration, c , for different C_8E_4 concentrations.

mated molecular weight of lysozyme is from 17.9 kDa to 46.8 kDa, much higher than that obtained in the absence of C_8E_4 . Apart from the approximation in dn/dc values as discussed above, this large value of estimated molecular weight can be attributed to the adsorption of surfactant monomers on protein molecules. This adsorption has been confirmed by the surface tension measurements, which will be discussed later. The molecular weight of C_8E_4 monomer is 306 Da (16). The increase of the molecular weight of protein indicates that ~70 C_8E_4 monomers are adsorbed. This number is less than the aggregation number of C_8E_4 micelle, which has been measured by pulsed-gradient nuclear magnetic resonance (PGNMR) and found to be 85 (17). The binding of nonionic surfactant on protein has been verified for bovine serum albumin (BSA) and is attributed to the hydrophobic interaction (18). It has been found that the strength of binding depends inversely on the number of oxyethylene units in the surfactant. The high number of C_8E_4 adsorbed on lysozyme may be because of the four oxyethylene units of C_8E_4 .

The second virial coefficients obtained from slopes are summarized in Table 3. For lysozyme in buffer, 0.5% and 5% C_8E_4 , the B_{22} values were positive and increased with C_8E_4 concentration, from 4.08×10^{-4} to 233×10^{-4} ml mol/g². This indicates an increase in repulsive interactions between protein molecules in the presence of small amounts of C_8E_4 , which stabilizes the protein solution. When the C_8E_4 concentration increases to 10%, B_{22} starts decreasing. In 20% and 30% C_8E_4 , negative values were obtained for B_{22} , indicating an attractive interaction between protein molecules. Comparing with crystallization results, it can be inferred that for the noncrystallization condition the interaction is repulsive, whereas for crystallization conditions the interaction is attractive and falls in the crystallization window defined for B_{22} . The amorphous aggregation observed in some protein samples in 10% and 20% C_8E_4 cannot be explained by the B_{22} values. However, a possible

TABLE 3 Variations with C₈E₄ concentration of parameters obtained from SLS and DLS according to Eqs. 1–5

Solvent	B_{22} (10^{-4} ml mol/g ²)	Crystallization result	M_w (kDa)	D_0 (10^{-7} cm ² /s)	k_d (ml/g)	R_H (nm)
Buffer	4.08	CS	13.7	12.08	−2.85	1.85
0.5% C ₈ E ₄	4.11	CS	17.9	8.40	23.15	2.66
5% C ₈ E ₄	233	CS	32.1	8.17	61.90	2.73
10% C ₈ E ₄	130	CS/AA	31.3	7.07	8.10	3.16
20% C ₈ E ₄	−2.64	CS/AA/crystal	31.4	7.62	−3.94	2.93
30% C ₈ E ₄	−4.72	crystal	46.8	8.42	−27.27	2.65

CS, clear solution; AA, amorphous aggregation.

explanation is that some free micelles may locally link protein molecules studded with surfactant monomers, causing amorphous aggregation.

Dynamic light scattering

The findings of SLS measurements on interactions of lysozyme molecules in C₈E₄ were verified by determining the mutual diffusion coefficient, D_m , using DLS as shown in Fig. 2. The parameters obtained from SLS and DLS are summarized in Table 3. According to Eqs. 3 and 5, the positive slopes for 0.5%, 5%, and 10% C₈E₄ suggest that B_{22} is positive for these conditions, whereas negative slopes for 20% and 30% C₈E₄ indicate negative B_{22} and a resultant attractive force between lysozyme molecules. For pure lysozyme in buffer, the slope is slightly negative. This is because B_{22} has a small positive value as obtained in SLS (4.08×10^{-4} ml mol/g²). As shown in Eq. 5, k_d can be negative for small positive value of B_{22} because k_f and ν are always positive. The mutual diffusion coefficient values, D_m , for pure lysozyme found in this study are consistent with those found by other authors (13,19,20).

The intercept of D_m versus c gives the self-diffusion coefficient, D_0 , of the scatterers. The hydrodynamic radius, $R_H = 1.85$ nm, evaluated from the intercept D_0 using Eq. 4 agrees well with values reported in the literature for pure

lysozyme in the absence of C₈E₄ (11,13,20). However, in the presence of C₈E₄, irrespective of its concentration, the D_0 value is found to be $\sim 8 \times 10^{-7}$ cm²/s. The viscosity of C₈E₄ was found to almost double for every 10% increase in concentration. Equation 4 was used to extract the R_H value using solvent viscosity η as that of buffer. The R_H value evaluated in the presence of C₈E₄ was ~ 2.7 nm. The reason for this higher R_H value may be twofold: 1), the adsorption of C₈E₄ monomers on lysozyme molecules increases the size of scatterers, and 2), the light scattering of micelles dominates over that of lysozyme molecules. The critical micelle concentration (CMC) of C₈E₄ is 0.2%, as found by surface tension measurements. At all the surfactant concentrations studied here, micelles already exist in a large amount. Therefore, when the protein concentration is low, micelles are predominant scatterers in the solution. Since DLS measures the collective diffusion of macromolecules (protein and micelles), the intercept, D_0 , may correspond to the more prominent scatterers, namely, the micelles. The size of C₈E₄ micelles was measured to be 2.5 nm using PGNMR (17). If the size of protein molecules with adsorbed C₈E₄ monomers is the same as that of micelles, they may not be distinguished by light scattering. Similar observation has been made by Valster et al. for BSA-C₁₂E₈ complexes (12). They concluded that at high surfactant concentration, the hydrodynamic radius of scatterers obtained is the same as free micelles. The dispute over R_H notwithstanding, the variation of D_m with protein concentration gives supportive evidence for the B_{22} measurements of the SLS study.




FIGURE 2 Plot of mutual diffusion coefficient versus lysozyme concentration for various C₈E₄ concentrations.

Surface tension measurement

Composed of hydrophobic and hydrophilic amino acids, protein is surface active in aqueous solution (21). Thus surface tension measurement is an effective method to reveal the properties of surfactants and proteins. To explain the origin of changes in lysozyme molecular interactions in the presence of C₈E₄, surface tension measurement was performed.

The surface tension of pure C₈E₄ in buffer solution was determined before the measurements involving C₈E₄/lysozyme mixtures. Although the measurement of the surface tension of C₈E₄ in pure water has been reported before (22), a surface tension profile for C₈E₄ in sodium acetate buffer

under the specific ionic strength of 0.1 M is not available in the literature. Fig. 3 shows the surface tension plot of C_8E_4 in buffer, together with the profile measured for the mixture under fixed lysozyme concentrations of 0.2 mg/ml and 2 mg/ml. The CMC of C_8E_4 in acetate buffer remains the same (0.2%) as that in pure water—an expected result for nonionic surfactant in salt solution.

Compared with pure C_8E_4 in buffer, the surface tension profiles for the mixed C_8E_4 /lysozyme system are shifted slightly upward for C_8E_4 concentrations < CMC. This is suggestive of the adsorption of C_8E_4 monomers on lysozyme molecules at these concentrations. According to Gibbs's equation (23),

$$\Gamma = -\frac{1}{RT} \frac{d\gamma}{d \ln a}, \quad (6)$$

where Γ is the adsorption amount of surfactant at the interface, R is the gas constant, T is the absolute temperature, γ is the surface tension, and a is the surfactant activity. The surface tension decreases in proportion to the amount of surfactant adsorbed at the interface. When some surfactant monomers are adsorbed on the protein, the activity of surfactant will decrease as the number being adsorbed on the solution surface is less. Therefore, the surface tension will be higher than that without protein. The larger the protein concentration, the greater will be the surface tension, as observed for 0.2 and 2 mg/ml protein and depicted in Fig. 3. Once the hydrophobic sites on the protein are occupied by the monomers, the surface tension profile follows the same path as in the absence of protein. This can be seen at C_8E_4 concentrations \geq CMC as shown in Fig. 3.

A similar result has been obtained for the system of mixed human serum albumin and dodecyl dimethyl phosphine oxide, HAS/ C_{10} DMDO, solutions (24). The quantitative fluorescence studies by Velev et al. also revealed the surfactant adsorption on the protein crystal matrices (25).




FIGURE 3 Plot of equilibrium surface tension of C_8E_4 at different concentrations with and without mixed lysozyme.

Mechanism of protein crystallization with nonionic surfactant

As discussed in the above sections, C_8E_4 monomers tend to adsorb on hydrophobic parts of lysozyme molecules. This adsorption screens the hydrophobic attractive interactions between protein molecules and increases the hydrophilicity of protein surface. Solvation becomes more favorable and results in a higher repulsive protein-protein interaction. As C_8E_4 concentration is increased to 20% or 30% v/v, the interaction becomes attractive and falls in the crystallization window. According to Loll et al., the crystallization slot for membrane protein in the presence of nonionic surfactant lies near the cloud point of the surfactant (6). However, the cloud points for all the solution conditions used by us were above 35°C, far higher than the crystallization temperature, 20°C. These excluded the clouding and phase separation of micelles as a reason for proteins to partition into aqueous phase and crystallize.

Asakura and Oosawa proposed a model to describe the “depletion attraction” between colloidal particles on addition of nonadsorbing polymers (26). Using this model, the globular protein can be approximated as a sphere of radius R , and the free micelle in solution as a sphere of radius R_g . There is a shell of thickness R_g around the protein into which the micelle cannot penetrate (Fig. 4 *a*). When far apart, a uniform osmotic pressure is exerted on protein molecules. As protein molecules come closer, micelles no longer penetrate the shaded region in Fig. 4 *b*. As a consequence, the osmotic pressure is unbalanced and drives protein molecules close together. Thus a net attractive potential is set up between protein molecules. When the concentration of free micelles is large, this excluded volume effect becomes predominant, enhancing the depletion attractive forces, which are short ranged.




FIGURE 4 Depletion mechanism. (*a*) When far apart, a uniform osmotic pressure is exerted on the protein molecules of radius R . (*b*) Micelles of radius R_g cannot enter the region between the closely spaced molecules of radius R . The result is net attraction between protein molecules of radius R .

This model explains why at higher C_8E_4 concentration, the interaction between lysozyme molecules becomes attractive. The depletion force in the region $R_g > R$ in aqueous protein-poly(ethylene glycol) (PEG) solution has been investigated by Kulkarni et al. (27). It was found that the change in B_{22} of protein solution is nonmonotonic as the concentration of PEG is increased. The radii of lysozyme and C_8E_4 micelle studied by us are ~ 1.85 nm and 2.5 nm, respectively, i.e., $R_g > R$. Our results also show a change from initial repulsive interaction to an attractive interaction with increasing C_8E_4 concentration. In a certain range, $\sim 20\%$ – 30% C_8E_4 , the B_{22} falls in the crystallization window resulting in the protein crystallization.

From the above analysis, a mechanism of protein crystallization in the presence of nonionic surfactant can be proposed. As illustrated in Fig. 5 *a*, protein molecules dissolved in buffer have certain hydrophobic patches on the otherwise hydrophilic surface. Small amounts of nonionic surfactant added will adsorb on the hydrophobic parts of the protein (Fig. 5 *b*). Thus the protein solution is further stabilized by screening off the attractive hydrophobic force and increasing the repulsive hydration force. When the amount of added surfactant is large and the number of free micelles is high enough, the attractive depletion force becomes appreciable and B_{22} falls in the crystallization window (Fig. 5 *c*). Therefore, the protein molecules with adsorbed nonionic surfactant are brought closer and form crystals (Fig. 5 *d*). The short range of the depletion force enables a slow nucleation process and hence crystal growth.

CONCLUSION

The nonionic surfactants found useful in membrane protein crystallization can also promote soluble protein crystalliza-

tion under certain conditions. Several experimental techniques were used to investigate the crystallization of the model soluble protein, lysozyme, in the presence of nonionic surfactant, C_8E_4 . Our results suggest the following mechanism for protein crystallization. The adsorption of surfactant on hydrophobic parts of protein molecules screens the hydrophobic attraction between protein molecules, increases the solvation, and results in an increase in repulsive interaction. However, at high surfactant concentration, the attractive depletion force due to the presence of free micelles becomes dominant and enables the protein molecules to assemble in crystalline order.

We thank Dr. Claire Lesieur Chungkham for enlightening discussions. We also thank Ms. Wang Lin and the Institute of Materials Research and Engineering for the help in surface tension and viscosity measurements.

REFERENCES

1. Michel, H., and D. Oesterhelt. 1980. Three-dimensional crystals of membrane proteins: bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA*. 77:1283–1285.
2. Garavito, R. M., and J. P. Rosenbush. 1980. Three-dimensional crystals of an integral membrane protein: an initial x-ray analysis. *J. Cell Biol.* 86:327–329.
3. Mustafa, A. O., J. P. Derrick, G. J. T. Tiddy, and R. C. Ford. 1998. A novel approach for the crystallization of soluble proteins using non-ionic surfactants. *Acta Crystallogr. D*. 54:154–158.
4. McPherson, A., S. Koszelak, H. Axelrod, J. Day, L. Robinson, M. McGrath, R. Williams, and D. Cascio. 1986. The effects of neutral detergents on the crystallization of soluble proteins. *J. Cryst. Growth*. 76:547–553.
5. George, A., and W. W. Wilson. 1994. Predicting protein crystallization from a dilute solution property. *Acta Crystallogr. D*. 50:361–365.
6. Loll, P. J., M. Allaman, and J. Wiencek. 2001. Assessing the role of detergent-detergent interactions in membrane protein crystallization. *J. Cryst. Growth*. 232:432–438.
7. Brown, W. 1996. *Light Scattering: Principles and Development*. Oxford University Press, New York.
8. Brown, W., and T. Nicolai. 1993. *Dynamic Light Scattering: The Method and Some Applications*. W. Brown, editor. Oxford University Press, New York.
9. Teraoka, I. 2002. *Polymer Solutions: An Introduction to Physical Properties*. Wiley, New York.
10. Huglin, M. B. 1972. *Light Scattering from Polymer Solutions*. M. B. Huglin, editor. Academic Press, London and New York.
11. Narayanan, J., and X. Y. Liu. 2003. Protein interactions in under-saturated and supersaturated solutions: a study using light and x-ray scattering. *Biophys. J.* 84:523–532.
12. Valstar, A., M. Almgren, and W. Brown. 2000. The interaction of bovine serum albumin with surfactants studied by light scattering. *Langmuir*. 16:922–927.
13. Valstar, A., W. Brown, and M. Almgren. 1999. The lysozyme-sodium dodecyl sulfate system studied by dynamic and static light scattering. *Langmuir*. 15:2366–2374.
14. Gimel, J. C., and W. Brown. 1996. A light scattering investigation of the sodium dodecyl sulfate-lysozyme system. *J. Chem. Phys.* 104: 8112–8117.
15. Jones, M. N., and P. Manley. 1979. Binding of n-alkyl sulphates to lysozyme in aqueous solution. *J. Chem. Soc., Faraday Trans. 1*. 75: 1736–1744.




FIGURE 5 Mechanism of protein crystallization in the presence of nonionic surfactant. (*a*) Protein in solution; (*b*) hydrophobic tails of nonionic surfactant monomers adsorb on the hydrophobic parts of protein molecules; (*c*) when the micelle concentration is high enough, the depletion force is predominant and fine tunes the crystallization; and (*d*) protein crystal with adsorbed nonionic surfactant.

16. van Os, N. M., J. R. Haak, and L. A. M. Rupert. 1993. *Physico-Chemical Properties of Selected Anionic, Cationic and Nonionic Surfactants*. Elsevier, New York.
17. Faucompre, B., and B. Lindman. 1987. Self-association of zwitterionic and nonionic surfactants. NMR self-diffusion studies. *J. Phys. Chem.* 91:383–389.
18. Sukow, W. W., H. E. Sandberg, E. A. Lewis, D. J. Eatough, and L. D. Hansen. 1980. Binding of the triton X series of nonionic surfactants to bovine serum albumin. *Biochemistry*. 19:912–917.
19. Eberstein, W., Y. Georgalis, and W. Saenger. 1994. Molecular interactions in crystallizing lysozyme solutions studied by photo correlation spectroscopy. *J. Cryst. Growth*. 143:71–78.
20. Muschol, M., and F. Rosenberger. 1995. Interactions in undersaturated and supersaturated lysozyme solutions: static and dynamic light scattering results. *J. Chem. Phys.* 103:10424–10432.
21. Magdassi, S. 1996. *Surface Activity of Proteins: Chemical and Physicochemical Modifications*. Marcel Dekker, New York.
22. Degiorgio, V., and M. Corti. 1985. *Physics of Amphiphiles-Micelles, Vesicles, and Microemulsions: Proceedings of the International School of Physics, Enrico Fermi, Course Xc*. Elsevier Science, Amsterdam.
23. Adamson, A. W., and A. P. Gast. 1997. *Physical Chemistry of Surfaces*. John Wiley & Sons, New York.
24. Miller, R., V. B. Fainerman, A. V. Makievski, J. Krägel, D. O. Grigoriev, V. N. Kazakov, and O. V. Sinyachenko. 2000. Dynamics of protein and mixed protein/surfactant adsorption layers at the water/fluid interface. *Adv. Colloid Interface Sci.* 86:39–82.
25. Velev, O. D., E. W. Kaler, and A. M. Lenhoff. 2000. Surfactant diffusion into lysozyme crystal matrices investigated by quantitative fluorescence microscopy. *J. Phys. Chem. B*. 104:9267–9275.
26. Asakura, S., and F. Oosawa. 1958. Interaction between particles suspended in solutions of macromolecules. *J. Polym. Sci. [B]*. XXXIII:183–192.
27. Kulkarni, A. M., A. P. Chatterjee, K. S. Schweizer, and C. F. Zukoski. 1999. Depletion interactions in the protein limit: effects of polymer density fluctuations. *Phys. Rev. Lett.* 83:4554–4557.