

other methods to determine the degree of structural perturbation caused by mutation of the EF-hands. Inhibition of  $\text{Ca}^{2+}$  binding to the N-domain in CaM12 does not affect  $\text{Ca}^{2+}$  binding to the C-domain, however, inhibition of  $\text{Ca}^{2+}$  binding to the C-domain in CaM3 and CaM34 significantly increases the  $\text{Ca}^{2+}$ -binding affinity of the N-domain by decreasing the  $k_{\text{off}}$  for  $\text{Ca}^{2+}$ . This was associated with increased exposure of hydrophobic regions in the N-domain as detected by ANS fluorescence. Significantly, 1H-15N HSQC spectra collected in the absence of  $\text{Ca}^{2+}$  show large structural perturbations in the C-domain of CaM3, CaM4 and especially CaM34 relative to apo-CaM. This was observed as resonance broadening and a loss of dispersion. These data indicate that conversion of Asp 93 and 129 to Ala destabilizes the C-domain of apo-CaM.

#### 598-Pos Board B477

##### Interactions of the Anti-Psychotic Drug Trifluoperazine with Calmodulin

Michael D. Feldkamp, Madeline A. Shea.

University of Iowa, Iowa City, IA, USA.

Calmodulin (CaM) is a  $\text{Ca}^{2+}$ -sensing protein essential to eukaryotic signal transduction pathways. It has two homologous domains (N and C), each binding two  $\text{Ca}^{2+}$  ions. The anti-psychotic drug trifluoperazine (TFP; Stelazine) is a CaM antagonist known to bind hydrophobic clefts of CaM that are exposed upon  $\text{Ca}^{2+}$  binding.

Equilibrium  $\text{Ca}^{2+}$  titrations monitored by changes in steady-state fluorescence of intrinsic Phe and Tyr residues were used to evaluate the effect of TFP on the  $\text{Ca}^{2+}$  affinity of full length CaM (CaM<sub>1-148</sub>), N-domain (CaM<sub>1-80</sub>) and C-domain (CaM<sub>76-148</sub>) over a range of TFP:CaM ratios. Low levels of TFP (1:1, 2:1 ratios) decreased the  $\text{Ca}^{2+}$  affinity of CaM. TFP had the greatest effect on  $\text{Ca}^{2+}$  binding to sites III and IV, in the C-domain of CaM<sub>1-148</sub>, but affected both domains. At an 8:1 ratio of TFP:CaM, the effect reversed and the  $\text{Ca}^{2+}$  affinity of CaM increased.

<sup>1</sup>H-<sup>15</sup>N-HSQC NMR showed that resonances assigned to apo and  $\text{Ca}^{2+}$ -saturated C-domain were the most perturbed during TFP titration, while a smaller subset of N-domain resonances were affected. The stoichiometry of TFP binding to apo-CaM<sub>1-148</sub> was determined to be 2:1, and 4:1 for (Ca<sup>2+</sup>)<sub>4</sub>-CaM.

Crystallographic structures of TFP bound to (Ca<sup>2+</sup>)<sub>4</sub>-CaM<sub>1-148</sub> indicate two possible orientations of TFP when bound in 1:1 vs 2:1 and 4:1 TFP:CaM ratios. A new structure of a (Ca<sup>2+</sup>)<sub>2</sub>-CaM<sub>76-148</sub>-TFP complex showed the trifluoromethyl group of TFP in both positions seen previously; distinct conformation of Met 144 correlated with orientation of TFP. NMR of apo-CaM<sub>76-148</sub> will be used to determine whether apo CaM-TFP complex adopts the semi-open conformation of apo CaM bound to a myosin peptide.

#### 599-Pos Board B478

##### Integration of Extracellular and Intracellular Calcium Signals via Calcium-Sensing Receptor (CaSR)

Yun Huang<sup>1</sup>, Yubin Zhou<sup>1</sup>, Adriana Castiblanco<sup>1</sup>, Hing-Cheung Wong<sup>1</sup>, Yangyi Chen<sup>1</sup>, Wei Yang<sup>1</sup>, Edward M. Brown<sup>2</sup>, Jenny J. Yang<sup>1</sup>.

<sup>1</sup>Georgia State University, Atlanta, GA, USA, <sup>2</sup>Brigham and Women's Hospital, Boston, MA, USA.

$\text{Ca}^{2+}$ , both as a first and a second messenger, is closely involved in the modulation and regulation of numerous important cellular events, such as cell proliferation, differentiation and cell death. Fine-tuned  $\text{Ca}^{2+}$  signaling is achieved by its reversible or irreversible binding to a repertoire of  $\text{Ca}^{2+}$  signaling molecules. Among them, the extracellular calcium sensing receptor (CaSR) senses  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_{\text{e}}$ ) in the milieu outside of cells where  $\text{Ca}^{2+}$  serves as a first messenger. An array of naturally-occurring mutations in CaSR has been found in patients with inherited disorders of  $\text{Ca}^{2+}$  homeostasis, leading to abnormal intracellular responses toward  $[\text{Ca}^{2+}]_{\text{e}}$ .

In the present study, we have computationally predicted and experimentally characterized the metal-binding properties of five  $\text{Ca}^{2+}$ -binding pockets within the extracellular domain of CaSR. Two complementary methods of grafting approach and the subdomain approach were used to probe site specific and cooperative metal binding as well as metal induced conformational change. Based on our results, a model has been proposed to explain the distinct CaSR-mediated responses toward diseases related-abnormally "high" or "low" extracellular  $\text{Ca}^{2+}$  levels. We here further demonstrate that the cytosolic terminal is essential for proper intracellular  $\text{Ca}^{2+}$  response to external signals.

#### 600-Pos Board B479

##### Altered Calcium Handling Between Healthy And Atherosclerotic Vascular Smooth Muscle

Marie-Ann Ewart<sup>1</sup>, Susan Currie<sup>2</sup>, John G. McCarron<sup>2</sup>, Simon Kennedy<sup>1</sup>.

<sup>1</sup>University of Glasgow, Glasgow, United Kingdom, <sup>2</sup>University of Strathclyde, Glasgow, United Kingdom.

In order for smooth muscle (SM) contraction and relaxation to proceed efficiently,  $\text{Ca}^{2+}$  handling is under tight regulation. The cyclic strain associated with hypertension is thought to play an initiating role in atherosclerosis, suggesting dysregulation of SM  $\text{Ca}^{2+}$  handling may be a contributing factor. Peroxynitrite (ONOO<sup>-</sup>), the reaction product of superoxide and nitric oxide, forms in diseased vessels and has been demonstrated to induce SM cell relaxation. In this study, we assessed function and expression levels of sarcoplasmic reticulum  $\text{Ca}^{2+}$  handling proteins; the inositol 1,4,5-trisphosphate receptor (IP<sub>3</sub>R) and the  $\text{Ca}^{2+}$ -ATPase (SERCA) in both healthy and atherosclerotic aorta. ONOO<sup>-</sup> dose-dependently relaxed U46619 pre-contracted aorta from both control and atherosclerotic ApoE<sup>-/-</sup> mice (2 months high fat diet) [ $51.2 \pm 4.7\%$  and  $78.5 \pm 4.3\%$  maximal relaxation, respectively ( $3 \times 10^{-5}$  ONOO<sup>-</sup>)]. This relaxation was antagonised in both C57 and ApoE<sup>-/-</sup> by the addition of either 3  $\mu\text{M}$  thapsigargin (TG), a SERCA inhibitor, or 60  $\mu\text{M}$  2-aminoethoxydiphenyl borate (2-APB), an IP<sub>3</sub>R blocker. In control aorta, relaxation was  $4.3 \pm 5\%$  (TG),  $p < 0.001$ ;  $n=7$  and  $14.6 \pm 6.2\%$  (2-APB)  $p=0.001$ ;  $n=9$ . In ApoE<sup>-/-</sup> aorta, % relaxation was  $22.09 \pm 3.1$  (TG)  $p < 0.001$ ;  $n=8$  and  $7 \pm 5.9\%$  (2-APB)  $p < 0.001$ ;  $n=7$ . There was no significant difference between endothelial denuded or intact vessels. These data indicate an alteration in the effect of  $\text{Ca}^{2+}$  handling protein inhibitors between control and ApoE<sup>-/-</sup> mice. This has been further correlated to expression of both SERCA and IP<sub>3</sub>R proteins. Studies with the potassium channel blocker tetraethylammonium (TEA) indicate plasma membrane hyperpolarisation is an effector of ONOO<sup>-</sup> induced relaxation [ $31.2\%$  relaxation reduction with TEA in C57 aorta,  $p=0.002$  vs  $35.5\%$  reduction in ApoE<sup>-/-</sup>,  $p=0.022$ ]. We provide additional evidence, through myography and biochemical analysis, of a time-dependent correlation between atherosclerotic development and SM  $\text{Ca}^{2+}$  handling machinery modulation.

#### 601-Pos Board B480

##### PEP-19 is an Intrinsically Disordered, Acidic/IQ Motif Regulator of Calmodulin Signaling

Quinn K. Kleerekoper, Liangwen Xiong, Xu Wang, John A. Putkey.

University of Texas Medical School, Houston, TX, USA.

PEP-19 is a small calmodulin (CaM) binding protein that inhibits apoptosis and protects cells against  $\text{Ca}^{2+}$ -toxicity. It binds to either apo or  $\text{Ca}^{2+}$ -CaM and greatly increases the  $k_{\text{on}}$  and  $k_{\text{off}}$  of  $\text{Ca}^{2+}$  binding but does not affect  $K_{\text{Ca}}$ . Here we investigate the molecular basis for modulation of  $\text{Ca}^{2+}$  binding to CaM by PEP-19. First, we identified an extended IQ motif that includes an N-terminal acidic sequence that is necessary for modulation of  $\text{Ca}^{2+}$  binding to CaM, and show the acidic/IQ motif is present in a variety of proteins from different species. Although PEP-19 binds to apo and  $\text{Ca}^{2+}$ -CaM with similar affinity, the  $k_{\text{off}}$  and  $k_{\text{on}}$  for binding to apo CaM are at least 50-fold slower than for  $\text{Ca}^{2+}$ -CaM, however, simulations show that these differences would not inhibit transfer of CaM from PEP-19 to a  $\text{Ca}^{2+}$ -dependent target protein during a  $\text{Ca}^{2+}$  pulse. Sequence analysis, CD and NMR show that PEP-19 is an intrinsically disordered protein, but with residual structure localized to its acidic/IQ motif. We also show that PEP-19 persists in a partially folded state when bound to either apo or  $\text{Ca}^{2+}$ -CaM, a feature of protein-protein interactions that has been called a fuzzy complex. These data show PEP-19 to be a representative of a class of acidic/IQ regulators of CaM signaling. They also support models in which intrinsic disorder confers plasticity that allows PEP-19 to bind to either apo or  $\text{Ca}^{2+}$  CaM, and that complex formation may be facilitated by conformational selection of residual structure in the acidic/IQ sequence. Moreover, conformational exchange of bound PEP-19 in a fuzzy complex with CaM could exert an allosteric effect that modulates or gates the  $k_{\text{on}}$  and  $k_{\text{off}}$  rates for binding  $\text{Ca}^{2+}$  to the C-domain of CaM.

#### 602-Pos Board B481

##### Preferential Binding and Orientation of Recoverin to Phospholipid Monolayers

Philippe Calvez, Julie Boucher, Philippe Desmeules, Christian Salesse.

Centre de recherche du CHUQ, Québec, QC, Canada.

Recoverin is a 201 amino acids calcium-myristoyl switch protein that is responsible for the regulation of the phosphorylation of the visual pigment rhodopsin. Calcium binding to myristoylated recoverin leads to a conformational change, which exposes its hydrophobic residues and its myristoyl moiety. We have previously demonstrated that the myristoyl group highly accelerates the membrane binding of recoverin in the presence of calcium. However, it is still unknown whether recoverin shows preferential membrane binding towards highly polyunsaturated phospholipids such as those found in photoreceptor membranes. In this study, we performed monolayer measurements to analyze the affinity of recoverin for different phospholipids that are representative of these membranes. We concluded that the affinity of recoverin increases with fatty acyl chain length and unsaturation of the phospholipids. In addition, we observed a preferential binding of recoverin for didocosahexaenoyl phosphatidylethanolamine

with a maximum insertion pressure of 30 mN/m, which is in the range of the lateral pressure postulated for biological membranes. Moreover, results show that the size and the charge of the polar head group of phospholipids are also implicated in recoverin adsorption in monolayers. Furthermore, polarization-modulated infrared reflection absorption spectroscopy (PM-IRRAS) has been used to determine the secondary structure and orientation of recoverin in monolayers. PM-IRRAS spectra indicated the prevalence of  $\alpha$ -helices in the secondary structure of recoverin, which is consistent with its known structure. In contrast, non-myristoylated recoverin is quickly denatured after its adsorption in monolayers. Finally, the amide I/amide II ratio allowed to determine that the  $\alpha$ -helices of myristoylated recoverin are oriented perpendicular to the plane of the monolayer.


#### 603-Pos Board B482

##### Characterization Of $\text{Ca}^{2+}$ Photo-release From DM-nitrophen Using Photothermal Beam Deflection

Gangadhar Dhulipala, Jaroslava Miksovská.

Florida International University, Miami, FL, USA.

DM-nitrophen is a popular caged calcium compound that allows for a rapid increase in calcium concentration from hundred nanomolar to tens to hundred micromolar level on the microsecond timescale. To fully understand the mechanism of calcium photo-release from DM-nitrophen, we have used photothermal beam deflection technique to investigate time-profiles of volume and enthalpy changes associated with DM-nitrophen photodissociation. Our data show that the photodissociation of calcium loaded DM-nitrophen occurs as a two-step process. The first step takes place within  $\sim 10 \mu\text{s}$  upon photolysis and is associated with a volume decrease of  $-7 \text{ mL mol}^{-1}$  and enthalpy change of  $66 \text{ kcal mol}^{-1}$ . On the longer timescale ( $\tau = 200 \mu\text{s}$ ), the second event with a positive volume change of  $7 \text{ mL mol}^{-1}$  and enthalpy change of  $8 \text{ kcal mol}^{-1}$  was detected. These data are in agreement with the previous fluorescence studies showing that calcium release from DM-nitrophen occurs as a two step process.



#### 604-Pos Board B483

##### Altered Structure Of The Cerebellar Granule Cell Layer Of Mice Lacking Calretinin

Patrick Bishop, Céline Roussel, Serge Schiffrmann, David Gall.

Université Libre de Bruxelles, Bruxelles, Belgium.

Calcium binding proteins, such as calretinin, are abundantly expressed in distinctive patterns in the central nervous system, but their physiological functions remain poorly understood. Calretinin is highly expressed in cerebellar granule cells and calretinin deficient mice exhibit alterations in motor coordination. Using confocal microscopy, we demonstrate that the cerebellar cortex of calretinin deficient mice exhibit a significantly decreased granule cells density. Interestingly, it has been shown that the migration of granule cells is tightly associated with intracellular calcium oscillations. Therefore, we hypothesize that an alteration of these calcium oscillations in calretinin deficient mice could be involved in the observed morphological alterations. To test this assumption, we are currently developing two strategies. First, using confocal microscopy and cerebellar microexplant culture, we are studying calcium oscillations occurring during granule cell migration in the wild type control and calretinin knock-out mice. This allow us to characterize the impact of variations in calcium buffering capacity over neuronal development and on the generation of the calcium oscillations observed during the granule cell migration. On the other hand, we are developing a theoretical model to study the impact of calcium buffering modifications on the dynamics underlying the observed calcium oscillations. This dedicated computational model will shed light on the possible mechanism responsible for the modulation, by calretinin, of calcium oscillations during the granule cell migration.

#### 605-Pos Board B484

##### Characterization Of Zebrafish (Danio Rerio) NCX4: A Novel Na/Ca Exchanger With Distinct Electrophysiological Properties

Glen F. Tibbitts.

Simon Fraser University, Burnaby, BC, Canada.

Members of the  $\text{Na}^{+}$ - $\text{Ca}^{2+}$  exchanger (NCX) family are important regulators of cytosolic  $\text{Ca}^{2+}$  in myriad tissues and are highly conserved across a wide range of species. Three distinct NCX genes and numerous splice variants exist in mammals, many of which have been characterized in a variety of heterologous expression systems. Recently, however, we discovered a fourth NCX gene (NCX4) which is found exclusively in teleost, amphibian and reptilian genomes. The zebrafish (Danio rerio) NCX4a encodes for a protein of 939 amino acids and shows a high degree of identity with known  $\text{Na}^{+}$ - $\text{Ca}^{2+}$  exchangers. Although knock down of NCX4a activity in zebrafish embryos has been shown to alter left-right patterning, it has not been demonstrated that NCX4a functions as a  $\text{Na}^{+}$ / $\text{Ca}^{2+}$  exchanger. In this study, we: 1) demonstrate for the first time that this gene encodes for a novel NCX; 2) characterize the tissue distribution of zebrafish NCX4a and 3) evaluate its kinetic and transport properties. While ubiquitously expressed, the highest levels of NCX4a expression occur in the brain and eyes. NCX4a exhibits modest levels of  $\text{Na}^{+}$ -dependent inactivation and requires much higher levels of regulatory  $\text{Ca}^{2+}$  to activate outward exchange currents. NCX4a also exhibited extremely fast recovery from  $\text{Na}^{+}$ -dependent inactivation of outward currents, faster than any previously characterized wild-type exchanger. While this result suggests that the II inactive state of NCX4a is far less stable than in other NCX family members. We have demonstrated that a new putative member of the NCX gene family NCX4a encodes for a  $\text{Na}^{+}$ / $\text{Ca}^{2+}$  exchanger with unique properties including an extremely rapid recovery from  $\text{Na}^{+}$ -dependent inactivation. These data will be useful in understanding the role that NCX4a plays in embryological development as well as in the adult where it is expressed ubiquitously.

## Calcium Signaling Pathways

#### 606-Pos Board B485

##### Modeling $[\text{Na}^{+}]$ in PM-SR Nanodomains of Vascular Smooth Muscle Cells

Nicola Fameli<sup>1</sup>, Cornelis van Breemen<sup>1,2</sup>.

<sup>1</sup>University of British Columbia, Vancouver, BC, Canada, <sup>2</sup>Child and Family

Research Institute, Vancouver, BC, Canada.

We present a stochastic computational model aimed at elucidating the mechanism of site-specific signalling between a source and a target ionic transporter, both of which are localized on the plasma membrane (PM) and are part of a nanodomain: nanometer-scale subplasmalemmal signalling compartments comprising the PM, the sarcoplasmic reticulum (SR),  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  transporters (channels, exchangers and pumps), and the intervening cytosol. In this chain of events, the physical and functional link between non-selective cation channels (NSCC) and  $\text{Na}^{+}$ / $\text{Ca}^{2+}$  exchangers (NCX) needs to be elucidated in view of two interesting recent findings: the identification of the TRPC6 as the NSCC in VSM cells and the observation of localized  $\text{Na}^{+}$  transients following purinergic stimulation of these cells. Having previously helped clarify the signalling step between NCX and SERCA behind sarcoplasmic reticulum (SR)  $\text{Ca}^{2+}$  refilling, this quantitative approach now allows us to make inroads into this important signalling step. We have implemented a random walk (RW) Monte Carlo (MC) model with simulations mimicking a  $\text{Na}^{+}$  diffusion process originating at the NSCC within PM-SR junctions. Physical features of the system (junctional dimensions, diffusivity of  $\text{Na}^{+}$  in cytosol, channel capacity, etc.) were obtained in our laboratory and from the existing literature. The model calculates the average  $[\text{Na}^{+}]$  in the junction and also produces iso-concentration profiles for  $[\text{Na}^{+}]$  as a function of distance from the  $\text{Na}^{+}$  source. It also analyzes the influence of the junctional geometry on the signalling ability of the nanodomain. Our results emphasize the necessity of a strategic juxtaposition of the relevant signalling channels and organelles to form nanodomains that permit adequate  $[\text{Na}^{+}]$  build-up to provoke NCX reversal and  $\text{Ca}^{2+}$  influx via NCX eventually to refill the SR during asynchronous  $\text{Ca}^{2+}$  waves.

#### 607-Pos Board B486

##### A 3d Pseudo-stochastic Model Of Intercellular Calcium Signaling In Smooth Muscle

Dumitru A. Iacobas, Sylvia O. Suadicani, David C. Spray.

Albert Einstein College of Medicine, Bronx, NY, USA.

We present a general 3D pseudo-stochastic model of intercellular calcium signaling (ICS) in smooth muscle, composed of independently tunable units aggregated through a flexible convolution procedure. ICS ensures propagation and synchronicity of myocytes' contraction within muscular syncytia from