

IGF-1 effect, supporting the involvement of a serine-threonine phosphatase in IGF-1 activity (De Luca et al., *Br. J. Pharmacol.* 1998). Angiotensin receptor antagonists and other tools are in use to gain further insight in the mechanisms involved in inflammation-sensitive CIC-1 impairment in muscular dystrophy (Telethon-Italy GGP05130).

#### 2423-Pos Board B393

##### Evaluation of the Effects of Statin and Fibrate Treatment on Rat Skeletal Muscle: Biophysics, Genetic and Proteomic Studies

Giulia M. Camerino<sup>1</sup>, Sabata Pierno<sup>1</sup>, Claudio Digennaro<sup>1</sup>, Maria Antonietta Pellegrino<sup>2</sup>, Alfred L. George<sup>3</sup>, Roberto Bottinelli<sup>2</sup>, Diana Conte Camerino<sup>1</sup>.

<sup>1</sup>University of Bari, Bari, Italy, <sup>2</sup>University of Pavia, Pavia, Italy, <sup>3</sup>Vanderbilt University, Nashville, TN, USA.

Resting chloride conductance (gCl) sustained by the CIC-1 chloride channel have a crucial role in muscle physiology by maintaining the sarcolemma electrical stability. Its reduction can produce myotonia-like symptoms. CIC-1 channel has been shown to be a target of therapeutic molecules such as statins and fibrates. We previously demonstrated that lipophilic statins and fibrates affect skeletal muscle function by reducing resting chloride conductance (gCl) (Pierno et al, *Br J Pharmacol* 149:909, 2006). Here we studied the time course changes of fluvastatin (20 mg/kg) and fenofibrate (60 mg/kg) effects on Extensor Digitorum Longus (EDL) muscle gCl measured by two-microelectrode current clamp method. The gCl decreased in a time-dependent manner, being significantly lower after 1 week ( $2464 \pm 66 \mu\text{S}/\text{cm}^2$ ,  $n=46$  and  $2510 \pm 53 \mu\text{S}/\text{cm}^2$ ,  $n=50$  in fluvastatin and fenofibrate treated rats, with respect to  $2706 \pm 83 \mu\text{S}/\text{cm}^2$ ,  $n=27$  of control). To investigate the causes of gCl reduction we analyzed the CIC-1 gene expression by real-time quantitative PCR. The results showed a marked decrease in CIC-1 mRNA expression in both fluvastatin and fenofibrate chronically treated animals which contributes to gCl reduction. To study the involvement of other proteins essential for muscle function we analysed the proteomic map of EDL muscle from rat treated with fluvastatin (20mg/kg), atorvastatin (10mg/kg), fenofibrate (60mg/kg) and with combined fluvastatin (5mg/kg) plus fenofibrate (30mg/kg) by two-dimensional gel electrophoresis (Gelfi et al, *J Proteome Res* 5:1344, 2006). Fluorescent stained proteomic map showing ~500 spots were obtained and 40, 74, 60 and 76 differently expressed proteins were found in the above mentioned treated groups, with respect to control. The identification of each spot will allow to identify the protein targets of the myopathic process.

#### 2424-Pos Board B394

##### Anomalous Mole Fraction Effect in CIC-2 Chloride Channel Pore

Jorge E. Sanchez, Jose A. de Santiago-Castillo, Jorge Arreola. Physics Institute, Univ. Autonoma de San Luis Potosi, San Luis Potosi, Mexico.

The  $\text{Cl}^-$  pathway of *E. Coli*  $\text{ClC H}^+/\text{Cl}^-$  exchanger has three binding sites that can be occupied by  $\text{Cl}^-$ , however  $\text{SeCN}^-$  (an anion similar to  $\text{SCN}^-$ ) occupies only one site. This pathway serves as a model for pore structure of CIC channels whose gating is coupled to permeation. Thus CIC channels may gate differently in the presence of  $\text{SCN}^-$ . We analyzed the relationship between gating and pore properties by performing  $\text{SCN}^-$  mole-fraction experiments in CIC-2 channels expressed in HEK cells. Internal and external solutions with different  $\text{SCN}^-$  fractions were prepared mixing  $[\text{Cl}^-] + [\text{SCN}^-] = 140 \text{ mM}$ . Replacing 100%  $\text{Cl}^-$  with  $\text{SCN}^-$  on both sides of the membrane shifted to the left the voltage ( $V_m$ ) vs channel  $P_0$  curve by ~18 mV without changing the slope. In contrast, the protopore gate  $P_p$  vs  $V_m$  curve was shallower and shifted rightward (+5 mV). Extracellular  $\text{SCN}^-$  mole-fractions produced negative shifts on reversal potential ( $E_R$ ) values which are not described by GHK equation with a constant  $P_{\text{SCN}}/P_{\text{Cl}}$  ratio. The  $P_{\text{SCN}}/P_{\text{Cl}}$  ratios increased from 1.4 to 2.5 when the mole fraction increased, suggesting extracellular  $\text{SCN}^-$  enters the pore better than  $\text{Cl}^-$ . In addition, the slope of  $P_0$  vs  $V_m$  curve was steeper and the mid-point voltage ( $V_{0.5}$ ) did not change. Intracellular  $\text{SCN}^-$  mole fractions produced nonlinear negative shifts on  $E_R$  suggesting that from this side  $\text{Cl}^-$  enters the pore better than  $\text{SCN}^-$ . The slope of the  $P_0$  vs  $V_m$  curve was shallower and  $V_{0.5}$  and slope conductance vs mole fraction displayed the classical anomalous behaviour. Interestingly, the protopore gate located in the  $\text{Cl}^-$  pathway also displays the same behaviour. Our data show that the CIC-2 channel has indeed a multi-ion pore and that  $\text{SCN}^-$  enters this pore preferably from the extracellular side.

Supported by CONACyT.

#### 2425-Pos Board B395

##### Pharmacological Characterization of GaTx2, a Peptide Inhibitor of CIC-2 Chloride Channels

Christopher H. Thompson<sup>1</sup>, Cody S. Freeman<sup>1</sup>, Robert J. French<sup>2</sup>, Nael A. McCarty<sup>3</sup>.

<sup>1</sup>Georgia Institute of Technology, Atlanta, GA, USA, <sup>2</sup>University of Calgary, Calgary, AB, Canada, <sup>3</sup>Emory University School of Medicine, Atlanta, GA, USA.

CIC-2 chloride channels are voltage-gated ion channels that are expressed in neuronal and epithelial cells where they are critical mediators for the passive diffusion of  $\text{Cl}^-$  across the plasma membrane. Although CIC-2 is nearly ubiquitously expressed in mammalian cells, many details regarding channel biophysics and the physiological role that CIC-2 channels play remain undefined. We recently isolated the first peptide toxin active against CIC channels: Georgia anion toxin 2 (GaTx2), a 3.2 kDa peptide composed of 29 residues with three disulfide bonds. Here, we describe the basic pharmacological features of the inhibitory activity of GaTx2 against CIC-2, including affinity, mechanism of inhibition, and specificity. Using two-electrode voltage-clamp, we created a dose-response curve for inhibition of CIC-2 by GaTx2 at  $V_m = -100 \text{ mV}$ , and calculated a  $K_D$  of  $22 \pm 10 \text{ pM}$ . This value was very similar to the value obtained from dose-response curves created from multi-channel patches, which gave a  $K_D$  of  $12 \pm 5 \text{ pM}$ . Additionally, from TEVC recordings we measured  $k_{\text{on}} = 43 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ , and  $k_{\text{off}} = 0.0034 \text{ s}^{-1}$ , which is consistent with rate constants for other peptide inhibitors. Single channel recording showed that the latency to first opening is increased nearly 8 fold in the presence of 20 pM GaTx2. Also, outside-out macropatches revealed that GaTx2 is unable to inhibit open CIC-2 channels; thus, this toxin may act as a gating modifier. Finally, we found that GaTx2 is specific for CIC-2, being unable to inhibit other CIC channels or transporters, other major classes of  $\text{Cl}^-$  channels, or voltage-dependent  $\text{K}^+$  channels. This high affinity, highly specific inhibitor of CIC-2 will provide an excellent tool for studies designed to understand the function and regulation of this channel, and will help define its physiological role(s).

#### 2426-Pos Board B396

##### Slow Gating in CIC Chloride Channels: Normal Mode Analysis

Gennady V. Miloshevsky<sup>1</sup>, Ahmed Hassanein<sup>1</sup>, Peter C. Jordan<sup>2</sup>.

<sup>1</sup>Purdue University, West Lafayette, IN, USA, <sup>2</sup>Brandeis University, Waltham, MA, USA.

All-atom NMA is used to explore possible mechanisms for slow gating in CIC  $\text{Cl}^-$  channels. As the "double-barreled" architecture is well established throughout the CIC family, both channels and transporters (Miller, 2006. *Nature*. 440:484), we use the high-resolution (2.5 Å) X-ray structure of an *E. coli* CIC transporter (pdb entry 1OTS) as a template, describe it with the CHARMM22 force field and carry out standard all-mode NMA. The slowest, intrinsic motions encoded in the structure are determined by protein shape. Perturbing the system in either direction along the 7<sup>th</sup> all-atom NM leads to slow relative swinging of the subunits, perpendicular to the membrane plane. The in-plane swivel axis lies at the subunit interface, near the protein's center. The intracellular interfacial domain is the region most affected. Here the two halves of the protein oscillate, separating and then nearly touching. The R and A helices execute large scale swaying, alternately increasing and decreasing their cytoplasmic ends' separations, motion in agreement with FRET experiments (Bykova et al., 2006. *Nat. Struct. Biol.* 13:1115). The ion-occupied intracellular pores behave as almost rigid units. As the subunits separate, the intracellular pore tilt relative to the membrane plane changes notably. In contrast, the extracellular portion of the subunit interface is significantly less affected, although small interfacial structural changes are clearly observable. Those extracellular regions structurally affected by the subunits' slow sway are localized near the extracellular  $\text{Cl}^-$  pathways. As the subunits separate, these regions compress, possibly shutting the extracellular pores. As they approach, the extracellular regions near the  $\text{Cl}^-$  conduction pathways relax, possibly opening them.

#### 2427-Pos Board B397

##### Ab Initio Calculations Of Structural Rearrangements and Energetic of Glutamate<sup>148</sup> Site Chain of the Ec-CIC H<sup>+</sup>/Cl<sup>-</sup> Exchanger

Pablo G. Nieto, Ricardo Guirado, Jorge Arreola.

Instituto de Fisica UASLP Mexico, San Luis Potosi, Mexico.

$\text{ClC H}^+/\text{Cl}^-$  exchangers are homodimers with three  $\text{Cl}^-$ -binding sites within each monomer. Previous studies suggest that transport cycles are triggered by protonation of the lateral chain of a glutamate residue (E148) located in the middle of the  $\text{Cl}^-$  pathway. In this work, we performed *ab initio* density functional theory calculations using the ultra soft pseudopotential approach in order to gain insight about microscopic movements induced by H and/or Cl during transport activity. For these computations the 16 amino-acids that line the  $\text{Cl}^-$  pathway were spatially aligned according to the X-ray structure. We found that the orientation of the unprotonated E148 carboxy group was influenced when Cl was in close proximity. Attaching a single H to this  $\text{COO}^-$  group displaced the lateral chain (with respect to the unprotonated structure) towards the extracellular side and led to the formation of a sizable hole in the entryway. When Cl was placed at 2 Å from the protonated  $\text{COO}^-$  group ( $\text{COOH}$ ) (in both intra- and extra-cellular positions) the conformational