

obtained by knocking down PKC $\alpha$  with small interfering RNA duplexes. Taken together, these data demonstrate that PtdIns(4,5)P<sub>2</sub> itself targets PKC $\alpha$  to the plasma membrane through the polybasic cluster located in the C2 domain, with this interaction being critical in the signaling network involved in neural differentiation.

#### 489-Pos Board B368

**Probing Phosphoinositide Kinetics With A Voltage-sensitive Phosphatase**  
**Bjoern H. Falkenburger**, Jill B. Jensen, Byung-Chang Suh, Bertil Hille.  
 University of Washington, Seattle, WA, USA.

Voltage-sensitive phosphatases (VSPs) have a voltage sensor linked to a phosphoinositide (PI) 5-phosphatase, which hydrolyzes plasma membrane PI-(4,5)-bisphosphate (PIP<sub>2</sub>) to PI(4)P [Iwasaki, PNAS 105, 7970]. We used PIP<sub>2</sub> hydrolysis by VSP from *Ciona intestinalis* (ci-VSP) and zebrafish (dr-VSP) to better understand PIP<sub>2</sub> binding and resynthesis. PIP<sub>2</sub> was monitored using the PIP<sub>2</sub>-sensitive M-current (KCNQ2/3) and FRET between a pair of PIP<sub>2</sub>-binding probes (PH-PLC $\delta$ 1-CFP & PH-PLC $\delta$ 1-YFP). Depolarizations to +100 mV lasting >50 ms reduced M-current and PH-probe FRET. PIP<sub>2</sub> depletion was saturated by depolarizations lasting 500-1000 ms. Evidently PH-probe FRET and M-current respond quickly to changes in plasma membrane PIP<sub>2</sub>.

After repolarization, PH-probe FRET and M-current relaxed to baseline values with time constants of ~10 s in a wortmannin-insensitive manner. This reflects endogenous PIP 5-kinase converting PI(4)P back to PI(4,5)P<sub>2</sub>. Overexpression of PIP 5-kinase increased the length of depolarization required to deplete PIP<sub>2</sub>, and speeded PIP<sub>2</sub> recovery after repolarization.

Recovery of PIP<sub>2</sub> after VSP activation is ~10x faster than after PLC activation. However, it only requires PIP 5-kinase, whereas recovery after PLC activation requires PI 4-kinase and PIP 5-kinase in series. Thus PI 4-kinase must be the slower enzyme. To estimate the rate of PI 4-kinase, we compared translocation of a fluorescent probe that reports plasma membrane PI(4)P (PH-OSH2, T.Balla) to translocation of the PIP<sub>2</sub>-binding probe in confocal time-lapse imaging. Upon PLC activation through M1 receptors, the plasma membrane PI(4)P signal decreased 20 s later than the PIP<sub>2</sub> signal. Both probes recovered with similar time courses. FRET photometry between either PIP<sub>2</sub>-probes or PI(4)P-probes showed comparable results. The simultaneous recovery of both probes is consistent with the hypothesis that recovery of PIP<sub>2</sub> is governed by rate-limiting synthesis of PI(4)P by the PI 4-kinase, followed by rapid conversion of PI(4)P into PI(4,5)P<sub>2</sub> by the PIP 5-kinase.

NIH-NS008174&HFSP

#### 490-Pos Board B369

**Interaction Of PTEN<sub>1-21</sub> Peptide With Phosphatidylinositol-4,5-Bisphosphate: A <sup>31</sup>P NMR Relaxation Study.**

**Edgar E. Kooijman<sup>1</sup>**, Avigdor Leftin<sup>2</sup>, Michael F. Brown<sup>2</sup>, Arne Gericke<sup>1</sup>.  
<sup>1</sup>Kent State University, Kent, OH, USA, <sup>2</sup>University of Arizona, Tucson, AZ, USA.

Phosphatidylinositol-4,5-bisphosphate (PI(4,5)P<sub>2</sub>) is by far the most abundant of all phosphoinositides (about 1% of all membrane phospholipids) and is found primarily at the cytoplasmic leaflet of the plasma membrane. PI(4,5)P<sub>2</sub> can be converted by class I PI 3-kinases to PI(3,4,5)P<sub>3</sub>, which is a second messenger molecule affecting processes like cell survival and proliferation. PTEN is an important tumor suppressor protein that converts PI(3,4,5)P<sub>3</sub> back to PI(4,5)P<sub>2</sub>, thereby keeping the basal levels of PI(3,4,5)P<sub>3</sub> low. We have demonstrated recently that PTEN is allosterically activated by PI(4,5)P<sub>2</sub>, which interacts with the N-terminal end of PTEN. Here we study the interaction of the PI(4,5)P<sub>2</sub> binding domain of PTEN (PTEN1-21) with model membranes containing PI(4,5)P<sub>2</sub> using <sup>31</sup>P-NMR. Using both magic angle spinning (MAS) and static solid state NMR we probe the interaction of PTEN1-21 with lipid phosphates. We determined the intrinsic T<sub>1</sub> and T<sub>2</sub> relaxation times of the phosphomonoester groups of PI(4,5)P<sub>2</sub> and the phosphodiester group of dioleoyl-phosphatidylcholine, which form the lipid matrix into which PI(4,5)P<sub>2</sub> was dispersed. PTEN1-21 is highly basic, containing several Lys and Arg residues which are thought to give rise to a largely electrostatic PTEN/PI(4,5)P<sub>2</sub> interaction. We show here that the binding of PTEN1-21 to PI(4,5)P<sub>2</sub> bilayers dramatically affects the membrane structure, indicating that the PTEN1-21/PI(4,5)P<sub>2</sub> interaction is likely more than a simple electrostatic interaction. This is in accordance with our recent findings that PTEN/phosphoinositide interaction is specific for PI(4,5)P<sub>2</sub>. Lys13 is crucial for this specific interaction and this study explores the interaction of the Lys13 of PTEN with the phosphomonoester groups of PI(4,5)P<sub>2</sub>.

#### 491-Pos Board B370

**Plasma Membrane Order In T Cell Signalling**

**Jelena Dinic**, Jeremy Adler, Ingela Parmryd.

Wenner-Gren Institute, Stockholm University, Stockholm, Sweden.

Plasma membrane nanodomains, referred to as lipid rafts, more ordered than the bulk membrane play an important role in T cell signalling by forming signalling platforms in activated T cells. However, the existence of lipid rafts in resting T cells is contentious. Using laurdan, a membrane probe whose peak emission wavelength depends on the lipid environment, evidence is presented for the existence of ordered nanodomains in resting T cells.

T cell signalling can be initiated by stimulating the T cell receptor (TCR), crosslinking the lipid raft markers GM1 (sphingolipid) or glycosylphosphatidylinositol (GPI) anchored proteins. The aggregation of lipid raft components induces the same response in Jurkat T cells as the ligation of an antigen to the TCR. Changes in membrane order linked with reorganization of the plasma membrane upon Jurkat T cell activation were followed at 37°C. Fluorescent images were analyzed for generalised polarisation values - a measure of the relative abundance of liquid ordered and liquid disordered domains. TCR patching does not increase the overall membrane order suggesting that membrane domains of high order are brought together in the patches. This supports the existence of small ordered membrane domains in resting T cells that aggregate upon activation. Patching of GM1, the GPI-anchored protein CD59 and the non lipid raft marker CD45 significantly increases the overall membrane order. So does general crosslinking of membrane components with Concanavalin A. Remodelling of the actin cytoskeleton is an integral part of TCR signaling and T cell activation. Disrupting actin polymerization using latrunculin B decreases membrane order and stabilizing actin filaments with jasplakinolide increases membrane order. An increase in membrane order appears to be a general effect of plasma membrane component patching and is likely due to a global induction of actin polymerization at the plasma membrane.

#### 492-Pos Board B371

**Adsorption Of Bar-domain Proteins To Charged Lipid Membranes Causes Deformations And Lipid Demixing**

**George Khelashvili<sup>1</sup>**, Daniel Harries<sup>2</sup>, Harel Weinstein<sup>1</sup>.

<sup>1</sup>Weill Medical College, New York, NY, USA, <sup>2</sup>The Hebrew University of Jerusalem, Jerusalem, Israel.

Many proteins participating in cellular processes contain BAR domains that have been implicated in membrane shaping and deformation. These BAR domains can either induce significant membrane curvatures or sense high-curvature regions on cell membranes, but the mechanism for this action is still not well understood. One suggestion is that BAR domains work collectively and achieve significant bilayer deformations only through a suitable organization at membrane interfaces. In contrast, evidence from some atomistic simulations suggests that a single BAR can substantially deform a lipid membrane locally. Here we present results from a self-consistent mean-field model of BAR domain association on membranes, suggesting that a single Amphiphysin BAR is capable of producing a steady state, where the initially near-planar membrane curves significantly. However, using our approach we predict that such deformation will occur only for membrane patches that have the propensity to attain high spontaneous curvature, and that such favorable preconditioning may be the result of either local lipid demixing, or of a preceding insertion of the BAR domain's amphiphatic N-helix. Both events have been predicted to bring about asymmetry in the two membrane monolayers. To contrast, our simulations also show that local segregation of charged lipids under the influence of the adsorbing BAR domain alone cannot produce high enough asymmetry between bilayer leaflets, and that in the absence of additional energetic sources that favor membrane asymmetry, the membrane will remain near-flat within fluctuations upon BAR adsorption. Thus, we conclude that N-helix insertions may have a critical mechanistic role in the function of the BAR domain, and that the electrostatic interactions between BAR and membrane are essential for sensing and stabilization of bilayer curvature.

#### 493-Pos Board B372

**Inducing and Reversing Anesthesia with Temperature Variation - Experiments on an Excised Frog Sciatic Nerve**

**Bineyam Kassahun**, Martin Bier, Alexander Murashov.

East Carolina University, Greenville, NC, USA.

The Meyer-Overton Rule and other more recent experimental observations suggest that the fluidity of the lipid membrane is involved in nerve propagation and in mechanisms behind anesthesia. In other words, Hodgkin-Huxley may not be