

- in the millisecond time region. *Proc. Natl. Acad. Sci. USA* 78:3318–3322.
4. Aoshima, H., D. J. Cash, and G. P. Hess. 1981. Mechanism of inactivation (desensitization) of acetylcholine receptor. Investigations by fast reaction techniques with membrane vesicles. *Biochemistry*. 20:3467–3474.
 5. Hess, G. P., D. J. Cash, and H. Aoshima. 1983. Acetylcholine receptor-controlled ion translocation: chemical kinetic investigations of the mechanism. *Annu. Rev. Biophys. Bioeng.* 12:443–473.
 6. Walker, J. W., M. G. McNamee, E. Pasquale, D. J. Cash, and G. P. Hess. 1981. Acetylcholine receptor inactivation in *Torpedo californica* electroplax membrane vesicles. Detection of two processes in the millisecond and record time regions. *Biochem. Biophys. Res. Comm.* 100:86–90.
 7. Hess, G. P., E. B. Pasquale, J. W. Walker, and M. G. McNamee. 1982. Comparison of acetylcholine receptor-controlled cation flux in membrane vesicles from *Torpedo californica* and *Electrophorus electricus*: chemical kinetic measurements in the millisecond region. *Proc. Natl. Acad. Sci. USA.* 79:963–967.
 8. Hess, G. P., D. J. Cash, and H. Aoshima. 1980. Kinetic mechanism of acetylcholine receptor-controlled ion flux: flow quench kinetic measurements of acetylcholine-induced flux in membrane vesicles. *Neurochemistry International*. 2:233–242.
 9. Hess, G. P., H. Aoshima, D. J. Cash, and B. Lenchitz. 1981. Specific reaction rate of acetylcholine receptor-controlled ion translocation: a comparison of measurements with membrane vesicles and with muscle cells. *Proc. Natl. Acad. Sci. USA.* 78:1361–1365.
 10. Neher, E., and B. Sakmann. 1976. Single-channel currents recorded from membranes of denervated frog muscle fibers. *Nature (Lond.)*. 260:779–802.

ACETYLCHOLINE-ACTIVATED CHANNEL CURRENT-VOLTAGE RELATIONS IN SYMMETRICAL Na⁺ SOLUTIONS

JOHN A. DANI AND GEORGE EISENMAN

Department of Physiology, University of California Los Angeles Medical School, Los Angeles, California 90024

A primary function of the acetylcholine receptor-channel complex (AChR) is transporting ions. The channel is impermeable to anions but passes divalent cations and a wide variety of monovalent cations with diameters less than ~7–8 Å (1, 2). A simple two-barrier, one-site (2B1S) model has been used to describe the transport of the simplest and biologically most important monovalent cations (3–6). In each case certain restrictions were applied to the model, such as arbitrarily setting the position or affinity of the site. In addition, because of experimental limitations the model was applied to data obtained at one symmetrical concentration or by changing only the external concentration.

Certainly permeation through ion channels is very complex. A simple barrier model may be viewed as a crude approximation of reality most useful as a phenomenological description of transport. The model can serve a functional role in the analysis of new data; any extensions or revisions necessary to describe transport of other permeants can give clues regarding the different interactions between the ions and channel. The model is also useful when considering the effect of electrolyte composition on other properties of the channel such as gating (4). Exciting hypotheses have recently been proposed relating the primary amino acid sequences to the structure of the AChR subunits, and relating subunit organization to the formation of the ion channel (7, 8). A simple barrier model can provide the first step toward integrating the details of permeation to the overall structure and function.

Here we test if a 2B1S model, applied without restricting the location or energies of the barriers and site, can

describe current-voltage (*I-V*) relations obtained over a wide concentration range of symmetrical Na⁺ solutions. The shortcomings of such an absolute rate theory model, like those that arise when the energy barriers are small (9), are not addressed. Although we shall conclude that a general 2B1S model without additional provisions is not completely adequate to describe the data, it does provide a surprisingly good first approximation and gives information about the energy profile that may be useful in further investigation of AChR permeation, structure, and function.

RESULTS

Single AChR events from rat myotubes were studied by voltage clamping membrane patches in symmetrical Na⁺ concentrations (10). To obtain *I-V* relations, voltage ramps were applied to the membrane (11). The legend of Fig. 1 describes the solutions, methods, and procedure for obtaining *I-V* relations from single-channel events. Fig. 1 C shows a leak-corrected single-channel response to a voltage ramp. Fig. 2 is a plot of the zero-voltage conductance (G_0) vs. activity. A Michaelis-Menten curve having a dissociation constant of 74 mM (activity) is drawn through the high concentration data. The insert is an Eadie-Hofstee plot (G_0 vs. $G_0/\text{activity}$) that shows the deviation of the low concentration data from an expected straight line. Fig. 3 shows the *I-V* relations obtained in symmetrical Na⁺ solutions of 45 mM, 150 mM, and 450 mM. The smooth curves result from the best theoretical fit of the 2B1S model. Although the fit to the *I-V* relations is quite good,




FIGURE 1 Standard patch-clamp techniques were used. Our rate of obtaining gigaohm seals increased following James Rae's suggestion to use Corning's Kovar 7052 pipettes (Corning Glass Works, Corning, NY). The Pt wire used in fire polishing should be frequently coated with Kovar glass. We believe the glass vaporizes from the hot wire, depositing a thin film on the patch pipette. The sealing rate of electrically superior pipettes can be improved by this procedure (19). The pipette was usually filled with 45, 150, or 450 mM NaCl, osmolarity unadjusted; 5 mM HEPES, pH 7.4; 0.1 to 0.6 μ M ACh and 0.5 mM BaCl₂ was added to the 45-mM and occasionally the 150-mM solution. After a patch was obtained it was placed deeply into an inflow port containing an equivalent Cl⁻ or F⁻ solution. *A*, raw record of a single channel response to a 51 ms, -170 to +175 mV ramp voltage clamp, filtered at 600 Hz with 8-pole Bessel, 21°C, in 150 mM Na⁺. *B*, average of two selected nearby blank records that accurately overlap the baseline of *A*, leak correction was made by subtracting *B* from *A*. Rarely, a second order polynomial was used to correct short events if blank records were unavailable or did not fit the event record baseline. *C*, leak-corrected event record. Many digitized current points were averaged by computer at their corresponding voltage point to obtain the *I-V* (11).

the model does not accurately predict G_0 . The dashed line in Fig. 2 shows the 2B1S model poorly predicts the G_0 data, especially at low concentrations.

The dissociation constant is in reasonable agreement with values obtained at 100 mV from myotubes (12), and from *Torpedo* AChR reconstituted into liposomes (13). These values are about four times smaller than those estimated in frog or toad endplates (14, 15).

The deviation of the low concentration data from a single site binding curve in Fig. 2 suggests possible shortcomings of the 2B1S model. Such a deviation would result if the channel had more than one nonequivalent site or if negative dipoles or charges increased the concentration of Na⁺ near the channel entrance. Lewis and Stevens (5) extended the 2B1S model to include an external surface




FIGURE 2 Zero-voltage conductance vs. activity fitted by $G_0 = G_{max} [Na]/([Na] + K_D)$ with $G_{max} = 68$ pS and $K_D = 74$ mM, where $[Na]$ is activity. *Insert*, an Eadie-Hofstee plot of G_0 vs. $G_0/[Na]$. The solid line is drawn through the high concentration data using the same G_{max} and K_D as above. The dashed line is the prediction of the 2B1S model with $G_{max} = 68$ pS and $K_D = 90$ mM. The x is taken from reference 12.

potential to fit their data. The surface potentials they used are larger than suggested by our results. Given that the channel extends 55 Å from the outer surface of the membrane and that the channel entrance is ~25 Å in diameter (16, 17), it seems likely that the effective field of




FIGURE 3 Shows the *I-V* relations at 45, 150, and 450 mM Na⁺. The vertical and horizontal lines mark $V = 0$ and $I = 0$ positions. The smooth curves are the result of a nonlinear least-squares Levenberg-Marquardt fit (algorithm developed by C. Clausen) to all the data using the 2B1S model. The net flux of ions is given by $J = [Na](k_{-1}k_{-2} - k_1k_2)/[Na](k_1 + k_{-2}) + k_{-1} + k_2$. The upper insert shows the free energy profile of the best fit labeled with the rate constant used in Eq. 1. The marks on the *Y*-axis represent + or - 10 kT. Unlike previous findings, this fit predicts the outer barrier to be larger than the inner.

dipoles and charges within this large vestibule will vary with ionic strength and will influence ion loading. The introduction of "surface potentials" into the barrier model using the Grahame equation (18) is an approximation that does not deal with these intricacies, but extends the model to handle the overall effect. Additional data may allow us to determine whether "surface potentials" can adequately diminish the shortcomings of a 2B1S model, whether a more complex barrier model is needed, or whether other types of models are more appropriate.

We thank Dr. R. Horn for helpful discussions.

This work was supported by grant PCM 76-20605 from the National Science Foundation and grant GM 24749 from the United States Public Health Service.

Received for publication 6 May 1983.

REFERENCES

- Adams, D. J., T. M. Dwyer, and B. Hille. 1980. The permeability of endplate channel to monovalent and divalent metal cations. *J. Gen. Physiol.* 75:493-510.
- Dwyer, T. M., D. Adams, and B. Hille. 1980. The permeability of the endplate-channel to organic cations in frog muscle. *J. Gen. Physiol.* 75:469-492.
- Adams, P. R. 1979. A completely symmetrical barrier model for endplate channels. *Biophys. J.* 25(2,Pt.2):70 a. (Abstr.)
- Marchais, D., and A. Marty. 1979. Interaction of permeant ions with channels activated by acetylcholine in *Aplysia* neurones. *J. Physiol. (Lond.)* 297:9-45.
- Lewis, C. A., and C. F. Stevens. 1979. Mechanism of ion permeation through channels in a postsynaptic membrane. In *Membrane Transport Processes*. C. F. Stevens and R. W. Tsien, editors. Raven Press, New York. 3:133-151.
- Horn, R., and M. S. Brodwick. 1980. Acetylcholine-induced current in perfused rat myoballs. *J. Gen. Physiol.* 75:297-321.
- Deville-Thierry, A., J. Giraudat, M. Bentaboulet, and J.-P. Changeux. 1983. Complete mRNA coding sequence of the acetylcholine binding α -subunit of *Torpedo marmorata* acetylcholine receptor: a model for the transmembrane organization of the polypeptide chain. *Proc. Natl. Acad. Sci. USA.* 80:2067-2071.
- Noda, M., H. Takahashi, T. Tanabe, M. Toyosato, S. Kikuyotani, Y. Furutani, T. Hirose, H. Takashima, S. Inayama, T. Miyata, and S. Numa. 1983. Structural homology of *Torpedo californica* acetylcholine receptor subunits. *Nature (Lond.)* 302:528-531.
- Montroll, E. W., and K. E. Shuler. 1959. The application of the theory of the stochastic processes to chemical kinetics. *Adv. Chem. Phys.* 1:361-399.
- Hamill, O. P., A. Marty, E. Neher, B. Sakmann, and F. J. Sigworth. 1981. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch. Eur. J. Physiol.* 391:85-100.
- Yellen, G. 1982. Single Ca^{2+} -activated nonselective cation channels in neuroblastoma. *Nature (Lond.)* 296:357-359.
- Horn, R., and J. Patlak. 1980. Single channel currents from excised patches of muscle membrane. *Proc. Natl. Acad. Sci. USA.* 77:6930-6934.
- Tank, D. W., R. L. Huganir, P. Greengard, and W. W. Webb. 1983. Patch-recording of purified and reconstituted *Torpedo* acetylcholine receptors. *Biophys. J.* 41(2,Pt.2):135a. (Abstr.)
- Adams, D. J., W. Nonner, T. M. Dwyer, and B. Hille. 1981. Block of endplate channels by permeant cations in frog skeletal muscles. *J. Gen. Physiol.* 78:593-615.
- Takeda, K., P. H. Barry, and P. W. Gage. 1982. Effects of extracellular sodium concentration on null potential, conductance and open time of endplate channels. *Proc. R. Soc. Lond. B Biol. Sci.* 216:225-251.
- Kistler, J., R. Stroud, M. Klymkowsky, R. Lalancette, and R. Fairclough. 1982. Structure and function of an acetylcholine receptor. *Biophys. J.* 37:371-383.
- Klymkowsky, M., and R. Stroud. 1979. Immunospecific identification and three-dimensional structure of a membrane-bound acetylcholine receptor from *Torpedo californica*. *J. Mol. Biol.* 128:319-334.
- Grahame, D. 1947. The electric double layer and the theory of electrocapillarity. *Chem. Rev.* 41:441-501.
- Horn, R., C. Vandenberg, and K. Lange. 1984. Statistical analysis of single sodium channels: effects of *N*-bromoacetamide. *Biophys. J.* In press.