

3. Tzaphlidou, M., J. A. Chapman, and M. H. Al-Samman. 1982. A study of positive staining for electron microscopy using collagen as a model system. II. Staining by uranyl ions. *Micron*. 13:133-145.
4. Steven, A. C., and M. A. Navia. 1982. Specificity of stain distribution in electron micrographs of protein molecules contrasted with uranyl acetate. *J. Microscopy*. 128:145-155.
5. Mannella, C. A., and J. Frank. 1982. Effects of divalent metal ions and chelators on the structure of outer mitochondrial membranes from *Neurospora crassa*. *Biophys. J.* 37:3-4.
6. Mannella, C. A., B. Cognon, and M. Colombini. 1982. Immunological evidence for identity between the subunits of the crystalline arrays in *Neurospora* outer mitochondrial membranes and the pore-forming components of these membranes. *J. Cell Biol.* 95:256 a.
7. Frank, J., and C. A. Mannella. 1983. Multivariate classification of diffraction patterns: effects of different stains on images of mitochondrial pore arrays. Proc. 41st Mtg. Electron Microsc. Soc. America. In press.
8. Frank, J., W. Goldfarb, D. Eisenberg, and T. S. Baker. 1978. Reconstruction of glutamine synthetase using computer averaging. *Ultramicroscopy*. 3:283-290.
9. Mannella, C. A., and J. Frank. 1982. Negative-staining characteristics of pore-like subunits of the mitochondrial outer membrane: implications for surface-charge distribution. Proc. 10th Intl. Congress on Electron Microsc. 3:41-43.
10. Hallermayer, G., and W. Neupert. 1974. Lipid composition of mitochondrial outer and inner membranes of *Neurospora crassa*. *Hoppe-Seyler's Z. Physiol. Chem.* 355:279-288.
11. Seimiya, T., and S. Ohki. 1973. Ionic structure of phospholipid membranes, and binding of calcium ions. *Biochim. Biophys. Acta*. 298:546-561.
12. Habeeb, A. F. S. A., H. G. Cassidy, and S. J. Singer. 1958. Molecular structural effects produced in proteins by reaction with succinic anhydride. *Biochim. Biophys. Acta*. 29:587-593.
13. Döring, C., and M. Colombini. 1984. On the nature of the molecular mechanism underlying the voltage dependence of the channel-forming protein, VDAC. *Biophys. J.* 45:44-46.

FLUORESCENCE PROBES FOR THE STUDY OF ACETYLCHOLINE RECEPTOR FUNCTION

M. MARTINEZ-CARRION, J. M. GONZALEZ-ROS, J. R. MATTINGLY, J. A. FERRAGUT,
M. C. FARACH, AND D. DONNELLY
*Department of Biochemistry, Medical College of Virginia, Virginia Commonwealth University,
Richmond, Virginia 23298*

INTRODUCTION

Our group initiated the use of pyrene derivatives for functional and structural studies of the membrane-bound Acetylcholine Receptor (AChR) system. Our studies have taken two approaches. We have recently used a hydrophobic derivative (pyrene-1-sulfonyl azide) that is able to bind covalently to regions of the AChR molecule accessible from the membrane lipid (1) to monitor ligand-induced effects at those hydrophobic regions of the AChR protein exposed to the membrane matrix (2). The structural effects induced by ligand binding are interpreted in relation to the AChR "desensitization" process. We have also used a hydrophilic pyrene derivative (1,3,6,8-pyrene tetra sulfonate [PTSA]) to develop a fast spectroscopic method for measuring AChR functionality, i.e., the formation of a short-lived cation channel, within the physiological time scale. The procedure is based on the fluorescence quenching of intravesicularly entrapped PTSA by externally added Tl^+ which substitutes for the physiologically occurring Na^+ . Addition of cholinergic agonist to preparations of AChR membranes containing entrapped PTSA promotes the formation of the AChR cation chan-

nel, which allows the added Tl^+ into the internal volume of the vesicles where collisional quenching between the cation and the entrapped fluorophore occurs. This Tl^+ influx is spectroscopically monitored as a time-dependent decrease of the fluorescence emitted by PTSA in the millisecond time scale.

In this communication, we use the latter, "stopped-flow/fluorescence quenching," procedure to examine several functional aspects of the membrane bound AChR, including the effect of alkaline extraction of peripheral proteins from the AChR membranes, and the effect of binding of specific anti-AChR antibodies, which serves as a molecular model for the autoimmune disease *Myasthenia gravis* (4, 5).

MATERIALS AND METHODS

AChR-enriched membranes were purified for *Torpedo californica* electric tissue. The electroplax was minced and homogenized 1:2 (wt/vol) in ice-cold buffer (10 mM Tris, pH 7.4, 5 mM EDTA, 5 mM iodoacetamide, 0.5 mM PMSF and 0.02% $NaNO_3$). The homogenate was centrifuged at 3,500 rpm for 10 min in a GS-3 Sorvall rotor (DuPont Instruments-Sorvall, Newton, CT). The supernatant was filtered through cheesecloth and centrifuged at 30,000 rpm for 30 min in a Beckman 35 rotor (Beckman Instruments, Inc., Fullerton, CA). The resulting pellet, referred to as crude AChR-membranes, was resuspended in HEPES buffer (10 mM HEPES, pH 7.4, 100 mM $NaNO_3$) at ~10 mg of protein/ml. This crude AChR had specific activity values of 10 μ g of α -Bungarotoxin (α -Bgt) bound/mg of protein. Routinely, three-fourths of that membrane preparation was submitted to alkaline extraction of the peripheral membrane proteins (5).

Abbreviations used in this paper: AChR, acetylcholine receptor; PTSA, 1,3,6,8-pyrene tetrasulfonate; EDTA, ethylenediaminetetraacetic acid; PMSF, phenylmethanesulfonyl fluoride; α -Bgt, α -bungarotoxin; DPH, 1,6-diphenyl-1,3,5-hexatriene; TMA-DPH, 1-[4-(trimethylammonium)phenyl]-6-phenyl-1,3,5-hexatriene.

Protein concentration and α -Bgt binding of the membrane fractions were determined (1). The extraction, fractionation and characterization of lipids from the membranes were performed as previously described (6). Lipid vesicles from the different lipid extracts were formed through a detergent dialysis method (6). Steady-state fluorescence measurements were obtained as described (2, 6) using a SLM 4800-series subnanosecond spectrofluorometer.

Fab fragments from polyclonal anti-AcChR IgG from immunized goats suffering from experimental autoimmune *Myasthenia gravis* were prepared following published procedures (3). For experiments involving the study of the anti-AcChR Fab on AcChR function, the AcChR membranes were incubated overnight at 4°C with a 30-fold molar excess of Fab over the α -Bgt binding sites. Unbound Fab fragments were removed by centrifugation of the membranes prior to resuspension in HEPES buffer for submission to the fluorophore-loading procedure described below.

Loading of the fluorophore PTSA (final concentration 4.3 mM) into AcChR membranes or lipid vesicles was accomplished through a rapid freezing/slow thawing technique (7). The external fluorophore was removed from the fluorophore vesicles by chromatography on a Sepharose 6 MB column.

Fast kinetic measurements of Ti^+ "influx" through the membranes were obtained using a Durrum D-110 stopped-flow spectrophotometer set up in the fluorescence mode. Excitation wavelength was 378 nm and emission of fluorescence was monitored using a Corning 3-75 filter (Corning Glass Works, Corning, NY). The flux process was initiated by rapid mixing of the fluorophore-loaded vesicles (dead time was ~3-4 ms) with an equal volume of a 10 mM HEPES, pH 7.4, 30 mM TiNO_3 , 70 mM NaNO_3 buffer containing various concentrations of cholinergic agonists. The spectroscopic data was stored in a Physical Data model 514A transient recorder (Physical Data, Inc., Beaverton, OR) and visualized in a Tektronix oscilloscope (Tektronix, Inc., Beaverton, OR). Permanent storage and data analysis were performed by an interfaced Apple micro-computer. The spectroscopic traces were computer-fitted to the Stern-Volmer equation describing the collisional quenching process (8).

RESULTS AND DISCUSSION

Since the development of the alkaline extraction procedure (9) to remove peripheral proteins associated with electroplax AcChR membranes, most researchers have utilized the alkali treatment to increase the AcChR-specific activities of their membrane preparations. As originally described, the alkaline treatment leads to AcChR-rich membranes having normal binding capabilities for cholinergic agonists and local anesthetics and having at least a qualitative retention of the cation permeability control by the AcChR molecule (9). However, further investigations revealed that the aggregation state and the distribution of the AcChR protein within the membrane were changed, probably as a consequence of the extraction of the 43,000 M_r polypeptide associated with the AcChR membranes. Also a greater rotational mobility of the AcChR protein following alkaline extraction has been reported.

Despite the data available on the effect of alkaline extraction on the AcChR protein, very little has been learned about the effect of this treatment on the properties of the membrane matrix. Thus we have studied the effect of the alkaline treatment on (i) the membrane lipid composition; (ii) the apparent "fluidity" of the native membranes and of lipid vesicles formed from plain lipid

extracts; (iii) the permeability of those lipid vesicles and membranes for monovalent cations.

The lipid compositional features of AcChR-membranes before and after alkaline treatment (Table I) reveals that although an almost twofold increase in the lipid/protein ratio is produced by extracting peripheral proteins, the molar ratios of phospholipid/cholesterol remain unchanged. However, the proportions of the major phospholipid components are somewhat altered, revealing what could be a "selective loss" of phosphatidyl ethanolamine during alkaline extraction. In consequence, phosphatidylcholine becomes the major phospholipid component of the alkaline treated membranes. No significant changes could be detected in the degree of unsaturation of the acyl components of either individual phospholipids or the membranes as a whole.

The selective extraction of peripheral proteins as well as the slight loss of phosphatidyl ethanolamine subsequent to alkaline extraction does not promote significant changes in the apparent "fluidity" of the membranes. This is measured by the temperature-dependent fluorescence depolarization of two fluorophores serving as probes for different membrane domains, diphenylhexatriene (DPH) and its trimethyl ammonium derivative (TMA-DPH). Fluorescence polarization values appear to indicate only a slightly less fluid environment for the alkaline membranes at the DPH domain. Even those small differences are absent when plain lipid vesicles formed from lipid extracts of crude and alkaline-treated membranes are compared, suggesting total similarity in "fluidity" of both lipid media.

Despite these similarities, the extraction of peripheral proteins appears to perturb the passive cation permeability of the membranes. We measured rate-constant values corresponding to the pseudo-first-order fluorescence decay process due to slow Ti^+ leakage through the membranes upon rapid mixing in the stopped-flow spectrophotometer. A faster leakage of Ti^+ into alkaline-treated membranes was always observed even when measurements were done at different temperatures for both crude and alkaline-treated membranes. Nevertheless, when "leak" rates were

TABLE I
COMPOSITIONAL FEATURES OF THE LIPIDS FROM
ACChR MEMBRANES BEFORE AND AFTER ALKALINE
EXTRACTIONS*

	Crude AcChR-Membranes	Alkaline-treated AcChR-Membranes
Lipid/protein (wt/wt)	0.86	1.40
Phospholipids/cholesterol (molar ratio)	1.11	1.04
Fatty acid composition	identical (see ref. 6)	
Phospholipid composition† (molar percentage)	PC, 41%; PE, 44%	PC, 45%; PE, 39%

*Averages for two different preparations.

†PC, phosphatidyl choline; PE, phosphatidyl ethanolamine.




FIGURE 1 Dependence of the apparent TI^+ flux rate on the concentration of carbamylcholine. Apparent rate constants were obtained from computer-fitting of the kinetic traces from the stopped-flow experiments to the Stern-Volmer equation. AcChR-enriched membranes with (x) or without (○, control) preincubation with anti-AcChR polyclonal Fab were used.

measured in plain lipid vesicles obtained from lipid extracts of both membrane types, there was a similarity of values for the rate constants, indicating that the observed increased permeability to cations of the alkaline treated membranes is due to a destabilizing effect of the absence of the peripheral proteins per se, and not to the perturbation of the lipid matrix.

In additional experiments we used the fast stopped-flow/fluorescence quenching technique to study the effect of binding of polyclonal anti-AcChR antibodies on the cation translocation properties of the AcChR (Fig. 1). Binding of anti-AcChR antibodies inhibits the AcChR-mediated cation flux. The specificity of the inhibition is undetermined because polyclonal Fab were used; the degree of saturation of possible selected sites in the AcChR molecule cannot be ascertained under these conditions where competition for antigenic sites on the AcChR has been shown to occur (4). Specific monoclonal antibodies will be used in the near future to achieve selectivity of

binding. Meanwhile, the observed inhibition patterns apparently follow noncompetitive behavior between Fab and cholinergic ligands, in agreement with previous observations based on equilibrium binding studies (4).

We thank Tina Fehskens for typing the manuscript.

This work was supported by grant BNS-77-24715 from the National Science Foundation, awarded to M. Martinez-Carrion, and by grant NS-17029 from the National Institutes of Health, awarded to J. M. Gonzalez-Ros.

Received for publication 29 April 1983.

Note Added in Proof: Hill coefficients, maximum rate constants of the flux and apparent dissociation constants for cholinergic agonists were determined as described by Neubig et al. (10).

REFERENCES

1. Sator, V., J. M. Gonzalez-Ros, P. Calvo, and M. Martinez-Carrion. 1979. Pyrenesulfonyl azide: a marker of acetylcholine receptor subunits in contact with membrane hydrophobic environment. *Biochemistry*. 18:1200-1206.
2. Gonzalez-Ros, J. M., M. C. Farach, and M. Martinez-Carrion. 1983. Ligand induced effects at regions of acetylcholine receptor accessible to membrane lipids. *Biochemistry*. 22:3807-3811.
3. Mihovilovic, M., and M. Martinez-Carrion. 1979. Purification of high affinity Fab fragments from experimental autoimmune *Myasthenia gravis* rabbits and their effect on isolated acetylcholine receptors. *Biochemistry*. 18:4522-4528.
4. Farach, M. C., M. Mihovilovic, A. Paraschos, and M. Martinez-Carrion. 1982. Effect of anti-acetylcholine receptor Fab fragments from goats on functional membrane bound receptor. *Arch. Biochem. Biophys.* 214:140-154.
5. Farach, M. C., and M. Martinez-Carrion. 1983. A differential scanning calorimetry study of acetylcholine receptor rich membranes from *Torpedo californica*. *J. Biol. Chem.* 258:4166-4170.
6. Gonzalez-Ros, J. M., M. Llanillo, A. Paraschos, and M. Martinez-Carrion. 1982. The lipid environment of acetylcholine receptor from *Torpedo californica*. *Biochemistry*. 21:3467-3474.
7. Pick, V. 1981. Liposomes with a large trapping capacity prepared by freezing and thawing of sonicated phospholipid mixtures. *Arch. Biochem. Biophys.* 212:186-194.
8. Moore, H. P., and M. A. Raftery. 1980. Direct spectroscopic studies of cation translocation by *Torpedo* acetylcholine receptor on a time scale of physiological relevance. *Proc. Natl. Acad. Sci. USA*. 77:4509-4513.
9. Neubig, R. R., E. K. Krodel, N. D. Boyd, and J. B. Cohen. 1979. Acetylcholine and local anesthetics binding of *Torpedo* nicotinic postsynaptic membranes after removal of non-receptor peptides. *Proc. Natl. Acad. Sci. USA*. 76:690-694.
10. Neubig R. R., N. D. Boyd, and J. B. Kohen. 1982. Conformations of *Torpedo* acetylcholine receptor associated with ion transport and desensitization. *Biochemistry*. 21:3460-3467.