

A Disulfide Crosslink Between Cys⁹⁸ of Troponin-C and Cys¹³³ of Troponin-I Abolishes the Activity of Rabbit Skeletal Troponin

Hye-Shin Park,* Bang-Jian Gong,* and Terence Tao**§

*Muscle Research Group, Boston Biomedical Research Institute, Boston, Massachusetts 02114; †Department of Neurology, Harvard Medical School, Boston, Massachusetts 02115; and §Department of Biochemistry, Tufts University School of Medicine, Boston, Massachusetts 02111 USA

ABSTRACT Various thio-reactive bifunctional crosslinkers as well as 5,5'-dithiobis(2-nitrobenzoate)-mediated disulfide bond formation were used to crosslink troponin-C and troponin-I, the Ca²⁺-binding and inhibitory subunits of troponin, respectively. In all cases, substantial crosslinking was obtained when the reactions were carried out in the absence of Ca²⁺. No disulfide crosslinking occurred if either Cys⁹⁸ of TnC, or Cys¹³³ of TnI were blocked, indicating that these thiols are involved in the crosslinking. Troponin containing the disulfide crosslink is no longer capable of regulating actomyosin ATPase activity in a Ca²⁺-dependent manner. Our results suggest that the relative movement between the Cys⁹⁸ region of TnC and the Cys¹³³ region of TnI is required for the Ca²⁺-regulatory process in skeletal muscle.

INTRODUCTION

Troponin (Tn) is the Ca²⁺-binding thin filament protein complex that, in concert with tropomyosin (Tm), mediates regulation of vertebrate striated muscle contraction by Ca²⁺ (Ebashi and Endo, 1968). Tn is composed of three subunits, viz. TnC, TnI, and TnT, the Ca²⁺-binding, inhibitory, and Tm-binding subunits, respectively. X-ray crystallography revealed that TnC is composed of an N- and a C-terminal domain well separated by a single central helix (Herzberg and James, 1985; Sundaralingam et al., 1985). The N-terminal domain contains the two low affinity Ca²⁺-specific sites (designated as sites I and II), whereas the C-terminal domain contains the two high affinity Ca²⁺/Mg²⁺ sites (designated as sites III and IV). There are four Cys's in Tn; Cys⁹⁸ in TnC, Cys⁴⁸, Cys⁶⁴, and Cys¹³³ in TnI; and none in TnT. It is currently thought that binding of Ca²⁺ to the low affinity triggering sites of TnC induces successive changes in the mode of interaction between TnC and TnI, TnI and actin, Tm and actin, and finally, myosin head and actin (reviewed by El-Saleh et al., 1986). In view of its crucial role in Ca²⁺ regulation, the interaction between TnC and TnI has been

studied by many workers (reviewed by Leavis and Gergely, 1984; Zot and Potter, 1987; Grabarek et al., 1992). In particular, the region of TnC containing Cys⁹⁸ has been shown to interact with the inhibitory region of TnI by studies employing proteolytic fragments (Grabarek et al., 1981), NMR (Dalgarno et al., 1982), and photocrosslinking (Leszyk et al., 1987, 1988). Furthermore, crosslinking between Cys⁹⁸ of TnC and Cys¹³³ of TnI by FNB was reported by Dobrovol'sky et al. (1984), whereas resonance energy transfer studies showed that the distance between these two residues decreases by ~5 Å in response to the binding of Ca²⁺ to TnC (Tao et al., 1989; Wang and Cheung, 1984).

In this study, we used a variety of crosslinkers to characterize further the proximity relationship between Cys⁹⁸ of TnC and Cys¹³³ of TnI in the ternary Tn complex. We found that the two residues can be crosslinked by sulfhydryl-specific bifunctional reagents with linker regions of various lengths and rigidity. Furthermore, a disulfide bond between them can be formed rapidly when Nbs₂ is used as a catalyst. The ternary Tn complex containing this disulfide bond is no longer capable of regulating actomyosin ATPase activity in a Ca²⁺-dependent manner. The significance of these results with respect to the mechanism of action of Tn will be discussed.

MATERIALS AND METHODS

Chemicals

DMSDS was purchased from Molecular Probes (Junction City, OR). BMME was from Boehringer Mannheim (Indianapolis, IN). HEPES was from Research Organics (Cleveland, OH). Materials for polyacrylamide gel electrophoresis were from Bio-Rad (Richmond, CA). All other reagents were from Sigma Chemical Co. (St. Louis, MO).

Protein preparation

Tn, Tn subunits, and Tm were prepared from rabbit skeletal ether powder as described in Greaser and Gergely (1973). Actin was extracted from rabbit

Received for publication 20 December 1993 and in final form 17 March 1994.

Address reprint requests to Dr. Terence Tao, Boston Biomedical Research Institute, 20 Staniford St., Boston, MA 02114. Tel: 617-742-2010; Fax: 617-523-6649; E-mail: tao@bbri.eri.harvard.edu.

H.-S. Park's present address: Lucky Research Institute, Science Town, Tae Jeon, Chung-Nam, Korea.

Abbreviations used: Tn, Tm, and S1: rabbit skeletal troponin, tropomyosin, and myosin subfragment 1, respectively. TnC, TnI, and TnT: the Ca²⁺-binding, inhibitory, and Tm-binding subunits of Tn, respectively; SDS-PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis; DTT, dithiothreitol; Nbs₂, 5,5'-dithio-bis-2-nitrobenzoic acid; TNB⁻, thionitrobenzoic acid ion; *p*-PDM, *N,N'*-*p*-phenyldimaleimide; *o*-PDM, *N,N'*-*o*-phenyldimaleimide; BMME, bis-(maleimide)-methyl ether; DMSDS, 4,4'-dimaleimidylstilbene-2,2'-disulfonic acid; FNB, 1,3-difluoro-4,6-dinitrobenzene.

© 1994 by the Biophysical Society

0006-3495/94/06/2062/04 \$2.00

skeletal muscle acetone powder according to Spudich and Watt (1971). Myosin was prepared according to Balint et al. (1975), and chymotryptic S1 was prepared from myosin by the method of Weeds and Pope (1977).

Tn reconstitution

Tn was routinely reconstituted from the individual components shortly before crosslinking experiments. Equimolar amounts of TnC, TnI, and TnT were mixed in 6 M urea, 10 mM HEPES, pH 7.0 containing 0.1 M NaCl, 5 mM EDTA, and 5 mM DTT. After incubation for 1 h at room temperature, the proteins were slowly renatured and reconstituted by dialysis against 10 mM HEPES, pH 7.0 containing 0.1 M NaCl, 0.1 mM CaCl_2 , and 1 mM DTT, followed by further dialysis against the same buffer, but without DTT.

Crosslinking by bismaleimides

Molar excess 1.2-fold of bismaleimide derivatives were added to the reduced and reconstituted Tn in 10 mM HEPES, pH 7.0, 0.1 M NaCl in the presence or absence of Ca^{2+} . After 30 min of incubation, the reaction was stopped by adding excess DTT, and the reaction mixture was analyzed by SDS-PAGE.

Sulphydryl blocking

Cys⁹⁸ of TnC was alkylated by adding iodoacetamide to denatured TnC at a molar ratio of 3:1 (in 4 M guanidinium chloride, 5 mM EDTA, 10 mM HEPES, pH 7.0) followed by incubation for 1 hour at room temperature. The reaction was stopped by the addition of excess DTT, and the unreacted reagents were removed by dialysis. Alkylation of Cys¹³³ in TnI exploits the finding that only this TnI thiol is reactive in the ternary Tn complex (Chong and Hodges, 1982). Following a procedure that has been shown by Strasburg et al. (1985) to modify only Cys¹³³ of TnI, reconstituted Tn was incubated with *N*-ethylmaleimide at a molar ratio of 5:1 and allowed to react for 4 h at room temperature. The reaction was quenched with excess DTT, and unreacted reagents were removed by dialysis.

Miscellaneous

Mg²⁺-ATPase activity of regulated actomyosin was measured according to White (1982). SDS-PAGE was performed by the method of Laemmli (1970).

RESULTS

When reconstituted Tn was treated with a stoichiometric amount of Nbs₂ in the presence of Ca^{2+} , the absorbance at 412 nm (A_{412}) rose rapidly, indicating the release of TNB⁻ (Fig. 1). The increase in A_{412} leveled off when approximately 1 mol of TNB⁻ per mol of Tn was liberated. SDS-PAGE of this reaction mixture revealed the formation of only a small amount of high molecular weight-crosslinked material (Fig. 2A, lane 2). Addition of EGTA at this stage resulted in another rapid rise in A_{412} , indicating the release of a second mol of TNB⁻. SDS-PAGE of the final reaction mixture revealed a high molecular weight band whose mobility is consistent with the molecular mass of the TnC-TnI complex (Fig. 2A, lane 3). Addition of DTT to this reaction mixture before SDS-PAGE completely removed the high molecular weight band (Fig. 2A, lane 4), indicating that the complex was formed via a disulfide bond. When the high molecular weight band was excised from the gel, treated with DTT, and rerun on SDS-PAGE under reducing conditions, bands cor-




FIGURE 1 Time course of Nbs₂-catalyzed disulfide crosslinking between TnC and TnI monitored by the absorbance at 412 nm (A_{412}). An equimolar amount of Nbs₂ was added to freshly reconstituted Tn (20 μM) in a buffer containing 0.1 M NaCl, 0.1 mM CaCl_2 , 10 mM HEPES, pH 7.0, at room temperature. At the time indicated by the arrow, 2 mM EGTA was added. Calculated A_{412} value for one mol of TNB⁻ ion released per mol of Tn is 0.272, using $\epsilon_{412} = 13,600 \text{ cm}^{-1} \text{ M}^{-1}$.

responding to TnC and TnI were observed (Fig. 2B, lane 3), confirming the notion that the crosslink is between TnC and TnI. If either Cys⁹⁸ of TnC or Cys¹³³ of TnI was pre-blocked with alkylating reagents, no crosslinking was observed (Fig. 2C), indicating that the disulfide crosslink is between these two Cys's.

The activity of the Tn containing this disulfide bond was assayed by measuring its ability to mediate Ca^{2+} regulation of acto-S1 ATPase in an in vitro system. As shown in Table 1, with increasing molar ratio of Nbs₂ to Tn the Ca^{2+} -regulatory capacity of the system decreased, indicating that formation of the TnC-TnI disulfide bond rendered the modified Tn incapable of inhibiting acto-S1 ATPase in the absence of Ca^{2+} .

A variety of bifunctional thio-reactive reagents were also used to crosslink TnC and TnI; these include FNB, *p*-PDM, *o*-PDM, BMME, and DMSDS. It can be seen that in the absence of Ca^{2+} , all these reagents are capable of mediating the crosslinking of TnC and TnI (Fig. 3).

DISCUSSION

It has been shown that of the four sulphydryl groups in Tn, Cys⁴⁸, and Cys⁶⁴ of TnI are buried and are unreactive regardless of Ca^{2+} , Cys¹³³ of TnI is exposed and is reactive regardless of Ca^{2+} , and Cys⁹⁸ of TnC is reactive only in the absence of Ca^{2+} (Chong and Hodges, 1982; Potter et al., 1976). Based on this information, we can reasonably propose the following scheme for our disulfide crosslinking reaction: 1) Incubation of Tn with Nbs₂ first in the presence of Ca^{2+}




FIGURE 2 SDS-PAGE analysis of Nbs_2 -catalyzed disulfide crosslinking between TnC and TnI. In panel A, lane 1 is untreated Tn complex; lane 2 is Tn treated with Nbs_2 in the presence of Ca^{2+} just before the addition of EGTA (see Fig. 1); lane 3 is material in lane 2 after addition of EGTA and allowed to react for ~ 30 min; lane 4 is material in lane 3 treated with DTT; lane 5 are molecular weight markers. In panel B, lanes 1 and 2 are TnI and TnC, respectively, excised from a polyacrylamide gel, then reelectrophoresed; lane 3 is the high molecular weight band that has been excised from lane 3, panel A, treated with DTT, then reelectrophoresed in the presence of DTT. In panel C, lane 1 is ternary Tn complexed composed of TnI, TnT, and TnC alkylated at Cys⁹⁸; lane 2 is the material in lane 1 treated with Nbs_2 first in the presence then in the absence of Ca^{2+} as described in the legend for Fig. 1; lane 3 is Tn complex composed of TnC, TnT, and TnI alkylated at Cys¹³³; lane 4 is the material in lane 3 treated with Nbs_2 first in the presence then in the absence of Ca^{2+} as described in the legend for Fig. 1. All samples were treated with excess iodoacetamide to block free sulphydryls before SDS-PAGE.

TABLE 1 Ca^{2+} -dependent ATPase activity of a reconstituted system composed of S1, Tm, actin, and Tn that has been treated with varying amounts of Nbs_2 to induce varying extents of disulfide crosslinking*

[Nbs_2]/[Tn]	ATPase activity (s^{-1})		Ca^{2+} sensitivity† (%)
	- Ca^{2+}	+ Ca^{2+}	
0	0.05	0.24	80
0.2	0.10	0.21	52
0.4	0.17	0.22	23
0.8	0.20	0.20	0
1.0	0.22	0.22	0

* Concentrations of materials in the assay solution were: [S1] = 2 μM , [F-actin] = 3 μM , [Tm] = [Tn] = 0.5 μM , [EGTA] = 4 mM, or [Ca^{2+}] = 0.1 mM. Temperature was 25°C.

† Defined as $[1 - \text{ATPase}(-\text{Ca}^{2+})/\text{ATPase}(+\text{Ca}^{2+})] \times 100$.




FIGURE 3 Crosslinking of TnC and TnI by thio-specific bifunctional reagents. Ternary Tn complex (lane 1) was treated with *o*-PDM (lane 2), *p*-PDM (lane 3), FNB (lane 4), BMME (lane 5), and DMSDS (lane 6) in the absence of Ca^{2+} as described in Materials and Methods. Lane 7 is Tn treated with Nbs_2 as described in the legend for Fig. 1.

modifies Cys¹³³ of TnI because it is the only reactive sulphydryl under this condition. One mol of TNB^- ion is released, giving rise to the observed rise in A_{412} ; 2) addition of EGTA thereafter results in Cys⁹⁸ of TnC becoming reactive.

Crosslinking occurs via disulfide exchange; a second mol of TNB^- is released giving rise to the further increase in A_{412} .

It was noted that at the end of the reaction more TnC remains than TnI (Fig. 1 A, lane 3), whereas our reaction scheme would predict concomitant removal of TnC and TnI. This may be due to inaccuracies in the amounts of the added proteins, although the same result was obtained using purified instead of reconstituted Tn (data not shown). A more likely explanation is that Tn aggregates to a small extent and that Nbs_2 induces disulfide crosslinking between TnI subunits in the aggregates. It is clear, however, that the major reaction product is the disulfide-crosslinked TnC-TnI complex.

Aside from disulfide crosslinking, Dobrovol'sky et al. (1984) showed that FNB can crosslink the two Cys's, an observation that is confirmed here. We found further that these two Cys's can be crosslinked by four bismaleimide compounds whose maleimidyl groups are separated by a methyl ether group (BMME), phenyl group in the ortho configuration (*o*-PDM), phenyl group in the para configuration (*p*-PDM), and stilbene group (DMSDS). Except for BMME, the inter-maleimide linker groups of all these compounds are rigid. All of the crosslinking results suggest that the distance between these two Cys's is not fixed, and can span from 2 Å (to form a disulfide bond) to ~ 18 Å (to accommodate the stilbene group of DMSDS). It is interesting to note that our previous energy transfer studies also revealed some heterogeneity in the distance separating these two Cys's (Tao et al., 1989), suggesting that the protein regions containing these two Cys's possess considerable flexibility, a feature that may be required for the regulatory function of these proteins.

It was noted that whereas the disulfide crosslinked TnC-TnI complex comprises a single band, the bismaleimide crosslinked product is composed of two bands. Because it is known that the maleimide moiety is less specific than Nbs_2 for sulphydryls, it is possible that the second (lower) band

corresponds to crosslinking between Cys¹³³ of TnI and a non-Cys residue in TnC.

Our results show that the formation of a disulfide bond between Cys¹³³ of TnI and Cys⁹⁸ of TnC abolished the capacity of the Tn complex to regulate acto-S1 ATPase activity in a Ca²⁺-dependent manner; specifically, Tn containing this crosslink is incapable of inhibiting acto-S1 ATPase in the absence of Ca²⁺. We believe that this observation is relevant to the mechanism of thin filament-based regulation in the following manner: it has been proposed that activation of striated muscle contraction occurs when Ca²⁺ binds to the triggering sites of TnC (sites I and II), whereupon the affinity between TnC and TnI is increased, resulting in TnI detaching from actin. This in turn gives rise to the movement of Tm to a position in the thin filament that allows the cyclic interaction between myosin heads and actin to take place (El-Saleh et al., 1986).

Previous reports showing that upon the binding of Ca²⁺ to TnC, Cys¹³³ of TnI moves away from Cys³⁷⁴ of actin (Tao et al., 1990), and towards Cys⁹⁸ of TnC (Tao et al., 1989; Wang and Cheung, 1984) are in support of the above model. Our present work is also in support of this model because the disulfide crosslink between Cys¹³³ of TnI and Cys⁹⁸ of TnC can be expected to inhibit the movement of the Cys¹³³ region of TnI away from TnC and towards actin when Ca²⁺ is removed; the crosslink, therefore, locks the thin filament in the activated state, giving rise to our observed constitutively active acto-S1 ATPase.

It should be noted that in the absence of the high resolution structure of TnI, the location of Cys¹³³ in the TnI molecule is not known. We can conjecture, however, that it is in the vicinity of the so-called "inhibitory region" (residues 96–116), the region that contains the inhibitory peptide sequence (Syska et al., 1976), and most likely undergoes the Ca²⁺-dependent switching between TnC and actin. This is based on our findings that Cys⁹⁸ of TnC can be crosslinked to the inhibitory region of TnI via an attached photocrosslinker (Leszyk et al., 1987, 1988), and to Cys¹³³ of TnI via a disulfide bond (this work). Taken together, our findings suggest that a portion of the TnI molecule comprising both the inhibitory and the Cys¹³³ region undergoes switching between TnC and actin during Ca²⁺ activation of skeletal muscle contraction.

We thank Dr. Gale Strasburg for carrying out some of the initial crosslinking studies. We thank Drs. John Gergely and Zenon Grabarek for critical comments and reviewing the manuscript.

This work was supported by National Institutes of Health grant AR21673. Hye-Shin Park was a postdoctoral fellow of the American Heart Foundation, Massachusetts affiliate.

REFERENCES

Balint, M., F. A. Sreter, I. Wolf, B. Nagy, and J. Gergely. 1975. Substructure of heavy meromyosin-effect of Ca²⁺ and Mg²⁺ on tryptic fragmentation of heavy meromyosin. *J. Biol. Chem.* 250:6168–6177.

Chong, P. C. S., and R. S. Hodges. 1982. Proximity of sulfhydryl groups to the sites of interaction between components of the troponin complex from rabbit skeletal muscle. *J. Biol. Chem.* 257:2549–2555.

Dalgarno, D. C., R. J. A. Grand, B. A. Levine, A. J. G. Moir, G. M. M. Scott, and S. V. Perry. 1982. Interaction between troponin I and troponin C. Definition of the topography by proton magnetic resonance spectroscopy. *FEBS Lett.* 150:54–58.

Dobrovol'sky, A. B., N. B. Gusev, and P. Friedrich. 1984. Crosslinking of troponin complex with 1,3-difluoro-4,6-dinitrobenzene. Identification of the crosslink formed between troponin C and troponin I in the absence of Ca²⁺. *Biochim. Biophys. Acta.* 789:144–151.

Ebashi, S., and M. Endo. 1968. Calcium ion and muscle contraction. *Prog. Biophys. Mol. Biol.* 18:123–183.

El-Saleh, S., K. D. Warber, and J. D. Potter. 1986. The role of tropomyosin-troponin in the regulation of skeletal muscle contraction. *J. Muscle Res. Cell Motil.* 7:387–404.

Grabarek, Z., W. Drabikowski, P. C. Leavis, S. S. Rosenfeld, and J. Gergely. 1981. Proteolytic fragments of troponin C: interactions with the other troponin subunits and biological activity. *J. Biol. Chem.* 256:13121–13127.

Grabarek, Z., T. Tao, and J. Gergely. 1992. Molecular mechanism of troponin-C function. *J. Muscle Res. Cell Motil.* 13:383–393.

Greaser, M. L., and J. Gergely. 1973. Purification and properties of the components from troponin. *J. Biol. Chem.* 248:2125–2133.

Herzberg, O., and M. N. G. James. 1985. Structure of the calcium regulatory muscle protein troponin-C at 2.8 Å resolution. *Nature.* 313:653–659.

Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227:680–685.

Leavis, P. C., and J. Gergely. 1984. Thin filament proteins and thin filament-linked regulation of vertebrate muscle contraction. *CRC Crit. Rev. Biochem.* 16:235–305.

Leszyk, J., J. H. Collins, P. C. Leavis, and T. Tao. 1987. Cross-linking of rabbit skeletal muscle troponin with the photoactive reagent 4-maleimido-benzophenone: identification of residues in troponin I that are close to cysteine-98 of troponin C. *Biochemistry.* 26:7042–7047.

Leszyk, J., J. H. Collins, P. C. Leavis, and T. Tao. 1988. Cross-linking of rabbit skeletal muscle troponin subunits: labeling of cysteine-98 of troponin C with 4-maleimido-benzophenone and analysis of products formed in the binary complex with troponin T and the ternary complex with troponins I and T. *Biochemistry.* 27:6983–6987.

Potter, J. D., J. C. Seidel, P. Leavis, S. S. Lehrer, and J. Gergely. 1976. Effect of Ca²⁺ binding on troponin C. Changes in spin label mobility, extrinsic fluorescence, and sulfhydryl reactivity. *J. Biol. Chem.* 251:7551–7556.

Spudich, J. A., and S. Watt. 1971. The regulation of rabbit skeletal muscle contraction. Biochemical studies of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of myosin. *J. Biol. Chem.* 246:4866–4871.

Strasburg, G. M., P. C. Leavis, and J. Gergely. 1985. Troponin-C-mediated calcium-sensitive changes in the conformation of troponin I detected by pyrene excimer fluorescence. *J. Biol. Chem.* 260:366–370.

Sundaralingam, M., R. Bergstrom, G. Strasburg, S. T. Rao, M. Greaser, and B. C. Wang. 1985. Molecular structure of troponin C from chicken skeletal muscle at 3-Angstrom Resolution. *Science.* 227:945–948.

Syska, H., J. M. Wilkinson, R. J. A. Grand, and S. V. Perry. 1976. The relationship between biological activity and primary structure of troponin I from white skeletal muscle of the rabbit. *Biochem. J.* 153:375–387.

Tao, T., B.-J. Gong, and P. C. Leavis. 1990. Calcium-induced movement of troponin-I relative to actin in skeletal muscle thin filaments. *Science.* 247:1265–1372.

Tao, T., E. Gowell, G. M. Strasburg, J. Gergely, and P. C. Leavis. 1989. Ca²⁺ dependence of the distance between Cys-98 of troponin C and Cys-133 of troponin I in the ternary troponin complex. Resonance energy transfer measurements. *Biochemistry.* 28:5902–5908.

Wang, C.-K., and H. C. Cheung. 1984. Proximity relationship in the binary complex formed between troponin I and troponin C. *J. Mol. Biol.* 190:509–521.

Weeds, A. G., and B. Pope. 1977. Studies on the chymotryptic digestion of myosin. Effects of divalent cations on proteolytic susceptibility. *J. Mol. Biol.* 111:129–157.

White, H. D. 1982. Special instrumentation and techniques for kinetic studies of contractile systems. *Methods Enzymol.* 85:698–708.

Zot, A. S., and J. D. Potter. 1987. Structural aspects of troponin-tropomyosin regulation of skeletal muscle contraction. *Annu. Rev. Biophys. Biophys. Chem.* 16:535–559.