

The Electrostatic Basis for the Interfacial Binding of Secretory Phospholipases A₂

David L. Scott,* Arthur M. Mandel,† Paul B. Sigler,* and Barry Honig†

*Department of Molecular Biophysics and Biochemistry and the Howard Hughes Medical Institute, Yale University, New Haven, Connecticut 06511, and †Department of Biochemistry and Molecular Biophysics, Columbia University, New York, New York 10032 USA

ABSTRACT Biochemical and structural data suggest that electrostatic forces play a critical role in the binding of secretory phospholipases A₂ to substrate aggregates (micelles, vesicles, monolayers, and membranes). This initial binding (adsorption) of the enzyme to the interface is kinetically distinct from the subsequent binding of substrate to the buried active site. Thus, in the absence of specific active-site interactions, electrostatic forces operating at the molecular surface may orient and hold the enzyme at the interface. We have calculated the electrostatic potentials for 10 species of secretory phospholipases A₂ whose atomic coordinates have been determined by x-ray crystallography. Most of these enzymes show a marked electrostatic sidedness that is accentuated to a variable degree by the presence of the essential cofactor calcium ion. This asymmetry suggests a discrete interfacial binding region on the protein's surface, the location of which is in general agreement with proposals derived from the results of chemical modification, mutational, and crystallographic experiments.

INTRODUCTION

Phospholipases A₂ (PLA₂) specifically hydrolyze the 2-ester bond of 1,2-diacyl-3-*sn*-phosphoglycerides. The large family of small (14 kDa) homologous proteins derived from extracellular sources (e.g., exocrine pancreas, reptile and insect venoms, and human synovial fluid) possesses a distinct set of functional features: (1) high specific activities (V_m of up to 5000 IU), (2) alkaline pH optima, (3) full enzymatic activity only in the presence of organized lipid-water interfaces (micelles, monolayers, vesicles, or membranes), and (4) calcium dependence (Waite, 1987; Achari et al., 1987; Dennis, 1983; Verheij et al., 1981).

Secretory PLA₂ (sPLA₂) hydrolyze phospholipids that contain one of several naturally occurring polar head groups (e.g., choline, ethanolamine, inositol, serine, and glycerol). High affinity binding of sPLA₂ to substrate aggregates occurs without any significant deformation in the gross organization of the bilayer or leakage of vesicle-trapped components (Jain and Rogers, 1989). Because adsorption of the enzyme to the substrate interface can be kinetically distinguished from the binding of a phospholipid molecule to the catalytic site (Fig. 1), it is possible to analyze the qualities of an interface separately from the process of substrate selection and the chemistry of catalysis (Ghomashchi et al., 1991; Berg et al., 1991; Jain et al., 1991; Ramirez and Jain, 1991). A crucial role for electrostatic interactions in interfacial adsorption is likely given that the affinity of sPLA₂ for anionic bilayers is generally orders of magnitude larger than for zwitterionic bilayers regardless of the precise phospholipid composition of

the bilayer (Volwerk et al., 1986; Berg et al., 1991). Anionic and cationic additives (e.g., detergents) to the substrate interface increase and decrease, respectively, the catalytic efficiency of several sPLA₂ species (Volwerk et al., 1986; Jain and Jahagirdar, 1985; Apitz-Castro et al., 1982; Jain and Cordes, 1973). The binding of calcium ion to sPLA₂ also increases the affinity for aggregated substrate in some, but not all, enzyme species (van den Bergh et al., 1989; Menashe et al., 1986; Wells, 1972). Indeed, the adsorption of sPLA₂ to highly anionic vesicles can be sufficiently strong to make the enzyme completely processive; that is, the enzyme remains attached to the vesicle until virtually all of the phospholipid in the outer leaflet is hydrolyzed.

The three-dimensional structures of several sPLA₂s have been determined both in the absence (Freemont et al., 1993;




FIGURE 1 Schematic illustration of the scooting mode of catalysis by sPLA₂ (Berg et al., 1991). In this mechanism, which has substantial kinetic support (Ghomashchi et al., 1991; Berg et al., 1991; Jain et al., 1991; Ramirez and Jain, 1991; Jain and Rogers, 1989), adsorption of the enzyme (E) to the interface (E*) is a process that occurs before, and independent of, binding of substrate (S) to the active site (E*S). This permits the factors affecting adsorption and persistence of sPLA₂ at lipid-water interfaces to be selectively studied. (P = products, I = an inhibitor)

Received for publication 13 December 1993 and in final form 28 April 1994.

Address reprint requests to Paul B. Sigler, Department of Molecular Biophysics and Biochemistry, Yale University, Rm 147 Boyer Center for Molecular Medicine, 295 Congress Avenue, New Haven, CT 06510. Tel.: 203-737-4443; Fax: 203-776-3550.

© 1994 by the Biophysical Society

0006-3495/94/08/493/12 \$2.00

TABLE 1 Selected crystallographic and biophysical characteristics of the phospholipases A₂ species included in this study

| Phospholipase A ₂ source | PLA ₂ Class | pI | Aggregation state | Crystallization conditions | Charge groups |
|---------------------------------------------------|------------------------|-------|-------------------|---------------------------------------|---------------|
| 1. Porcine pancreas | I | 7.4 | Monomeric | pH 7.5(+) Ca ²⁺ /Organic | -15/+16 |
| 2. Bovine pancreas - zymogen | I | | Monomeric | pH 7.5(+) Ca ²⁺ /Organic | -15/+15 |
| 3. Bovine pancreas - enzyme | I | 9.2 | Monomeric | pH 7.5(+) Ca ²⁺ /Organic | -14/+14 |
| 4. <i>Naja naja atra</i> venom | I | 5.2 | Monomeric | pH 7.5(+) Ca ²⁺ /Organic | -10/+10 |
| 5. <i>Agkistridon piscivorus piscivorus</i> venom | II | 6.5 | Dimeric | pH 7.5(-) Ca ²⁺ /Organic | -17/+16 |
| 6. <i>Agkistridon piscivorus piscivorus</i> venom | II | 9.5 | Monomeric | pH 7.5(-) Ca ²⁺ /Organic | -13/+21 |
| 7. <i>Crotalus atrox</i> venom | II | 5.5 | Dimeric | pH 7.5(-) Ca ²⁺ /High Salt | -18/+12 |
| 8. <i>Crotalus adamanteus</i> venom | II | | Dimeric | pH 7.5(-) Ca ²⁺ /High Salt | -16/+14 |
| 9. Human nonpancreatic (synovial fluid) | II | >10.5 | Monomeric | pH 7.5(+) Ca ²⁺ /High Salt | - 8/+27 |
| 10. <i>Apis mellifera</i> venom | | >10.5 | Monomeric | pH 7.5(+) Ca ²⁺ /High Salt | -16/+25 |

Class distinctions (I or II) are made based upon the arrangement of disulfide bonds within the protein (Randolph et al., 1980). The bee venom PLA₂ is evolutionarily divergent but retains many of the core features of Class I/II enzymes (Scott et al., 1990b). The refined coordinates of these PLA₂ were derived from crystals obtained from a variety of solvent systems; however, it is unlikely that these distinctions, other than the presence or absence of calcium ion, are important to the electrostatic properties. "Charged residues" refers to a simple tabulation of the potential sources of negative charge [Asp, Glu] and the potential sources of positive charge [His, Arg, Lys].

TABLE 2 Amino acid sequences for the phospholipases A₂ included in this study

| | 1 | 5 | 10 | 15 | 20 | 25 | 29 | 33 |
|-----------------------|-----|-----|-----|-----|-----|-----|-----|----|
| Porcine pancreas | A | L | W | Q | F | R | S | M |
| Bovine pancreas | * | * | * | * | N | G | * | * |
| <i>Naja naja atra</i> | N | * | Y | * | K | N | * | Q |
| App-dimer | D | * | M | * | E | T | L | * |
| App-D49 | N | * | F | * | E | K | L | * |
| <i>C. atrox</i> | S | * | V | * | E | T | L | * |
| <i>C. adamanteus</i> | S | * | V | * | E | T | L | * |
| Human nonpanc. | N | * | V | N | * | H | R | * |
| | 35 | 40 | 45 | 50 | 55 | 60 | 65 | |
| Porcine pancreas | G | T | P | V | D | E | L | D |
| Bovine pancreas | * | * | * | * | D | * | * | * |
| <i>Naja naja atra</i> | * | * | * | * | D | * | * | * |
| App-dimer | * | R | * | Q | * | A | T | * |
| App-D49 | * | R | * | K | * | A | T | * |
| <i>C. atrox</i> | * | L | * | Q | * | A | T | * |
| <i>C. adamanteus</i> | * | R | * | Q | * | A | T | * |
| Human nonpanc. | * | K | * | K | * | * | * | * |
| | 70 | 75 | 80 | 85 | 90 | 95 | 100 | |
| Porcine pancreas | T | E | S | Y | S | Y | S | C |
| Bovine pancreas | * | N | N | * | * | * | * | * |
| <i>Naja naja atra</i> | F | K | T | * | * | * | * | * |
| App-dimer | L | D | S | * | T | * | V | E |
| App-D49 | M | D | I | * | T | * | V | E |
| <i>C. atrox</i> | * | V | I | * | T | * | E | E |
| <i>C. adamanteus</i> | * | V | I | * | T | * | E | E |
| Human nonpanc. | * | G | K | * | * | * | * | * |
| | 105 | 110 | 115 | 120 | 125 | 130 | | |
| Porcine pancreas | C | F | S | K | A | - | - | P |
| Bovine pancreas | * | * | * | * | * | * | * | * |
| <i>Naja naja atra</i> | * | A | G | * | - | - | * | N |
| App-dimer | * | R | D | N | K | V | T | * |
| App-D49 | * | R | D | N | L | K | T | * |
| <i>C. atrox</i> | * | R | D | N | I | P | S | * |
| <i>C. adamanteus</i> | * | R | D | N | I | P | S | * |
| Human nonpanc. | * | * | * | * | * | * | * | * |
| | 1 | 5 | 9 | 15 | 20 | 25 | 29 | 33 |
| <i>Apis mellifera</i> | I | I | Y | P | G | T | L | W |
| | 35 | 40 | 45 | 50 | 55 | 60 | 65 | |
| | D | H | C | P | D | V | M | S |
| | 70 | 75 | 80 | 85 | 90 | 95 | 100 | |
| | C | L | K | N | S | A | D | T |
| | 105 | 110 | 115 | 120 | 125 | 130 | 134 | |
| | C | G | E | R | T | E | G | R |

Asterisks are used to identify residues that are homologous to the prototype sequence (porcine pancreatic). The numbering system used is based upon the homologous core developed by Renetseder et al., 1985. Letters of residues bearing a potential positive charge (His, Arg, and Lys) or negative charge (Asp, Glu) are *italic* and **boldface**, respectively. Black rectangles appear above residues involved in binding the primary calcium ion; open rectangles appear above conserved residues of the catalytic network, and His48 appears with an open circle. The references for these sequences are porcine pancreatic (Puijk et al., 1977), bovine pancreatic (Fleer et al., 1978), *Naja naja atra* (Tsai et al., 1981), *A. piscivorus piscivorus* (App) D49 monomer (Maragnore and Heinrikson, 1993) and D49 dimer (Heinrikson, 1991), *C. atrox* (Randolph and Heinrikson, 1982), *C. adamanteus* (Heinrikson et al., 1977), and human nonpancreatic (Seilhamer et al., 1989). The sequence of the App-D49 monomer differs at position 78 (Asp→Glu) from the cited reference based on the interpretation of the high resolution (1.5 Å) crystal structure. The bee-venom PLA₂ sequence (Kuchler et al., 1989) is not aligned because of its low overall homology.

Figure 2(a)




Figure 2(b)




FIGURE 2 (a) Stereopair of the proposed interfacial binding surface—"0 degrees." The orientation of the C_α trace of this typical Class I/II PLA₂ (human nonpancreatic secretory) corresponds to the zero degree views used in subsequent figures displaying the calculated electrostatic potentials. The calcium ion cofactor (*yellow sphere*), the transition-state analog (colored *red*), the side chains of the hydrophobic channel (colored *green*) [Leu-2, Phe-5, His-6, Ile-9, Ala-19, Val-31, Cys-45, Tyr-52, Lys-69, and Phe-106], and additional surface residues also considered to be important to interfacial binding [Val-3, Glu-17, Leu-20, and Phe-24] are shown. (b) Stereopair of the proposed interfacial binding surface—"90 degrees." The molecule shown in *a* has been rotated 90° counter-clockwise (as viewed from above) around a vertical axis that lies in the plane of the figure.

Westerlund et al., 1992; Wery et al., 1991; Scott et al., 1990a; Brunie et al., 1985; Dijkstra et al., 1983, 1982, 1981) and presence of transition-state (Scott et al., 1992; White et al., 1991; Scott et al., 1991b) and substrate analogs (Thunnissen et al., 1990). These structures, along with data from mutational and chemical modification studies (Yang and Chang, 1989; van Oort et al., 1985; Yoshida et al., 1988; de Haas et al., 1987; Dijkstra et al., 1981b), have implicated a

discrete surface of the protein in the process of interfacial adsorption. This surface consists of a ring of positively charged side chains surrounding a hydrophobic channel that leads to the centrally located active site (Scott et al., 1990a). The binding, orientation, and persistence of sPLA₂ at the lipid-water interface is thought to be governed by the electrostatic and hydrophobic forces operative at this surface.







FIGURE 3 Electrostatic potentials calculated from the coordinates of crystalline phospholipases A₂: category I. Figures depict the electrical potentials of one or both standard orientations (0°/90°). An electrostatic potential of 1 kT/e (where k is the Boltzmann constant and T is the Kelvin temperature) is equal to 25 mV at 300 Kelvin. The contour levels of the maps were chosen to provide the maximum pictorial information possible about the respective electrostatic potential magnitudes and orientations. A contour level of 1 kT means that an atom of one unit of positive charge, sitting at a point on the contour grid, would have an interaction energy of +1kT (approximately +0.6 kcal/mole at 300 Kelvin) with the electrostatic potential at that point. Blue is used to indicate positive potentials, and red is used to indicate negative potentials. In three cases (the *A. p. piscivorus* and *C. atrox* dimers, and the bee-venom sPLA₂), additional skeletal traces are provided to clarify the molecular orientation. a. *C. atrox* sPLA₂ (dimer): alpha-carbon trace—(linear numbering system). b. *C. atrox* sPLA₂ (dimer): 0 degrees—0.5 kT—0 calcium ions. c. *C. atrox* sPLA₂ (monomer created by using only one of the two molecules of the dimer): 90 degrees—2.0 kT—0 calcium ions. d. *C. adamantus* sPLA₂ (dimer): 0 degrees—0.5 kT—0 calcium ions. e. *A. p. piscivorus* sPLA₂ (dimer): alpha-carbon trace—(linear numbering system). f. *A. p. piscivorus* sPLA₂ (dimer): 0 degrees—1.0 kT—0 calcium ions. g. *A. p. piscivorus* sPLA₂ (monomer created by using only one of the two molecules of the dimer): 0 degrees—1.0 kT—0 calcium ions. h. *Apis mellifera* sPLA₂: alpha-carbon trace—(linear numbering system). i. *Apis mellifera* sPLA₂ (the glycosyl group of Asn13 was not included in the electrostatics calculations): 0 degrees—0.25 kT—1 calcium ion.



(3e) *A. p. piscivorus* dimer: alpha-carbon trace - (linear-numbering system).



(3f) *A. p. piscivorus* dimer
0 degrees - 1.0 kT - 0 calcium ions.



(3g) *A. p. piscivorus* dimer (created by using the coordinates for only one of the two subunits of the dimer)
0 degrees - 1.0 kT - 0 calcium ions.

FIGURE 3—continued

In this paper, we examined the surface electrostatic potentials of a diverse group of sPLA₂ that includes members of both Class I and Class II sPLA₂, the pancreatic pro-phospholipase (Class I), as well as the evolutionarily distant bee-venom enzyme. The electrostatic potentials calculated for these enzymes demonstrate a marked molecular sidedness, with the proposed interfacial binding surface generally lying in or adjacent to the most positively charged regions. This is consistent with the enhanced affinity and increased processivity exhibited by sPLA₂ on more negatively charged substrate aggregates. The essential, but weakly bound ($K_d > 10^{-4}$ M), calcium ion accentuates the protein's electrostatic

sidedness and might play a particularly important functional role in those enzymes whose apo- state is only marginally asymmetric. In contrast, enzyme dimerization (Myatt et al., 1991) as seen in the sPLA₂ species from *Crotalus* and *Agkistridon* venoms confers no obvious electrostatic advantages.

MATERIALS AND METHODS

Coordinates for the sPLA₂s were obtained from either the Brookhaven Protein Data Bank or from recently completed studies at Yale. Secretory PLA₂ species were chosen for inclusion based on the availability of well




FIGURE 3—continued

refined crystallographic coordinates, a suitable degree of biochemical and kinetic characterization, and the presence of unique structural/functional features (e.g., dimerization). The relevant biophysical properties and amino acid sequences are provided in Tables 1 and 2, respectively.

Electrostatic calculations were carried out using the DelPhi computer program (Nicholls and Honig, 1991; Klapper et al., 1986) and displayed using INSIGHT II (Biosym Technologies Inc., San Diego, CA). No corrections were made for the overestimation of electrostatic charge potentials by the linear Poisson-Boltzmann equation in the few cases where the potentials were very high. An ionic strength of 145 mM was chosen for the solvent region, and the temperature was set to 298 K for all calculations. The dielectric constant was set to 4 for the protein interior. The crystallographically observed water molecules were not treated explicitly but were instead included as part of the solvent region that was assigned a dielectric constant of 80 (Gilson et al., 1987; Sharp and Honig, 1990). In most cases, pK_a calculations were made for individual side chains and incorporated into the determination of electrical potentials. The effects of these calculations on the global results were, however, minor.

RESULTS

The high overall structural homology of these sPLA₂s permitted the choice of a single molecular orientation as the standard for comparison of electrostatic potential maps (Fig. 2). The view chosen as the standard ("0 degrees") is similar to that depicted in recent papers describing the structures of several sPLA₂ transition-state analog complexes (Scott et al., 1992, 1991a, b; White et al., 1991). The advantage of this view is that it looks directly down the hydrophobic channel while remaining perpendicular to the plane of the proposed interfacial binding surface. The calculated electrostatic potential maps (Figs. 3-5) are oriented according to this system. An additional view ("90 degrees" of rotation around the in-page vertical axis, Fig. 2 b) is provided where the standard view failed to illustrate electrical potential differences adequately between the front and back of the molecule.

DISCUSSION

The 10 sPLA₂ included in this study can be divided into three categories based on their calculated electrostatic potentials. Members of the first category, which includes the enzymes from the venoms of *C. atrox* (Western diamondback rattlesnake), *C. adamanteus* (Eastern diamondback rattlesnake), *A. p. piscivorus* (American cottonmouth water moccasin)-dimeric D49, and *Apis mellifera* (the honey bee), show distinct molecular sidedness in their distribution of positive and negative electrical potentials, but the scope and magnitude of the positive potential is relatively small (Fig. 3). The *C. atrox* sPLA₂ is the most striking in this regard, with the intact dimer having only a small patch of positive potential (Fig. 3 b). Calculation of the electrical potential for a theoretical monomer (only the structure of the crystallized dimer is known), however, reveals that the dimer interface quenches part of the charge distribution intrinsic to the dissociated subunits (Fig. 3 c). The magnitude and location of this quenched charge suggests that on the surface of a negatively charged substrate aggregate, the monomeric form is preferentially stabilized. An analogous prediction can be made for the dimeric enzyme from *A. p. piscivorus* venom where most of the positive electrostatic potential also resides at the shielded dimer interface (Fig. 3, f and g). The oligomerization state of the active enzyme at the interface remains controversial despite recent evidence supporting the monomer (Jain et al., 1991; Ferreira et al., 1993).

The second category of electrostatic distributions includes the enzymes from the venom of *Naja naja atra* (Taiwanese cobra) and from bovine pancreas (Fig. 4). In both of these sPLA₂s, the most prominent positive electrostatic potential envelops a surface that is adjacent, but 45–90° away from the proposed interfacial binding surface (Scott et al., 1990a). The electrostatic asymmetry of this category of enzymes is quite






FIGURE 4 Electrostatic potentials calculated from the coordinates of crystalline phospholipases A₂: category II. Figures depict the electrical potentials of one or both standard orientations (0°/90°). An electrostatic potential of 1 kT/e (where k is the Boltzmann constant and T is the Kelvin temperature) is equal to 25 mV at 300 Kelvin. The contour levels of the maps were chosen to provide the maximal pictorial information possible about the respective electrostatic potential magnitudes and orientations. A contour level of 1 kT means that an atom of one unit of positive charge, sitting at a point on the contour grid, would have an interaction energy of +1 kT (approximately +0.6 kCal/mole at 300 Kelvin) with the electrostatic potential at that point. Blue is used to indicate positive potentials, and red is used to indicate negative potentials. a. *Naja naja atra* sPLA₂: 0 degrees—1.0 kT—2 calcium ions. b. *Naja naja atra* sPLA₂: 90 degrees—1.0 kT—2 calcium ions. c. Bovine pancreatic sPLA₂: 0 degrees—0.5 kT—1 calcium ion. d. Bovine pancreatic sPLA₂: 90 degrees—0.5 kT—1 calcium ion. e. Bovine pancreatic sPLA₂ (trans-aminated): 0 degrees—0.5 kT—1 calcium ion. f. Bovine pro-enzyme (the positions of the seven N-terminal residues that are disordered in the crystal structure [Glu-Ala-Gly-Leu-Asn-Ser-Arg] were modeled based on probable stereochemistry and packing considerations within the crystallographic unit cell): 0 degrees—0.5 kT—1 calcium ion.



(4e) Bovine pancreatic trans-aminated sPLA₂
0 degrees - 0.5 kT - 1 calcium ion.



(4f) Bovine pro-enzyme (the positions of the seven N-terminal residues which are disordered in the crystal structure [Glu-Ala-Gly-Leu-Asn-Ser-Arg] were modelled based on probable stereochemistry and packing considerations within the crystallographic unit cell) 0 degrees - 0.5 kT - 1 calcium ion.

FIGURE 4—continued

distinctive, with obvious implications for the orientation of the molecule in its interactions with negatively charged substrate aggregates. The bovine pancreatic sPLA₂ is a particularly interesting case because the availability of the refined coordinates for the transaminated enzyme and the pro-enzyme permit the impact of an intact amino-terminus on the global electrostatics to be evaluated (Fig. 4 c-f). Modifications of the amino-terminus appear to only modestly alter the global electrostatic potentials. This argues against a simple electrostatic explanation for the failure of the transaminated and pro-enzymes to hydrolyze organized substrate aggregates efficiently (Jain et al., 1986).

The electrostatic potentials of the third category of enzymes grossly resemble those of the second category, but the maximum development of the positive potential shifts to envelop the proposed interfacial adsorption surface (Fig. 5). The three enzymes in this group, the D49 monomer from *A. p. piscivorus* venom, the human nonpancreatic secretory sPLA₂ (synovial fluid), and the porcine pancreatic enzyme, preserve a striking molecular asymmetry despite the predominance of positive electrical potential. In the case of the synovial sPLA₂, demonstration of this asymmetry requires the electrostatic maps to be displayed at very high contour levels (14 kT). The influence of the binding of the essential cofactor calcium ion on global charge distributions is particularly evident in the case of the porcine sPLA₂. The porcine enzyme has been shown by crystallography and by so-

lution kinetics to bind two calcium ions per molecule. One of these ions participates as the essential cofactor in catalysis, the second calcium ion appears to be critical for the binding of the enzyme to aggregated substrate at alkaline pH (van den Bergh, 1989). From Fig. 5, e and f it is obvious that calcium ion markedly enhances the porcine enzyme's positive electrostatic potentials. A similar effect of the binding of calcium ion would be expected on the electrostatic potential distributions of other sPLA₂, especially for those enzymes with acidic isoelectric points (e.g., *C. atrox*).

In general, the electrostatic potentials calculated for these sPLA₂ show clear molecular sidedness with the maximum positive potential adjacent to or incorporating the face of the protein that contains the opening to the hydrophobic channel. These results correlate well with biochemical and genetic evidence for a specific surface on the enzyme that interacts with lipid aggregates (e.g., vesicles, monolayers, and membranes) (Jain and Vaz, 1987; Jain et al., 1986; Mao et al., 1986; Pieterse et al., 1974). The efficiency of this surface in promoting binding of the enzyme to the lipid-water interface is critically dependent on its charge composition (Ghomashchi et al., 1991; Berg et al., 1991; Jain et al., 1991; Ramirez and Jain, 1991; Thuren et al., 1987; de Haas et al., 1987; Volwerk et al., 1986). Spectroscopic experiments with the porcine pancreatic sPLA₂ have directly correlated the degree of interfacial adsorption with the mole fraction of the anionic component in the lipid aggregate. Highly anionic




FIGURE 5 Electrostatic potentials calculated from the coordinates of crystalline phospholipases A₂: category III. Figures depict the electrical potentials of one or both standard orientations (0°/90°). An electrostatic potential of 1 kT/e (where k is the Boltzmann constant and T is the Kelvin temperature) is equal to 25 mV at 300 Kelvin. The contour levels of the maps were chosen to provide the maximal pictorial information possible about the respective electrostatic potential magnitudes and orientations. A contour level of 1 kT means that an atom of one unit of positive charge, sitting at a point on the contour grid, would have an interaction energy of +1 kT (approximately +0.6 kcal/mole at 300 Kelvin) with the electrostatic potential at that point. No correction was made for the over-estimation of electrostatic charge potentials by the linear Poisson-Boltzmann equation for *c* and *d*. Blue is used to indicate positive potentials, and red is used to indicate negative potentials. a. *A. p. piscivorus* sPLA₂ (49 monomer): 90 degrees—0.5 kT—0 calcium ions. b. *A. p. piscivorus* sPLA₂ (D49 monomer): 90 degrees—0.5 kT—1 calcium ion. c. Human nonpancreatic (synovial) sPLA₂: 0 degrees—4.0 kT—2 calcium ions. d. Human nonpancreatic (synovial) sPLA₂: 90 degrees—14.0 kT—2 calcium ions. e. Porcine pancreatic sPLA₂: 90 degrees—1.0 kT—0 calcium ions. f. Porcine pancreatic sPLA₂: 90 degrees—1.0 kT—2 calcium ions.




FIGURE 5—continued

mixed vesicles permit the efficient catalysis of substrate with little discrimination among phospholipid headgroups. The high efficiency of this “scooting mode” form of catalysis results, at least in part, from the persistence of the enzyme at the interface ($K_d < 10^{-13}$ M).

Mutational and chemical modification studies with the bovine pancreatic sPLA₂ suggested that the interfacial binding domain was a discrete protein face whose center lay at the opening to the hydrophobic channel (Fig. 2). Substitution of positively charged residues (e.g., arginine) for uncharged residues contributing to this surface (e.g., Asn-6) resulted in a two- to threefold higher value of K_{cat} on substrate aggregates (de Haas et al., 1987). Removal of sources of positive charge in this region either by specific mutations or by chemical modification of the protein lead to dramatic reductions in K_{cat} . Spectroscopic studies have correlated adsorption of the enzyme to the interface with the desolvation of uncharged residues forming the opening to the hydrophobic channel (Jain and Maliwal, 1993). The recently completed crystal structures of three sPLA₂ transition-state analog complexes (*Naja naja atra*, White et al., 1991; bee-venom, Scott et al., 1991b; and human nonpancreatic, Scott et al., 1992) and a substrate-based inhibitor complex (bovine pancreatic, Thunnissen et al., 1990) also clearly implicate this surface of the protein in interfacial adsorption.

Although the results described here are compatible with delocalized molecular electrostatics playing a key role in the orientation and persistence of sPLA₂ at water-lipid interfaces, there is also evidence for a strong hydrophobic effect (Tomasselli et al., 1989; Van der Weile et al., 1988; Cho et al., 1988; Drainas and Lawrence, 1978). Lugtigheid et al. (1993) observed a 60-fold increase in the apparent K_m of the porcine sPLA₂ for micellar substrate after acylation of

Lys-56. Specific acylation of the bee-venom sPLA₂ improved the enzyme's K_{cat} by up to two orders of magnitude (Drainas and Lawrence, 1978). Presumably, the covalent attachment of hydrophobic groups activates sPLA₂ in a manner analogous to the addition of positive charge to the interfacial binding surface; i.e., increasing the affinity and persistence of the enzyme at the interface.

Clearly, the degree to which electrostatic forces affect interfacial catalysis can only be approximated by the techniques used here. Although an adequate description can be made of the enzyme's electrostatic field potentials, describing the environment at the surface of the bilayer in similar terms is at best difficult (Cevc, 1990). The 10 crystalline enzymes used in this study reveal a common, electrostatically favorable surface for potential interaction with substrate aggregates. Small variations in the location of this surface, such as seen in the Category II enzymes, suggest some flexibility in the initial adsorption that is presumably either optimized for catalysis by other forces or is inconsequential. Additional mutational and chemical modification studies should prove helpful in determining the precise boundaries and chemistry of the interfacial adsorption surface.

The research at Yale was supported by a grant from the United States Public Health Service (GM24324 and NS 25867) and the Howard Hughes Medical Institute. The work at Columbia was funded by a grant from the National Science Foundation (MCB9304127). A. M. Mandel is the recipient of a Medical Scientist Training Program grant (5T32GM07367) and gratefully acknowledges helpful discussions with Marilyn Gunner and An-Suei Yang. Coordinates for the crystalline sPLA₂ can be obtained from the Brookhaven Protein Data Bank or from the authors.

REFERENCES

- Achari, A., D. Scott, P. Barlow, J. C. Vidal, Z. Otwinowski, S. Brunie, and P. B. Sigler. 1987. Facing up to membranes: structure/function relation-

- ships in phospholipases. *Cold Spring Harbor Symp. Quant. Biol.* 52: 441-452.
- Apitz-Castro, R. J., M. K. Jain, and G. H. de Haas. 1982. Origin of the latency phase during the action of phospholipase A₂ on unmodified phosphatidylcholine vesicles. *Biochim. Biophys. Acta.* 688:349-356.
- Berg, O. G., B.-Z. Zhu, J. Rogers, and M. K. Jain. 1991. Interfacial catalysis by phospholipase A₂: determination of the interfacial kinetic rate constants. *Biochemistry.* 30:1283-1291.
- Brunie, S., J. Bolin, D. Gewirth, and P. B. Sigler. 1985. The refined crystal structure of dimeric phospholipase A₂ at 2.5 Å: access to a shielded catalytic site. *J. Biol. Chem.* 260:9742-9749.
- Cevc, G. 1990. Membrane electrostatics. *Biochim. Biophys. Acta.* 1031: 311-382.
- Chettibi, S., F. Lyall, and A. J. Lawrence. 1990. Rapid activation of the non-toxic basic isoform of phospholipase A₂ from *Naja mossambica* (spitting cobra) by long-chain fatty acylation. *Toxicon.* 28:953-961.
- Cho, W., A. G. Tomasselli, R. L. Heinrikson, and F. J. Keady. 1988. The chemical basis for interfacial activation of monomeric phospholipases A₂. *J. Biol. Chem.* 263:11237-11241.
- de Haas, G. H., G. J. M. van Scharrenburg, and A. J. Slotboom. 1987. Interaction of pancreatic phospholipases A₂ and semisynthetic mutants with anionic substrates and substrate analogues. *Biochemistry.* 26: 3402-3408.
- Dennis, E. A. 1983. Phospholipases A₂. *The Enzymes.* 16:307-353.
- Dijkstra, B. W., J. Drenth, and K. H. Kalk. 1981a. Active site and catalytic mechanism of phospholipase A₂. *Nature.* 289:604-606.
- Dijkstra, B. W., K. H. Kalk, W. G. J. Hol, and J. Drenth. 1981b. Structure of bovine pancreatic phospholipase A₂ at 1.7 Å resolution. *J. Mol. Biol.* 147:97-123.
- Dijkstra, B. W., R. Renetseder, K. H. Kalk, W. G. J. Hol, and J. Drenth. 1983. Structure of porcine pancreatic phospholipase A₂ at 2.6 Å resolution and comparison with bovine phospholipase A₂. *J. Mol. Biol.* 168:163-179.
- Dijkstra, B. W., G. J. H. van Nes, K. H. Kalk, N. P. Brandenberg, W. G. J. Hol, and J. Drenth. 1982. The structure of bovine pancreatic phospholipase A₂ at 3.0 Å resolution. *Acta Cryst.* B38:793-799.
- Drinas, D., and A. J. Lawrence. 1978. Activation of bee venom phospholipase A₂ by oleoyl imidazole. *Eur. J. Biochem.* 91:131-138.
- Fleer, E. A. M., H. M. Verheij, and G. H. de Haas. 1978. The primary sequence of bovine pancreatic phospholipase A₂. *Eur. J. Biochem.* 82: 261-269.
- Freemont, D. H., D. H. Anderson, I. A. Wilson, and E. A. Dennis. 1993. Crystal structure of phospholipase A₂ from Indian cobra reveals a trimeric association. *Proc. Natl. Acad. Sci. USA.* 90:342-346.
- Ghomashchi, F., B.-Z. Yu, O. Berg, M. K. Jain, and M. H. Gelb. 1991. Interfacial catalysis by phospholipase A₂: substrate specificity in vesicles. *Biochemistry.* 30:7318-7329.
- Gilson, M., K. Sharp, and B. Honig. 1987. Calculating the electrostatic potentials of molecules in solution: method and error assessment. *J. Comp. Chem.* 9:327-335.
- Heinrikson, R. L. 1991. Dissection and sequence analysis of phospholipases A₂. *Methods Enzymol.* 197:201-214.
- Heinrikson, R. L., E. T. Krueger, and P. S. Keim. 1977. The amino acid sequence of phospholipase A₂ from the venom of *Crotalus adamanteus*. *J. Biol. Chem.* 252:4913-4921.
- Jain, M. K., and E. H. Cordes. 1973. Phospholipase I. Effect of n-alkanols on the rate of enzymatic hydrolysis of egg phosphatidylcholine. *J. Membr. Biol.* 14:101-118.
- Jain, M. K., and D. V. Jahagirdar. 1985. Action of phospholipases A₂ on bilayers: effects of inhibitors. *Biochim. Biophys. Acta.* 814:319-326.
- Jain, M. K., and B. P. Maliwal. 1993. Spectroscopic properties of the states of pig pancreatic phospholipase A₂ at interfaces and their possible molecular origin. *Biochemistry.* 32:11838-11846.
- Jain, M. K., B. P. Maliwal, G. H. de Haas, and A. J. Slotboom. 1986. Anchoring of phospholipase A₂: the effect of anions and deuterated water, and the role of the n-terminus region. *Biochim. Biophys. Acta.* 860: 448-461.
- Jain, M. K., G. Ranadive, B.-Z. Yu, and H. M. Verheij. 1991. Interfacial catalysis by phospholipase A₂: monomeric enzyme is fully catalytically active at the bilayer interface. *Biochemistry.* 30:7330-7340.
- Jain, M. H., and J. Rogers. 1989. Substrate specificity for interfacial catalysis by phospholipase A₂ in the scooting mode. *Biochim. Biophys. Acta.* 1003:91-97.
- Jain, M. K., and W. L. C. Vaz. 1987. Dehydration of the lipid-protein microinterface on binding of phospholipase A₂ to lipid bilayers. *Biochim. Biophys. Acta.* 905:1-8.
- Klapper, I., R. Hagstrom, R. Fine, K. Sharp, and B. Honig. 1986. Focusing the electrostatic fields in the active site of Cu, Zn superoxide dismutase: effects of ionic strength and amino acid modification. *Proteins.* 1:47-59.
- Kuchler, K., M. Gmachl, M. J. Sippl, and G. Kreil. 1989. Analysis of the cDNA for phospholipase A₂ from honeybee venom glands. *Eur. J. Biochem.* 184:249-254.
- Lugtighed, R. B., G. A. F. Nicolaes, E. J. A. Veldhuizen, A. J. Slotboom, H. M. Verheij, and G. H. de Haas. 1993. Acylation of porcine pancreatic phospholipase A₂ influences penetration and substrate head-group binding, depending on the position of the acylated lysine in the enzyme molecule. *Eur. J. Biochem.* 216:519-525.
- Mao, S.-Y., A. H. Maki, and G. H. de Haas. 1986. Optically detected magnetic resonance studies of porcine pancreatic phospholipase A₂ binding to a negatively charged substrate analogue. *Biochemistry.* 25:2781-2786.
- Maragnone, J. M., and R. L. Heinrikson. 1993. Erratum to "The lysine-49 phospholipase A₂ from the venom of *Agkistrion piscivorus piscivorus*. Relation of structure and function to other phospholipases A₂". *J. Biol. Chem.* 268:6064.
- Menashe, M., G. Romero, R. L. Biltonen, and D. Lichtenberg. 1986. Hydrolysis of dipalmitoylphosphatidylcholine small unilamellar vesicles by porcine pancreatic phospholipase A₂. *J. Biol. Chem.* 261:5328-5333.
- Myatt, E. A., F. J. Stevens, and P. B. Sigler. 1991. Effects of pH and calcium ion on self-association properties of two dimeric phospholipases A₂. *J. Biol. Chem.* 266:16331-16335.
- Nicholls, A., and B. Honig. 1991. A rapid finite difference algorithm, utilizing successive over-relaxation to solve the Poisson-Boltzmann equation. *J. Comp. Chem.* 12:435-445.
- Pieterse, W. A., J. C. Vidal, J. J. Volwerk, G. H. de Haas. 1974. Zymogen-catalyzed hydrolysis of monomeric substrates and the presence of a recognition site for lipid-water interfaces to phospholipase A₂. *Biochemistry.* 13:1455-1460.
- Puijk, W. C., H. M. Verheij, and G. H. de Haas. 1977. The primary structure of the phospholipase A₂ from porcine pancreas: a reinvestigation. *Biochim. Biophys. Acta.* 492:254-259.
- Ramirez, F., and M. H. Jain. 1991. Phospholipase A₂ at the bilayer interface. *Proteins Struct. Funct. Genet.* 9:229-239.
- Randolph, A., and R. L. Heinrikson. 1982. *Crotalus atrox* phospholipase A₂. *J. Biol. Chem.* 257:2155-2161.
- Randolph, A., T. P. Sakmar, and R. L. Heinrikson. 1980. Phospholipase A₂: structure, function, and evolution. In *Frontiers in Protein Chemistry*. A. B. Liu, C. D. Mamiya, and E. F. Yasunoku, editors. Elsevier/North Holland Inc., Amsterdam. 297-312.
- Renetseder, R., S. Brunie, B. W. Dijkstra, J. Drenth, and P. B. Sigler. A comparison of the crystal structures of phospholipase A₂ from bovine pancreas and *Crotalus atrox* venom. *J. Biol. Chem.* 260:11627-11634.
- Scott, D. L., Z. Otwinowski, M. H. Gelb, and P. B. Sigler. 1990b. Crystal structure of bee-venom phospholipase A₂ in a complex with a transition-state analogue. *Science.* 250:1563-1566.
- Scott, D. L., S. P. White, J. L. Browning, J. J. Rosa, M. H. Gelb, and P. B. Sigler. 1992. Crystal structures of free and inhibited human secretory phospholipase A₂ from inflammatory exudate. *Science.* 254:1007-1010.
- Scott, D. L., S. P. White, Z. Otwinowski, W. Yuan, M. H. Gelb, and P. B. Sigler. 1990a. Interfacial catalysis: the mechanism of phospholipase A₂. *Science.* 250:1541-1546.
- Seilhamer, J. J., W. Pruzanski, P. Vadas, S. Plant, J. A. Miller, J. Kloss, and L. K. Johnson. 1989. Cloning and recombinant expression of phospholipase A₂ present in rheumatoid arthritic synovial fluid. *J. Biol. Chem.* 264:5335-5338.
- Sharp, K. A., and B. Honig. 1990. Electrostatic interactions in macromolecules: theory and applications. *Annu. Rev. Biophys. Biophys. Chem.* 19: 301-332.
- Thunnissen, M. M. G. M., A. B. Elso, K. H. Kalk, J. Drenth, B. W. Dijkstra, O. P. Kuipers, R. Dijkman, G. H. de Haas, and H. M. Verheij. 1990. X-ray

- structure of phospholipase A_2 complexed with a substrate-derived inhibitor. *Nature*. 347:689-691.
- Thuren, T., J. A. Virtanen, R. Verger, and P. K. J. Kinnunen. 1987. Hydrolysis of 1-palmitoyl-2-[6-(pyren-1-yl)]hexanoyl-sn-glycero-3-phospholipids by phospholipase A_2 : effect of the polar head-group. *Biochim. Biophys. Acta*. 917:411-417.
- Tomasselli, A. G., J. Hui, J. Fisher, H. Zurcher-Neely, I. M. Reardon, E. Oriaku, F. J. Kezdy, and R. L. Heinrikson. 1989. Dimerization and activation of porcine pancreatic phospholipase A_2 via substrate level acylation of lysine 56. *J. Biol. Chem.* 264:10041-10047.
- Tsai, I. H., S. H. Wu, and T. B. Lo. 1981. The complete amino acid sequence of the phospholipase A_2 from the venom of *Naja naja atra* (Taiwan cobra). *Toxicon*. 19:141-152.
- van den Bergh, C. J., A. C. A. P. A. Bekkers, H. M. Verheij, and G. H. de Haas. 1989. Glutamic acid 71 and aspartic acid 66 control the binding of the second calcium ion in porcine pancreatic phospholipase A_2 . *Eur. J. Biochem.* 182:307-313.
- Van der Weile, F. C., W. Atsma, R. Dijkman, A. M. M. Schreurs, A. J. Slotboom, and G. H. de Haas. 1988. Site-specific e-NH₂ monoacylation of pancreatic phospholipase A_2 . 1. Preparation and properties. *Biochemistry*. 27:1683-1688.
- van Oort, M. G., R. Dijkman, J. D. R. Hille, and G. H. de Haas, G. H. 1985. Kinetic behavior of porcine pancreatic phospholipase A_2 on zwitterionic and negatively charged single-chain substrates. *Biochemistry*. 24:7987-7993.
- Verheij, H. M., A. J. Slotboom, and G. H. de Haas. 1981. Structure and function of phospholipase A_2 . *Rev. Physiol. Biochem. Pharmacol.* 91:91-203.
- Volwerk, J. J., P. C. Jost, G. H. de Haas, and O. H. Griffith. 1986. Activation of porcine pancreatic phospholipase A_2 by the presence of negative charges at the lipid-water interface. *Biochemistry*. 25:1726-1733.
- Waite, M., editor. 1987. The Phospholipases, Handbook of Lipid Research 5. Plenum Press, New York.
- Wells, M. A. 1972. A kinetic study of the phospholipase A_2 (*Crotalus adamanteus*) catalyzed hydrolysis of 1,2-dibutyl-sn-glycero-3-phosphorylcholine. *Biochemistry*. 11:1030-1041.
- Wery, J. P., R. W. Schevitz, D. K. Clawson, J. L. Bobbitt, E. R. Dow, G. Gamboa, T. Goodson, Jr., R. B. Hermann, R. M. Kramer, D. B. McClure, E. D. Mihelich, J. E. Putnam, J. D. Sharp, D. H. Stark, C. Teater, M. W. Warrick, and N. D. Jones. 1991. Structure of recombinant human rheumatoid arthritic synovial fluid phospholipase A_2 at 2.2 Å resolution. *Nature*. 352:79-82.
- Westerlund, B., P. Nordlund, U. Uhlin, D. Eaker, and H. Eklund. 1992. The three-dimensional structure of notexin, a presynaptic neurotoxic phospholipase A_2 at 2.0 Å resolution. *FEBS Lett.* 301:159-164.
- White, S. P., D. L. Scott, Z. Otwinowski, M. H. Gelb, and P. B. Sigler. 1990. Crystal structure of cobra-venom phospholipase A_2 in a complex with a transition-state analogue. *Science*. 250:1560-1563.
- Yang, C.-C., L.-S. Chang. 1989. Studies on the status of lysine residue in phospholipase A_2 from *Naja naja atra* (Taiwan cobra) snake venom. *Biochem. J.* 262:855-860.
- Yoshida, M., T.-C. Shieh, N. Oda, H. Kihara, C.-C. Chang, and M. Ohno. 1988. Tryptophan residue essential for activity of *Naja naja atra* phospholipase A_2 . *J. Biochem.* 103:156-161.