

A Spectrally Silent Transformation in the Photolysis of Octopus Rhodopsin: A Protein Conformational Change Without Any Accompanying Change of the Chromophore's Absorption

Yoshinori Nishioku,* Masashi Nakagawa,[†] Motoyuki Tsuda,[†] and Masahide Terazima*

*Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan; and [†]Department of Life Science, Himeji Institute of Technology, Harima Science Garden City, Kamigori, Akou-gun, Hyogo 678-1297, Japan

ABSTRACT A spectrally silent transformation in the photolysis of octopus rhodopsin was detected by the time-resolved transient grating method. Our results showed that at least two photointermediates, which share the same chromophore absorption spectrum, exist after the final absorption changes. Previously, mesorhodopsin was thought to decay to the final photoproduct, acid metarhodopsin with a lifetime of 38 μ s at 15°C, but the present results show that there is at least one intermediate species (called transient acid metarhodopsin) with a lifetime of 180 μ s at 15°C, before forming acid metarhodopsin. This indicates that the parts of the protein distant from the chromophore are still changing even after the changes in microenvironment around the chromophore are over. From the signal intensity detected by the transient grating method, the volume change of the spectrally silent transformation was found to be $\Delta V = 13$ ml/mol. The activation energy of the spectrally silent transformation is much lower than those of other transformations of octopus rhodopsin. Since stable acid metarhodopsin has not been shown to activate the G protein, this transient acid metarhodopsin may be responsible for G protein activation.

INTRODUCTION

Rhodopsins consist of 11-*cis* retinal chromophore covalently attached to the opsin via a protonated Schiff base linkage. Light isomerizes the 11-*cis* retinal to its all-*trans* form which initiates a series of protein conformational changes, called the photointermediates of bleaching, each usually characterized by their different absorption spectra. One of the photointermediates interacts with the G protein, resulting in the electrical excitation of a photoreceptor cell (Stryer, 1986; Tsuda, 1987). Although the photoreaction processes of both vertebrate and invertebrate rhodopsins resemble each other for several μ s after photoexcitation, eventually they begin to differ. After metarhodopsin II, which activates the G protein, the vertebrate rhodopsins are finally hydrolyzed to free all-*trans* retinal and opsin. On the other hand, photolysis of the invertebrate pigments leads to a stable species containing all-*trans* retinal. In octopus rhodopsin, the stable acid metarhodopsin can be converted back to the original rhodopsin upon orange-light illumination (Tsuda, 1979, 1987). In a previous paper, Nakagawa et al. (1998) showed that stable final photoproduct could not activate the G protein, but a novel intermediate that lies between mesorhodopsin and stable acid metarhodopsin is a candidate for the species that can activate the G protein. Spectroscopic methods have been extensively applied to

the study of the reactions following light excitation (Tsuda, 1979; Tsuda et al., 1980; Pande et al., 1987; Ohtani et al., 1988; Bagley et al., 1989; Deng et al., 1991; Taiji et al., 1992; Masuda et al., 1993; Nakagawa et al., 1997; Nishimura et al., 1997). Transient absorption spectroscopy showed that transformation of mesorhodopsin to acid metarhodopsin (with a lifetime of 31 μ s at 15°C) is the final spectral transformation in the photolysis of octopus rhodopsin. Previously, Nakagawa et al. (1998) studied the photointermediate of octopus rhodopsin responsible for G protein activation by a GTP γ S binding assay and concluded that a novel intermediate between mesorhodopsin and stable acid metarhodopsin, but not stable acid metarhodopsin, is the most likely candidate for the activating species. The presence of a precursor to stable acid metarhodopsin was suggested by transient absorption changes at alkaline conditions (Tsuda 1979), Fourier-transform infrared spectroscopy (FTIR) (Masuda et al., 1993) and light scattering measurements (Nakagawa et al., 1998). However, this postulated intermediate has not been yet identified kinetically. To study kinetics of a process which is not accessible by any optical absorption changes, the transient grating technique (Richard et al., 1992; Hara et al., 1996; Takeshita et al., 2000) or the photoacoustic technique (van Brederode et al., 1995) could be powerful methods; they monitor the reaction volume change. We examined the kinetics of the activation of octopus rhodopsin by using the time-resolved transient grating technique and found a spectrally silent transition accompanying the protein structure changes after the absorption changes are complete. Based on these investigations, formation of the acid metarhodopsin, which has previously been assigned by absorp-

Received for publication 25 September 2000 and in final form 21 March 2001.

Address reprint requests to Masahide Terazima, Department of Chemistry, Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan. Tel: 81-78-753-4026; Fax: 81-75-753-4000; E-mail: mterazima@kuchem.kyoto-u.ac.jp.

© 2001 by the Biophysical Society

0006-3495/01/06/2922/06 \$2.00

tion spectroscopy, is shown in this paper to be the formation of transient acid metarhodopsin.

EXPERIMENTAL

Octopus rhodopsin was prepared as described previously (Tsuda et al., 1986). Briefly, microvillar membranes from the octopus retina (*Octopus dofleini*) were isolated by sucrose floatation and then washed with buffer A [400 mM KCl, 10 mM MgCl₂, 10 mM 2-(*N*-morpholine)ethanesulfonic acid (pH 6.5), 1 mM dithiothreitol, 1 mM benzamidinium-HCl, 20 μ M 4-(amidinophenyl)methanesulfonyl fluoride hydrochloride (APMSF)] and then with buffer B [10 mM 4-morpholinepropanesulfonic acid (pH 6.5), 1 mM dithiothreitol, 1 mM benzamidinium-HCl, 20 μ M APMSF]. After the microvillar membranes were solubilized in 1% (w/v) sucrose monolaurate (SM 1200), 10 mM Tris-HCl buffer (pH 7.4) containing 1 mM dithiothreitol, 1 mM benzamidinium-HCl, and 20 μ M APMSF, octopus rhodopsin was affinity purified by concanavalin-A (Con-A) sepharose (Amersham Pharmacia Biotech, Uppsala, Sweden). Finally, the sample (~ 80 μ M rhodopsin) for the transient grating and transient absorption measurements was prepared in 10 mM 3-(*N*-morpholinopropanesulfonic acid (pH 7.4) and 1% SM 1200.

The transient grating setup was similar to that previously described (Terazima and Hirota, 1993; Hara et al., 1996). Briefly, the beam of a XeCl excimer laser-pumped dye laser (Lambda Physik Compex 102xc, Lumomics Hyper Dye 300; λ , 465 nm, Göttingen, Germany) was divided by a beam splitter and the beams crossed inside a quartz sample cell (optical path-length, 2 mm). The laser power of the excitation was < 8 μ J/pulse. The interference pattern (transient grating) created in the sample was probed by a He-Ne laser (633 nm) or a diode laser (840 nm) as a Bragg diffracted signal (transient grating signal). The transient grating signal was detected by a photo-multiplier (Hamamatsu R928, Iwata, Japan) and averaged by a digital oscilloscope (Tektronix 2430A, Toyko, Japan). A transient absorption signal was also detected by a photo-multiplier and averaged by a digital oscilloscope using 543.5 nm (He-Ne laser) probe light. The repetition rate of the excitation laser was 1-Hz. The sample of solution was gently stirred to prevent an accumulation of photoproduct in the photoexcitation region. The photolyzed sample was irradiated with orange light (a tungsten lamp with a cutoff filter ($\lambda > 590$)) to convert the photoproduct back to rhodopsin. Using these manipulations, the decrease of signal intensity due to pigment bleaching can be prevented. Experiments were performed at between 0 to 35°C, with the temperature controlled by a thermal bath. Bromocresol purple (Nacalai Tesque, Kyoto, Japan) was used for a calorimetric reference. The value of q was determined from the decay rate of the thermal grating signal of this reference sample (see below).

PRINCIPLE

The transient grating method can be used to detect protein dynamics without utilizing the optical absorption of a chromophore. The transient grating is created by the interference of the two excitation pulsed laser beams. There are several origins for the transient gratings. After optical excitation with a nanosecond laser pulse, one of the dominant contributions is the temperature change of the medium induced by the energy from the radiationless decay of excited states and by the enthalpy change of the reaction (thermal grating $\delta n_{th}(t)$). Another important contribution originates from the change in the absorbance of the solution induced by light. The newly formed species and the depletion of the original species contribute to the gratings through the absorption changes. We have called this component the population grating ($\delta n_{pop}(t)$). The partial molecular volume (V) difference between the reactant and products (ΔV) induces a change of the average density of the medium, and this change causes the refractive index change δn_{vol} (volume grating). In this work, the term "species grating" is used to describe both of these contributions ($\delta n_{spe} = \delta n_{pop} + \delta n_{vol}$). If the contribution of absorbance is negligible and the diffraction efficiency of the probe light is small, the transient grating signal intensity is proportional to the square of the peak-null difference of the refractive index created by the interference pattern of the excitation light intensity (Eichler et al., 1986; Miller, 1989; Terazima and Hirota, 1993; Hara et al., 1996).

$$I_{TG}^{1/2}(t) = A|\delta n_{th}(t) + \delta n_{pop}(t) + \delta n_{vol}(t)| \quad (1)$$

where A is a constant. The refractive index change of the volume grating component is given by

$$\delta n_{vol} = V \frac{dn}{dV} \Delta V \Delta N \quad (2)$$

where Vdn/dV is the refractive index change by the molecular volume change, and ΔN is the number of the reacting molecules in a unit volume. If the protein structure changes, the dynamics should be apparent in the transient grating signal through this component (even if the optical absorption spectrum does not change). By a quantitative measurement of δn_{vol} and the known solvent properties (Vdn/dV), we can determine the absolute value of ΔV .

The temporal profile of the refractive index grating due to the thermal grating is given by

$$\delta n_{th}(t) = (dn/dT)/\rho C_p [dQ(t)/dt * \exp(-D_{th}q^2t)] \quad (3)$$

where $*$ is the convolution integral, $Q(t)$ is the thermal energy coming out from the sample, dn/dT is the temperature dependence of the refractive index, ρ is the density, C_p is the heat capacity at a constant pressure, and D_{th} is the thermal diffusivity. The time dependence of the species grating is determined by the kinetics of the reaction and the molecular diffu-

sion process. If we can neglect the reaction kinetics in the molecular diffusion time-region, the time dependence is given by (Terazima and Hirota, 1993; Hara et al., 1996)

$$\delta n_{\text{spe}}(t) = \delta n_r \exp(-D_r q^2 t) + \delta n_p \exp(-D_p q^2 t) \quad (4)$$

where q is a grating wave number, and δn_r and δn_p represent refractive index changes by the reactant and product, respectively. D_r and D_p are the mass diffusion coefficients of the reactant and product, respectively. Because the magnitude of D_{th} is ~ 2 – 3 orders of magnitude larger than D_r and D_p , we can easily separate the thermal and species grating contributions (Hara et al., 1996; Takeshita et al., 2000). Furthermore, the reaction kinetics can be separated from the diffusion process by measuring the transient grating dynamics at different q^2 , because the diffusion process depends on q^2 , while the reaction kinetics should not.

RESULTS AND DISCUSSION

Fig. 1 shows the time course of the transient absorption changes at 543.5 nm after photolysis at 15°C. The kinetics can be described with two lifetimes, 520 ns and 38 μs , at this probe wavelength. These processes have been attributed to the lumirhodopsin \rightarrow mesorhodopsin \rightarrow acid metarhodopsin transformations. The absorption changes were essentially complete within 100 μs . These observations are consistent with previous findings (Nakagawa et al., 1997). The species formed with 38 μs (or 31 μs as reported by Nakagawa et al., 1997) has previously been considered to be the final photoproduct and has been called acid metarhodopsin (Nakagawa et al., 1997). However, hereafter in this paper, we call it the transient acid metarhodopsin because we discovered new dynamics for the creation and decay of this species.

Fig. 2 shows the early time profile of the transient grating signal of the photolyzed octopus rhodopsin probed at 840 nm at 15°C. We found that the signal over this fast time




FIGURE 1 The kinetics of the light-induced transient absorption changes of octopus rhodopsin probed at 543.5 nm. The inset is the time profile with a longer time scale to show that there are no absorbance changes after about 100 μs (mesorhodopsin \rightarrow transient acid metarhodopsin).




FIGURE 2 The kinetics of the transient grating signal of octopus rhodopsin probed at 840 nm on a fast time scale ($t < 20 \mu\text{s}$). Assignment of each kinetic component is labeled.

scale ($t < 100 \mu\text{s}$) can be fitted by a sum of four exponential functions with lifetimes of 130 ns, 520 ns, 3.4 μs , and 38 μs at $q^2 = 2.0 \times 10^{12} \text{ m}^{-2}$. One of the lifetimes, 3.4 μs , depends on q^2 , but the others do not. Considering the reaction scheme (Scheme 1) and the lifetimes, we can identify the q^2 -independent lifetimes to the bathorhodopsin \rightarrow lumirhodopsin, lumirhodopsin \rightarrow mesorhodopsin, and mesorhodopsin \rightarrow transient acid metarhodopsin processes, respectively. These lifetimes agree well with those observed from the transient absorption measurements discussed just above, and support the reaction scheme (Nakagawa et al., 1997).

Besides these q^2 -independent lifetimes, a q^2 -dependent lifetime is observed. The q^2 -dependent dynamic implies that this dynamic is governed by a diffusion process. It is easily assigned to a thermal diffusion process (eq. (3)), because the decay-rate constant agrees with $D_{\text{th}} q^2$ calculated from the reported D_{th} . Therefore, this signal is the thermal grating signal. It is created by the thermal energy from a nonradiative transition or an enthalpy change (ΔH) accompanying the reaction. [This signal is not apparent under weak excitation conditions, but its intensity increases with increasing laser power (data not shown). The relative enhancement of the thermal grating intensity can be explained in terms of the multiphoton excitation of either rhodopsin or the intermediates within the excitation pulse width.] From the thermal grating signal intensity we can determine the enthalpies of these intermediates, and these results will be published elsewhere. Interestingly, even after the absorption changes are over, there are several components in the transient grating signal (Fig. 3). The longer time-temporal profile can be reproduced by three exponential functions, one with a lifetime of 180 μs and two with lifetimes that depend on q^2 (4.5 ms and 16 ms at $q^2 = 2.0 \times 10^{12} \text{ m}^{-2}$). The slower q^2 -dependent kinetics were assigned to the diffusion processes of rhodopsin and the final species in this time range (Eq. 4). The most interesting and important component is the one with the 180- μs lifetime. Because this rate does not depend on q^2 , this




FIGURE 3 (a) The kinetics of the transient grating signal probed at 840 nm in a slow time scale ($t > 20 \mu\text{s}$) (dotted line). The kinetics can be well-fitted by three exponential functions with lifetimes of $150 \mu\text{s}$ and two lifetimes that depend on q (diffusion terms) (solid line). The inset is the time profile with a longer time scale to show the diffusion dynamics of rhodopsin. (b) In order to show the $150\text{-}\mu\text{s}$ component clearly, the diffusion terms are subtracted from the transient grating signal.

dynamic is not due to a molecular diffusion process. We will examine this component further in detail.

First, because the final photoproduct of octopus rhodopsin can be converted back to rhodopsin upon light irradiation, the excitation laser pulse may excite this photoproduct, which was created by the previous pulses and initiate the back reaction. In such a case, kinetics of intermediates in the back reaction could contribute to the transient grating signal. This possibility was excluded by transient grating measurements with different excitation repetition rates. We confirmed that the transient grating signal becomes weak, and almost disappears, by successive excitation with a faster repetition rate and without stirring the solution. Therefore, the contribution of the back reaction from the final product in the signal can almost be neglected. Secondly, with increasing laser power, the thermal grating signal intensity increases, whereas the species grating signal intensity is saturated. This implies that all of the species in the transient grating signal come from the photoexcitation of rhodopsin, and none come from the sequential multiphoton excitation of the rhodopsin. Thirdly, the origin of the signal was examined by changing the probe wavelength. The refractive index change due to the absorption spectrum change (population grating) depends on the probe wavelength as pre-




FIGURE 4 The kinetics of the transient grating signals probed at 840 nm at 0, 10.2, 24.8, and 33.9°C.

dicted by the Kramers-Kronig relationship (Longhurst, 1967), whereas the thermal grating and the volume grating intensities are less sensitive to the probe wavelength. If the refractive index change, which gives rise to the population grating signal, comes from an absorption change of the chromophore, the population grating signal intensity is expected to increase at shorter wavelengths (closer to the absorption bands). It was found that the relative intensities of the population grating signals, except for the $180\text{-}\mu\text{s}$ component, were indeed enhanced at 633 nm compared with those at 840 nm. Since the $180 \mu\text{s}$ component was less sensitive to the probe wavelength, the origin of this signal is not due to an absorption change of the chromophore, but due to a volume change. Since this component is silent for absorption spectroscopy, the component that induces the volume change originates from protein conformational changes without any accompanying absorption changes of the chromophore. From quantitative measurements of the signal intensity, we determined the volume change to be 13




FIGURE 5 Arrhenius plots of three transformation process; the lumirhodopsin \rightarrow mesorhodopsin (circles), and the mesorhodopsin \rightarrow transient acid metarhodopsin (squares) and transient acid metarhodopsin \rightarrow acid metarhodopsin formation (diamonds).

TABLE 1 ΔH^\ddagger and ΔS^\ddagger of the formation of acid metarhodopsin, the lumirhodopsin \rightarrow mesorhodopsin, and the mesorhodopsin \rightarrow transient acid metarhodopsin processes

	$E_{\text{act}}/\text{kJ mol}^{-1}$	$\ln A$	$\Delta H^\ddagger/\text{kJ mol}^{-1}$	$\Delta S^\ddagger/\text{J mol}^{-1} \text{ K}^{-1}$	$\Delta G^\ddagger/\text{kJ mol}^{-1}$
Lumi \rightarrow meso	59.8	39.2	57.3	72.9	35.6
Meso \rightarrow transient acid meta	79.5	43.3	77.1	107.3	45.1
Transient acid meta \rightarrow acid meta	32.4	21.9	30.0	-70.6	51.0

ΔG^\ddagger , Gibbs free energy at $T = 298.15 \text{ K}$.

ml/mol (volume expansion). We attributed this new kinetic component to a protein conformational change.

A long-lived intermediate was found to exist before stable acid metarhodopsin by time-resolved Fourier-transform infrared spectroscopy (Masuda et al., 1993). During illumination of microvillar membranes by blue light, bands in the 1800 to 1700 cm^{-1} region changed, which could be associated with the change in the state of carboxylic group of Asp-81. A preliminary observation of a light-induced fluorescence quenching (Nakagawa and Tsuda, to be published) showed that the lifetime of the movement of the fluorescence probe attached at Cys345 in the C terminus of octopus rhodopsin was $\sim 1 \text{ ms}$ at 15°C .

The results of the present transient grating, FTIR and the fluorescence quenching experiments show that parts of the protein distant from the chromophore are still changing even after the changes in microenvironment around the chromophore are over. The observed kinetics (180 μs at 15°C) in the transient grating signal could represent the transformation from the transient acid metarhodopsin to the final product (acid metarhodopsin), or to another intermediate species, which should eventually transform to the acid metarhodopsin with a slower rate. We could examine whether there are other transient intermediate species with much longer lifetimes, or not, using the transient grating signal method on much longer time scale. However this longer time scale measurement is very difficult because the dynamics, which represent the molecular diffusion process, becomes dominant on


such a later time scale. Thus, the species created with 180 μs is tentatively called acid metarhodopsin.

To further understand this process, we measured the transient grating signal at various temperatures from 0 to 35°C . Fig. 4 shows the kinetics of the transient grating signals probed at 840 nm for 0, 10.2, 24.8, and 33.9°C . From this temperature dependence of the rate (k), the enthalpy of activation (ΔH^\ddagger) and the entropy of activation (ΔS^\ddagger) were obtained from activated complex theory,

$$\ln(kh/k_B T) = \Delta S^\ddagger/R - \Delta H^\ddagger/RT,$$

where h is the Planck constant, k_B is the Boltzmann constant and R is the gas constant. Fig. 5 shows the Arrhenius plots of the formation rate of transient acid metarhodopsin together with the rates of the lumirhodopsin \rightarrow mesorhodopsin, and the mesorhodopsin \rightarrow transient acid metarhodopsin processes. Each of these kinetics can be described by the relation with a single ΔH^\ddagger . The ΔH^\ddagger and ΔS^\ddagger that we determined are listed in Table 1. It is interesting that, although ΔH^\ddagger and ΔS^\ddagger for the lumirhodopsin \rightarrow mesorhodopsin and mesorhodopsin \rightarrow transient acid metarhodopsin processes are similar, the values for the creation of acid metarhodopsin are much lower. The different values of ΔH^\ddagger and ΔS^\ddagger imply that the activation processes for lumirhodopsin \rightarrow mesorhodopsin and mesorhodopsin \rightarrow transient acid metarhodopsin are very different from those for the transient acid metarhodopsin \rightarrow acid metarhodopsin process. The kinetics of the lumirhodopsin to transient acid metarhodopsin form should reflect conformational changes of the protein moiety around the chromophore. Therefore, this different activation process is consistent with our conclusion that the structure change to acid metarhodopsin is not the dynamics around the chromophore.

In bovine rhodopsin, the formation of metarhodopsin II induces helix changes in the pigment so that this species can activate the G protein (Cohen et al., 1993). The structural change of octopus rhodopsin observed in this study may correspond to a similar change on this pigment. The low activation energy of the new dynamics probably reflects that this large amplitude motion is less hindered. The molecular volume expansion of 13 ml/mol may result from the deactivation process of the transient acid metarhodopsin to form acid metarhodopsin, or initially contacted area of the helix is open for the water, which may be responsible for the activation of the G protein by a possible intermediate species.



SCHEME 1

CONCLUSION

A new product, transient acid metarhodopsin, in the photolysis of octopus rhodopsin was detected by the time-resolved transient grating method. The lifetime at 15°C is 180 μ s, which is about 5 times slower than the final absorption change. The protein structure around the chromophore should be very similar in both species, but the structures apart from the chromophore must be different. The volume change of $\Delta V = 13$ ml/mol was obtained for transient acid metarhodopsin \rightarrow acid metarhodopsin transform. The activation energy for this movement is much smaller than those for the structural movement around the chromophore. Since stable acid metarhodopsin has not been shown to activate the G protein, this transient acid metarhodopsin may be responsible for G protein activation.

REFERENCES

- Bagley, K. A., L. Eisenstein, T. G. Ebrey, and M. Tsuda. 1989. A comparative study of the infrared difference spectra for octopus and bovine rhodopsins and their bathorhodopsin photointermediates. *Biochemistry*. 28:3366–3373.
- Cohen, G. B., T. Yang, P. R. Robinson, and D. D. Oprian. 1993. Constitutive activation of opsin: influence of charge at position 134 and size at position 296. *Biochemistry*. 32:6111–6115.
- Deng, H., D. Manor, G. Weng, P. Rath, Y. Koutalos, T. Ebrey, R. Gebhard, J. Lugtenburg, M. Tsuda, and R. H. Callender. 1991. A resonance Raman study of octopus bathorhodopsin with deuterium labeled retinal chromophores. *Photochem. Photobiol.* 54:1001–1007.
- Eichler, H. J., P. Günter, and D. W. Pohl. 1986. Laser Induced Dynamic Gratings. Springer-Verlag, Berlin.
- Ernst, O. P., K. P. Hofmann, and T. P. Sakmar. 1995. Characterization of rhodopsin mutants that bind transducin but fail to induce GTP nucleotide uptake: classification of mutant pigments by fluorescence, nucleotide release, and flash-induced light scattering assays. *J. Biol. Chem.* 270:10580–10586.
- Franke, R. R., T. P. Sakmar, R. M. Graham, and H. G. Khorana. 1992. Structure and function in rhodopsin: studies of the interaction between the rhodopsin cytoplasmic domain and transducin. *J. Biol. Chem.* 267:14767–14774.
- Hara, T., N. Hirota, and M. Terazima. 1996. New application of the transient grating method to a photochemical reaction: the enthalpy, reaction volume change, and partial molar volume measurements. *J. Phys. Chem.* 100:10194–10200.
- Hashimoto, S., H. Takeuchi, M. Nakagawa, and M. Tsuda. 1996. Ultraviolet resonance Raman evidence for the absence of tyrosinate in octopus rhodopsin and the participation of Trp residues in the transformation to acid metarhodopsin. *FEBS Lett.* 398:239–242.
- König, B., A. Arendt, J. H. McDowell, M. Kahlert, P. A. Hargrave, and K. P. Hofmann. 1989. *Proc. Natl. Acad. Sci. U.S.A.* 86:6878–6882.
- Longhurst, R. S. 1967. Geometrical and Physical Optics, 2nd ed. John Wiley, New York.
- Masuda, S., E. Morita, M. Tasumi, T. Iwasa, and M. Tsuda. 1993. Infrared studies of octopus bathorhodopsin: existence of a long-lived intermediate and the states of the carboxylic group of Asp81 in rhodopsin and its photoproducts. *FEBS Lett.* 317:223–227.
- Miller, R. J. D. 1989. In Time Resolved Spectroscopy. R. J. H. Clark, and R.E. Hester, editors. John-Wiley and Sons.
- Nakagawa, M., T. Iwasa, S. Kikkawa, M. Tsuda, and T. G. Ebrey. 1999. How vertebrate and invertebrate visual pigments differ in their mechanism of photoactivation. *Proc. Natl. Acad. Sci. U.S.A.* 96:6189–6192.
- Nakagawa, M., S. Kikkawa, T. Iwasa, and M. Tsuda. 1997. Light-induced protein conformational changes in the photolysis of octopus rhodopsin. *Biophys. J.* 72:2320–2328.
- Nakagawa, M., S. Kikkawa, K. Tominaga, N. Tsugi, and M. Tsuda. 1998. A novel photointermediate of octopus rhodopsin activates its G-protein. *FEBS Lett.* 436:259–262.
- Nishimura, S., H. Kandori, M. Nakagawa, M. Tsuda, and A. Maeda. 1997. Structural dynamics of water and the peptide backbone around the Schiff base associated with the light-activated process of octopus rhodopsin. *Biochemistry*. 36:864–870.
- Ohtani, H., M. Kobayashi, M. Tsuda, and T. G. Ebrey. 1988. Primary processes in photolysis of octopus rhodopsin. *Biophys. J.* 53:17–24.
- Pande, C., A. Pande, K. T. Yue, R. H. Callender, T. G. Ebrey, and M. Tsuda. 1987. Resonance Raman spectroscopy of octopus rhodopsin and its photoproducts. *Biochemistry*. 26:4941–4947.
- Richard, L., L. Genberg, J. Deak, H. Chiu, and R. J. D. Miller. 1992. Picosecond phase grating spectroscopy of hemoglobin and myoglobin: energetics and dynamics of global protein motion. *Biochemistry*. 31:10703–10715.
- Stryer, L. 1986. Cyclic GMP cascade of vision. *Annu. Rev. Neurosci.* 9:87–119.
- Taiji, M., K. Bryl, M. Nakagawa, M. Tsuda, and T. Kobayashi. 1992. Femtosecond studies of primary photoproduct in octopus rhodopsin. *Photochem. Photobiol.* 56:1003–1011.
- Takeshita, K., N. Hirota, Y. Imamoto, M. Kataoka, F. Tokunaga, and M. Terazima. 2000. Temperature dependent volume change of initial step of the photoreaction of photoactive yellow protein (PYP) studied by the transient grating. *J. Am. Chem. Soc.* 122:8524–8528.
- Terazima, M., and N. Hirota. 1993. Translational diffusion of a transient radical studied by the transient grating method, pyrazinyl radical in 2-propanol. *J. Chem. Phys.* 98:6257–6262.
- Tsuda, M. 1979. Transient spectra of intermediates in the photolytic sequence of octopus rhodopsin. *Biochim. Biophys. Acta.* 545:537–546.
- Tsuda, M. 1987. Photoreception and phototransduction in invertebrate photoreceptor. *Photochem. Photobiol.* 45:915–931.
- Tsuda, M., T. Tsuda, Y. Terayama, Y. Fukada, T. Akino, G. Yamanaka, L. Stryer, T. Katada, M. Ui, and T. G. Ebrey. 1986. Kinship of cephalopod photoreceptor G protein with vertebrate transducin. *FEBS Lett.* 198:5–10.
- van Brederode, M. E., T. Gensch, W. D. Hoff, K. J. Hellingwerf, and S. E. Braslavsky. 1995. Photoinduced volume change and energy storage associated with the early transformations of the photoactive yellow protein from *ectothiorhodospira halophila*. *Biophys. J.* 68:1101–1109.