

Molecular Dynamics Simulations of Lipid Bilayers: Major Artifacts Due to Truncating Electrostatic Interactions

M. Patra,* M. Karttunen,* M. T. Hyvönen,^{†‡} E. Falck,[‡] P. Lindqvist,[‡] and I. Vattulainen[‡]

*Biophysics and Statistical Mechanics Group, Laboratory of Computational Engineering, Helsinki University of Technology, FIN-02015 HUT, Finland; [†]Wihuri Research Institute, FIN-00140 Helsinki, Finland; and [‡]Laboratory of Physics and Helsinki Institute of Physics, Helsinki University of Technology, FIN-02015 HUT, Finland

ABSTRACT We study the influence of truncating the electrostatic interactions in a fully hydrated pure dipalmitoylphosphatidylcholine (DPPC) bilayer through 20 ns molecular dynamics simulations. The computations in which the electrostatic interactions were truncated are compared to similar simulations using the particle-mesh Ewald (PME) technique. All examined truncation distances (1.8–2.5 nm) lead to major effects on the bilayer properties, such as enhanced order of acyl chains together with decreased areas per lipid. The results obtained using PME, on the other hand, are consistent with experiments. These artifacts are interpreted in terms of radial distribution functions $g(r)$ of molecules and molecular groups in the bilayer plane. Pronounced maxima or minima in $g(r)$ appear exactly at the cutoff distance indicating that the truncation gives rise to artificial ordering between the polar phosphatidyl and choline groups of the DPPC molecules. In systems described using PME, such artificial ordering is not present.

INTRODUCTION

One of the great challenges in biophysics is to understand the basic principles that govern lipid bilayer mixtures. Lipid bilayers, or membranes, govern and mediate various biologically relevant processes on the cellular level. Transfer of ions through membranes and the function of enzymes attached to membranes provide two examples of these situations. Besides cellular membranes, lipid membranes are present in various man-made applications such as liposomes used in novel drug delivery techniques (Langer, 1998) and in many natural entities such as lipoproteins. The variety of situations where lipid bilayers play an important role is truly fascinating and is discussed in a number of review articles (Bloom et al., 1991; Lipowsky and Sackmann, 1995; Merz and Roux, 1996; Nagle and Tristram-Nagle, 2000; Katsaras and Gutberlet, 2001).

The characteristics of membranes have been extensively investigated for many decades, and experiments have provided substantial information about the intriguing physicochemical aspects of membrane systems (Bloom et al., 1991; Lipowsky and Sackmann, 1995; Merz and Roux, 1996; Nagle and Tristram-Nagle, 2000; Katsaras and Gutberlet, 2001). However, although the experimental approach is the cornerstone of membrane research, it is often difficult or even impossible to obtain a thorough understanding of the phenomena taking place in lipid bilayers by experiments only. Therefore, atomistic computer simulation techniques such as classical molecular dynamics (MD) have become a standard tool for studies of biomembrane systems

at the molecular level (Merz and Roux, 1996; Tieleman et al., 1997; Feller, 2000; Saiz and Klein, 2002).

One drawback of the computational approach is that its success depends on various methodological issues such as force fields, constraints, and the accuracy of integration schemes for the equations of motion (Tieleman et al., 1997; van Gunsteren and Mark, 1998; Chiu et al., 2000; Besold et al., 2000; Vattulainen et al., 2002). In particular, the treatment of electrostatic interactions deserves special attention, since biomembrane systems are highly charged: lipid molecules are either polar or charged and they interact with each other, the polar water environment, counterions (Pandit and Berkowitz, 2002), proteins (Ibragimova and Wade, 1998), and DNA (Bandyopadhyay et al., 1999). Proper treatment of electrostatic interactions in MD simulations is therefore one of the most important issues in this field and it continues to pose significant challenges for computer simulations.

The calculation of electrostatic interactions is typically based on solving the Poisson equation for the electrostatic potential such that all charged particles and their periodic images are taken into account in some systematic fashion. The Ewald summation method, its variants (Sagui and Darden, 1999), and the fast multipole method (Greengard and Rokhlin, 1987; Frenkel and Smit, 2002) are commonly used techniques that exploit this idea. In particular, the particle-mesh Ewald (PME) technique has been used increasingly often in lipid bilayer simulations (Venable et al., 2000; Saiz and Klein, 2001; Feller et al., 2001; Tobias, 2001; Pandit and Berkowitz, 2002).

Alternatively, one can neglect the long-range Coulombic tail and truncate the interactions at some suitable distance, a typical choice being 1.5–2.0 nm. This technique leads to considerable savings in the computational load and hence is widely used. Due to the speedup, it is particularly useful in studies of large systems over long times (Lindahl and Edholm, 2000; Marrink et al., 2001; Chiu et al., 2002), and

Submitted November 27, 2002, and accepted for publication February 21, 2003.

Address reprint requests to Ilpo Vattulainen, E-mail: Ilpo.Vattulainen@csc.fi.

© 2003 by the Biophysical Society

0006-3495/03/06/3636/10 \$2.00

when the computational requirements are demanding due to, e.g. long timescales associated with complex processes such as membrane fusion (Marrink and Tieleman, 2002). The discontinuities in the potential and forces at the cutoff radius are typically not considered to be a major issue, since they can be handled using various shifting and switching techniques (Steinbach and Brooks, 1994; Leach, 2001).

One might expect the artifacts due to truncation, if any, to become smaller as the cutoff distance r_{cut} is increased, and that for reasonably large cutoffs the system should not be influenced by truncation. In practice, however, this is not the case. The classical example is water: its bulk properties (Alper and Levy, 1989; Feller et al., 1996) and properties at the surfaces of lipid monolayers have been found to be affected by truncation (Alper et al., 1993a,b; Feller et al., 1996). Other cases where direct effects due to truncation have been observed include peptides, proteins, and DNA (Smith and Pettitt, 1991; Schreiber and Steinhäuser, 1992; York et al., 1993, 1995; Ibragimova and Wade, 1998; Norberg and Nilsson, 2000).

Given these findings, it is rather surprising that only Venable et al. have considered the effects of truncation in the context of lipid bilayers (Venable et al., 2000). They compared the areas per molecule in a DPPC bilayer in the gel phase using systems in which the electrostatic interactions had been treated using PME and a truncation of 1.2 nm. They found the results to differ by $\sim 4\%$. To the best of our knowledge, further systematic studies of truncation effects in lipid bilayers have not been reported. The lack of information is particularly striking in the case of the liquid-crystalline (L_α) phase, which is highly relevant from a physiological point of view. Instead, the general impression seems to be that truncation may lead to artifacts, but they are minor, or even negligible, if the cutoff is longer than ~ 1.8 nm (Alper et al., 1993a,b; Feller et al., 1996; Jakobsson et al., 1996).

In this article, we show through an extensive set of 20-ns MD simulations for a fully hydrated pure lipid bilayer of 128 dipalmitoylphosphatidylcholine (DPPC) molecules in the liquid-crystalline phase that the truncation of electrostatic interactions can have significant consequences on the properties of lipid bilayer systems. We consider several truncation distances from 1.8 to 2.5 nm and compare them to a case where the particle-mesh Ewald technique has been applied. We find that the simulations where PME has been used lead to an area per lipid molecule consistent with experiments, whereas the truncation of electrostatic interactions leads to 5–14% smaller values. This dramatic result is reflected in various properties of the lipid bilayer, including the probability distribution of the area per lipid, the density profile across the membrane, and the ordering of acyl chains. In addition to these, truncation leads to prominent artifacts in the electrostatic potential across the bilayer. We interpret the artifacts in terms of radial distribution functions $g(r)$ of molecules and molecular groups in the plane of the

membrane. The radial distribution functions reveal without doubt that truncation leads to artificial ordering in the head-groups of lipid molecules.

We conclude that the truncation of electrostatic interactions may lead to profound artifacts in the properties of lipid bilayer systems, and should be used with great care, if at all.

SYSTEM

Model and simulation details

We have simulated a lipid bilayer system consisting of 128 DPPC molecules (shown schematically in Fig. 1), fully hydrated by 3655 water molecules. The united atom model description used in this work has been validated previously (Tieleman and Berendsen, 1996).

We used a system available at <http://moose.bio.ucalgary.ca/files/dppc128.pdb> as our initial configuration. This system corresponds to the final structure of run E discussed elsewhere (Tieleman and Berendsen, 1996). The bilayer is in the x - y plane.

The parameters for bonded and nonbonded interactions were taken from a rather recent study on a DPPC bilayer system (Berger et al., 1997), available in electronic form at <http://moose.bio.ucalgary.ca/files/lipid.itp>. The partial charges were obtained from the underlying model description (Tieleman and Berendsen, 1996) and can be found at <http://moose.bio.ucalgary.ca/files/dppc.itp>. For water, the SPC model (Berendsen et al., 1981) was used.

Lennard-Jones interactions were cut off at 1.0 nm without shift or switch functions. Electrostatic interactions within 1.0 nm were calculated at each time step, whereas interactions beyond this range were determined every 10 time steps. These choices follow the parameterization of DPPC (Tieleman and Berendsen, 1996). This choice further corresponds to the scheme referred to as twin-range cutoff, to distinguish it from the multiple time step schemes. They both have certain practical advantages, and there are cases where the multiple time step would probably be more useful. However, since basically all DPPC simulations reported so far have used a twin-range scheme and our objective was to use an approach that is common on this field, we decided to follow the same idea.

Long-range electrostatics was handled either by using a cutoff at $r_{\text{cut}} = 1.8$ nm, 2.0 nm, or 2.5 nm, or by means of the particle-mesh Ewald (Essman et al., 1995b) method to




FIGURE 1 Representation of a DPPC molecule showing the numbering of carbons in the acyl chains discussed in the text.

RESULTS AND DISCUSSION

System dimensions

One of the central quantities in describing lipid bilayers is the average area per molecule $\langle A \rangle$. For DPPC it has been experimentally determined to be $\langle A \rangle = 0.64 \text{ nm}^2$ (Nagle and Tristram-Nagle, 2000). It provides a measure for tuning the force fields and other parameters for lipid systems as one aims for a quantitative description of lipid bilayers through MD simulations. In the present work, the average area per lipid was computed using the size of the simulation box in the x - y plane. Since we employ pressure coupling, the simulation box is allowed to fluctuate during the simulation. The temporal behavior of the area per lipid $A(t)$ is shown in Fig. 2. The simulations have equilibrated after 10 ns and for the analysis we discarded the first 10 ns, and used the second 10-ns period only.

Fig. 2 shows that $A(t)$ depends strongly on the treatment of electrostatic interactions. The simulations using PME yielded $\langle A \rangle = (0.645 \pm 0.010) \text{ nm}^2$ consistent with recent experiments (Nagle and Tristram-Nagle, 2000). Truncation at both 2.0 nm and 2.5 nm lead to $\langle A \rangle = (0.615 \pm 0.010) \text{ nm}^2$, which deviates $\sim 5\%$ from the PME result. Further decrease of the cutoff distance to 1.8 nm leads to $\langle A \rangle = (0.555 \pm 0.010) \text{ nm}^2$. This is $\sim 14\%$ smaller than the reference value of 0.645 nm^2 .

We verified that the large differences in the average area per lipid are indeed due to electrostatics and not a consequence of initial conditions. To this end, we chose an equilibrated configuration from the simulation with $r_{\text{cut}} = 1.8 \text{ nm}$ (at 10 ns) as the initial configuration for a new simulation. In this new simulation, the electrostatic interactions were computed using PME instead of a cutoff. This transition point is marked by an arrow in Fig. 2. As seen from Fig. 2, the area per lipid quickly approaches the value of 0.645 nm^2 . We can thus conclude that the different areas per lipid




FIGURE 2 The area per lipid $A(t)$ as a function of time for truncated and PME electrostatics. Cutoff radii are shown in the plot. In addition, $A(t)$ is shown for the case where the electrostatics was switched from a cutoff at 1.8 nm to PME at 10 ns (marked by an arrow).

reported here are solely due to the treatment of electrostatic interactions.

For the purpose of comparison, let us note that our results for the average area per lipid at short times are in agreement with earlier studies by Tieleman et al. (Tieleman and Berendsen, 1996), whose model description we follow in the present work. They used a cutoff at 2.0 nm and found $\langle A \rangle \approx 0.60 \text{ nm}^2$ for the area per molecule. A full comparison is not meaningful, however, since the timescale in their studies was 0.5 ns and the analysis was done over the last 100 ps only.

We have extended the above studies by considering the probability distributions for the area per lipid molecule $P(A)$. This quantity is of interest for a number of processes in lipid bilayers, e.g. the lateral diffusion of lipids in the bilayer plane. Results for $P(A)$ are shown in Fig. 3. The distributions reveal that the minimum area per lipid is $\sim 0.3 \text{ nm}^2$. Further, we find that the shapes of the distributions are similar and scaled by the average area per molecule. However, it is worth pointing out that even if the cutoff distance is increased to a value close to the maximal one (i.e., half the linear dimension of the system), the artifacts in $P(A)$ persist. This indicates that cutoff distances even as large as 2.5 nm are not sufficient for a proper quantitative treatment of electrostatics.

Next we estimate the density distribution and the dimensions along the bilayer normal. The mass density profile for water and lipid molecules is shown in Fig. 4. As before, we find that the smallest cutoff leads to major artifacts as compared to the PME description, whereas the other two truncation distances perform better. The results obtained using any truncation distance are distinctly different from those obtained using PME, however. To demonstrate this, we consider the thickness of the bilayer defined as the point where the densities of water and lipids are equal. The half thicknesses of the bilayers were found to be 2.03 nm for PME, 2.15 nm for $r_{\text{cut}} = 2.0 \text{ nm}$ and 2.5 nm, and 2.33 nm for $r_{\text{cut}} = 1.8 \text{ nm}$. It is noteworthy that the deviations in these results are of similar size as the deviations found in the average area per molecule. However, although truncation




FIGURE 3 Distribution of the area per lipid, $P(A)$, computed by the Voronoi analysis.




FIGURE 4 Mass density profile across the bilayer computed separately for lipids (*left*) and water (*right*). $z = 0$ corresponds to the center of the bilayer and both sides of the bilayer have been folded toward positive z .

leads to a decreased area per lipid, it also leads to an increased height of the membrane. Thus these two artifacts compensate each other, and one finds that the average volume per lipid obtained by truncation methods is consistent with the value found by PME. This result was confirmed by the Voronoi analysis for the lipid bilayers in three dimensions (data not shown).

It should be noted that the density of the bulk water phase is essentially independent of the simulation parameters. This is in contrast to the density of the lipid phase. These observations confirm that the differences between the simulations are caused by the electrostatic interactions between the lipids and/or lipids and water, but not by the interactions among the water molecules.

Order parameters

We have computed S_{CD} for all carbon atoms in both chains (*sn-1* and *sn-2*) by averaging over all equivalent atoms in all DPPC molecules. The results are shown in Fig. 5. We find that PME yields an order parameter profile which is in good agreement with experimental data (Brown et al., 1979; Douliez et al., 1995; Petrache et al., 2000). Note that $S_{CD} \approx 0.20$ close to the glycerol group, and tends toward zero toward the end of a tail. The acyl chains are therefore reasonably ordered close to the headgroup, whereas conformational disorder becomes more and more apparent toward the center of the bilayer.

The results with different cutoff distances differ significantly from those obtained using PME. The situation close to the glycerol group clearly demonstrates the problem. The truncation of electrostatic forces at $r_{cut} = 2.0$ nm and 2.5 nm yields order parameters that deviate $\sim 13\%$ from the values obtained using PME. In the case of $r_{cut} = 1.8$ nm, the deviation is even larger, being of the order of 40% close to the glycerol group. Differences are expected since the ordering of acyl chains must be affected by the packing of lipids as discussed in the previous section. A reduction in the




FIGURE 5 The order parameter $|S_{CD}|$ separately for the two acyl chains of DPPC: *sn-1* chain and *sn-2* chain. For numbering of carbons, see Fig. 1.

area per molecule leads to an enhanced ordering of the acyl chains and correlates with an increased thickness of the bilayer.

We have also considered the ordering of water in the vicinity of the bilayer-water interface by calculating the time averaged projection of the water-dipole unit vector onto the interfacial normal. As revealed by Fig. 6, the water molecules prefer to order themselves in such a way that the dipoles are oriented toward the bilayer. Ordering persists up to the height where the density of the lipids approaches zero.

This ordering can be explained by the orientational behavior of water molecules around phosphoryl groups. The radial distribution of the oxygens and hydrogens of the water molecules around the phosphatidyl and choline groups were determined (data not shown), revealing that the water molecules are strongly oriented around the phosphorus atom. This is due to the hydrogen bonding between the oxygen atoms of the phosphatidyl group and the hydrogen atoms of the water molecules. Only a weak orientation is found around the nitrogen atom, most likely due to the hydrophobic nature of the surrounding methyl groups, which results in hydrogen bonding among the water molecules. This behavior is very similar for the different treatments of electrostatic interactions and also to earlier MD simulation studies of phosphatidylcholine systems (Alper et al., 1993a; Damodaran and Merz, 1994; Essman et al., 1995a; Hyvönen et al., 1997). As the




FIGURE 6 $P(z) = \langle \cos \theta \rangle$ describing the ordering of water close to the bilayer-water interface. $z = 0$ corresponds to the center of the bilayer. The noise at small z is due to the small number of water molecules inside the bilayer.

number of hydrogen-bonded water molecules around the phosphatidyl group is larger on the side of the water region, the dominating orientation of the water dipoles is toward the bilayer center. Only a few water molecules penetrate into the hydrocarbon region, resulting in poor statistics and a spiky profile.

To conclude this section, our results indicate that the ordering of fatty acyl chains is strongly affected by the method by which electrostatic interactions are treated. In addition, we find that the use of a relatively large cutoff in electrostatic interactions does not give rise to major artifacts in the properties of water molecules themselves or their radial distribution around the headgroups. At the surface of the bilayer, however, packing of lipids affects the properties of the interfacial water layer. These conclusions support the view of previous research on the properties of water close to a water-lipid monolayer interface (Alper et al., 1993a,b; Feller et al., 1996). In these studies it was found that the artifacts were reduced by an increase in the cutoff, but were not eliminated for cutoff distances as large as $r_{\text{cut}} = 1.8$ nm. The work by Feller et al. provides a particularly interesting example of this issue, since they considered the radial distribution of oxygen-oxygen pairs in bulk water and found minor peaks close to the cutoff distance (Feller et al., 1996). This is consistent with our results for RDFs close to the water-bilayer interface. We studied the radial distribution functions for O-N and O-P pairs (where O stands for oxygen in water) and found very weak but systematic ordering effects at r_{cut} . In RDFs found by PME, such ordering effects were not present.

Electrostatic potential

Based on the results discussed above, it seems obvious that the truncation of Coulombic interactions plays an important role in the electrostatic properties of the bilayer. To verify this and to quantify the magnitude of possible artifacts due to

truncation, we studied the electrostatic potential $V(z)$. The results are shown in Fig. 7.

The general behavior agrees with previous simulation studies of PC bilayers (Tieleman and Berendsen, 1996; Shinoda et al., 1998b; Smondyrev and Berkowitz, 1999, 2000). The lipid molecules contribute with a large positive potential, which is compensated by the contribution due to the water molecules. The total potential was determined to be -570 mV. For comparison, the experimental values for the potential range from -200 mV to -575 mV for different PC/water interfaces (Flewelling and Hubbel, 1986; Simon and McIntosh, 1989; Gawrisch et al., 1992; McIntosh et al., 1992).

As what comes to the differences between the curves in Fig. 7, we note that the profiles of $V(z)$ are correlated with the mass density profiles in Fig. 4. The differences in the packing of lipids are also reflected in the electrostatic potential: The strongest orientation of water molecules is found just above the peak in the density of lipids, and the orientation of the water ranges just as far as the headgroups of the lipids prevail.

Radial distribution of lipids

We first consider RDFs between the two charged groups in a DPPC molecule. The positively charged choline group is at the top and the negatively charged phosphate group at the lower part of the headgroup (see Fig. 1). However, the average orientation of the P-N vector is almost parallel to the plane of the bilayer (data not shown). The details of the calculation are described in the section for data analysis.

The RDFs for the two pairs of P and N atoms in the headgroup are shown in Fig. 8. The RDF of N-N pairs serves as a good example of our findings. The application of PME yields a radial distribution function which has a hard core at small distances, a rather narrow peak around 0.8 nm, and essentially no structure beyond $r = 1.0$ nm. This behavior is expected since we are dealing with a liquid-crystalline phase in the absence of translational order in the bilayer plane. The




FIGURE 7 Total electrostatic potential $V(z)$ across the bilayer, where $z = 0$ corresponds to the center of the bilayer.




FIGURE 8 Radial distribution function (RDF) $g(r)$ between the two central atoms in the headgroup of a DPPC molecule: RDFs for N-N and P-P pairs. Cutoff distances are indicated by arrows.

RDFs from the simulations in which a cutoff was used are dramatically different. In all of these cases we find that there is a wildly oscillating long-range component which has a local maximum exactly at the cutoff distance. In addition to this, the oscillations persist for distances far beyond r_{cut} . Although the details are slightly different for P-P (as well as N-P) pairs, similar conclusions on artificial ordering can be drawn.

These structural artifacts have a very strong character. This is evident from the RDF for the DPPC center of mass positions presented in Fig. 9. The $g(r)$ given by the PME method is consistent with the assumption of a fluid-like phase, having just a small peak around 1.2 nm. The truncation methods, on the other hand, give rise to further structure manifested as artificial maxima precisely at the cutoff distance.

Results of similar nature have been reported for ionic systems (van Gunsteren and Mark, 1998). For an aqueous NaCl solution van Gunsteren and Mark found that the truncation of Coulombic interactions gave rise to an artificial peak around the cutoff distance. In our work the DPPC molecules are polar, though the behavior of the bilayer is not expected to be solely dictated by polar groups. In this sense, the pronounced artifacts in Fig. 9 for lipid molecules may come as a surprise.




FIGURE 9 Radial distribution function $g_{2d}(r)$ for the center of mass positions of the DPPC molecules. Note the relatively soft core of $g(r)$ in the vicinity of $r = 0$. This is due to the fact that the lipids may be entangled around each other such that the x,y coordinates of their center of mass positions may lie close to each other.

We can conclude that the truncation of electrostatic interactions gives rise to artificial order in the plane of the membrane. Truncation changes the phase behavior of the bilayers, and consequently affects thermodynamic properties such as the compressibility of the bilayer. It is clear that this is a matter of serious concern. It suggests that the truncation of Coulombic interactions in lipid bilayer systems may not only influence the short-range order of the system, but also the long-range behavior. Various intriguing phenomena such as organization of bilayer-protein systems involve scales of several molecular diameters. The artificial ordering observed in this work persists over these distances, and care should be taken when treating these systems.

SUMMARY AND CONCLUSIONS

We have investigated the effects of truncating Coulombic interactions on the properties of a DPPC bilayer and water in the vicinity of a bilayer-water interface. We have found that the truncation of electrostatic interactions can have serious consequences for the structural and electrostatic properties of lipid bilayer systems. The reduced area per molecule and the corresponding changes in the ordering of acyl chains provide two examples of the problems associated with truncation. Preliminary studies of dynamic quantities such as the lateral diffusion coefficient have shown that they can also be sensitive to truncation. However, a thorough analysis of this issue requires simulations over much longer timescales (Patra et al., unpublished results). As shown in this work, these artifacts are due to the fact that truncation gives rise to artificial ordering in the bilayer plane, which in turn implies that the lipid bilayer is no longer in a truly fluid-like state. Instead, the artificial order persists over a long range, which for typical system sizes studied by molecular dynamics simulations may exceed the actual size of the system. Furthermore, since the artificial order due to truncation affects the

phase behavior of the system, we may conclude that there is indeed a reason for major concern.

The particle-mesh Ewald (PME) technique, on the other hand, has performed very well in this study. Besides providing results consistent with experimental data, it has given no reason for concern with respect to pair correlation behavior in the plane of the bilayer. The current trend of using PME in MD simulations of lipid membrane systems seems to be justified and should be encouraged.

We would like to stress, however, that even PME and its variants may lead to artifacts unless great care is taken. These artifacts are related to the periodicity of the system, as periodic boundary conditions are used to eliminate finite size effects in simulations of small systems. Hünenberger et al. have observed (Hünenberger and McCammon, 1999; Weber et al., 2000) that the artificial periodicity used in Ewald techniques may indeed affect the conformational equilibria of e.g. peptides and proteins by stabilizing the most compact conformations of the molecules. Despite the fact that a substantial amount of work has already been done (Sagui and Darden, 1999; Martyna and Tuckerman, 1999; Zhou et al., 2001; Mináry et al., 2002; Karttunen et al., 2002), it is clear that more attention is called for to develop more reliable and efficient techniques for dealing with electrostatic interactions in simulations of biomolecular systems.

APPENDIX A: EFFECT OF OTHER SIMULATION PARAMETERS

We have complemented the main body of research by further simulations to check the importance of other simulation parameters in the present work. For example, we have checked the importance of constraints by keeping the DPPC bonds flexible, investigated the role of pressure coupling by a study in which the time constant of the barostat was set to a value of 5 ps, and repeated all the simulations at a temperature of $T = 325$ K. In all cases we have found that both the quantitative results and the conclusions remain the same. However, changing the cutoff of the van der Waals interaction r_{vdW} appears to have effects worth noticing. Although the parameterization of




FIGURE 10 Radial distribution function for N-N pairs in the headgroups of DPPC molecules based on simulations with a van der Waals cutoff $r_{vdW} = 1.4$ nm (cf. Fig. 8 with $r_{vdW} = 1.0$ nm). Cutoff distances are indicated by arrows.

DPPC was done with a cutoff of $r_{vdW} = 1.0$ nm, and was therefore used in the major part of this study, the value of 1.4 nm is also a common choice. This is motivated in particular by the Gromos96 force field which assumes $r_{vdW} = 1.4$ nm.

Simulations similar to those with $r_{vdW} = 1.0$ nm have been run with $r_{vdW} = 1.4$ nm. The average area per lipid using PME is reduced by 0.051 nm^2 , whereas the average areas per lipid for the three different cutoffs— $r_{cut} = 1.8$ nm, 2.0 nm, or 2.5 nm—are reduced by 0.017, 0.046, and 0.067 nm^2 , respectively. This trend is understandable since an increase in r_{vdW} effectively increases the attractive interaction between acyl chains, thus reducing the area per molecule. Interestingly, also the average volume per lipid is reduced by about 0.03 nm^3 , as was revealed by the Voronoi analysis of lipid bilayers in three dimensions. Despite these quantitative differences between the different cutoff distances for the van der Waals interactions, the conclusions of our work remain intact. This is demonstrated in Fig. 10, which presents data for the RDFs between the N-N pairs with a cutoff $r_{vdW} = 1.4$ nm. The data indicate that the truncation of electrostatic interactions still gives rise to artificial ordering, which is not observed in the case of PME.

This work has, in part, been supported by the Academy of Finland through its Center of Excellence Program (E.F., P.L., and I.V.), the National Graduate School in Materials Physics (E.F.), the Academy of Finland Grant Nos. 54113 (M.K.) and 80246 (I.V.), the Jenny and Antti Wihuri Foundation (M.H.), and the Federation of Finnish Insurance Companies (M.H.).

REFERENCES

- Alper, H. E., D. Bassolino, and T. R. Stouch. 1993a. Computer simulation of a phospholipid monolayer-water system: The influence of long range forces on water structure and dynamics. *J. Chem. Phys.* 98:9798–9807.
- Alper, H. E., D. Bassolino-Klimas, and T. R. Stouch. 1993b. The limiting behavior of water hydrating a phospholipid monolayer: A computer simulation study. *J. Chem. Phys.* 99:5547–5559.
- Alper, H. E., and R. M. Levy. 1989. Computer simulations of the dielectric properties of water: Studies of the simple point charge and transferable intermolecular potential models. *J. Chem. Phys.* 91:1242–1251.
- Bandyopadhyay, S., M. Tarek, and M. L. Klein. 1999. Molecular dynamics study of a lipid-DNA complex. *J. Phys. Chem. B.* 103:10075–10080.
- Berendsen, H. J. C., J. P. M. Postma, W. F. van Gunsteren, A. DiNola, and J. R. Haak. 1984. Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* 81:3684–3690.
- Berendsen, H. J. C., J. P. M. Postma, W. F. van Gunsteren, and J. Hermans. 1981. Interaction models for water in relation to protein hydration. In *Intermolecular Forces*. B. Pullman, editor, Reidel, Dordrecht, The Netherlands. 331–342.
- Berger, O., O. Edholm, and F. Jahnig. 1997. Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophys. J.* 72: 2002–2013.
- Besold, G., I. Vattulainen, M. Karttunen, and J. M. Polson. 2000. Towards better integrators for dissipative particle dynamics simulations. *Phys. Rev. E.* 62:R7611–R7614.
- Bloom, M., E. Evans, and O. G. Mouritsen. 1991. Physical properties of the fluid lipid-bilayer component of cell membranes: A perspective. *Q. Rev. Biophys.* 24:293–397.
- Brown, M. F., J. Seelig, and U. Häberlen. 1979. Structural dynamics in phospholipid bilayers from deuterium spin-lattice relaxation time measurements. *J. Chem. Phys.* 70:5045–5053.
- Chiu, S.-W., M. Clark, S. Subramaniam, and E. Jacobsson. 2000. Collective motion artifacts arising in long-duration molecular dynamics simulations. *J. Comput. Chem.* 21:121–131.

- Chiu, S. W., E. Jacobsson, R. J. Mashl, and H. L. Scott. 2002. Cholesterol-induced modifications in lipid bilayers: A simulation study. *Biophys. J.* 83:1842–1853.
- Damodaran, K. V., and J. K. M. Merz. 1994. A comparison of DMPC- and DLPE-based lipid bilayers. *Biophys. J.* 66:1076–1087.
- Douliez, J.-P., A. Leonard, and E. J. Dufourc. 1995. Restatement of order parameters in biomembranes: Calculation of C–C bond order parameters from C-D quadrupolar splittings. *Biophys. J.* 68:1727–1739.
- Essman, U., L. Perera, and M. L. Berkowitz. 1995a. The origin of the hydration interaction of lipid bilayers from MD simulation of dipalmitoylphosphatidylcholine membranes in gel and liquid crystalline phases. *Langmuir*. 11:4519–4531.
- Essman, U., L. Perera, M. L. Berkowitz, H. L. T. Darden, and L. G. Pedersen. 1995b. A smooth particle mesh Ewald method. *J. Chem. Phys.* 103:8577–8592.
- Feller, S. E. 2000. Molecular dynamics simulations of lipid bilayers. *Curr. Opin. Colloid Interface Sci.* 5:217–223.
- Feller, S. E., K. Gawrisch, and A. D. MacKerell, Jr. 2001. Polyunsaturated fatty acids in lipid bilayers: Intrinsic and environmental contributions to their unique physical properties. *J. Am. Chem. Soc.* 124:318–326.
- Feller, S. E., R. W. Pastor, A. Rojnuckarin, S. Bogusz, and B. R. Brooks. 1996. Effect of electrostatic force truncation on interfacial and transport properties of water. *J. Phys. Chem.* 100:17011–17020.
- Flewelling, R. F., and W. L. Hubbel. 1986. The membrane dipole potential in a total membrane potential model. *Biophys. J.* 49:541–552.
- Frenkel, D., and B. Smit. 2002. *Understanding Molecular Simulation: From Algorithms to Applications*, 2nd edition. Academic Press, San Diego.
- Gawrisch, K., D. Ruston, J. Zimmerberg, V. A. Parsegian, R. P. Rand, and N. Fuller. 1992. Membrane dipole potentials, hydration forces, and the ordering of water at membrane surface. *Biophys. J.* 61:1213–1223.
- Greengard, L., and V. Rokhlin. 1987. A fast algorithm for particle simulation. *J. Comput. Phys.* 73:325–348.
- Hess, B., H. Bekker, H. J. C. Berendsen, and J. G. E. M. Fraaije. 1997. LINCS: A linear constraint solver for molecular simulations. *J. Comput. Chem.* 18:1463–1472.
- Hünenberger, P. H., and J. A. McCammon. 1999. Effect of artificial periodicity in simulations of biomolecules under Ewald boundary conditions: A continuum electrostatics study. *Biophys. Chem.* 78:69–88.
- Hyyönen, M., T. T. Rantala, and M. Ala-Korpela. 1997. Structure and dynamic properties of diunsaturated 1-palmitoyl-2-linoleoyl-sn-glycero-3-phosphatidylcholine lipid bilayer from molecular dynamics simulation. *Biophys. J.* 73:2907–2923.
- Ibragimova, G. T., and R. C. Wade. 1998. Importance of explicit salt ions for protein stability in molecular dynamics simulation. *Biophys. J.* 74:2906–2911.
- Jacobsson, E., S. Subramaniam, and H. L. Scott. 1996. Strategic issues in molecular dynamics simulations of membranes. In *Biological membranes—A Molecular Perspective from Computation And Experiment*, K. M. Merz, Jr. and B. Roux, editors. Dordrecht Birkhäuser, Boston. 105–123.
- Karttunen, M., A. L. Pakkanen, P. K. Kinnunen, and K. Kaski. 2002. Computational modeling of DNA-cationic lipid complexation. *Cell. Mol. Biol. Lett.* 7:238–239.
- Katsaras, J., and T. Gutberlet. 2001. *Lipid bilayers: Structure and Interactions*. Springer-Verlag, Berlin.
- Langer, R. 1998. Drug delivery and targeting. *Nature*. 392(Suppl.):5–10.
- Leach, A. R. 2001. *Molecular Modelling: Principles and Applications*, 2nd edition. Prentice Hall, Harlow.
- Lindahl, E., and O. Edholm. 2000. Mesoscopic undulations and thickness fluctuations in lipid bilayers from molecular dynamics simulations. *Biophys. J.* 79:426–433.
- Lindahl, E., B. Hess, and D. van der Spoel. 2001. GROMACS 3.0: A package for molecular simulation and trajectory analysis. *J. Mol. Model.* 7:306–317.
- Lipowsky, R., and E. Sackmann. 1995. *Structure and dynamics of membranes: From cells to vesicles*. Elsevier, Amsterdam.
- Marrink, S. J., E. Lindahl, O. Edholm, and A. E. Mark. 2001. Simulation of the spontaneous aggregation of phospholipids into bilayers. *J. Am. Chem. Soc.* 123:8638–8639.
- Marrink, S. J., and D. P. Tieleman. 2002. Molecular dynamics simulation of spontaneous membrane fusion during a cubic-hexagonal phase transition. *Biophys. J.* 83:2386–2392.
- Martyna, G. J., and M. E. Tuckerman. 1999. A reciprocal space based method for treating long range interactions in ab initio and force-field-based calculations in clusters. *J. Chem. Phys.* 110:2810–2821.
- McIntosh, T. J., S. A. Simon, D. Needham, and C.-H. Huang. 1992. Inter-bilayer interactions between sphingomyelin and sphingomyelin/cholesterol bilayers. *Biochemistry*. 31:2020–2024.
- Merz, K. M., Jr., and B. Roux. 1996. *Biological membranes: A molecular perspective from computation and experiment*. Birkhäuser, Boston.
- Mináry, P., M. E. Tuckerman, K. A. Pihakari, and G. J. Martyna. 2002. A new reciprocal space based treatment of long range interactions on surfaces. *J. Chem. Phys.* 116:5351–5362.
- Nagle, J. F., and S. Tristram-Nagle. 2000. Structure of lipid bilayers. *Biochim. Biophys. Acta*. 1469:159–195.
- Norberg, J., and L. Nilsson. 2000. On the truncation of long-range electrostatic interactions in DNA. *Biophys. J.* 79:1537–1553.
- Pandit, S. A., and M. L. Berkowitz. 2002. Molecular dynamics simulation of dipalmitoylphosphatidylserine bilayer with Na⁺ counterions. *Biophys. J.* 82:1818–1827.
- Petrache, H. I., S. W. Dodd, and M. F. Brown. 2000. Area per lipid and acyl length distributions in fluid phosphatidylcholines determined by ²H NMR spectroscopy. *Biophys. J.* 79:3172–3192.
- Sagui, C., and T. A. Darden. 1999. Molecular dynamics simulations of biomolecules: Long-range electrostatic effects. *Annu. Rev. Biophys. Biomol. Struct.* 28:155–179.
- Saiz, L., and M. L. Klein. 2001. Structural properties of a highly polyunsaturated lipid bilayer from molecular dynamics simulations. *Biophys. J.* 81:204–216.
- Saiz, L., and M. L. Klein. 2002. Computer simulation studies of model biological membranes. *Acc. Chem. Res.* 35:482–489.
- Schreiber, H., and O. Steinhauser. 1992. Cutoff size does strongly influence molecular dynamics results on solvated polypeptides. *Biochemistry*. 31:5856–5860.
- Shinoda, W., and S. Okazaki. 1998a. A Voronoi analysis of lipid area fluctuation in a bilayer. *J. Chem. Phys.* 109:1517–1521.
- Shinoda, W., M. Shimizu, and S. Okazaki. 1998b. Molecular dynamics study on electrostatic properties of a lipid bilayer: Polarization, electrostatic potential, and the effects on structure and dynamics of water near the interface. *J. Phys. Chem. B*. 102:6647–6654.
- Simon, S. A., and T. J. McIntosh. 1989. Magnitude of the solvation pressure depends on dipole potential. *Proc. Natl. Acad. Sci. USA*. 86:9263–9267.
- Smith, P. E., and B. M. Pettitt. 1991. Peptides in ionic solutions: A comparison of the Ewald and switching function techniques. *J. Chem. Phys.* 95:8430–8441.
- Smondyrev, A. M., and M. L. Berkowitz. 1999. Structure of dipalmitoylphosphatidylcholine/cholesterol bilayer at low and high cholesterol concentrations: Molecular dynamics simulation. *Biophys. J.* 77:2075–2089.
- Smondyrev, A. M., and M. L. Berkowitz. 2000. Molecular dynamics simulation of dipalmitoylphosphatidylcholine membrane with cholesterol sulfate. *Biophys. J.* 78:1672–1680.
- Steinbach, P. J., and B. R. Brooks. 1994. New spherical-cutoff methods for long-range forces in macromolecular simulations. *J. Comput. Chem.* 15:667–683.
- Tieleman, D. P., and H. J. C. Berendsen. 1996. Molecular dynamics simulations of a fully hydrated dipalmitoylphosphatidylcholine bilayer with different macroscopic boundary conditions and parameters. *J. Chem. Phys.* 105:4871–4880.

- Tieleman, D. P., S. J. Marrink, and H. J. C. Berendsen. 1997. A computer perspective of membranes: Molecular dynamics studies of lipid bilayer systems. *Biochim. Biophys. Acta*. 1331:235–270.
- Tobias, D. J. 2001. Electrostatics calculations: Recent methodological advances and applications to membranes. *Curr. Opin. Struct. Biol.* 11:253–261.
- van Gunsteren, W. F., and A. E. Mark. 1998. Validation of molecular dynamics simulation. *J. Chem. Phys.* 108:6109–6116.
- Vattulainen, I., M. Karttunen, G. Besold, and J. M. Polson. 2002. Integration schemes for dissipative particle dynamics: From softly interacting systems towards hybrid models. *J. Chem. Phys.* 116:3967–3979.
- Venable, R. M., B. R. Brooks, and R. W. Pastor. 2000. Molecular dynamics simulations of gel L_{β} phase lipid bilayers in constant pressure and constant surface area ensembles. *J. Chem. Phys.* 112:4822–4832.
- Weber, W., P. H. Hünenberger, and J. A. McCammon. 2000. Molecular dynamics simulations of a polyalanine octapeptide under Ewald boundary conditions: Influence of artificial periodicity on peptide conformation. *J. Phys. Chem. B*. 104:3668–3675.
- York, D. M., T. A. Darden, and L. G. Pedersen. 1993. The effect of long-range electrostatic interactions in simulations of macromolecular crystals: A comparison of the Ewald and truncated list methods. *J. Chem. Phys.* 99:8345–8348.
- York, D. M., W. Yang, H. Lee, T. Darden, and L. G. Pedersen. 1995. Toward the accurate modeling of DNA: The importance of long-range electrostatics. *J. Am. Chem. Soc.* 117:5001–5002.
- Zhou, R., E. Harder, H. Xu, and B. J. Berne. 2001. Efficient multiple time step method for used with Ewald and particle mesh Ewald for large biomolecular systems. *J. Chem. Phys.* 115:2348–2358.