

Increased Concentration of Polyvalent Phospholipids in the Adsorption Domain of a Charged Protein

Emir Haleva,* Nir Ben-Tal,* and Haim Diamant†

*Department of Biochemistry, George S. Wise Faculty of Life Sciences, and †School of Chemistry, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978, Israel

ABSTRACT We studied the adsorption of a charged protein onto an oppositely charged membrane, composed of mobile phospholipids of differing valence, using a statistical-thermodynamical approach. A two-block model was employed, one block corresponding to the protein-affected region on the membrane, referred to as the adsorption domain, and the other to the unaffected remainder of the membrane. We calculated the protein-induced lipid rearrangement in the adsorption domain as arising from the interplay between the electrostatic interactions in the system and the mixing entropy of the lipids. Equating the electrochemical potentials of the lipids in the two blocks yields an expression for the relations among the various lipid fractions in the adsorption domain, indicating a sensitive dependence of lipid fraction on valence. This expression is a result of the two-block picture but does not depend on further details of the protein-membrane interaction. We subsequently calculated the lipid fractions themselves using the Poisson-Boltzmann theory. We examined the dependence of lipid enrichment, i.e., the ratio between the lipid fractions inside and outside the adsorption domain, on various parameters such as ionic strength and lipid valence. Maximum enrichment was found for lipid valence in the range between -3 and -4 in physiological conditions. Our results are in qualitative agreement with recent experimental studies on the interactions between peptides having a domain of basic residues and membranes containing a small fraction of the polyvalent phosphatidylinositol 4,5-bisphosphate (PIP₂). This study provides theoretical support for the suggestion that proteins adsorbed onto membranes through a cluster of basic residues may sequester PIP₂ and other polyvalent lipids.

INTRODUCTION

Some membrane-associated proteins are known to bind to membranes nonspecifically through electrostatic interactions (Murray et al., 1997, 2002; Resh, 1999; McLaughlin et al., 2002). These interactions result from the attraction between a cluster of charged residues in the protein and the oppositely charged membrane lipids. As the charged protein approaches the membrane, it changes the local membrane composition in its vicinity. We refer to this protein-affected region on the membrane as the adsorption domain. The lateral fluidity of the membrane allows oppositely charged lipids to migrate toward the adsorption domain to minimize the interaction free energy. Evidence for such redistribution was reported in experimental (Heimburg et al., 1999; Rauch et al., 2002; Wang et al., 2002) and theoretical (Harries et al., 1998; May et al., 2000, 2002; Fleck et al., 2002) studies. Lipid redistribution was observed also in monolayers (Lee et al., 1994; Lee and McConnell, 1995) and bilayers (Groves et al., 1997, 1998) that were subjected to external electric fields.

Local changes in lipid concentration may have biological significance. For example, PIP₂, a polyvalent phospholipid with valence in the range from -3 to -5 (Toner et al., 1988; McLaughlin et al., 2002; Rauch et al., 2002; Wang et al., 2002) participates in signal transduction (Czech, 2000; Payraastre et al., 2001; Simonsen et al., 2001). Its average

fraction in plasma membranes is very low, $\sim 1\%$, and it is known to be concentrated in specific regions of the membrane (Liu et al., 1998; Stauffer et al., 1998). The PIP₂ lipid serves as a substrate for phospholipase C (PLC), which cleaves it to two secondary messengers (Katan and Williams, 1997). Another component, the myristoylated alanine-rich C kinase substrate protein (MARCKS), containing an amino acid segment of 13 basic and no acidic residues (Blackshear, 1993; McLaughlin and Aderem 1995), is believed to form a PIP₂ “reservoir” in its adsorption domain. As long as PIP₂ is concentrated in the MARCKS adsorption domain, PLC is inhibited and cannot catalyze the PIP₂ hydrolysis (Wang et al., 2001). It is assumed that upon demand, by phosphorylating the MARCKS effector segment, these lipids are freed for signaling (McLaughlin et al., 2002). Thus the capability of MARCKS to sequester PIP₂ potentially affects intracellular signaling.

Lipids of various valences are attracted to the adsorption domain to different extents. This electrostatically induced enrichment is partially balanced by entropy effects that favor homogeneous lipid distribution. Theoretical studies of bilayers composed of neutral and monovalent lipids, where the lipid mobility was taken into account, showed that the formation of a charged lipid domain due to the adsorbed protein is energetically favorable and outweighs entropy effects (e.g., Heimburg et al., 1999; May et al., 2000). Recently, Fleck et al. (2002) presented a detailed formulation for the interactions between charged objects and a membrane composed of lipids of various valences. These studies demonstrate the importance of lipid redistribution in the thermodynamics of protein-membrane adsorption.

Submitted August 13, 2003, and accepted for publication October 17, 2003.

Address reprint requests to Haim Diamant, School of Chemistry, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Ramat Aviv 69978, Israel. Tel.: +972-3-640-6967; Fax: +972-3-640-9293; E-mail: hdiamant@tau.ac.il.

© 2004 by the Biophysical Society

0006-3495/04/04/2165/14 \$2.00

Based on the findings of these theoretical studies, we focus in this work on a simplified model for the redistribution of different-valence lipids in the adsorption domain of a charged protein. We start by deriving an expression that relates the fractions of the various lipids in the adsorption domain to their values in the unperturbed membrane. The general expression is restricted, however, to *relations* between concentrations of different lipids. To obtain the actual concentration values for each lipid type, we use the Poisson-Boltzmann (PB) theory (e.g., Andelman, 1995). The model is then applied to peptide segments, such as a polylysine chain, interacting with a membrane composed of uncharged, monovalent, and trivalent lipids, corresponding to zwitterionic PC, PS, and PIP₂, respectively. Finally, the model is evaluated and the biological implications of its results are discussed.

MODEL

A schematic view of the protein-membrane system under study is provided in Fig. 1. In the model, the membrane is considered to be an infinite surface composed of k phospholipid species. Each phospholipid type i is ascribed a fraction (in the unperturbed membrane) ϕ_i and a fixed valence z_i . (We do not consider pH-dependent dissociation of charged lipids, which was found to have a minor effect compared to lipid mobility; Fleck et al., 2002.) The indices $i = 0$ and $i = 1$ are assigned to neutral lipids ($z_0 = 0$) and monovalent anionic ones ($z_1 = -1$), respectively, which are always present in biological membranes. For simplicity, all phospholipids are ascribed the same headgroup area a . (In a more detailed model, one can treat different headgroup areas; Andelman et al., 1994.)

We assume that the adsorption domain is a “patch” of finite area A and uniform charge density σ_m , whose size is much larger than the Debye screening length of the solution κ^{-1} , i.e., $\kappa^2 A \gg 1$. The screening provided by the surrounding ionic solution ensures a cutoff for the effect of the adsorbed protein on the membrane, justifying the finite-area assumption. The uniform charge density, employed merely for simplicity, can be thought of as an effective or average domain charge density. We thus neglect effects related to charge discreteness. (This assumption becomes invalid in certain circumstances; we shall return to it in the Discussion.) The membrane outside the domain serves as a large reservoir, assumed to be unaffected by the protein. Thus, in this study (similar to the description of May et al., 2002) we regard the membrane as composed of two blocks, the protein-affected adsorption domain, with lipid fractions ψ_i , and the unaffected remainder of

the membrane, with fractions ϕ_i . Both blocks are assumed to contain enough molecules to be considered, to a good approximation, as macroscopic phases. (A treatment of finite-size effects can be found in May et al., 2002.) The charge densities $\bar{\sigma}$ and σ_m , in the protein-free and protein-bound regions, respectively, are defined as:

$$\bar{\sigma} = \frac{e}{a} \sum_i z_i \phi_i, \quad \sigma_m = \frac{e}{a} \sum_i z_i \psi_i, \quad (1)$$

where e is the elementary charge.

The free energy per unit area of the bare (protein-free) membrane, $F^{(0)}$, is

$$F^{(0)} = \frac{T}{a} \sum_i \phi_i \ln \phi_i + F_{es}^{(0)}(\bar{\sigma}). \quad (2)$$

The first term is the mixing entropy contribution, where T is the temperature in energy units (taking the Boltzmann constant as unity). We neglect short-range, nonelectrostatic interactions between lipid molecules (May et al., 2002) except for excluded-volume effects. The second term accounts for the electrostatic contribution. Similarly, the free energy per unit area of the protein-bound domain, F , is

$$F = \frac{T}{a} \sum_i \psi_i \ln \psi_i + F_{es}(\sigma_m), \quad (3)$$

where F_{es} accounts for the electrostatic interactions among the phospholipids and between them and the protein. Note that $F_{es}^{(0)}$ and F_{es} are functions of ϕ_i and ψ_i , respectively, only via the charge densities $\bar{\sigma}$ and σ_m as defined in Eq. 1. These free energies can be calculated using various theories, e.g., the commonly used PB theory (see Appendix 2). However, at this stage of our formulation, we need not specify the expressions for $F_{es}^{(0)}$ and F_{es} at all.

The adsorption domain and the rest of the membrane are at thermodynamic equilibrium, thus the electrochemical potentials μ_i of each lipid type are equal in the two regions. In addition, the membrane incompressibility adds two constraints:

$$\sum_i \phi_i = 1, \quad \sum_i \psi_i = 1. \quad (4)$$

The electrochemical potential of phospholipid i in the protein-free membrane is

$$\mu_i^{(0)} = a \frac{\partial F^{(0)}}{\partial \phi_i} = T \ln \frac{\phi_i}{\phi_0} + z_i e \frac{\partial F_{es}^{(0)}}{\partial \bar{\sigma}}, \quad (5)$$

and similarly, in the adsorption domain,

$$\mu_i = a \frac{\partial F}{\partial \psi_i} = T \ln \frac{\psi_i}{\psi_0} + z_i e \frac{\partial F_{es}}{\partial \sigma_m}. \quad (6)$$

In Eqs. 5 and 6 the dependencies on ϕ_0 and ψ_0 arise from the incompressibility constraint, Eq. 4. They can be viewed as partial surface pressures exerted by the uncharged species ($i = 0$) due to excluded-volume effects. (Their surface pressure is equal to $-T \ln \phi_0$ and $-T \ln \psi_0$, respectively, in the bare membrane and in the adsorption domain.) We still have not specified explicit expressions for $F_{es}^{(0)}$ and F_{es} . As a particular example, one may assume a mean electric potential (as in PB), having a value $\Psi^{(0)}(0)$ at the bare membrane, and then $\partial F_{es}^{(0)} / \partial \sigma_m = \Psi^{(0)}(0)$. If, in addition, we set $\phi_0 \simeq 1$ (low membrane charge), then the familiar expression




FIGURE 1 Schematic view of the protein-membrane system. The protein region interacting with the membrane is modeled as a planar surface of charge density σ_p , hovering parallel to the membrane at a distance h . The unperturbed membrane is composed of various lipids, each having valence z_i and area fraction ϕ_i . The interaction region on the membrane, the adsorption domain, has area A and lipid fractions ψ_i . The whole system is embedded in an ionic solution characterized by a dielectric constant ϵ , Debye screening length κ^{-1} , and temperature T .

for the electrochemical potential is recovered, $\mu_i^{(0)} = T \ln \phi_i + z_i e \Psi^{(0)}(0)$ (and similarly for μ_i). Note, however, that the validity of our formulation is more general than this specific example.

Equating $\mu_i^{(0)} = \mu_i$ we get

$$\frac{e}{T} \left(\frac{\partial F_{es}^{(0)}}{\partial \bar{\sigma}} - \frac{\partial F_{es}}{\partial \sigma_m} \right) = \frac{1}{z_i} \ln \frac{\psi_i \phi_0}{\phi_i \psi_0}. \quad (7)$$

Importantly, the left-hand side of Eq. 7 is independent of i . We can therefore compare the right-hand side of the equation for a certain species i with the same expression for the monovalent species ($i = 1$). This gives a set of equations relating the enrichment ratios for the various lipids,

$$\frac{\psi_i}{\phi_i} = \left(\frac{\psi_1}{\phi_1} \right)^{z_i/z_1} \left(\frac{\phi_0}{\psi_0} \right)^{z_i/z_1 - 1}. \quad (8)$$

We refer to Eq. 8 as the “relative enrichment equation.” It is a set of $(k - 2)$ equations for every $i \neq 0, 1$. The relative enrichment equation is one of the key results of this work. In the limit of low charge density, $\phi_0, \psi_0 \simeq 1$, and assuming a mean electric potential having the values $\Psi^{(0)}(0)$ and $\Psi(0)$ at the unperturbed membrane and the adsorption domain, respectively, Eq. 8 is directly related to a Boltzmann relation,

$$\psi_i \simeq \phi_i \exp[-e z_i (\Psi(0) - \Psi^{(0)}(0))/T]. \quad (9)$$

As derived above, however, the applicability of Eq. 8 is more general; it is more accurate and is valid for highly charged membranes and beyond the mean-field approximation. In particular, Eq. 9 implies that the protein does not perturb the local concentration of the neutral lipid, $\psi_0/\phi_0 \simeq 1$. Yet, as we shall see below, in the biologically relevant case, where the membrane contains a large fraction of monovalent lipid, the neutral lipid is significantly depleted from the adsorption domain. Hence, Eq. 8, rather than Eq. 9, will be used throughout our analysis.

In the absence of protein, the phospholipid composition would not change and both sides of Eq. 8 are trivially equal to 1. Another consequence of Eq. 8 is that perturbation of one lipid fraction necessarily entails perturbation in others. In cases where the charge density in the adsorption domain increases (in absolute value), $|\sigma_m| > |\bar{\sigma}|$, neutral phospholipids will be depleted from the domain to allow the entrance of charged ones, $\phi_0/\psi_0 > 1$. The relative enrichment of lipid i , ψ_i/ϕ_i , is then at least that of the monovalent one raised to the power $|z_i|$.

As an example, let us consider the binding of a basic protein to a membrane containing uncharged zwitterionic lipids (e.g., PC), monovalent lipids (e.g., PS), and trivalent ones (e.g., PIP₂). According to Eq. 8, the polyvalent enrichment ratio will be stronger than that of the monovalent fraction by at least a power of $z_i/z_1 = 3$. This is a strong effect. If the monovalent concentration increases twofold, the trivalent concentration will increase by a factor of $2^3 = 8$. Similarly, a slight decrease in the negative charge of the adsorption domain leads to a significant decrease in polyvalent fraction. This entropy-driven enhancement of high-valence ion concentration is an example of a more general phenomenon manifest, e.g., in the favorable accumulation of dissolved polyvalent ions near charged surfaces or polyelectrolytes (e.g., Rouzina and Bloomfield, 1996). Note again that the relative enrichment equation is independent of the specific expressions for $F_{es}^{(0)}(\bar{\sigma})$ and $F_{es}(\sigma_m)$. It does not depend explicitly on details such as the distance between the protein and membrane, or the protein charge.

Equation 8 provides us only with a relation between the different fractions ψ_i . To calculate the actual values of ψ_i we need to derive explicit expressions for the electrochemical potentials $\mu_i^{(0)}$ and μ_i . To this end, we

must introduce details of the protein. It is treated, for simplicity, as a flat surface of uniform charge density σ_p located at a distance h parallel to the membrane (see Fig. 1). This schematic description may be relevant to proteins that have a flat cluster of basic residues facing the membrane at close proximity, whereas the rest of the charged residues are further away, screened by the ionic solution. We regard the protein-membrane distance h as an external parameter determined by other interactions (e.g., desolvation effects), which are not taken into account in our theory. We further assume that h is small enough ($h \ll \sqrt{A}$), such that the induced adsorption domain on the membrane and the interacting cluster on the protein can be taken to have roughly the same area A .

We apply the commonly used mean-field Poisson-Boltzmann (PB) theory (e.g., Andelman, 1995; Gilson, 1995; Honig and Nicholls, 1995) to calculate the electrochemical potentials $\mu_i^{(0)}$ and μ_i . The applicability of this theory to solutions containing polyvalent ions is questionable (e.g., Netz, 2001). However, in the systems discussed here the polyvalent ions (phospholipids) are restricted to the membrane and their mol fraction is much smaller than that of the monovalent lipids, in accord with biological conditions. The more mobile salt ions in the aqueous solution are assumed to be monovalent. Polyvalent ions thus enter the PB calculation merely as a (minor) contribution to the membrane surface charge. As such they should not induce strong correlations, and the PB theory should be applicable. (An exception, where the polyvalent lipid is the majority charge in the membrane and simple electrostatic models indeed seem to fail, will be presented in the Discussion.)

We derive $\mu_i^{(0)}$ and μ_i using three alternative levels of approximation, all of which are discussed in detail in Appendix 1. In presenting the three methods we wish to demonstrate that even a much simplified approach, involving minimum computation, still gives useful results. The nonlinear problem (NLPB) resulting from the PB theory can only be treated numerically. Subsequently, we examine a further approximation where the PB expressions are linearized. This approximation is valid when the electrostatic potential Ψ is much smaller than T/e everywhere (Andelman, 1995). Although this condition is not fulfilled in the relevant biological systems, the two derivations give similar results for reasons that are discussed in Appendix 1. The LPB approximation allows us to derive analytical expressions for μ_i , yet solutions for the various lipid fractions (i.e., for the equations $\mu_i^{(0)} = \mu_i$) still cannot be obtained in closed form.

The fact that, in most relevant systems, the electrostatic interactions dominate over entropy effects (May et al., 2000) led us to examine one last approximation, in which the derivation is divided into two stages. In the first, we neglect entropy when equating $\mu_i^{(0)} = \mu_i$. This implies that the electrostatic potential is uniform along the membrane (i.e., the membrane behaves as a perfect conductor). The resulting membrane charge density in the adsorption domain is

$$\sigma_m = \frac{\sinh(\kappa h) \bar{\sigma} - \sigma_p}{\cosh(\kappa h)}. \quad (10)$$

In the second stage, this value of σ_m is substituted in Eq. 1. Equations 1, 4, and 8 (the latter incorporating the effect of entropy) thus provide a closed set of k polynomial equations that can be easily solved for the k lipid fractions ψ_i . We refer to this scheme as the simplified linear Poisson-Boltzmann method (SLPB).

RESULTS

To study the effects of protein adsorption on a mixed membrane, we calculated the lipid fractions for several representative conditions. Our aim was to examine the redistribution of different-valence lipids in the adsorption domain of a membrane-adsorbed protein as a function of

several parameters: protein-membrane distance, protein charge, and the valence of the most charged lipid species. Unless otherwise stated, we used physiological values for the Debye length ($\kappa^{-1} = 10 \text{ \AA}$), temperature (300 K), dielectric constant of water ($\epsilon = 80$), and lipid headgroup area ($a = 70 \text{ \AA}^2$). Throughout the text, the notation 69%/30%/1% uncharged/monovalent/polyvalent was used to describe the unperturbed membrane composition (ϕ_i values). For consistency, the results presented in this section were all obtained using the more elaborate NLPB method.

Enrichment as a function of protein-membrane distance

The enrichment ratio, ψ_i/ϕ_i , was calculated for the association of a charged protein with a membrane composed of 69% neutral (zwitterionic), 30% monovalent, and 1% trivalent lipids (in the absence of a protein). This implies an average charge density $\bar{\sigma} = -0.33e$ per lipid headgroup area. We present the enrichment arising from a protein that is slightly more charged than the membrane, $\sigma_p = -1.3\bar{\sigma}$. (These values are typical to protein-membrane interactions, as demonstrated below.) Fig. 2 shows the enrichment ratios as a function of protein-membrane distance.

Far from the membrane ($h \gg \kappa^{-1}$) the protein charge is screened and its effect on the membrane is weak. As the charged protein approaches the membrane, oppositely charged lipids move into the adsorption domain whereas the neutral lipids are depleted from it. The choice of similar charge densities (in absolute value) for the protein and




FIGURE 2 Protein-membrane distance effect. Enrichment ratios (ψ_i/ϕ_i) of uncharged (dotted), monovalent (dashed), and trivalent (solid) lipid fractions in the adsorption domain of a protein as a function of the protein-membrane distance h , scaled by the Debye length κ^{-1} . The unperturbed membrane contains 69%/30%/1% uncharged/monovalent/trivalent lipid fractions. The charge density of the protein is 1.3-fold that of the unperturbed membrane and of opposite sign. The inset shows the actual lipid fractions. Although quite similar in charge density to the membrane, the protein causes a marked change in lipid composition. Note the relatively large increase in the trivalent lipid fraction.

membrane leads to minor changes in the fractions of the abundant (neutral and monovalent) lipids, even at distances smaller than κ^{-1} . Notably, the fraction of trivalent lipids changes by a much larger factor of ~ 3 . This result is a consequence of the exponential dependence of the enrichment on lipid valence as seen in Eq. 8.

The nonmonotonic behavior at small distances, shown in Fig. 2, is a delicate point that deserves further discussion. If the membrane charge density had a fixed value $\sigma_m \neq -\sigma_p$, then, at a sufficiently short distance, the mutual attraction between the surfaces would turn into repulsion (Parsegian and Gingell, 1972). This is caused by the increased concentration of the salt ions, which are bound to remain in the confined volume between the protein and membrane to neutralize the system. In our case, however, the system has the additional freedom to change σ_m . As a result, the electrostatic contribution to the free energy of the protein-membrane interaction decreases monotonously with decreasing distance, i.e., the interaction is purely attractive (cf. Fig. 11). (The effect of charge mobility on the interaction between two membranes has been studied in detail recently; Russ et al., 2003.) As long as the two objects are not too close, it may become favorable to overcharge the membrane and gain attraction energy. This is what happens in the system of Fig. 2 for $\kappa h < 1$. For example, at $\kappa h = 0.5$, we find $|\sigma_m/\sigma_p| \simeq 1.17$. In such a case of overcharging, as the distance is further reduced, the osmotic pressure of the salt ions at short distances causes the membrane to decrease its charge density to lower the energetic penalty of further compression (Fig. 2 in the range $\kappa h \lesssim 0.5$). At contact ($\kappa h = 0$) we have $\sigma_m = -\sigma_p$, such that the system is neutral without mobile ions. Thus, the ability to redistribute the lipids allows the system to avoid high concentration of ions in the solution.

Effect of protein charge

In Fig. 3 we present the enrichment ratios (ψ_i/ϕ_i) of the different lipid species as a function of the protein charge density for a given protein-membrane distance and membrane composition. As expected, when the protein is highly charged, the adsorption domain is strongly enriched with oppositely charged lipids. On the other hand, at low protein charge, depletion of charged lipids is observed. Remarkably, in both cases, the trivalent species exhibits a much stronger effect than the monovalent one. This is again a consequence of the sensitive dependence of the enrichment ratio on lipid valence (Eq. 8). In between the strongly charged and weakly charged limits, there is a value of σ_p for which the membrane is unperturbed (see arrow in Fig. 3). This point does not correspond to $|\sigma_p| = |\bar{\sigma}|$, as might have been expected. As discussed in the previous subsection, for a nonzero protein-membrane distance the membrane may become overcharged. As a result, this special point where $\psi_i/\phi_i = 1$ is obtained for




FIGURE 3 Protein charge effect. Enrichment ratios (ψ_i/ϕ_i) of the uncharged (dotted), monovalent (dashed), and trivalent (solid) lipids in the adsorption domain of a protein as a function of its charge density (in units of elementary charge per 1000 \AA^2). The inset shows the actual lipid fractions. The unperturbed membrane contains 69%/30%/1% uncharged/monovalent/trivalent lipid fractions, corresponding to $\bar{\sigma} = -4.7e$ per 1000 \AA^2 and the protein-membrane distance is $\kappa h = 0.3$. The enrichment in charged lipids increases with protein charge. The arrow indicates the σ_p value for which the membrane is unperturbed.

$|\sigma_p| < |\bar{\sigma}|$. For example, for the parameters of Fig. 3 it occurs at $|\sigma_p/\bar{\sigma}| \simeq 0.57$.

Effect of lipid valence

Equation 8 implies that, the higher the valence $|z_2|$ of the lipid, the stronger its enrichment, ψ_2/ϕ_2 , relative to that of the monovalent species, ψ_1/ϕ_1 . This does not imply that ψ_2/ϕ_2 per se (not relative to ψ_1/ϕ_1) increases monotonously with $|z_2|$. In fact, there is a competition between two opposing effects. The first, which is entropy driven, favors charging of the adsorption domain by high-valence lipids to minimize the perturbation of membrane composition. On the other hand, from simple stoichiometry, only a small concentration of a high-valence lipid is needed to attain a given membrane charge density. The competition should result in a maximum of ψ_2/ϕ_2 at a certain value of valence $z_2 = z_2^*$. This is confirmed in Fig. 4, where we present ψ_2/ϕ_2 as a function of z_2 for a wide range of protein charge densities. (To obtain smoother curves we calculated ψ_2/ϕ_2 also for artificial, noninteger values of z_2 .) For high protein charge, where electrostatic interactions are strong, the entropy effect is negligible and $|z_2^*|$ is small; for unphysically high protein charge $|z_2^*|$ eventually tends to $|z_1| = 1$. At low protein charge, entropy dominates and $|z_2^*|$ increases.

As seen in Fig. 4, $|z_2^*|$ does not drastically change with protein charge. For reasonable, physiological charge densities (σ_p of up to $\sim 10 e/1000 \text{\AA}^2$) we find $|z_2^*| \sim 3$ –4. Thus, lipid enrichment due to the adsorption of an oppositely charged protein will be most effective for a certain lipid




FIGURE 4 Valence effect. The enrichment ratio of polyvalent lipids (ψ_2/ϕ_2) in the adsorption domain of a protein as a function of the lipid valence z_2 . The different curves correspond to different protein charges, the number on each curve indicating the number of elementary charges per 1000 \AA^2 . Maximum enrichment is marked by a box. The unperturbed membrane composition and protein-membrane distance are the same as in Fig. 3.

valence which, within our idealized model assumptions, seems to be in the range between -3 and -4 .

APPLICATION TO PEPTIDE-MEMBRANE INTERACTIONS

In this section we compare the qualitative results of our model with recent experimental studies of the lateral sequestration of the polyvalent lipid PIP₂ by adsorbed basic peptides. It should be borne in mind that our simple model can only provide an approximate description of such systems. Treating the interaction region of the peptide as a large flat surface is a particularly severe simplification. We return to this and other weaknesses of the model in the Discussion below.

We focus on two peptides for which there are available experimental data: FA-MARCKS(151–175) and a polylysine chain of 13 residues, (Lys)₁₃. The former corresponds to the basic effector segment of the MARCKS protein, where five alanine residues were substituted for the original phenylalanine ones (Gambhir et al., 2004; Wang et al., 2004). We avoid dealing with the MARCKS peptide, because experiments indicate that its hydrophobic phenylalanine residues pull the peptide into the membrane such that its backbone penetrates the membrane (Qin and Cafiso, 1996; Zhang et al., 2003). This kind of interaction is expected to be sensitive to specific molecular details, which are not encompassed by our model.

To apply the model, we need an estimate for the area of the peptide that interacts with the membrane to determine the peptide charge density. In addition, this area is assumed to be equal to that of the adsorption domain, A (Fig. 1). We built

extended peptides (MOE software 2002, Chemical Computing Group, Montreal, Quebec, Canada), similar to the ones used by Wang et al. (2004). We then defined the effective peptide area as the area of its projected backbone plus an envelope of width κ^{-1} around it. (This somewhat arbitrary definition will be further examined at the end of this section.) Dividing the number of charged residues in the peptide by this area, we got the estimates $\sigma_p \simeq 13 \text{ e}/2120 \text{ \AA}^2$ and $13 \text{ e}/1060 \text{ \AA}^2$ for FA-MARCKS(151–175) and (Lys)₁₃, respectively. Note that for these calculated areas and typical peptide–membrane distances h of a few angstroms, the basic assumption of the model, $h \ll \sqrt{A}$, is well satisfied.

We used these estimated σ_p values to produce Fig. 5, plotting the trivalent lipid fraction in the adsorption domain of each peptide as a function of its distance from the membrane. In accordance with experiments (Gambhir et al., 2004), we took the unperturbed membrane composition to be: $\phi_0 = 82\%$ (corresponding to the uncharged zwitterionic PC lipid), $\phi_1 = 17\%$ (monovalent PS), and $\phi_2 = 1\%$ (trivalent PIP₂). Fig. 5 shows that the PIP₂ fraction rises to 18% in the (Lys)₁₃ adsorption domain, whereas only 6% PIP₂ is obtained in the case of FA-MARCKS(151–175). This is caused by the higher charge density of (Lys)₁₃, roughly double that of FA-MARCKS(151–175). However, FA-MARCKS(151–175) has twice the effective area of (Lys)₁₃; thus, if we examine the average number of PIP₂ molecules per peptide adsorption domain, N_2 , the difference is less significant—whereas $N_2 \sim 2.7$ for (Lys)₁₃ at $\kappa h \lesssim 0.3$, for FA-MARCKS(151–175) at the same distance $N_2 \sim 1.8$. (The value of $\kappa h \approx 0.3$ corresponds to $h \approx 3 \text{ \AA}$ at 100 mM salt, which is the approximate peptide–membrane distance; Ben-Tal et al., 1996; Murray et al., 2002. Consequences of this small value will be addressed in the Discussion below.)




FIGURE 5 Comparison between lipid rearrangement induced by FA-MARCKS(151–175) and (Lys)₁₃. Trivalent lipid fraction in the adsorption domain of the FA-MARCKS(151–175) (dashed) and (Lys)₁₃ (solid) peptides as a function of their distance from the membrane. The unperturbed membrane is composed of 82%/17%/1% uncharged/monovalent/trivalent lipid fractions. The enrichment caused by (Lys)₁₃ is much stronger than that achieved by FA-MARCKS(151–175) due to its higher (roughly double) charge density.

Next, we examined the dependence of the trivalent lipid (PIP₂) fraction in the adsorption domain on ϕ_1 , the monovalent lipid fraction in the unperturbed membrane. Fig. 6 presents the results obtained for FA-MARCKS(151–175) and (Lys)₁₃ at various distances. The smaller the value of ϕ_1 , the stronger the enrichment in trivalent lipids. When there is a little amount of monovalent lipids, the membrane charge density induced by the peptide–membrane interaction is attained primarily by the trivalent species. Therefore, at $\phi_1 = 0$, the enrichment in trivalent lipid is maximum. At that limit, the number of PIP₂ molecules per adsorption domain can be simply approximated (at short distances) as the number of charges on the peptide divided by the lipid valence, $N_2 \approx |A\sigma_p/(z_2e)|$. (In practice, however, this example of the polyvalent lipid being the majority charge is problematic, as will be presented in the Discussion.) Fig. 6 shows that, under physiological conditions of 30% monovalent lipid fraction and only 1% PIP₂, both peptides sequester PIP₂—roughly one molecule per FA-MARCKS(151–175) peptide and approximately two molecules per (Lys)₁₃ peptide. This result is in qualitative agreement with experiments (Gambhir et al., 2004), where (Lys)₁₃ was found to attract PIP₂ more strongly than FA-MARCKS(151–175).

To emphasize the strong sequestration of the trivalent lipid, consider the case of a 73%/17%/0 PC/PS/PIP₂ membrane interacting with (Lys)₁₃. At a short distance the peptide charge will be neutralized by ~ 13 PS molecules. For a 72%/17%/1% membrane, as shown by the dashed curve in Fig. 6, there are roughly two to three sequestered PIP₂ lipids per (Lys)₁₃, providing six to nine elementary charges. Thus, introducing 1% PIP₂ into a membrane of 17% PS leads to replacement of about one half of the PS lipids in the adsorption domain by PIP₂. Indication of such an exchange of PS for PIP₂ upon addition of a small amount of PIP₂ was found in a recent experiment (S. McLaughlin, personal communication).

Interestingly, the concentration, on average, of about two PIP₂ molecules per (Lys)₁₃ would not be possible if PIP₂ were of much different valence. Fig. 7 shows the average number of PIP₂ molecules per adsorption domain of both peptides as a function of the PIP₂ hypothetical valence. Similar to the results shown in Fig. 4, we find a nonmonotonic behavior as a function of valence with a maximum at $|z_2| \sim 3$ –4. It is stressed again that, in view of our simplified model, one should pay more attention to the existence of a competition mechanism, leading to an optimum valence, than to the exact value obtained for that valence.

In Fig. 8 we show the dependence of the trivalent lipid fraction in the adsorption domain on its value in the unperturbed membrane. As expected, ψ_2 increases with ϕ_2 . This calculation shows that one needs $\phi_2 \gtrsim 1\%$ to get an average stoichiometry of 1:1 between FA-MARCKS(151–175) and PIP₂. As we have seen above, (Lys)₁₃ sequesters PIP₂ more effectively. Hence, as demonstrated in Fig. 8, a ϕ_2 value of only 0.1% is sufficient to obtain a 1:1 (Lys)₁₃:PIP₂




FIGURE 6 Effect of monovalent lipid concentration. Trivalent lipid fraction (left ordinate) and number (right ordinate) in the adsorption domain of (A) FA-MARCKS(151–175) and (B) $(\text{Lys})_{13}$ peptides as a function of the peptide–membrane distance. Different curves correspond to membranes of different lipid compositions: 69%/30%/1% (solid), 72%/17%/1% (dashed), and 99%/0%/1% (dotted). The enrichment in the polyvalent lipid increases with decreasing monovalent-lipid fraction.

ratio. That is, $(\text{Lys})_{13}$ can sequester an appreciable amount of PIP_2 even when the membrane contains 300-fold more PS lipids than PIP_2 . This value is in very good agreement, perhaps fortuitously, with the measurements of Gambhir et al. (2004).

Finally, we examined the dependence of the PIP_2 enrichment on the ionic strength, i.e., the concentration of mobile salt ions in the solution, n_0 . This parameter enters into the model through the Debye screening length κ^{-1} , which both scales the distance h and affects the amplitude of the electrostatic potential (see, e.g., Eq. 21 in Appendix 1). Fig. 9 shows that ψ_2 decreases with ionic strength. This is a result of the increased screening of the electrostatic attraction between the membrane and protein. The changes are not dramatic up to quite high n_0 values. The reason is the very close proximity of the two objects (3 Å) for which $\kappa h < 1$ in the entire n_0 range examined.




FIGURE 7 Effect of hypothetical PIP_2 valence on sequestration. Average number of polyvalent lipids per adsorption domain of FA-MARCKS(151–175) and $(\text{Lys})_{13}$ as a function of PIP_2 valence. Concentration is a maximum for $|z_2| \sim 3$ –4. Both calculations were performed using a membrane composition of 69%/30%/1% uncharged/monovalent/polyvalent lipid fractions in the unperturbed membrane, and a peptide–membrane distance of $\kappa h = 0.3$.

Effect of approximated peptide size

As discussed above, we define the effective area of the peptide as the area of its projected backbone plus an envelope of width κ^{-1} around it. This definition, however, is somewhat arbitrary. We therefore examine the effect of relaxing the effective-area definition on the values obtained for N_2 .

Fig. 10 shows the average number of trivalent lipids N_2 per adsorption domain of a 20-amino-acid-long peptide for a range of $1/A$ values extending to $\pm 50\%$ of our original estimate. In this range the alteration in N_2 is limited to ± 1 lipid molecules.

Interaction free energy

From the Poisson-Boltzmann theory, as applied to our model, we can calculate the contribution to the free energy of peptide-membrane association coming from electrostatics and entropy. The derivation is given in Appendix 2. The results for the case of FA-MARCKS(151–175) interacting with a membrane of different compositions are presented in Fig. 11. Some of these compositions have also been studied by Gambhir et al. (2004) and Wang et al. (2004). It should be recalled that our model does not take into account repulsive interactions, such as the Born desolvation effect (see Wang et al., 2004). Lipid demixing effects are included in the model, so the local membrane charge density may change due to the peptide. As a result, the electrostatic free energy in our model decreases monotonously as FA-MARCKS(151–175) approaches the membrane. To get the total free energy of association, one ought to add a repulsive interaction at short distances (not included in our model), that would yield




FIGURE 8 Effect of ϕ_2 on enrichment. Trivalent lipid fraction (left ordinate) and average number (right ordinate) in the adsorption domain of (A) FA-MARCKS(151–175) and (B) (Lys)₁₃ peptides as a function of the peptide–membrane distance. The two curves correspond to membrane compositions of 69%/30%/1% PC/PS/PIP₂ (solid) and 69.9%/30%/0.1% PC/PS/PIP₂ (dashed).

a free-energy minimum at a distance of a few angstroms (see, e.g., Wang et al., 2004). Thus, the free-energy values presented in Fig. 11 are probably more negative than the actual binding free energy. More detailed models provide values for the total binding free energy that are roughly one-half the contribution presented here (Wang et al., 2004).

As expected, electrostatic attraction between the peptide and the membrane is proportional to the charge density of the membrane (Fig. 11). The interaction free energy is not sensitive to the specific lipid composition of the membrane, but rather to its average charge density. This result stems from the minor contribution of mixing entropy to the free energy (fourth term in Eq. 23) and the approximate behavior of the membrane in our model as a surface of constant electric potential, which is determined by the average charge density $\bar{\sigma}$ (May et al., 2000). The calculated contribution to the free energy, therefore, is mainly the work required to bring a charged object (protein) into such a potential,




FIGURE 9 Ionic strength effect. Trivalent lipid fraction in the adsorption domain of FA-MARCKS(151–175) (dashed) and (Lys)₁₃ (solid) peptides as a function of salt concentration. The two curves for each peptide were calculated using membrane compositions of 69%/30%/1% (upper), and 82%/17%/1% (lower). A peptide–membrane distance of $h = 3$ Å was used.

regardless of the lipid redistribution. This may also be the reason why finite difference Poisson-Boltzmann (FDPB) calculations (e.g., Ben-Tal et al., 1996; Murray et al., 2002), although ignoring lipid redistribution, often give a good estimate of the binding free energy.

DISCUSSION

Model evaluation

A simplified theoretical model for the effect of an adsorbed charged protein on the distribution of phospholipids in the




FIGURE 10 Dependence of polyvalent lipid number on estimated peptide area. Average number of trivalent lipids (N_2) as a function of the inverse area of the adsorption domain $1/A$. A is also taken as the effective peptide area. The arrow marks the $1/A$ value according to the definition used throughout this study. Results were obtained for a 10 (dotted), 15 (dashed), and 20 (solid) charged residues in a 20-amino-acid-long peptide interacting with a 69%/30%/1% uncharged/monovalent/trivalent membrane at a distance of $\kappa h = 0.3$.




FIGURE 11 Contribution from electrostatics and entropy within the smeared charge model to the free energy of peptide-membrane interaction. Free energy in units of T as a function of the peptide-membrane distance. The curves correspond to FA-MARCKS(151–175) and membrane compositions of 69%/30%/1% (solid), 77%/18%/5% (dash-dotted), 72%/17%/1% (dashed), and 90%/10%/0% (dotted). The vertical dotted line indicates a distance of $\kappa h = 0.3$. The presented contribution to the free energy decreases monotonously with distance. The solid and dash-dotted curves correspond to membranes of very different lipid compositions but of the same charge density, $-0.33 e/a$.

membrane has been presented. One of the key theoretical results is the relative enrichment equation, Eq. 8, relating the concentrations of various lipid species in the adsorption domain. It should be emphasized that this result is almost model independent. Within a two-block picture, it should hold for any protein-membrane system governed by electrostatics, regardless of particular details of the protein, the strength of the electrostatic interactions, and the validity of a mean-field (Poisson-Boltzmann) assumption. The main physical effect described by this relation is the sensitivity of lipid enrichment to valence, i.e., the increased concentration of polyvalent lipids compared to that of monovalent ones in the adsorption domain.

We have demonstrated how simple Poisson-Boltzmann calculations can be added to Eq. 8 to obtain further details of the membranal interaction region. In most of the biologically relevant cases, including the peptides discussed here, the peptide-membrane distance h is of the order of a few angstroms only (Ben-Tal et al., 1996, 1997; Murray et al., 2001, 2002). In the context of our model, the elaborate Poisson-Boltzmann theory is unnecessary for such a thin water layer between the two charged objects. Nevertheless, it should be recalled that, at any rate, our calculations converge to the correct membrane charge density at small h due to charge neutrality, $\sigma_m = -\sigma_p$ (cf. Fig. 12 in Appendix 1).

The model is focused on the qualitative behavior of electrostatically dominated protein-membrane systems. Our description of the protein-membrane interaction is evidently crude, as it omits the molecular details of this complex system; see the atomistic description used in the companion

report of Wang et al. (2004). Thus, effects related to charge discreteness and three-dimensional structure are ignored. Such effects, for example, may play an important role in the membrane adsorption of the MARCKS protein, whose backbone is believed to penetrate the membrane (Zhang et al., 2003) and whose charges are not evenly spaced. Furthermore, we considered only electrostatic and entropy effects while neglecting other interactions (e.g., desolvation). The model is not intended to reproduce such data as the total binding free energy or the optimum protein-membrane distance (e.g., Ben-Tal et al., 1996; Arbuzova et al., 2000). Similarly, the values that have been obtained for the various lipid concentrations should be regarded as rough estimates.

Treating a peptide as a large, flat, uniformly charged surface is probably the gravest simplification of the model, deserving further discussion. The assumption of flatness may be applicable to proteins interacting with the membrane through such effector regions as those described in Wang et al. (2004) and Gambhir et al. (2004). The assumption regarding the lateral extent of the peptide is reasonable in cases where the peptide-membrane distance h is much smaller than both lateral dimensions. For example, in the case of the FA-MARCKS(151–175) discussed above, we estimated the peptide surface area facing the membrane as a rectangle of dimensions $106 \text{ \AA} \times 20 \text{ \AA}$. The smaller lateral dimension is still significantly larger than the typical value of $h \sim 3 \text{ \AA}$. In addition, the fact that this smaller lateral dimension ($\sim 20 \text{ \AA}$) is comparable to the screening length may lead, in principle, to significant finite-size effects. Yet, as already noted above, because of the small value of h in the relevant systems, the charge density in the adsorption domain should become insensitive to these details and be determined, to a good approximation, merely by charge neutrality.

The small value of h , on the other hand, raises a difficulty with respect to the smeared-charge simplification. Because h is similar to or smaller than the typical distance between charged groups on the peptide, neglecting charge discreteness is clearly questionable. Spreading the charges of the lipid headgroups evenly over the membrane is problematic as well. Nonetheless, several recent experiments might help us indicate the limits of validity of models based on smeared electrostatics, such as ours. The affinity of FA-MARCKS(151–175) to PC:PS membranes was found to depend linearly on PS fraction (Murray et al., 1999). Adding a small amount of PIP₂ to such membranes, having a significant PS fraction, did not change the binding significantly (S. McLaughlin, personal communication). These two observations are in line with simple electrostatic considerations. However, the binding affinity of the same peptide to a membrane of relatively low charge density, composed of 99:1 PC:PIP₂, is surprisingly large—similar to that of a 5:1 PC:PS membrane—even though the membrane charge densities in the two cases differ by a factor of ~ 5 (Wang et al., 2002). Clearly, the latter observation cannot be ac-

counted for by smeared electrostatics. We therefore believe that our model gives reliable results regarding lipid redistribution only in cases where the membrane has a high “background” charge density, i.e., a large fraction of monovalent lipid, which is the biologically relevant case. For such membranes, as discussed in the previous section, the model predictions are in qualitative agreement with available experiments.

Apart from these assumptions, the model contains another implicit simplification, namely, that all the electric field lines are contained within the aqueous spacing between the membrane and protein. This commonly used assumption is strictly correct in the limit where the objects are either infinitely thick or of a vanishing dielectric constant. As will be reported elsewhere, we find that this approximation is, in fact, still good for objects of $\epsilon = 2$ and thickness as small as one-third the Debye length.

On the positive side, the simplified model presented here provides new insights into lipid redistribution caused by protein adsorption. Although most of the results presented in this work were obtained using a numerical solution of the nonlinear Poisson-Boltzmann equation, we have demonstrated how one can get the same qualitative results using a much simpler calculation, the SLPB method, involving merely a set of polynomial equations (see Appendix 1). Such a scheme may serve as a better starting point for more detailed numerical calculations, e.g., FDPB (Honig et al., 1993). In most of the current FDPB calculations, a predefined membrane composition identical to that of the bare membrane is used (Ben-Tal et al., 1996, 1997; Murray et al., 1999; Wang et al., 2004). This arbitrary description of the adsorption domain may be improved if one uses a preliminary analysis of the type presented here to produce an approximate lipid configuration.

Future extensions of this work may include phenomena such as elastic deformation of the membrane (Dan et al., 1993; May, 2000), adsorption of multiple proteins (May et al., 2000, 2002), nonuniform charge density in the adsorption domain (May et al., 2000), and acid dissociation at different pH values (Fleck et al., 2002).

Biological implications

ENTH, FYVE, PX, and other membrane-association domains use predefined stereochemistry to recognize polyphosphoinositides (McLaughlin et al., 2002; Lemmon, 2003). These domains, which are commonly found in proteins involved in intracellular signaling, bind tightly to the poly-phosphoinositides, often via ion pairs, anchoring the protein firmly and irreversibly to membrane surfaces. The binding specificity is reflected in the evolutionary conservation across the homologous domains comprising the family; usually the amino acid residues that mediate the poly-phosphoinositides binding are strictly conserved and

can often serve as sequence signatures to recognize these domains using sequence analysis tools.

Here we dealt with a much less specific, and often reversible, mode of membrane recognition via a cluster of basic residues on the membrane-facing region of the protein. These residues interact electrostatically with acidic lipids in the adsorption domain on the bilayer surface. Our model showed that in such cases membrane association induces a preference for polyvalent lipids such as PIP₂ to sequester in the adsorption domain. The number of sequestered polyvalent lipids may be regulated by the charge density and size of the membrane-interaction region on the protein. This result supports the suggestion, advocated in the accompanying papers (Gambhir et al., 2004; Wang et al., 2004), that membrane-associated proteins such as adducin (Matsuoka et al., 2000), DAKAP200 (Rossi et al., 1999), GAP43 (Laux et al., 2000), MARCKS (Wang et al., 2002), and MacMARCKS (Blackshear, 1993), which contain a cluster of basic residues, may create a reservoir of PIP₂ molecules in their adsorption domain (McLaughlin et al., 2002). It may further imply that the membrane-interaction region of such proteins does not have to be strictly conserved evolutionarily; it should only preserve a specific charge density, as in C2 domains (Murray and Honig, 2002). This speculation, naturally, needs to be checked in future studies.

We found that, when all other parameters are held fixed, there is an optimum value of lipid valence that yields maximum enrichment in the adsorption domain (Figs. 4 and 7). This value, resulting from a competition between entropy and stoichiometry, is found to be at reasonable valence values (e.g., $|z_2^*| = 3-4$ in the examples above). Thus, if in a certain biological scenario there is a need to increase the local concentration of a phospholipid by electrostatic interactions, a polyvalent lipid of valence larger than 1 but not too large (say, ~ -3) would be advantageous. The valence of PIP₂, considered to be between -3 and -5 (Toner et al., 1988; McLaughlin et al., 2002; Rauch et al., 2002; Wang et al., 2002) appears to be in line with this criterion.

Assuming a certain valence for PIP₂ and a certain distance for peptide association with the membrane, our model enables the derivation of a set of approximate rules relating the number of basic residues on the peptide to the average number of sequestered PIP₂ molecules per peptide. For example, assuming a trivalent PIP₂ and association distance of ~ 3 Å, under physiological conditions, each segment of seven consecutive lysine residues of an adsorbed poly-lysine peptide such as (Lys)₁₃ (Fig. 6 B) would sequester on the average approximately one PIP₂ molecule when the membrane composition is 69%/30%/1% uncharged/monovalent/trivalent, and roughly one-and-a-half PIP₂ molecules when the membrane composition is 82%/17%/1%. These results are in good agreement with the detailed calculations reported in the companion report of Wang et al. (2004).

APPENDIX 1: METHODS FOR CALCULATING LIPID CONCENTRATIONS

In this section we present the method for calculating the values of ψ_i . We present three levels of approximation and compare their results.

In thermodynamic equilibrium the electrochemical potentials of each lipid species in the protein-free membrane and in the protein adsorption domain should be equal, $\mu_i^{(0)} = \mu_i$. Within a mean-field approximation, this condition can be written as

$$z_i e \Psi^{(0)}(0) + T \ln \frac{\phi_i}{\phi_0} = z_i e \Psi(0) + T \ln \frac{\psi_i}{\psi_0}, \quad (11)$$

where $\Psi^{(0)}(0)$ is the mean electrostatic potential at the bare membrane, and $\Psi(0)$ its value at the adsorption domain. Equation 11 is actually a set of $(k-1)$ equations for every species $i \neq 0$. The incompressibility condition, $\sum_i \psi_i = 1$, closes a set of k equations for the k unknown ψ_i . To solve these equations we need the surface potentials $\Psi^{(0)}(0)$ and $\Psi(0)$. We derive them using the Poisson-Boltzmann theory. Note that $\Psi(0)$ depends on the variables ψ_i that determine the charge density of the adsorption domain, σ_m .

Nonlinear Poisson-Boltzmann

To find the surface potentials we need to solve the Poisson-Boltzmann equation,

$$\frac{\partial^2 y}{\partial z^2} = \kappa^2 \sinh y, \quad (12)$$

with the appropriate boundary conditions. In Eq. 12 y is the local dimensionless potential at a distance z from the membrane, $y(z) \equiv e\Psi(z)/T$. Note that this equation holds only for salt comprised of monovalent ions.

For the bare membrane the boundary conditions are

$$\left. \frac{\partial y^{(0)}}{\partial z} \right|_{z=0} = -\frac{4\pi e \bar{\sigma}}{\epsilon T}, \quad (13)$$

and vanishing of the field at $z \rightarrow \infty$. This problem, of a single charged plate in an electrolyte, is analytically solvable in closed form (Andelman, 1995). The result is

$$y^{(0)}(0) = -4 \tanh^{-1} \gamma, \quad (14)$$

where γ is the positive root of the quadratic equation $\gamma^2 + \gamma \kappa \epsilon T / (\pi e |\bar{\sigma}|) = 1$. (We have assumed the membrane to be negatively charged.)

For the protein-membrane system, the boundary conditions are

$$\left. \frac{\partial y}{\partial z} \right|_{z=0} = -\frac{4\pi e \sigma_m}{\epsilon T}, \quad \left. \frac{\partial y}{\partial z} \right|_{z=h} = \frac{4\pi e \sigma_p}{\epsilon T}. \quad (15)$$

Integrating the Poisson-Boltzmann equation (Eq. 12) once while applying both boundary conditions, we get

$$\frac{\partial y}{\partial z} = \kappa \sqrt{2 \cosh y + C}, \quad C = \left(\frac{4\pi e \sigma_m}{\epsilon \kappa T} \right)^2 - 2 \cosh y(0). \quad (16)$$

$$\cosh y(0) - \cosh y(h) = 2 \left(\frac{2\pi e}{\epsilon \kappa T} \right)^2 (\sigma_m^2 - \sigma_p^2). \quad (17)$$

We then integrate Eq. 16 to get

$$h = \int_0^h dz = \frac{1}{\kappa \sqrt{2}} \int_{y(0)}^{y(h)} \frac{dy}{\sqrt{\cosh y + C/2}}. \quad (18)$$

Equations 17 and 18 are solved numerically for the two unknowns $y(0)$ and $y(h)$, the surface potentials of the membrane and protein. Finally, the values of $y(0)$ and $y^{(0)}(0)$ (Eq. 14) are used in Eq. 11 to calculate the values of ψ_i .

Linear Poisson-Boltzmann

The LPB approximation holds when the electrostatic interactions are weak compared to T , $y \ll 1$. (In fact, the systems relevant to our study are far from this limit; cf. Fig. 11.) In this limit we can linearize the Poisson-Boltzmann equation (Eq. 12),

$$\frac{\partial^2 y}{\partial z^2} = \kappa^2 y. \quad (19)$$

The surface potential of the unperturbed membrane then takes the form (Evans and Wennerstrom, 1994),

$$y^{(0)}(0) = \frac{4\pi e \bar{\sigma}}{\epsilon \kappa T}. \quad (20)$$

For the protein-membrane system we integrate Eq. 19 twice while applying the boundary conditions (Eq. 15) to obtain (Parsegian and Gingell, 1972)

$$y(0) = \frac{4\pi e}{\epsilon \kappa T \sinh(\kappa h)} [\sigma_p + \sigma_m \cosh(\kappa h)]. \quad (21)$$




FIGURE 12 Comparison of the different computational methods. Trivalent lipid fraction in the adsorption domain of a highly charged protein ($\sigma_p = 13 e/1000 \text{ \AA}$) as a function of the protein-membrane distance. Results were obtained using three methods: nonlinear Poisson-Boltzmann (solid curve), linear Poisson-Boltzmann (dashed curve), and the simplified linear Poisson-Boltzmann (dotted curve). At distances $\kappa h < 0.3$, where the membrane approaches charge matching, the three methods differ by $< 10\%$. The unperturbed membrane composition is the same as in Fig. 3.

The potential values of Eqs. 20 and 21 are then used in Eq. 11 to calculate the ψ_i values.

Simplified linear Poisson-Boltzmann

In this further approximation the entropy is neglected in $\mu_i^{(0)}$ and μ_i , and we are left with a uniform surface potential $y(0) = y^{(0)}(0)$. Note that the entropy contribution is not necessarily small compared to T . The strong enrichment in trivalent lipid, as demonstrated in Figs. 3 and 12, entails an entropy penalty of a few T . Yet, it is still much smaller than the electrostatic contribution, which amounts to tens of T (cf. Fig. 11).

Within the LPB approximation we can equate Eqs. 20 and 21 to obtain the charge density in the adsorption domain in closed form as given in Eq. 10 (May et al., 2000). Substituting σ_m in Eq. 1 yields a set of k equations, Eqs. 1, 4, and 8, which are easily solved for the k lipid fractions ψ_i .

Comparison of the methods

Here we compare the NLPB, LPB, and SLPB calculation methods for a specific example. In the limit of weak electrostatic interactions compared to the thermal energy T the LPB method should coincide with the NLPB one. To highlight the difference between the methods we therefore chose as an example a highly charged protein ($\sigma_p = 13 \text{ e}/1000 \text{ \AA}^2$). The trivalent lipid fractions in the adsorption domain, as calculated using the three methods, are plotted in Fig. 12. For both long and short distances all three curves match. In the long distance limit ($\kappa h \gg 1$), the protein-membrane interaction is weak and the linearization of the Poisson-Boltzmann equation is valid. At very short distances the membrane, as described by all three methods, is forced by charge neutrality to match the charge density of the protein (in opposite sign). In the intermediate range the methods differ, yet because of the two constraints at $\kappa h \ll 1$ and $\kappa h \gg 1$ the differences are mild. It is noteworthy that at distances $\kappa h \leq 0.3$ (typical to protein-membrane adsorption), the calculated ψ_i values obtained using the three methods differ by $<10\%$, and the difference in magnitude between the charge densities of the protein and membrane are also $<10\%$.

APPENDIX 2: FREE-ENERGY CALCULATION

In this appendix we calculate the free energy of protein-membrane interaction as arising from our model. It should be recalled that what is presented here is not the total binding free energy but only the contributions due to simple electrostatics and entropy. In particular, because charges are allowed to redistribute in the membrane, the interaction is purely attractive and the free energy decreases monotonously with distance, reaching its minimum at $h = 0$. This should be contrasted with the FDPB calculations (e.g., Ben-Tal et al., 1996; Wang et al., 2004), which take desolvation into account and yield a free-energy minimum at a finite distance.

The general form of the free energy per unit area within a mean-field approximation is

The first term in Eq. 22 is the energy associated with the mean electric field. The second term corresponds to the interaction of the mobile ions with the field, where n_+ and n_- are the local concentrations of the monovalent positive and negative ions. The second integral accounts for the ideal entropy of mixing of the mobile ions and their contact with ion reservoirs having chemical potentials μ_+ and μ_- . The last four terms correspond to the surface energy of the membrane and protein, including mixing entropy of the lipids and electrostatic interactions.

Setting the variations of F with respect to n_+ , n_- , and Ψ to zero, one properly recovers the Boltzmann relations for the mobile ions, the Poisson equation, and, hence, also the Poisson-Boltzmann equation. Using these results along with Eq. 16, and changing the integration variable from z to y , we obtain the following simplified expression for the free energy:

$$F(h) = \frac{2Tn_0}{\kappa} \int_{y(0)}^{y(h)} \frac{1 - 2 \cosh y - C/2}{\sqrt{2 \cosh y + C}} dy + \frac{T\sigma_p}{e} y(h) + \frac{T\sigma_m}{e} y(0) + \frac{T}{a} \sum_i \psi_i \ln \psi_i - \frac{1}{a} \sum_{i>0} \mu_i \psi_i, \quad (23)$$

where the constant C was defined in Eq. 16. We can now use the values obtained for $y(0)$, $y(h)$, μ_i , and ψ_i , as described in Appendix 1, to calculate the free energy of interaction $F(h)$.

We thank S. McLaughlin and D. Murray for providing us with draft copies of their papers before submission, and we are grateful to S. McLaughlin for comments on the manuscript and for valuable discussions. We benefited from discussions with D. Andelman, Y. Burak, M. Gutman, and M. Kozlov.

H.D. acknowledges support from the Israeli Council of Higher Education (Alon Fellowship).

REFERENCES

- Andelman, D., M. M. Kozlov, and W. Helfrich. 1994. Phase transitions between vesicles and micelles driven by competing curvatures. *Europhys. Lett.* 25:231–236.
- Andelman, D. 1995. Electrostatic properties of membranes: the Poisson Boltzmann theory. In *Structure and Dynamics of Membranes*, 2nd Ed, Vol 1B. R. Lipowsky and E. Sackmann, editors. Elsevier, Amsterdam, The Netherlands. 603–642.
- Arbuzova, A., L. Wang, J. Wang, G. Hangyas-Mihalyne, D. Murray, B. Honig, and S. McLaughlin. 2000. Membrane binding of peptides containing both basic and aromatic residues. Experimental studies with peptides corresponding to the scaffolding region of caveolin and the effector region of MARCKS. *Biochemistry*. 39:10330–10339.
- Ben-Tal, N., B. Honig, R. M. Peitzsch, G. Denisov, and S. McLaughlin. 1996. Binding of small basic peptides to membranes containing acidic

$$F(h) = \int_0^h \left[-\frac{\epsilon}{8\pi} \left(\frac{\partial \Psi}{\partial z} \right)^2 + e(n_+ - n_-) \Psi \right] dz + \int_0^h [T(n_+ \ln n_+ - n_+ + n_- \ln n_- - n_- - 2n_0 \ln n_0 + 2n_0) - \mu_+(n_+ - n_0) - \mu_-(n_- - n_0)] dz + \frac{T}{a} \sum_i \psi_i \ln \psi_i - \frac{1}{a} \sum_{i>0} \mu_i \psi_i + \sigma_p \Psi(h) + \sigma_m \Psi(0). \quad (22)$$

- lipids: theoretical models and experimental results. *Biophys. J.* 71: 561–575.
- Ben-Tal, N., B. Honig, C. Miller, and S. McLaughlin. 1997. Electrostatic binding of proteins to membranes. Theoretical predictions and experimental results with charybdotoxin and phospholipid vesicles. *Biophys. J.* 73:1717–1727.
- Blackshear, P. J. 1993. The MARCKS family of cellular protein kinase C substrates. *J. Biol. Chem.* 268:1501–1504.
- Czech, M. P. 2000. PIP₂ and PIP₃: complex roles at the cell surface. *Cell.* 100:603–606.
- Dan, N., P. Pincus, and S. A. Safran. 1993. Membrane-induced interactions between inclusions. *Langmuir.* 9:2768–2771.
- Evans, D. F., and H. Wennerstrom. 1994. The Colloidal Domain, Where Physics, Chemistry, Biology, and Technology Meet, 2nd Ed. VCH Publishers, New York.
- Fleck, C., R. R. Netz, and H. H. von Grünberg. 2002. Poisson-Boltzmann theory for membranes with mobile charged lipids and the pH-dependent interaction of a DNA molecule with a membrane. *Biophys. J.* 82: 76–92.
- Gambhir, A., G. Hangyás-Mihályiné, I. Zaitseva, D. Cafiso, J. Wang, D. Murray, S. Pentyala, S. Smith, and S. McLaughlin. 2004. Electrostatic sequestration of PIP₂ on phospholipid membranes by basic/aromatic regions of proteins. *Biophys. J.* 86:2188–2207.
- Gilson, M. K. 1995. Theory of electrostatic interactions in macromolecules. *Curr. Opin. Struct. Biol.* 5:216–223.
- Groves, J. T., S. G. Boxer, and H. M. McConnell. 1997. Electric field-induced reorganization of two-component supported bilayer membranes. *Proc. Natl. Acad. Sci. USA.* 94:13390–13395.
- Groves, J. T., S. G. Boxer, and H. M. McConnell. 1998. Electric field-induced critical demixing in lipid bilayer membranes. *Proc. Natl. Acad. Sci. USA.* 95:935–938.
- Harries, D., S. May, W. M. Gelbart, and A. Ben-Shaul. 1998. Structure, stability, and thermodynamics of lamellar DNA-lipid complexes. *Biophys. J.* 75:159–173.
- Heimburg, T., B. Angerstein, and D. Marsh. 1999. Binding of peripheral proteins to mixed lipid membranes: effect of lipid demixing upon binding. *Biophys. J.* 76:2575–2586.
- Honig, B., K. Sharp, and A. Suei-Yang. 1993. Macroscopic models of aqueous solutions: biological and chemical applications. *J. Phys. Chem.* 97:1101–1109.
- Honig, B., and A. Nicholls. 1995. Classical electrostatics in biology and chemistry. *Science.* 268:1144–1149.
- Katan, M., and R. L. Williams. 1997. Phosphoinositide-specific phospholipase C: structural basis for catalysis and regulatory interactions. *Semin. Cell Dev. Biol.* 8:287–296.
- Laux, T., K. Fukami, M. Thelen, T. Golub, D. Frey, and P. Caroni. 2000. GAP43, MARCKS, and CAP23 modulate PI(4,5)P₂ at plasmalemmal rafts, and regulate cell cortex actin dynamics through a common mechanism. *J. Cell Biol.* 149:1455–1472.
- Lee, K. Y. C., J. F. Klinger, and H. M. McConnell. 1994. Electric-field-induced concentration gradients in lipid monolayers. *Science.* 263: 655–658.
- Lee, K. Y. C., and H. M. McConnell. 1995. Effect of electric-field gradients on lipid monolayer membranes. *Biophys. J.* 68:1740–1751.
- Lemmon, M. A. 2003. Phosphoinositide recognition domains. *Traffic.* 4:201–213.
- Liu, Y., L. Casey, and L. J. Pike. 1998. Compartmentalization of phosphatidylinositol 4,5- bisphosphate in low-density membrane domains in the absence of caveolin. *Biochem. Biophys. Res. Commun.* 245:684–690.
- Matsuoka, Y., X. Li, and V. Bennett. 2000. Adducin: structure, function and regulation. *Cell. Mol. Life Sci.* 57:884–895.
- May, S. 2000. Theories on structural perturbations of lipid bilayers. *Curr. Opin. Coll. Interface Sci.* 5:244–249.
- May, S., D. Harries, and A. Ben-Shaul. 2000. Lipid demixing and protein-protein interactions in the adsorption of charged proteins on mixed membranes. *Biophys. J.* 79:1747–1760.
- May, S., D. Harries, and A. Ben-Shaul. 2002. Macroion-induced compositional instability of binary fluid membranes. *Phys. Rev. Lett.* 89:268102.
- McLaughlin, S., and A. Aderem. 1995. The myristoyl-electrostatic switch: a modulator of reversible protein-membrane interactions. *Trends Biochem. Sci.* 20:272–276.
- McLaughlin, S., J. Wang, A. Gambhir, and D. Murray. 2002. PIP₂ and proteins: interactions, organization, and information flow. *Annu. Rev. Biophys. Biomol. Struct.* 31:151–175.
- Murray, D., N. Ben-Tal, B. Honig, and S. McLaughlin. 1997. Electrostatic interaction of myristoylated proteins with membranes: simple physics, complicated biology. *Structure.* 5:985–989.
- Murray, D., A. Arbuzova, G. Hangyás-Mihályné, A. Gambhir, N. Ben-Tal, B. Honig, and S. McLaughlin. 1999. Electrostatic properties of membranes containing acidic lipids and adsorbed basic peptides: theory and experiment. *Biophys. J.* 77:3176–3188.
- Murray, D., S. McLaughlin, and B. Honig. 2001. The role of electrostatic interactions in the regulation of the membrane association of G protein beta gamma heterodimers. *J. Biol. Chem.* 276:45153–45159.
- Murray, D., A. Arbuzova, B. Honig, and S. McLaughlin. 2002. The role of electrostatic and nonpolar interactions in the association of peripheral proteins with membranes. *Curr. Top. Membr.* 52:271–302.
- Murray, D., and B. Honig. 2002. Electrostatic control of the membrane targeting of C2 domains. *Mol. Cell.* 9:145–154.
- Netz, R. R. 2001. Electrostatics of counter-ions at and between planar charged walls: from Poisson-Boltzmann to the strong-coupling theory. *Eur. Phys. J. E.* 5:557–574.
- Parsegian, V. A., and D. Gingell. 1972. On the electrostatic interaction across a salt solution between two bodies bearing unequal charges. *Biophys. J.* 12:1192–1204.
- Payraastre, B., K. Missy, S. Giuriato, S. Bodin, M. Plantavid, and M. Gratacap. 2001. Phosphoinositides: key players in cell signalling, in time and space. *Cell. Signal.* 13:377–387.
- Qin, Z. and D. S. Cafiso. 1996. Membrane structure of protein kinase C and calmodulin binding domain of myristoylated alanine rich C kinase substrate determined by site-directed spin labeling. *Biochemistry.* 35:2917–2925.
- Rauch, M. E., C. G. Ferguson, G. D. Prestwich, and D. Cafiso. 2002. Myristoylated alanine-rich C kinase substrate (MARCKS) sequesters spin-labeled phosphatidylinositol 4,5-bisphosphate in lipid bilayers. *J. Biol. Chem.* 277:14068–14076.
- Resh, M. D. 1999. Fatty acylation of proteins: new insights into membrane targeting of myristoylated and palmitoylated proteins. *Biochim. Biophys. Acta.* 1451:1–16.
- Rossi, E. A., Z. Li, H. Feng, and C. S. Rubin. 1999. Characterization of the targeting, binding, and phosphorylation site domains of an A kinase anchor protein and a myristoylated alanine-rich C kinase substrate-like analog that are encoded by a single gene. *J. Biol. Chem.* 274:27201–27210.
- Rouzina, I., and V. A. Bloomfield. 1996. Competitive electrostatic binding of charged ligands to polyelectrolytes: planar and cylindrical geometries. *J. Phys. Chem.* 100:4292–4304.
- Russ, C., T. Heimburg, and H. H. von Grünberg. 2003. The effect of lipid demixing on the electrostatic interaction of planar membranes across a salt solution. *Biophys. J.* 84:3730–3742.
- Simonsen, A., A. E. Wurmser, S. D. Emr, and H. Stenmark. 2001. The role of phosphoinositides in membrane transport. *Curr. Opin. Cell Biol.* 13:485–492.
- Stauffer, T. P., S. Ahn, and T. Meyer. 1998. Receptor-induced transient reduction in plasma membrane PI(4,5)P₂ concentration monitored in living cells. *Curr. Biol.* 8:343–346.
- Toner, M., G. Vaio, A. McLaughlin, and S. McLaughlin. 1988. Adsorption of cations to phosphatidylinositol 4,5-bisphosphate. *Biochemistry.* 27:7435–7443.

- Wang, J., A. Arbuzova, G. Hangyas-Mihalyne, and S. McLaughlin. 2001. The effector domain of myristoylated alanine-rich C kinase substrate binds strongly to phosphatidylinositol 4,5-bisphosphate. *J. Biol. Chem.* 276:5012–5019.
- Wang, J., A. Gambhir, G. Hangyas-Mihalyne, D. Murray, U. Golebiewska, and S. McLaughlin. 2002. Lateral sequestration of phosphatidylinositol 4,5-bisphosphate by the basic effector domain of myristoylated alanine-rich C kinase substrate is due to nonspecific electrostatic interactions. *J. Biol. Chem.* 277:34401–34412.
- Wang, J., A. Gambhir, S. McLaughlin, and D. Murray. 2004. Computational model for the electrostatic sequestration of $\text{PI}(4,5)\text{P}_2$ by membrane-adsorbed basic peptides. *Biophys. J.* 86:1969–1986.
- Zhang, W., E. Crocker, S. McLaughlin, and S. O. Smith. 2003. Binding of peptides with basic and aromatic residues to bilayer membranes: phenylalanine in the myristoylated alanine-rich C kinase substrate effector domain penetrates into the hydrophobic core of the bilayer. *J. Biol. Chem.* 278:21459–21466.