

Probing the Q-Proton Pathway of *ba*₃-Cytochrome *c* Oxidase by Time-Resolved Fourier Transform Infrared Spectroscopy

Constantinos Koutsoupakis,* Tewfik Soulimane,[†] and Constantinos Varotsis*

*Department of Chemistry, University of Crete, 71409 Heraklion, Crete, Greece; and [†]Paul Scherrer Institut, Life Sciences, OSRA/008, CH-5232 Villigen PSI, Switzerland

ABSTRACT In cytochrome *c* oxidase, the terminal respiratory enzyme, electron transfers are strongly coupled to proton movements within the enzyme. Two proton pathways (K and D) containing water molecules and hydrophobic amino acids have been identified and suggested to be involved in the proton translocation from the mitochondrial matrix or the bacterial cytoplasm into the active site. In addition to the K and D proton pathways, a third proton pathway (Q) has been identified only in *ba*₃-cytochrome *c* oxidase from *Thermus thermophilus*, and consists of residues that are highly conserved in all structurally known heme-copper oxidases. The Q pathway starts from the cytoplasmic side of the membrane and leads through the axial heme *a*₃ ligand His-384 to the propionate of the heme *a*₃ pyrrol ring A, and then via Asn-366 and Asp-372 to the water pool. We have applied FTIR and time-resolved step-scan Fourier transform infrared (TRS²-FTIR) spectroscopies to investigate the protonation/deprotonation events in the Q-proton pathway at ambient temperature. The photolysis of CO from heme *a*₃ and its transient binding to Cu_B is dynamically linked to structural changes that can be tentatively attributed to ring A propionate of heme *a*₃ (1695/1708 cm⁻¹) and to deprotonation of Asp-372 (1726 cm⁻¹). The implications of these results with respect to the role of the ring A propionate of heme *a*₃-Asp372-H₂O site as a proton carrier to the exit/output proton channel (H₂O pool) that is conserved among all structurally known heme-copper oxidases, and is part of the Q-proton pathway in *ba*₃-cytochrome *c* oxidase, are discussed.

INTRODUCTION

Hydrogen-bonded networks and regulated electron transfer pathways play the dominant role in the dual function of heme-copper oxidases to reduce O₂ to H₂O and pump protons (Ostermeier et al., 1996; Kannt et al., 1998; Soulimane et al., 2000; Than and Soulimane, 2001). Cytochrome *ba*₃ is a member of the large family of heme-copper oxidases, and in addition to activating O₂ and conserving the energy of the O₂ reduction for subsequent ATP synthesis, is able to catalyze the reduction of nitric oxide (NO) to nitrous oxide (N₂O) under reducing anaerobic conditions (Giuffrè et al., 1999a; Soulimane et al., 2000; Than and Soulimane, 2001). The crystal structure of the protein indicates that the conserved to all heme-copper oxidase subunit I consists of a low-spin heme *b*, and a high-spin heme *a*₃/Cu_B binuclear center, where the dioxygen and nitric oxide reactions take place (Soulimane et al., 2000; Than and Soulimane, 2001). Subunit II contains a mixed valence homodinuclear copper complex (Soulimane et al., 2000; Than and Soulimane, 2001). The *a*-type heme in *ba*₃ and its counterpart *caa*₃ contain a hydroxyethylgeranylgeranyl side chain instead of a hydroxyethylfarnesyl side chain as seen in most eukaryotic and bacterial oxidases (Iwata et al., 1995; Tsukihara et al., 1995; Ostermeier et al., 1997; Yoshikawa et al., 1998; Soulimane et al., 2000; Than and Soulimane, 2001). Three proton pathways were identified in

the crystal structure of *ba*₃ (Soulimane et al., 2000; Than and Soulimane, 2001). They originate at the cytoplasmic side of the membrane and serve for the transfer of protons to either the periplasmic side of the membrane or the active site. Two of these pathways correspond, with respect to their location in the enzyme, to the putative K and D pathways found in the *Paracoccus denitrificans* (Iwata et al., 1995; Ostermeier et al., 1997) and bovine oxidases (Tsukihara et al., 1995; Yoshikawa et al., 1998), despite the fact that most of the residues belonging to these pathways are not conserved. Importantly, Glu-278 (residue numbering of *P. denitrificans*) that is located at the end of the D-channel, and is highly conserved in heme-copper oxidases and has been implicated in redox-induced proton transfer reactions, is replaced by Ile in *ba*₃-cytochrome *c* oxidase (Soulimane et al., 2000; Than and Soulimane, 2001). The third pathway, called Q, starts from the cytoplasmic side of the membrane and leads through the axial heme *a*₃ ligand His-384 to the propionate of the heme *a*₃ pyrrol ring A, and then via Asn-366 and Asp-372 to an accumulation of water molecules, called water pool (Soulimane et al., 2000; Than and Soulimane, 2001). Although there are significant differences concerning the amino residues in the K and D channels between cytochrome *ba*₃ and heme-copper oxidases, the water pool, His-383, Asn-366, and Asp-372 are all highly conserved among all structurally known heme-copper oxidases (Iwata et al., 1995; Tsukihara et al., 1995; Ostermeier et al., 1997; Yoshikawa et al., 1998; Soulimane et al., 2000). Along these lines, it has been postulated that the water pool is part of the proton exit channel, and the primary acceptor for both pumped protons and H₂O molecules that are formed at the binuclear center of the enzyme (Soulimane et al., 2000; Than and Soulimane,

Submitted August 11, 2003, and accepted for publication December 16, 2003.

Address reprint requests to Constantinos Varotsis, Fax: 30-2810-393601; E-mail: varotsis@edu.uoc.gr.

Abbreviations used: FTIR, Fourier transform infrared; TRS²-FTIR, time-resolved step-scan Fourier transform infrared; MCT, mercury cadmium telluride; RR, resonance Raman.


© 2004 by the Biophysical Society

0006-3495/04/04/2438/07 \$2.00

2001). Therefore, it is essential to elucidate the protein dynamics near the H₂O-Asp-372-heme *a*₃-Cu_B site.

Fourier transform infrared difference spectroscopy (FTIR)

*ba*₃ follows that found in other heme-copper oxidases (Woodruff, 1993; Koutsoupakis et al., 2002, 2003a,b,c), and proceeds according to the following scheme:



is a powerful structure-specific technique for exploring changes that occur to individual amino acid residues in a protein as a result of changes to redox and ligation states. The FTIR difference approach has also been used to investigate the CO-photoproduct and the electrochemical oxidized-minus-reduced difference spectra of heme-copper oxidases (Hellwig et al., 1998, 1999, 2002; Iwase et al., 1999; Rich and Breton, 2001; Bailey et al., 2002; Heitbrink et al., 2002; Koutsoupakis et al., 2002, 2003a,b,c; Pinakoulaki et al., 2002a; Stavrakis et al., 2002; Tomson et al., 2002). In the latter case, the perturbation is the change in the redox state of the metal centers, whereas in the former it is the photodissociation of CO from the heme. Recently, it was demonstrated that although the exogenous ligand vibrations (CO) were essentially identical between the room- and low-temperature FTIR spectra of photodissociated CO-cytochromes *aa*₃ and *bo*₃, significant differences exist in the protein bands between these temperatures (Bailey et al., 2002). It was suggested that these differences originate from the fact that at room temperature, CO has dissociated from Cu_B, whereas at low temperature (80 K) the final state has CO coordinated to Cu_B. Moreover, with the ligand dissociation approach the highly conserved Glu-278 (*P. denitrificans* numbering) in the bovine, *P. denitrificans*, *Rhodobacter sphaeroides*, and *bo*₃-cytochrome oxidases, has been proposed to be involved in protonation/deprotonation reactions, and most recently in the protonation reactions during the catalytic cycle of cytochrome *c* oxidase from bovine (Iwaki et al., 2003), *P. denitrificans* (Iwaki et al., 2003), and *R. sphaeroides* (Nyquist et al., 2003).

Due to the unusual ligand-binding kinetic properties of its binuclear center, cytochrome *ba*₃ oxidase is unique among the heme-copper oxidases in being susceptible to a detailed analysis of its ligation dynamics in the heme *a*₃-Cu_B site (Goldbeck et al., 1992; Surerus et al., 1992; Woodruff, 1993; Giuffrè et al., 1999b; Koutsoupakis et al., 2002, 2003a,b,c). Resonance Raman (RR), electron nuclear double resonance (ENDOR), electron paramagnetic resonance (EPR) spectroscopies, in conjunction with permutations of ¹³C- and ¹⁵N-labeled cyanide have indicated that the reaction of oxidized *ba*₃ with cyanide yields heme *a*₃-CN-Cu_B(II)-CN complex (Surerus et al., 1992). The comparative kinetics data on CO photodissociation and rebinding of various heme-copper oxidases and the derived activation parameters have indicated that the CO-ligation/release mechanism in cytochrome

In contrast to the bovine *aa*₃ oxidase, Cu_B of cytochrome *ba*₃ has a relative high affinity for CO (*K*₁ > 10⁴), whereas the transfer of CO to heme *a*₃²⁺ is characterized by a small *k*₂ = 8 s⁻¹, and by a *k*₋₂ = 0.8 s⁻¹ that is 30-fold greater than that of the bovine *aa*₃ (Giuffrè et al., 1999b; Koutsoupakis et al., 2002).

In our previous cytochrome *ba*₃ work, we identified the equilibrium Cu_B¹⁺-CO complex, and concluded that the environment of the binuclear center is not altered in the pD-5.5–9.7 range. (Koutsoupakis et al., 2002). The time-resolved step-scan FTIR (TRS²) difference spectra revealed the dynamics of the binuclear center and showed protein conformational changes near the heme *a*₃ propionates (Koutsoupakis et al., 2002). In subsequent work we have demonstrated, that the ligand delivery channel is located at the Cu_B site, and the presence of a docking site near the heme *a*₃ propionates (Koutsoupakis et al., 2003a,b,c). In recent FTIR studies it has been noted that functional groups, including carboxyl groups of amino acids residues, are difficult to deuterate (Okuno et al., 2003). Therefore, we have investigated the CO-bound *ba*₃ complex in the pH 5.5–9.5 range, aiming to finalize the pH/pD sensitivity of the binuclear center by FTIR. We have also investigated the protein response subsequent to CO photolysis from heme *a*₃ by TRS² FTIR spectroscopy. On the basis of the tentative assignments of the 1695/1708 and 1726 cm⁻¹ modes, the TRS² data may reflect that Asp-372 undergoes deprotonation upon photodissociation of CO from heme *a*₃, and that there is a H-bonded connectivity between the ring A propionate of heme *a*₃-Asp-372-H₂O. By combining our results with those from a variety of other experiments, we postulate that the ring A propionate of heme *a*₃-Asp-372-H₂O site, which is conserved among all structurally known heme-copper oxidases, and is part of the Q-proton pathway in cytochrome *ba*₃, forms an output proton channel. This way, the ring A propionate of heme *a*₃-Asp-372-H₂O group may accept a proton, which in turn causes release of a proton to the exit channel, the so-called water pool.

MATERIALS AND METHODS

Cytochrome *ba*₃ was isolated from *Thermus thermophilus* HB8 cells according to previously published procedures (Soulimane et al., 2000; Than and Soulimane, 2001). The samples used for the FTIR measurements had an

enzyme concentration of ~ 1 mM and were placed in a desired buffer (pH 5.25–6.5, MES; pH 7.5, HEPES; pH 8.5–9.8, CHES). The pH solutions prepared in D_2O buffers were measured by using a pH meter and assuming $pD = pH(\text{observed}) + 0.4$. Dithionite reduced samples were exposed to 1 atm CO (1 mM) in an anaerobic cell to prepare the carbonmonoxy adduct and transferred to a tightly sealed FTIR cell with CaF_2 windows, under anaerobic conditions. CO gas (99.9%) was obtained from Messer (Frankfurt, Germany) and isotopic CO (91.6% $^{13}C^{16}O$ and 8.4% $^{13}C^{18}O$) was purchased from Isotec (Miamisburg, OH). FTIR measurements were performed on a Bruker (Newark, DE) Equinox IFS 55 spectrometer equipped with a mercury cadmium telluride (MCT) detector (Graseby Infrared D316, response limit 600 cm^{-1}). The experimental techniques used for generating and timing the green photolysis pulse (532 nm and 10 ns) and the IR probe beam to obtain time-resolved step-scan FTIR difference spectra have been reported (10, 17–19). Optical absorption spectra were recorded with a Perkin-Elmer (Fremont, CA) Lambda 20 ultraviolet-visible spectrometer before and after the FTIR measurements to ensure the formation and stability of the CO adducts.

RESULTS

Fig. 1 shows the FTIR spectra of CO-bound cytochrome *ba₃* in the pH 5.25–9.8 range. The spectra exhibit peaks at 1967, 1973, and 1982 cm^{-1} that have been assigned (Koutsoupakis

et al., 2002) to the C–O stretching modes of heme *a₃*-CO (complex **B** in Scheme 1) originating from three different conformers, and the C–O stretching mode of *Cu_B*-CO (complex **A**) located at 2053 cm^{-1} . These modes are downshifted to 1923, 1928, 1937, and 2007 cm^{-1} , respectively, when ^{12}CO is replaced by ^{13}CO . The intensity and bandwidth of all three heme *a₃*-CO modes remains unchanged in the pH 5.25–9.8 range. Similar observations have been reported for the bovine enzyme (Einarsdóttir et al., 1988; Iwase et al., 1999; Rich and Breton, 2001). The insensitivity of heme *a₃*-CO to pH demonstrates that the properties of the proximal to heme *a₃* His-384, that is part of the Q-proton pathway, and known to affect the frequencies of the heme *a₃* bound CO, remain unchanged in the pH 5.25–9.8 range (Pinakoulaki et al., 2002b). The insensitivity of complex **A** to pH changes indicates that the environment of the *Cu_B*-N(His) ligands that is distal to the bound heme *a₃*-CO also remains unchanged in the pH range 5.25–9.8.

In the oxidized-minus-reduced (electrochemical) FTIR difference spectra of *aa₃* oxidase from *P. denitrificans* the observation of a trough/peak pattern in the 1700 cm^{-1} region has been interpreted as an environmental change induced by the change in the redox state of the metal centers (Hellwig et al., 1998). In the FTIR difference spectra obtained upon CO-photolysis from the heme Fe, the appearance of a negative peak in the 1700 cm^{-1} region (protonated carboxylic acids) has been interpreted as deprotonation of a carboxyl group (Rich and Breton, 2001; Heitbrink et al., 2002; Nyquist et al., 2003; Okuno et al., 2003). With these approaches the properties of the highly conserved Glu-278 in heme-copper oxidases have been investigated. In *ba₃*-cytochrome *c* oxidase Glu is replaced by Ile, and no trough/peaks patterns were observed in the oxidized-minus-reduced FTIR difference spectra in the 1700 cm^{-1} region (Hellwig et al., 1999). This observation indicates that a change in the redox state of the metal centers is not observed as a change in the protonation/deprotonation of Glu and/or Asp residues in the enzyme.

Signals in the amide I region ($1620\text{--}1690\text{ cm}^{-1}$) can be attributed to changes of the C=O modes caused by perturbation in the polypeptide backbone and, to the C=O modes of Asn and Gln (Hellwig et al., 1998). Coupled CN stretching and NH bending modes and, the asymmetric COO^- modes from deprotonated heme propionates and Glu and Asp side chains are expected in the $1530\text{--}1590\text{ cm}^{-1}$ region (Hellwig et al., 1998). It has been established that the deprotonated symmetric COO^- vibrations of heme-propionates and Asp residues are expected at 1350 and 1450 cm^{-1} , respectively, whereas the C=O bonds of the protonated forms are at 1730 cm^{-1} (Hellwig et al., 1998, 1999, 2002). Also, the asymmetric vibrations of heme-propionates and Asp are located at 1530 and 1590 cm^{-1} , respectively (Hellwig et al., 1998, 1999, 2002). Fig. 2 collects TRS² FTIR difference spectra (coaveraged first 100 μs after photodissociation of CO) in the pH 6.5–9.35 and pD 7.5–10.1 range. The spectra represent




FIGURE 1 FTIR spectra of the cytochrome *ba₃*-CO complex at the indicated pH values, 293 K. The spectra at pH 5.25 and 9.60 correspond to the *ba₃*- ^{13}CO complex. Enzyme concentration was 1 mM and the pathlength $15\text{ }\mu m$. The spectral resolution was 2 cm^{-1} except for the spectra at pH 9.10 and 9.80, where it was 4 cm^{-1} .




FIGURE 2 Time-resolved step-scan FTIR difference spectra of the CO-bound form of fully reduced cytochrome *ba*₃ oxidase, at the indicated pH and pD values, after CO photolysis from heme *a*₃. Each spectrum is the average of 20 individual spectra from 5 to 100 μ s. The pathlength was 15 and 30 μ m for the pH and pD samples, respectively, and the spectral resolution 8 cm^{-1} . The time resolution was 5 μ s and 10 coadditions were collected and averaged per data point. The excitation wavelength was 532 nm (4 mJ/pulse) and three measurements were recorded and averaged for each data set. The spectra are normalized to the intensity of the 2053 cm^{-1} mode (Cu_B^{1+} -CO transient species).

the difference between the heme *a*₃-CO and the Cu_B -CO species because at 5–100 μ s subsequent to CO photolysis, the ligand is bound to Cu_B (Koutsoupakis et al., 2002, 2003a,b,c). The TRS² FTIR difference spectra in the 1690–1780 cm^{-1} region (Fig. 2 E) present an excellent “W” shape characteristic of substantial perturbation of carboxyl groups upon light-induced dissociation of CO from heme *a*₃ and subsequent ligation to Cu_B . The C=O stretching band that we tentatively assign to the ring A of heme *a*₃ propionate is seen as a derivative-shaped feature in the TR-FTIR difference spectrum with the trough/peak at 1708/1695 cm^{-1} that is 2–3 cm^{-1} higher than that observed in D₂O (Fig. 2 E). The frequency at 1695 cm^{-1} in the transient spectra means weaker C=O bond and therefore stronger H-bonding to surrounding groups. The other half of the “W” shape difference spectrum consists of a negative band at 1726 cm^{-1} . We also tentatively assign the 1726 cm^{-1} negative band to the C=O stretch of Asp-372 (*T. Thermophilus* sequencing number) because neither Glu nor other Asp residues are near the binuclear center, where the induced perturbation is expected to affect the structures of nearby residues. The TRS² FTIR difference spectra in the pH 6.47–9.35 range show little change. The insensitivity of the 1726 cm^{-1} mode to external pH indicates that the pK_a of Asp-372 must be higher than 9.4. The spectra obtained in the pD 7.5–10.1 range show that the 1726 cm^{-1} mode is absent (see below). The peak/trough of the propionate C=O stretching band observed at 1693/1705 cm^{-1} at pD 7.5 and 8.5 is similar to that obtained at pD 10.1 but with a noticeable intensity increase of the 1705 cm^{-1} trough in all the pD experiments. The rate of decay of the transient 1694(+)/1706(–) signals attributed to perturbation of the heme *a*₃ propionates (COOH) displays similar time constant as the transient Cu_B^{1+} -CO complex (Koutsoupakis et al., 2002). Although we have not been able to monitor the

kinetics of the 1726 cm^{-1} species accurately, due to interference from H₂O in this frequency range, the TRS²-FTIR spectra at times longer than 2 ms (data not shown) show a substantial decrease of the 1726 cm^{-1} mode, suggesting that there is a coupling between ligation dynamics in the binuclear center and the environment sensed by both the Asp-372 and the heme *a*₃ propionates (Koutsoupakis et al., 2002). Based on ¹³C labeling experiments in *aa*₃ oxidase from *P. denitrificans* (Behr et al., 2000) the modes at 1570 and 1538 have been assigned to $\nu(\text{COO}^-)^{\text{asym}}$ of heme propionates. Intensity changes and/or frequency shifts of the symmetric and asymmetric vibrations that could be attributed to both the deprotonated forms of heme-propionates and Asp-372 in *ba*₃ oxidase are observed (Hellwig et al., 1999). These include the peaks/troughs at 1390/1370 cm^{-1} ($\nu(\text{COO}^-)^{\text{sym}}$) of ring A propionate of heme *a*₃ and at 1456/1444 cm^{-1} ($\nu(\text{COO}^-)^{\text{sym}}$) of Asp-372. Furthermore, the negative band at 1529 cm^{-1} and the peak/trough at 1592/1582 cm^{-1} can be tentatively assigned to $\nu(\text{COO}^-)^{\text{asym}}$ of the ring A propionate of heme *a*₃ and of Asp-372, respectively. Comparison of the pH/pD spectra shows that there is noticeable downshift (3 cm^{-1}) of $\nu(\text{COO}^-)^{\text{asym}}$ of propionates at 1526 cm^{-1} in the pD 7.5–10.1 range. In addition, the 1456/1444 cm^{-1} $\nu(\text{COO}^-)^{\text{sym}}$ of Asp-372 has lost most of its intensity at pD 10.1 indicating alterations in the Asp-372 environment due to H/D exchange. It should be noted, however, that the deprotonated forms of both the heme-propionates $\nu(\text{COO}^-)^{\text{sym}}$ and Asp-372 persist up to pH 6.5. The appearance of COO(H) modes ascribed to heme-propionates (Fig. 2, A, C, and E) and Asp (Fig. 2 B, D, and E) in both the protonated (Fig. 2 E) and deprotonated (Fig. 2, A–D) spectral region of the TRS² FTIR-difference spectra indicates the presence of both conformations. Interestingly, in the oxidized-minus-reduced electrochemical FTIR difference spectra of *ba*₃ only the protonated forms of the propionates

were observed, and no modes ascribed to either protonated or deprotonated Glu and/or Asp were observed (Hellwig et al., 1999).

DISCUSSION

TRS²-FTIR spectroscopy has already proven to be a very powerful technique in studying transient changes at the level of individual amino acids during protein action. The intensity changes and frequency shifts of side chains and backbone structures observed in the TR-FTIR difference spectra is the result of the perturbation induced by the photodissociation of CO from heme *a*₃ and its subsequent binding to Cu_B, to structures near heme *a*₃ and Cu_B. The following discussion for the behavior of the ring A propionates and Asp-372 is based on our tentative assignments. The observation of the 1695/1708 cm⁻¹ peak/trough and its subsequent shift to 1693/1705 cm⁻¹ upon H/D exchange is similar to that observed in the oxidized-minus-reduced spectra from which it was concluded that the heme-propionates in the *ba*₃ oxidase from *T. thermophilus* are essentially protonated (Hellwig et al., 1999). The presence of deprotonated signals at 1390/1370 cm⁻¹ indicates, however, that this is not the case. Obviously there is an equilibrium of COO⁻ ↔ COOH. The shift of the 1529 cm⁻¹ mode to 1526 cm⁻¹ upon H/D exchange indicates a dependence on local environment and/or hydrogen bonding interactions. Similar conclusion can be drawn from the reduced intensity of the 1456/1444 cm⁻¹ modes in the D₂O experiments. To account for the lack of an observable negative peak at 1726 cm⁻¹ in the D₂O experiments we suggest that the loss of the H-bonding connectivity in the local environment of heme *a*₃-Asp-372-H₂O upon H/D exchanges do not alter the deuterated Asp-372, and thus, we do not observe a negative peak upon the induced perturbation (CO-photolysis from heme *a*₃). Consequently, the proton connectivity between the three groups is disrupted in the presence of D₂O, allowing Asp-372 to adopt a conformation that is significantly different from that observed in the pH experiments. Therefore, the detection of the deprotonated Asp-372 is not only the result of the induced perturbation, but rather a combination of the H-bonded connectivity of the three groups that is lost in the presence of D₂O. Taken together, the detection of both protonated and deprotonated forms of the ring A of heme *a*₃ propionate and the deprotonated Asp-372 in conjunction with the dependence of their deprotonated forms on the local environment suggests a protonic connectivity between the ring A propionate of heme *a*₃, Asp-372, and a H₂O molecule that is part of the Q-proton pathway.

One of the strongest interactions between different groups in heme-copper oxidases is that between the ring A propionate and Asp-372 because the two carboxyl groups are only 3.3 Å apart (Kannt et al., 1998). It was concluded that net protonation of the coupled system will depend on the interaction with the environment and that these two residues share a single proton over a pH 4–11.5 range (Kannt et al., 1998).

In addition, this network recently has been proposed as a part of the exit pathway for the pump protons (Soulimane et al., 2000; Than and Soulimane, 2001). To account for the presence of both the protonated and deprotonated forms of the ring A propionate and only the deprotonated form of Asp-372, we present in Fig. 3 a schematic view, based on the TRS² FTIR data presented here and the crystal structure (Soulimane et al., 2000; Than and Soulimane, 2001), that involves the Asp-372/propionate pair and a H₂O molecule. In the scheme, we invoke a specific role to the ring A propionate-Asp-372 to proton motion. This pair may accept a proton either in the oxidative or reductive phase (Verkhovsky et al., 1999), which in turn causes release of a proton to the water pool. The accumulation of H₂O molecules has been identified in the *P. denitrificans* oxidase and its involvement in proton exit channels has been demonstrated by mutagenesis experiments (Ostermeier et al., 1996; Kannt et al., 1998). It is important to note that in the scheme presented here only states B and D, in which a single proton is shared between the ring A propionate and Asp-372, may accept a proton that in turn causes the release of a proton to the water pool. We postulate that this pathway is blocked when both groups are protonated (state A) or deprotonated (state C). Although we do not know the source of the proton, our data strongly indicate that it is not from His-283 or any of the other Cu_B-His ligands (Koutsoupakis et al., 2002). This sequential or concerted H-bonded connectivity between the environments sensed by the ring A heme *a*₃ propionate-Asp-372-H₂O could have an activation energy for proton motion. The abovementioned protonic connectivity and the fast equilibrium of the water pool with bulk solvent suggest that the water pool may serve as a primary acceptor for both the H₂O molecules, formed during the catalytic turnover, and pumped protons.




FIGURE 3 Schematic view of the protonic connectivity between the ring A heme *a*₃ propionate-Asp-372-H₂O of the Q-proton pathway in *ba*₃-cytochrome *c* oxidase.

The exchangeable protons could play a vital role in the biological function of the enzyme. The first step in locating possible sites of proton motion requires identification of labile protons that could be either redox linked or perturbed by ligand motion; in this case, the CO photodissociation from heme *a₃*. The model discussed above postulates the role of the ring A heme *a₃* propionate-Asp-372-H₂O site in the Q-proton channel as a proton carrier to the water pool, demonstrating a facile pathway connecting the catalytic binuclear center and the exit/output proton channel. Based on the status of the ring A heme *a₃* propionate-Asp-372-H₂O site in all structurally known heme-copper oxidases we suggest that our data do not reflect only specific properties of the *ba₃*-cytochrome *c* oxides but rather are extended to the superfamily of heme-copper oxidases. The lack of similar observation in other heme-copper oxidases containing Glu-278 can be explained by the observation of difference peaks near 1734 cm⁻¹ that have been attributed to Glu-278. This way, strong overlap between the Glu and Asp modes in that frequency domain may have prevented the spectral isolation of the C=O stretching vibration of Asp that is located near the water pool. Experiments are in progress with emphasis on these questions.

We thank Stavros Stavrakis for excellent technical assistance and Dr. Eftychia Pinakoulaki for reading the manuscript.

This work was partially supported by the Greek Ministry of Education.

REFERENCES

- Bailey, J. A., F. L. Tomson, S. L. Mecklenburg, G. M. MacDonald, A. Katsonouri, A. Puustinen, R. B. Gennis, W. H. Woodruff, and R. B. Dyer. 2002. Time-resolved step-scan Fourier transform infrared spectroscopy of the CO adducts of bovine cytochrome *c* oxidase and of cytochrome *bo₃* from *Escherichia coli*. *Biochemistry*. 41:2675–2683.
- Behr, J., H. Michel, W. Mäntele, and P. Hellwig. 2000. Functional properties of the heme propionates in cytochrome *c* oxidase from *Paracoccus denitrificans*. Evidence from FTIR difference spectroscopy and site-directed mutagenesis. *Biochemistry*. 39:1356–1363.
- Einarsdóttir, Ó., M. G. Choc, S. Weldon, and W. S. Caughey. 1988. The site and mechanism of dioxygen reduction in bovine heart cytochrome *c* oxidase. *J. Biol. Chem.* 263:13641–13654.
- Giuffrè, A., G. Stubauer, P. Sart, M. Brunori, W. G. Zumft, G. Buse, and T. Soulimane. 1999a. The heme-copper oxidases of *Thermus thermophilus* catalyze the reduction of nitric oxide: evolutionary implications. *Proc. Natl. Acad. Sci. USA*. 96:14718–14723.
- Giuffrè, A., E. Forte, G. Antonini, E. D'Itri, M. Brunori, T. Soulimane, and G. Buse. 1999b. Kinetic properties of *ba₃* oxidase from *Thermus thermophilus*: effect of temperature. *Biochemistry*. 38:1057–1065.
- Goldbeck, R. A., O. Einarsdóttir, T. D. Dawes, D. B. O'Connor, K. K. Surerus, J. A. Fee, and D. S. Kliger. 1992. Magnetic circular dichroism study of cytochrome *ba₃* from *Thermus thermophilus*: spectral contributions from cytochromes *b* and *a₃* and nanosecond spectroscopy of carbon monoxide photodissociation intermediates. *Biochemistry*. 31:9376–9387.
- Heitbrink, D., H. Sigurdson, C. Bowlwien, P. Brzezinski, and J. Heberle. 2002. Transient binding of CO to Cu_B in cytochrome *c* oxidase is dynamically linked to structural changes around a carboxyl group: a time-resolved step-scan Fourier transform infrared investigation. *Biophys. J.* 82:1–10.
- Hellwig, P., J. Behr, C. Ostermeier, O.-M. H. Richter, U. Pfitzner, A. Odenwald, B. Ludwig, H. Michel, and W. Mäntele. 1998. Involvement of glutamic acid 278 in the redox reaction of the cytochrome *c* oxidase from *Paracoccus denitrificans* investigated by FTIR spectroscopy. *Biochemistry*. 37:7390–7399.
- Hellwig, P., T. Soulimane, G. Buse, and W. Mäntele. 1999. Electrochemical, FTIR, and UV/VIS spectroscopic properties of the *ba₃* oxidase from *Thermus thermophilus*. *Biochemistry*. 38:9648–9658.
- Hellwig, P., U. Pfitzner, J. Behr, B. Rost, R. P. Pasavento, W. V. Donk, R. B. Gennis, H. Michel, B. Ludwig, and W. Mäntele. 2002. Vibrational modes of tyrosines in cytochrome *c* oxidase from *Paracoccus denitrificans*: FTIR and electrochemical studies on Tyr-D₄-labeled and on Tyr280His and Tyr35Phe mutant enzymes. *Biochemistry*. 41:9116–9125.
- Iwaki, M., A. Puustinen, M. Wikström, and P. Rich. 2003. ATR-FTIR spectroscopy of the P_M and F intermediates of bovine and *Paracoccus denitrificans* cytochrome *c* oxidase. *Biochemistry*. 42:8809–8817.
- Iwase, T., C. Varotsis, K. Shinzawa-Itoh, S. Yoshikawa, and T. Kitagawa. 1999. Infrared evidence for Cu_B ligation of photodissociated CO of cytochrome *c* oxidase at ambient temperatures and accompanied deprotonation of a carboxyl side chain of protein. *J. Am. Chem. Soc.* 121:1415–1416.
- Iwata, S., C. Ostermeier, B. Ludwig, and H. Michel. 1995. Structure at 2.8 Å resolution of cytochrome *c* oxidase from *Paracoccus denitrificans*. *Nature*. 376:660–669.
- Kannt, A., C. R. D. Lancaster, and H. Michel. 1998. The coupling of electron transfer and proton translocation: electrostatic calculations on *Paracoccus denitrificans* cytochrome *c* oxidase. *Biophys. J.* 74:708–721.
- Koutsoupakis, K., S. Stavrakis, E. Pinakoulaki, T. Soulimane, and C. Varotsis. 2002. Observation of the equilibrium Cu_B-CO complex and functional implications of the transient heme *a₃* propionates in cytochrome *ba₃*-CO from *Thermus thermophilus*. Fourier transform infrared (FTIR) and time-resolved step-scan FTIR studies. *J. Biol. Chem.* 277:32860–32866.
- Koutsoupakis, K., S. Stavrakis, T. Soulimane, and C. Varotsis. 2003a. Oxygen-linked equilibrium Cu_B-CO species in cytochrome *ba₃* oxidase from *Thermus thermophilus*. Implications for an oxygen channel at the Cu_B site. *J. Biol. Chem.* 278:14893–14896.
- Koutsoupakis, K., S. T. Soulimane, and C. Varotsis. 2003b. Docking site dynamics of *ba₃*-cytochrome *c* oxidase from *Thermus thermophilus*. *J. Biol. Chem.* 278:36806–36809.
- Koutsoupakis, K., S. T. Soulimane, and C. Varotsis. 2003c. Ligand binding in a docking site of cytochrome *c* oxidase: a time-resolved step-scan Fourier transform infrared study. *J. Am. Chem. Soc.* 125:14728–14732.
- Nyquist, R. M., D. Heitbrink, C. Bolwien, R. B. Gennis, and J. Heberle. 2003. Direct observation of protonation reactions during the catalytic cycle of cytochrome *c* oxidase. *Proc. Natl. Acad. Sci. USA*. 100:8715–8720.
- Okuno, D., T. Iwase, K. Shinzawa-Itoh, S. Yoshikawa, and T. Kitagawa. 2003. FTIR detection of protonation/deprotonation of key carboxyl side chains caused by redox change of the Cu_A-heme *a* moiety and ligand dissociation from the heme *a₃*-Cu_B center of bovine heart cytochrome *c* oxidase. *J. Am. Chem. Soc.* 125:7209–7218.
- Ostermeier, C., S. Iwata, and H. Michel. 1996. Cytochrome *c* oxidase. *Curr. Opin. Struct. Biol.* 6:460–466.
- Ostermeier, C., A. Harrenga, U. Ermler, and H. Michel. 1997. Structure at 2.7 Å resolution of the *Paracoccus denitrificans* two-subunit cytochrome *c* oxidase complexed with an antibody F_v fragment. *Proc. Natl. Acad. Sci. USA*. 94:10547–10553.
- Pinakoulaki, E., T. Soulimane, and C. Varotsis. 2002a. Fourier transform infrared (FTIR) and step-scan time-resolved FTIR spectroscopies reveal a unique active site in cytochrome *caa₃* oxidase from *Thermus thermophilus*. *J. Biol. Chem.* 277:32867–32874.
- Pinakoulaki, E., U. Pfitzner, B. Ludwig, and C. Varotsis. 2002b. The role of the cross-link His-Tyr in the functional properties of the binuclear center in cytochrome *c* oxidase. *J. Biol. Chem.* 277:13563–13568.

- Rich, P. R., and J. Breton. 2001. FTIR studies of the CO and cyanide adducts of fully reduced bovine cytochrome *c* oxidase. *Biochemistry*. 40:6441–6449.
- Soulimane, T., G. Buse, G. P. Bourenkov, H. D. Bartunik, R. Huber, and M. E. Than. 2000. Structure and mechanism of the aberrant *ba*₃-cytochrome *c* oxidase from *Thermus thermophilus*. *EMBO J.* 19:1766–1776.
- Stavakis, S., K. Koutsoupakis, E. Pinakoulaki, A. Urbani, M. Saraste, and C. Varotsis. 2002. Decay of the transient Cu_B-CO complex is accompanied by formation of the heme Fe-CO complex of cytochrome *cbb*₃-CO at ambient temperature: evidence from time-resolved Fourier transform infrared spectroscopy. *J. Am. Chem. Soc.* 124:3814–3815.
- Surerus, K. K., W. A. Oertling, F. Chaoliang, R. J. Gurbel, O. Einarsdóttir, W. E. Antoline, R. B. Dyer, B. M. Hoffman, W. H. Woodruff, and J. A. Fee. 1992. Reaction of cyanide with cytochrome *ba*₃ from *Thermus thermophilus*: spectroscopic characterization of the Fe(II)_{a3}-CN-Cu(II)_B-CN complex suggests four ¹⁴N atoms are coordinated to Cu_B. *Proc. Natl. Acad. Sci. USA*. 89:3195–3199.
- Than, M. E., and T. Soulimane. 2001. *Ba*₃-cytochrome *c* oxidase from *Thermus thermophilus*. In *Handbook of Metalloproteins*. A. Messerschmidt, R. Huber, T. Poulos, and K. Wieghardt, editors. John Wiley and Sons, Ltd., Chichester, UK. 363–378.
- Tomson, F., J. A. Bailey, R. B. Gennis, C. J. Unkefer, Z. Li, L. A. Silks, R. A. Martinez, R. J. Donahoe, R. B. Dyer, and W. H. Woodruff. 2002. Direct infrared detection of the covalently ring linked His-Tyr structure in the active site of the heme-copper oxidases. *Biochemistry*. 41:14383–14390.
- Tsukihara, T., H. Aoyama, E. Yamashita, T. Tomizaki, H. Yamaguchi, K. Shinzawa-Itoh, R. Nakashima, R. Yaono, and S. Yoshikawa. 1995. Structures of metal sites of oxidized bovine heart cytochrome *c* oxidase at 2.8 Å. *Science*. 269:1069–1074.
- Verkhovsky, M. I., A. Jasaitis, M. L. Verkhovskaya, J. E. Morgan, and M. Wikström. 1999. Proton translocation by cytochrome *c* oxidase. *Nature*. 400:480–483.
- Woodruff, W. H. 1993. Coordination dynamics of heme-copper oxidases. The ligand shuttle and the control and coupling of electron transfer and proton translocation. *J. Bioenerg. Biomembr.* 25:177–188.
- Yoshikawa, S., K. Shinzawa-Itoh, R. Nakashima, R. Yaono, E. Yamashita, N. Inoue, M. Yao, M. J. Fei, C. P. Libeu, T. Mizushima, H. Yamaguchi, T. Tomizaki, and T. Tsukihara. 1998. Redox-coupled crystal structural changes in bovine heart cytochrome *c* oxidase. *Science*. 280:1723–1729.