

change the diminishing effect of the ablation on spontaneous activity ($n=3$). Epicardial imaging showed that spontaneous ectopic beats were manifested as concentric epicardial breakthrough patterns, located near spatiotemporal gradients of NADH fluorescence. These data strongly suggest that in un-paced hearts from healthy rats that are perfused with Tyrode's solution, the main mechanism of spontaneous ectopic activity associated with either ischemia, low-flow or full-flow reperfusion is activation of local Purkinje fibers.

1324-Pos Board B168

Acute effects of Lipopolysaccharide on L-type Ca^{2+} channel currents and Transient Outward K^{+} channel currents in Rat Ventricular Myocytes

Dushon DeVere Riley^{1,2}, W. Jonathan Lederer^{1,2}.

¹University of Maryland, Baltimore, Baltimore, MD, USA, ²Institute of Molecular Cardiology, University of Maryland Biotechnology Institute, Baltimore, MD, USA.

Septic Shock has accounted for nearly 2 percent of all hospitalizations and has a mortality rate of 40-50%. It is a progression of sepsis, which is caused by bacterial infection in the blood and is characterized by a whole body inflammatory state known as systemic inflammatory response syndrome (SIRS). Cardiac dysfunction is one of the principal pathologies of sepsis and septic shock, along with other hemodynamic changes and dysfunction in multiple organs including the lungs and brain. While cardiac dysfunction is typically associated with the late stages of septicemia in clinical observations, animal models of septicemia have shown that cardiac dysfunction can occur well before late stages of sepsis and the induction of septic shock. Sepsis is caused by exposure to lipopolysaccharides (LPS), an endotoxin found in the outer membrane of gram-negative bacteria. LPS is a known Toll-like receptor 4 (TLR4) agonist which is associated with signaling cascades that lead to cellular inflammatory events. Studies have shown that short term exposure to LPS cause a significant increase in the amount of phosphorylated NF- κ B in ARVCs. Using patch clamp single-cell electrophysiology we investigate acute effects of LPS exposure on membrane currents of adult rat ventricular cardiomyocytes (ARVC) *in vitro*. Given the necessity of L-type Ca^{2+} currents for proper myocardial function through excitation-contraction coupling (ECC) with links calcium membrane current with myocardial contraction, we investigate changes in L-type Ca^{2+} (I_{CaL}). Due to its role in shaping the early phase of cardiac ventricular action potential we also investigate alterations of the transient outward Ca^{2+} -independent K^{+} current (I_{TO}).

1325-Pos Board B169

Investigating Ion Channel Diseases With Dynamic Action Potential Clamp

Stefan A. Mann, Adam Hill, Jamie I. Vandenberg.

Victor Chang Cardiac Research Institute, Darlinghurst, Australia.

The number of ion channel mutations that are found in genetic screening of patients with cardiac arrhythmias is far outstripping our ability to functionally characterize all the mutations and assess the *in vivo* consequences of each mutant. Currently much time has to be invested in developing mathematical models of mutant ion channels in order to be able to predict their functional significance *in vivo*.

Recently the concept of a dynamic action potential clamp system was introduced that allows integration of electrophysiological recordings from mutant channels into *in silico* models of cellular systems. The current of interest is replaced in the model by current recorded from a mutant channel recorded using voltage-clamp electrophysiology, thereby bypassing the need to formulate a new mathematical model for the mutant ion channel. To simultaneously compute the action potential in the *in silico* model and record the ion current from a cell necessitates the use of a real time operating system so that the real cell can be clamped at the membrane potential of the virtual cell at all times (hence the term dynamic action potential clamp). The ionic current flow through the expressed channels would in turn contribute to changes in the membrane potential of the virtual cell model, so integrating the mutant channels into the virtual cell. We have developed a system where gating models are formulated using the graphical approach offered by Simulink / Matlab. Representation of the models in this way greatly simplifies the user interface compared to standard programming languages, making the system accessible to the less computer-savvy. In the future we will use this system to study the effects of ion channel mutations on the cardiac function cardiac action potential.

1326-Pos Board B170

Estimating Contribution Of Individual Ionic Components To The Cardiac Pacemaker Potential

Yukiko Himeno¹, Chae Young Cha¹, Takao Shimayoshi²,

Yasuhiro Nakamura¹, Jian-Wu Wang¹, Akinori Noma^{3,1}, Nobuya Inagaki¹.

¹Graduate School of Medicine, Kyoto University, Kyoto, Japan, ²ASTEM Research Institute of Kyoto, Kyoto, Japan, ³Faculty of Lifescience, Ritsumeikan University, Kusatsu, Japan.

Action potentials are generated by complicated interactions of various ionic channels and transporters through variations of membrane potential and/or the intracellular Ca^{2+} concentrations. It is still difficult to isolate the contribution of individual current from the secondary effect of modified channel activities. For example, if an ion channel is blocked, the pacemaker activity is drastically changed from the control time course. We have proposed a theoretical method to visualize the contribution of each current in the simulation study. Namely an instantaneous equilibrium potential, the lead potential (V_L), was calculated along the time course of pacemaker potential. V_L is given as,

$$V_L = (\Sigma G_X E_X - \Sigma I_{\text{pump}}) / \Sigma G_X,$$

where G_X and E_X are whole cell membrane conductance and reversal potential for ion X. Contribution of a given current system is evaluated by comparing V_L obtained by fixing the gating parameter of a given current system with the control V_L . In the present study, the gating parameters are fixed during slow diastolic depolarization and action potential repolarization.

It is revealed that the initial phase of the slow diastolic depolarization is mostly attributable to deactivation of the rapid component of the delayed rectifier K^{+} current (I_{Kr}). Then, the major inward currents, the hyperpolarization-activated current (I_f) and the sustained inward current (I_{st}) are activated to depolarize the membrane further to reach the threshold potential of the L-type Ca^{2+} channel current (I_{CaL}). Activation of I_{CaL} initiates generation of action potentials. Deactivation of I_{CaL} and activation of I_{Kr} and the slow component of the delayed rectifier K^{+} current (I_{Ks}) occur during repolarizing phase. Contribution of I_{st} activation to the repolarization is much larger than that of I_f . Activation of I_f , I_{st} , I_{CaL} and I_{Ks} through phosphorylation during β 1-adrenergic stimulation failed to modify each contribution dramatically.

1327-Pos Board B171

Extracellular Zinc Enhances Cardiomyocyte Relaxation Function in Diabetic Rats

Ting Yi, Zengyi Chen, Peter C. Van Buren, Bradley M. Palmer.

University of Vermont, Burlington, VT, USA.

Diabetes mellitus (DM) leads to a cardiomyopathy in humans and rodent models. Interestingly, the chronic infusion of zinc in DM mice prevents the development of the cardiomyopathy. To elucidate the possible mechanism underlying this observation, we examined the effects of extracellular zinc ion (Zn^{2+}) on cardiomyocyte function in DM rats ($n=5$) compared to non-DM controls (Ctrl, $n=5$). Both the DM and Ctrl rats were hypothyroid, which assured similar upregulation of β -myosin heavy chain (β -MHC) in both populations. Isolated cardiomyocytes were electrically stimulated at 2, 4, 6 and 6.5 Hz, maintained at 35°C and exposed to 1.2 mM extracellular Ca^{2+} . Sarcomere shortening and relengthening dynamics were monitored using a video-based Fourier-transform technique. Without extracellular Zn^{2+} peak shortening as a fraction of diastolic sarcomere length was statistically greater ($P<0.05$) in the DM ($6.56 \pm 0.99\%$; $n=16$) compared to Ctrl ($5.27 \pm 1.91\%$; $n=20$) at 2 Hz, but not at higher frequencies. Time to 50% return to diastolic sarcomere length was not statistically different between the groups at every pacing frequency. Exposure to 12 μM extracellular Zn^{2+} significantly reduced ($P<0.001$) peak shortening in both the DM and Ctrl at all frequencies. There was a strong trend ($P=0.070$) toward Zn^{2+} exposure significantly shortening the time to 50% return in the DM but not in the Ctrl, as revealed by repeated-measures ANOVA. Similar trends were found for time to peak shortening ($P=0.087$) and time to 10% return ($P=0.048$). These results suggest that extracellular Zn^{2+} inhibits cardiomyocyte contractile function independent of DM, perhaps by competing with intracellular Ca^{2+} , and enhances cardiomyocyte diastolic function in the DM. The enhanced sensitivity of the DM to the relaxing effects of Zn^{2+} may underlie the protective effects of Zn^{2+} against diabetic cardiomyopathy.

1328-Pos Board B172

Chronic Iron-overload Causes Sinus Bradycardia By Altering Electrical Activity In Sinoatrial Node Myocytes

Michael Sellan, Robert A. Rose, Carlo Cifelli, Scott P. Heximer,

Peter H. Backx.

University of Toronto, Toronto, ON, Canada.

Treatment of blood disorders such as thalassemias require constant blood transfusions that cause iron-overload leading to iron-mediated cardiomyopathy, which is characterized by contractile dysfunction and electrical disturbances, especially bradyarrhythmias. In this study we explored the cellular mechanisms underlying iron-mediated bradycardia by examining the effects of chronic iron-overload (CIO; 0.6 mg/g iron-dextran 3 days/week for 4 weeks by IP injection) on heart rate and sinoatrial node (SAN) function in mice. As expected, heart rate (assessed with telemetry electrocardiograms), was lower ($p<0.001$) in CIO mice (509 ± 21 beats/min; $n=5$) compared to controls (601 ± 12 beats/min; $n=6$, dextrose injected). To examine intrinsic SAN function, heart rate was further studied in anesthetized mice following autonomic nervous system blockade with propranolol (10 mg/kg IP) and atropine (1 mg/kg IP), as