We suggest, therefore, that 6-OHDA may release endogenous dopamine in an active form in the brain to cause the observed hypothermia.

M.A.S. was a Beit Memorial Research Fellow and N.J.U. held U.S. Public Health Service Post-doctoral Fellowship No. 5 FO 2 MH 35810-02PE.

REFERENCES

EVETTS, K. D. & IVERSEN, L. L. (1970). Effects of protriptyline on 6-hydroxydopamine induced depletion of catecholamines in the rat brain. *J. Pharm. Pharmac.*, in the Press.

URETSKY, N. J. & IVERSEN, L. L. (1970). Effects of 6-hydroxydopamine on catecholamine containing neurones in the rat brain. J. Neurochem., 17, 269-278.

Comparative assays using gastric acid secretion: a simple application of non-linear curve fitting (T)

J. W. BLACK, M. E. PARSONS and M. THOMPSON, Smith, Kline & French Research Institute, Welwyn Garden City, Hertfordshire

Mechanism of extraction of angiotensin II in coronary and renal circulations (T)

G. ANGLES D'AURIAC, P. MEYER, M. J. OSBORNE* and M. WORCEL (introduced by Y. S. BAKHLE), Hypertension Laboratory, Hôpital Broussais, Paris 14, France

Conversion of angiotensin I to angiotensin II in isolated strips of arterial smooth muscle (T)

J. W. AIKEN* and J. R. VANE, Department of Pharmacology, Institute of Basic Medical Sciences, Royal College of Surgeons of England, London

Angiotensin effect on biosynthesis of noradrenaline (T)

D. DAVILA and P. A. KHAIRALLAH* (introduced by J. R. VANE), Research Division, Cleveland Clinic, Cleveland, Ohio, U.S.A.

A method for measuring the effectiveness of drugs on platelet thrombus formation in vivo (T)

NICOLA BEGENT* and G. V. R. BORN, M.R.C. Thrombosis Research Group, Department of Pharmacology, Royal College of Surgeons, Lincoln's Inn Fields, London W.C.2