These investigations were supported in part by grants from the Medical Research Council and the Mental Health Research Fund.

REFERENCES

- ANDEN, N.-E. (1966). On the function of the nigrostriatal pathway. In: Mechanisms of Release of Biogenic Amines, ed. von Euler, U.S., Rosell, S. & Uvnas, B., pp. 357-359. Oxford: Pergamon
- ANDEN, N.-E., Roos, B.-E. & WERDINIUS, B. (1964). Effects of chlorpromazine, haloperidol, and reserpine on the levels of phenolic acids in rabbit corpus striatum. *Life Sci.*, Oxford, 3, 149-158.
- Arbuthnott, G. W., Crow, T. J., Fuxe, K. & Ungerstedt, U. (1970). Behavioural effects of
- stimulation in the region of the substantia nigra. J. Physiol., Lond., 210, 61-62P.

 HILLARP, N.-A., FUXE, K. & DAHLSTROM, A. (1966). Demonstration and mapping of central neurones containing dopamine, noradrenaline, and 5-hydroxytryptamine and their reactions to psycho-
- Containing dopamine, noradrenaline, and 5-hydroxytryptamine and their reactions to psychopharmaca. Pharmac. Rev., 18, 727-741.
 Jonas, W. & Scheel-Kruger, J. (1969). Amphetamine-induced stereotyped behaviour correlated with the accumulation of o-methylated dopamine. Arch. int. Pharmacodyn., 177, 379-389.
 O'Keefe, R., Sharman, D. F. & Vogt, M. (1970). Effect of drugs used in psychoses on cerebral dopamine metabolism. Br. J. Pharmac., 38, 287-304.
 Randrup, A. & Munkvad, I. (1967). Stereotyped activities produced by amphetamine in several animal species and Man. Psychopharmacologia, 11, 300-310.

Selective vasoconstrictor effect of atropine on the parametrial arteries in the pregnant guinea-pig (T)

C. Bell, Department of Pharmacology, Institute of Basic Medical Sciences, Royal College of Surgeons, Lincoln's Inn Fields, London, WC2A 3PN