

Reduction of locomotor activity in mice by dopamine- β -hydroxylase inhibitors — evidence against the involvement of non-specific irritation

A.C. DOLPHIN, P.N.C. ELLIOTT & P. JENNER

Department of Neurology, Institute of Psychiatry and King's College Hospital Medical School, Denmark Hill, London SE5 8AF

Dopamine- β -hydroxylase (DBH) inhibitors are widely used as research tools to distinguish between dopaminergic and noradrenergic mechanisms. However, it has been suggested (Moore, 1969; Thornburg & Moore, 1973) that the effects of DBH inhibitors on reducing spontaneous motor activity are not causally related to the depletion of brain noradrenaline which they produce, but to the stress provoked by local irritation which is reflected in the rise in blood glucose and plasma corticosterone which occurs after their intraperitoneal administration (Thornburg & Moore, 1971).

We have tested this hypothesis by comparing two DBH inhibitors with two known irritants, for their ability to produce irritation and their effect on motor activity. The two DBH inhibitors used were 1,1-dimethyl-3-phenyl-2-thiourea (U10,157) and bis(4-methyl - 1 - homopiperazinylthiocarbonyl)-disulphide (FLA-63) in doses which are known to inhibit cerebral DBH. The two irritants used were colloidal carrageenan and kaolin. Their effects were compared on the induction of mouse

paw oedema after subplantar administration, and in stimulation of peritoneal exudation after intraperitoneal administration as measured by leakage of intravenous Evans' Blue dye into the peritoneal cavity. Their effects were also compared on apomorphine and L-DOPA-induced motor activity. The results show that although FLA-63 and carrageenan in the concentrations used have a similar ability to produce inflammation and intraperitoneal irritation, carrageenan has no effect on motor activity produced by L-DOPA; neither the irritants nor the DBH inhibitors affect apomorphine-induced locomotor activity.

We conclude that although the DBH inhibitors tested are irritants, they are producing their effect on motor activity by a more specific direct mechanism which we believe to be related to depletion of cerebral noradrenaline.

References

- MOORE, K.E. (1969). Effects of disulfuran and diethyl-dithiocarbamate on spontaneous locomotor activity and brain catecholamine levels in mice. *Biochem. Pharmacol.*, **18**, 1627-1634.
- THORNBURG, J.E. & MOORE, K.E. (1971). Stress-related effects of various inhibitors of catecholamine synthesis in the mouse. *Arch. Int. Pharmacodyn. Thér.*, **194**, 158-167.
- THORNBURG, J.E. & MOORE, K.E. (1973). The relative importance of dopaminergic and noradrenergic neuronal systems for the stimulation of locomotor activity induced by amphetamine and other drugs. *Neuropharmacol.*, **12**, 853-866.

A pithed rabbit preparation for stimulation of different segments of the autonomic outflow

J.C. McGRATH & J.E. MacKENZIE

Department of Anaesthesia, University of Glasgow

Gillespie, MacLaren & Pollock (1970) described a method of stimulating different segments of the autonomic outflow from the spinal cord to various organs in the pithed cat and rat. We have modified this method for use in the rabbit. Since the geometry of the rabbit skull does not permit pithing via the orbit as in the cat and rat, this has been accomplished through a trephine-hole.

New Zealand white rabbits (3.0-3.5 kg) were anaesthetized with halothane (3%) and were decerebrated following placement of cannulae in

the trachea, both carotid arteries (left, for arterial and left ventricular pressure; right, for thermistor for cardiac output), and a jugular vein (for administration of drugs). Halothane was then discontinued, gallamine (1 mg/kg i.v.) was given and mechanical ventilation with 100% O₂ (Harvard Respirator) was adjusted to give an end-tidal CO₂ of 3-4% (Beckman Medical Gas Analyser LB-2). Pithing was carried out using a teflon covered stainless steel rod identical to that used for the cat by Gillespie *et al.* (1970). The resulting preparation had a resting arterial pressure of 48 ± 3 mmHg systolic and 26 ± 2 mmHg diastolic, and heart rate of 230 ± 4 min⁻¹ ($n = 9$), which together with blood gas values remained stable over a period of at least 6 hours. As in the previous study in the rat and cat (Gillespie *et al.*, 1970), reproducible cardiovascular responses could be obtained to stimulation of the autonomic

outflow (via the tip of the movable pithing rod) or to injected drugs.

In each animal, left ventricular pressure (and hence dP_v/dt) was also monitored via a cannula placed within that in the carotid artery. Cardiac output was measured by thermal dilution, enabling calculation of peripheral resistance. This preparation can assist in identifying the peripheral cardiovascular effects of drugs.

J.C.McG. received a Wellcome Trust Interdisciplinary Fellowship, and J.E.M. an M.R.C. Fellowship.

Reference

GILLESPIE, J.S., MacLAREN, A. & POLLOCK, D. (1970). A method of stimulating different segments of the autonomic outflow from the spinal column to various organs in the pithed cat and rat. *Br. J. Pharmac.*, **40**, 257-267.

The episcopes for large screen projection of tracings on opaque media during their recording for demonstrations

J.B.E. BAKER

Department of Pharmacology, Charing Cross Hospital Medical School, London W6

Measurement of inflammatory exudation and local blood flow changes in rabbit skin, using [131 I]-albumin and 133 Xe

M.J. PECK & T.J. WILLIAMS (introduced by G.P. LEWIS)

Department of Pharmacology, Institute of Basic Medical Sciences, Royal College of Surgeons of England, London WC2A 3PN

Measurement of blood flow in rabbit skin homografts and autografts using a 133 Xe-clearance technique

G.P. LEWIS, M.J. PECK, T.J. WILLIAMS & BEVERLEY A. YOUNG

Department of Pharmacology, Royal College of Surgeons, London WC2A 3PN

Prostaglandin production by rat polymorphonuclear leucocytes

ELAINE McCALL & L.J.F. YOULTEN

Department of Pharmacology, Royal College of Surgeons, London WC2A 3PN