

A sponge implantation test in the rat as a model for screening anti-inflammatory activity

J.P. BOLAM, A.W. FORD-HUTCHINSON,
M.J.H. SMITH & J.R. WALKER

Department of Biochemical Pharmacology, King's College Hospital Medical School, Denmark Hill, London SE5 8RX

Despite the large number of tests currently available to screen potential anti-inflammatory and antirheumatic drugs (Swingle, 1974) there are very few based on leucocyte migration into inflammatory exudates. The implanted sponge technique (Saxena, 1960) seems worthy of more general use since it is a simple procedure, allows a variety of measurements to be performed on exudates at various time intervals after implantation and is suitable for the assessment of inhibitory effects on granuloma formation (Ford-Hutchinson, Smith, Elliott, Bolam, Walker, Lobo, Badcock, Colledge & Billimoria, 1975; Wiener, Wiener, Urivetzky, Shafer, Isenberg, Janov & Meilman, 1975; Clarke, Vernon-Roberts & Currey, 1975).

The method may be of particular value in the assessment of substances which interfere with either the formation or action of leucotactic factors derived from complement since rats depleted of complement components by purified cobra venom factor showed a marked inhibition of cellular migration into the sponge exudate (Wiener, Lendvai, Rogers, Urivetsky & Meilman, 1973).

Some applications of the sponge method to the

study of the sites of action of conventional antirheumatic drugs and the relationship between the involvement of blood platelets, leucotaxis and prostaglandins in the inflammatory exudates will be presented.

This work was supported by the King's College Hospital and Medical School Research Committee, the Wates Foundation and the National Research Development Corporation.

References

- CLARKE, A.K., VERNON-ROBERTS, B. & CURREY, H.L.F. (1975). Assessment of anti-inflammatory drugs in the rat using subcutaneous implants of polyurethane foam impregnated with dead tubercle bacilli. *Ann. rheum. Dis.*, **34**, 326–331.
- FORD-HUTCHINSON, A.W., SMITH, M.J.H., ELLIOTT, P.N.C., BOLAM, J.P., WALKER, J.R., LOBO, A.A., BADCOCK, J.K., COLLEDGE, A.J. & BILLIMORIA, F.J. (1975). Effects of a human plasma fraction on leucocyte migration into inflammatory exudates. *J. Pharm. Pharmacol.*, **27**, 106–112.
- SAXENA, P.N. (1960). Effect of drugs on early inflammatory reactions. *Arch. int. Pharmacodyn. Ther.*, **126**, 228–237.
- SWINGLE, K.F. (1974). Evaluation for anti-inflammatory activity. In *Anti-inflammatory agents, chemistry and pharmacology*. Vol. II, ed. Scherrer, R.A. & Whitehouse, M.W. pp. 33–122. New York: Academic Press.
- WIENER, S., LENDVAI, S., ROGERS, B., URIVETSKY, M. & MEILMAN, E. (1973). Non-immune chemotaxis *in vivo*. *Am. J. Pathol.*, **73**, 807–816.
- WIENER, S.L., WIENER, R., URIVETSKY, M., SHAFER, S., ISENBERG, H.D., JANOV, C. & MEILMAN, E. (1975). The mechanism of action of a single dose of methylprednisolone on acute inflammation *in vivo*. *J. clin. Invest.*, **56**, 679–689.

Pethidine pharmacokinetics in dog: dose and time studies

M.R. BENDING, P.N. BENNETT &
M. ROWLAND

Department of Clinical Pharmacology, Royal Postgraduate Medical School, London and Department of Pharmacy, University of Manchester

Data in man suggest that pethidine is highly cleared predominantly by metabolism and may therefore exhibit a 'first pass' effect on oral dosing (Mather, Tucker, Pflug, Lindop & Wilkerson, 1975; Chan, Kendall, Wells & Mitchard, 1975). To examine and extend this possibility, the effect of dose and time on

the oral availability and disposition kinetics of pethidine were studied in three female greyhound dogs. In the dose-range study the following doses were administered on separate occasions—1.0, 2.0 and 3.0 mg/kg intravenously (i.v.) and 2.0, 3.0 and 4.0 mg/kg orally (p.o.) by stomach tube. In the time study, each dog received 2.0 mg/kg i.v. eight-hourly for seven doses and on another occasion 3.0 mg/kg p.o. eight-hourly for seven doses.

The terminal half-life of the blood-concentration-time curve increased with dose (means—50 and 72 min at the 1.0 and 3.0 mg/kg i.v. doses and 77 and 103 min at the 2.0 and 4.0 mg/kg p.o. doses). Comparison of the area under the blood-concentration-time curve (AUC) with dose suggested a non-linear relationship with AUC increasing dis-

proportionately at higher doses. Consequently total systemic blood clearance of pethidine (dose i.v./AUC i.v.) fell with increasing dose, the values for the three dogs at the 1.0 and 3.0 mg/kg i.v. doses being 59 and 34, 51 and 42, 65 and 52 ml min⁻¹ kg⁻¹ respectively. However a plot of β (the terminal disposition rate constant) \times AUC against dose (i.v.) was linear, indicating that volume of distribution was a constant (values for each dog—4.2, 4.4 and 3.6 l/kg). This indicates that changes in clearance are reflected in changes in β . In two dogs a plot of $\beta \times$ AUC against dose (p.o.) was linear and the systemic availability of pethidine in these dogs, calculated as the ratio of the slopes of the p.o. and i.v. regression lines was 0.41 and 0.38. In the third dog $\beta \times$ AUC against dose was non-linear: thus availability changed with dose in this dog between 0.57 and 0.64 at the 2.0 and 3.0 mg/kg p.o. doses.

The findings in single dose administrations of pethidine were reproducible when further single doses were given. The clearance of pethidine in these animals

approached liver blood flow and the low oral availability is compatible with significant 'first pass' hepatic metabolism of pethidine. In the repeated dose study for both i.v. and p.o. administration, no accumulation of pethidine occurred and the kinetics of pethidine did not change with time. The data suggest that while pethidine may exhibit dose-dependent disposition kinetics, there is no evidence for time dependent kinetics at the doses studied.

References

- CHAN, K., KENDALL, M.J., WELLS, W.D.E. & MITCHARD, M. (1975). Factors influencing the excretion and relative physiological availability of pethidine in man. *J. Pharm. Pharmacol.*, **27**, 235–241.
 MATHER, L.E., TUCKER, G.T. PFLUG, E.A., LINDOP, M.J. & WILKERSON, C. (1975). Meperidine kinetics in man. Intravenous injection in surgical patients and volunteers. *Clin. Pharmacol. Therap.*, **17**, 21–30.

Fluorimetric determination of ampicillin and cephalixin

R.H. BARBHAIYA & P. TURNER

Department of Clinical Pharmacology, St Bartholomew's Hospital, London EC1A 7BE

A fluorimetric method for quantitative analysis of two β -lactam antibiotics ampicillin and cephalixin has been developed, based upon the production of fluorescent derivatives formed by alkaline hydrolysis with 2N NaOH and heating at 100°C. The rate of formation of fluorescent material is enhanced by addition of formaldehyde in the buffer. The fluorescent compounds exhibit uncorrected excitation and emission spectra at 345 and 420 nm respectively. The structural similarities between the side chains of ampicillin and cephalixin and their identical fluorescent spectra suggest that both the antibiotics could be forming a similar fluorescent derivative. Although the precise chemical identity of the fluorescent derivative has not yet been determined, there is evidence to suggest that it may be the diketopiperazine derivative (Cohen, Funke & Puar, 1973; Indelicato, Narvilas, Pfeiffer, Wheeler & Wilham, 1974, 1974; Yamana, Tsuji, Kanayama & Nakono, 1974). An assay procedure

based on these observations permits detection of less than 0.1 μ g/ml of ampicillin and cephalixin in aqueous solution and plasma. Because of greater sensitivity, the proposed method is more suitable for clinical studies of these antibiotics than the previously described acid hydrolysis method for ampicillin (Jusko, 1971).

The results obtained with the fluorimetric assay have been compared with the results obtained in microbiological assay. The fluorimetric assay has advantage over microbiological assay in terms of ease and speed of performance, reproducibility and sensitivity.

References

- COHEN, A.I., FUNKE, P.T. & PUAR, M.S. (1973). Alkaline degradation product of cephradine. *J. Pharm. Sci.*, **62**, 1559–1561.
 INDELICATO, J.M., NARVILAS, T.T., PFEIFFER, R.R., WHEELER, W.J. & WILHAM, W.L. (1974). Substituent effect upon the base hydrolysis of penicillins and cephalosporins. *J. Med. Chem.*, **17**, 523–527.
 JUSKO, W.J. (1971). Fluorometric analysis of ampicillin in biological fluids. *J. Pharm. Sci.*, **60**, 728–732.
 YAMANA, T., TSUJI, A., KANAYAMA, K. & NAKONO, O. (1974). Comparative stabilities of cephalosporins in aqueous solutions. *J. Antibiot.*, **27**, 1000–1002.