

Assay of liver membrane insulin receptors in obese, hyperglycaemic (ob/ob) mice: stimulant effect of an oral antidiabetic sulphonylurea drug

B.D. GREENSTEIN

Department of Pharmacology, St. Thomas's Hospital Medical School, London SE1 7EH

A reliable *in vitro* assay of insulin receptors has been developed, and a preliminary study of the *in vivo* effects of an antidiabetic sulphonylurea drug has been attempted.

Purified membranes were prepared from livers of male ob/ob 6 week-old mice and from normal, lean littermates by the method of Cuatrecasas (1972), using a buffer consisting of (mM): Tris HCl 50, MgCl₂ 10, NH₄Cl 5, bovine serum albumin 0.1% (w/v), pH 7.9. The homogeneity of the preparation was checked by electron microscopy. The protein concentration of the microsomal fraction was 1.9 ± 0.06 mg/ml: $n = 9$ (mean \pm s.d.) for obese mice, and 1.38 ± 0.04 mg/ml: $n = 9$, for lean mice. Purified membranes were incubated at 13°C with 1.5 nM [¹²⁵I] insulin (Radiochemical Centre, Amersham) alone or in the presence of unlabelled insulin (50 µg/ml) to distinguish between receptor and non-saturably bound hormone. Incubates were layered over cushions of (0.3 M) sucrose in buffer in Beckman microfuge tubes and centrifuged. The supernatant was aspirated, tip of the tube cut off, and the radioactivity in the pellet counted.

Equilibrium between specifically bound and free [¹²⁵I] insulin was attained within 60 min of incubation at 13°C, and there was no apparent loss of binding sites during this period. Higher incubation temperatures appeared to destroy the binding sites. The binding affinity and capacity were determined from linear Lineweaver-Burke plots after incubating membranes with a range of concentrations of [¹²⁵I] insulin (0.02–1.5 M) alone or in the presence of excess unlabelled insulin. The molar dissociation constant (K_d) of the reaction and the binding capacity were, for ob/ob mice, $2.22 \pm 0.6 \times 10^{-10}$ M: $n = 4$ (mean \pm s.e. mean), and $6.23 \pm 3.6 \times 10^8$ binding sites per mg membrane protein respectively, and for lean mice, $3.94 \pm 0.7 \times 10^{-10}$ M: $n = 6$, and $5.47 \pm 1.3 \times 10^9$ binding sites respectively.

After i.p. injection of the antidiabetic sulphonylurea drug Gliquidon (4 µg/g bodyweight; Boehringer Ingelheim Ltd) twice daily in 0.1 ml of lactamide for 7 days, there was a five-fold increase in liver membrane insulin receptors from ob/ob mice, when compared with the effects of the solvent alone. Bodyweights were unchanged during the course of the treatment.

It is concluded that insulin receptors can be measured under conditions which maintain the stability of the complex. Also, preliminary experiments suggest that sulphonylurea derivatives may exert their antidiabetic effects, at least in part, by increasing liver insulin receptors.

Reference

CUATRECASAS, P. (1972). Isolation of the insulin receptor of liver and fat-cell membranes. *Proc. Nat. Acad. Sci. USA.*, **69**, 318–322.

Comparison of the binding characteristics of tritiated opiates and opioid peptides

MAUREEN G.C. GILLAN,
H.W. KOSTERLITZ &
S.J. PATERSON

Unit for Research on Addictive Drugs, University of Aberdeen, Aberdeen

Opiate receptor binding in homogenates of guinea-pig brain minus cerebellum was studied by a modification of the method of Pert & Snyder (1973), using [³H]-D-Ala²-D-Leu⁵-enkephalin (44 Ci/mmol); [³H]-D-Ala²-leucine-enkephalin amide (17.6 Ci/mmol); [³H]-D-Ala²-methionine-enkephalin amide (15 Ci/mmol); [³H]-etorphine (36 Ci/mmol); [³H]-dihydromorphine (81 Ci/mmol) and [³H]-morphine (28 Ci/mmol) as primary ligands. The brain tissue was homogenized in

Tris buffer (pH 7.4, 0°C), centrifuged at 49,000 g for 10 min; the pellet was resuspended in Tris buffer (pH 7.4, 37°C) and incubated at 37°C for 45 min. The homogenate was then centrifuged again and the pellet resuspended in Tris buffer (pH 7.4, 25°C) to give a final concentration of 1 g tissue/100 ml. Aliquots of the homogenate were incubated with various concentrations of the primary ligands, without addition of Na⁺, for 40 min at 25°C. Specific binding was the difference in counts obtained in the presence and absence of 50 nM of the antagonist (–)-2-(3-furylmethyl)-5,9-diethyl-2'-hydroxy-6,7 - benzomorphan (MR 2266). The kinetic parameters of the binding were determined by Scatchard analysis.

There is apparently only one binding site for D-Ala²-D-Leu⁵-enkephalin in guinea-pig brain homogenate. At this site, the ligand has a K_d of 1.27 ± 0.16 nM ($n = 5$) and the number of binding sites corresponds to 7.4 ± 0.31 pmol/g wet wt ($n = 5$). Under the same conditions, only one binding site was detected for D-Ala²-leucine-enkephalin amide or D-Ala²-