

The effect of metiamide on protracted anaphylaxis in the guinea-pig

M.A. CHADWICK & P. GOADBY

Department of Pharmacology, School of Pharmacy, Sunderland Polytechnic, Sunderland, SR1 3SD

The intravenous injection of antigen in actively sensitized guinea-pigs results in death due to respiratory failure which occurs within a few minutes of injection. However, the injection of antigen by intraperitoneal or subcutaneous route usually results in protracted anaphylactic shock a major effect of which is congestion and/or haemorrhaging of the gastrointestinal tract (Stone, 1959). In the present experiments we have studied the effect of the histamine H_2 -receptor antagonist, metiamide (Black, Duncan, Emmett, Ganellin, Hesselbo, Parsons & Wyllie, 1973) upon the reaction.

Guinea-pigs (500–700 g), sensitized to egg albumen at least 3 weeks previously, were pretreated with mepyramine (2 mg/kg) 1 h before challenge by the i.p. administration of egg albumen (40 mg). The animals showed some respiratory distress but did not die as a result of respiratory failure. After 30 min they became lethargic and showed piloerection. Those animals that died (7/15) did so between 30 and 120 min after the antigen injection and the remainder were killed after 3 hours.

Upon examination, little evidence of lung damage

was seen, but the stomach and large intestine showed signs of hyperaemia and localised haemorrhagic patches were apparent. The volume and pH of the contents were measured and the mucosa examined for damage using a dissection microscope. Mucosal damage varied in extent from small petechiae to larger haemorrhagic lesions located in the main body of the stomach. Rarely were lesions seen in the fundus or pyloric antrum. Pretreatment with metiamide (10 mg/kg s.c.) 30 min before challenge reduced the severity of the lesions but did not prevent them.

In the anaesthetized guinea-pig, histamine infusions ($0.5\text{--}20\text{ }\mu\text{g kg}^{-1}\text{ min}^{-1}$ i.v.) caused an increase in gastric acid secretion but the introduction of egg albumen, i.p. or i.v. in sensitized animals failed to increase gastric acid secretion. Thus protracted anaphylaxis following i.p. egg albumen results in gastric mucosal damage that is not associated with an increase in gastric acid secretion.

References

- BLACK, J.W., DUNCAN, W.A.M., EMMETT, J.C., GANELLIN, C.R., HESSELBO, T., PARSONS, M.E. & WYLLIE, J.H. (1973). Metiamide—an orally active histamine H_2 -receptor antagonist. *Agents and Actions*, **3**, 133–137.
STONE, S.H. (1959). Acute and protracted anaphylactic shock in guinea-pigs after subcutaneous eliciting injection of antigen. *J. Immunol.*, **82**, 138–145.

Evidence *in vivo* for a 5-hydroxytryptamine link in dopamine-receptor-mediated hypothermia in the rat

B. COX†, R.W. KERWIN, T.F. LEE† & C.J. PYCOCK

Department of Pharmacology, University of Manchester, Manchester M13 9PT† & Department of Pharmacology, University of Bristol, Bristol BS8 1TD

In rats, intrahypothalamic injection of dopamine agonists causes a fall in core temperature which appears to be mediated through tail skin vasodilation (Cox & Lee, 1977). An involvement of 5-hydroxytryptamine (5-HT) in this response has been suggested (Maj & Przewlocka, 1975; Przewlocki, 1977), but there is no general agreement (Grabowska, Michaluk & Antkiewicz, 1973). Therefore in this study, we have investigated the possibility of a 5-HT link.

Male Sprague-Dawley rats were used at the ambient temperature of 17°C. Drugs were injected

into the preoptic anterior hypothalamus through previously implanted guide cannulae in a dose volume of 1 μl . Core temperature was measured by means of a rectal probe and tail skin temperature by a small strap-on thermistor.

Intrahypothalamic injection of either dopamine or 5-HT caused a fall in core temperature which was blocked by intrahypothalamic pretreatment with either methysergide or cyproheptadine (Figure 1). Oxotremorine-induced hypothermia was unaffected by these antagonists. Haloperidol antagonized dopamine but not 5-HT (Figure 1). These findings support the suggestion of a 5-HT link.

Further evidence for 5-HT involvement came from studies with 5,6-dihydroxytryptamine (5,6-DHT), which was injected bilaterally into the preoptic area two weeks prior to the experiment in a dose of 8 μg in 2 μl and which significantly depleted hypothalamic 5-HT ($P < 0.01$) without affecting hypothalamic dopamine. This pretreatment antagonized dopamine, but was without effect on intrahypothalamic 5-HT (Figure 1). Furthermore, 5,6-DHT treated rats sub-