## British Journal of Pharmacology

## **PAPERS**

- 447 HERMANSSON, L.-E. & JOHANSSON, P. Postsynaptic α<sub>1</sub>- and α<sub>2</sub>-adrenoceptors in the vascular system of the herring gull, *Larus argentatus*
- HOLZER, P., HOLZER-PETSCHE, U. & LEANDER,
  S. A tachykinin antagonist inhibits gastric emptying and gastrointestinal transit in the rat
- 461 ANDREWS, P.L.R. & WOOD, K.L. Systemic baclofen stimulates gastric motility and secretion via a central action in the rat
- 469 ASHCROFT, S.J.H., HARRISON, D.E., POJE, M. & ROCIC, B. Structure-activity relationships of alloxanlike compounds derived from uric acid
- 473 HATA, F., ISHIDA H. & KONDO, E. Dopamineinduced amylase secretion from rat parotid salivary gland in vitro: an effect mediated via noradrenergic and cholinergic nerves
- 485 RICE, W.R. & SINGLETON, F.M. P<sub>2</sub>-purinoceptors regulate surfactant secretion from rat isolated alveolar Type II cells
- 493 LEFF, P., MARTIN, G.R. & MORSE, J.M. The classification of peripheral 5-HT<sub>2</sub>-like receptors using tryptamine agonist and antagonist analogues
- 501 CUMMINGS, S.A., GROSZMANN, R.J. & KAUMANN, A.J. Hypersensitivity of mesenteric veins to 5-hydroxytryptamine-and ketanserin-induced reduction of portal pressure in portal hypertensive rats
- 515 BALLEJO, G., CALIXTO, J.B. & MEDEIROS, Y.S. In vitro effects of calcium entry blockers, chlorpromazine and fenoterol upon human pregnant myometrium contractility
- 525 ALOSACHIE, I. & GODFRAIND, T. Role of cyclic GMP in the modulation by endothelium of the adrenolytic action of prazosin in the rat isolated aorta
- 533 FINET, M. & GODFRAIND, T. Selective inhibition by ethylisopropylamiloride of the positive inotropic effect evoked by low concentrations of ouabain in rat isolated ventricles
- 539 BOURUT, C., CHENU, E., GODENÈCHE, D., MADELMONT, J.-C., MARALT, R., MATHÉ, G. &

- MEYNIEL, G. Cytostatic action of two nitrosoureas derived from cysteamine
- 547 BLACK, J.W., GERSKOWITCH, V.P., LEFF, P. & SHANKLEY, N.P. Analysis of competitive antagonism when this property occurs as part of a pharmacological resultant
- 557 MARTIN, W., MORGAN, R.O., SMITH, J.A. & WHITE, D.G. Atriopeptin II-induced relaxation of rabbit aorta is potentiated by M & B 22,948 but not blocked by haemoglobin
- 563 MARTIN, W., SMITH, J.A. & WHITE, D.G. The mechanisms by which haemoglobin inhibits the relaxation of rabbit aorta induced by nitrovasodilators, nitric oxide, or bovine retractor penis inhibitory factor
- 573 ARIMURA, H. & IKEMOTO, Y. Action of enflurane on cholinergic transmission in identified *Aplysia* neurones
- 583 LARGE, W.A. & SIM, J.A. A comparison between mechanisms of action of different nicotinic blocking agents on rat submandibular ganglia
- 593 BLACKBURN, T.P., BORKOWSKI, K.R., FRIEND, J. & RANCE, M.J. On the mechanisms of κ-opioidinduced diuresis
- 599 FONTAINE, J., GRIVEGNEE, A.R. & ROBBERE-CHT, P. Evidence against VIP as the inhibitory transmitter in non-adrenergic, non-cholinergic nerves supplying the longitudinal muscle of the mouse colon
- 603 BARRON, E., MARSHALL, R.J., MARTORANA, M. & WINSLOW, E. Comparative antiarrhythmic and electrophysiological effects of drugs known to inhibit calmodulin (TFP, W7 and bepridil)
- 613 BOURA, A.L.A., HODGSON, W.C. & KING, R.G. Changes in cardiovascular sensitivity of alloxan-treated diabetic rats to arachidonic acid
- 619 GIBB, A.J. & MARSHALL, I.G. Nicotinic antagonists produce differing amounts of tetanic fade in the isolated diaphragm of the rat

## **MEETINGS CALENDAR**