



Erratum

Br. J. Pharmacol. (1996), **117**, 817–822.

L.A. Shaw & S.J. Coker Suppression of reperfusion-induced arrhythmias with combined administration of 5-HT₂ and thromboxane A₂ antagonists.

In the above article, Table 2 was duplicated and Table 1 was omitted due to a typesetting error. Table 1 appears below.

Table 1 Haemodynamics in the high dose reperfusion study: heart rate, systolic blood pressure (BP) and diastolic BP 1 min before and 10 min after drug or vehicle administration, i.e. immediately prior to coronary artery occlusion

	Heart rate (beats min ⁻¹)		Systolic BP (mmHg)		Diastolic BP (mmHg)	
	<i>Pre</i>	<i>Post</i>	<i>Pre</i>	<i>Post</i>	<i>Pre</i>	<i>Post</i>
Control	403 ± 11	371 ± 9	91 ± 3	95 ± 5	72 ± 5	74 ± 6
ICI 170,809 1 mg kg ⁻¹	394 ± 11	339 ± 12*	94 ± 5	94 ± 3	67 ± 5	65 ± 4*
ICI 192,605 1 mg kg ⁻¹ min ⁻¹	402 ± 12	360 ± 10	103 ± 6	100 ± 6	79 ± 7	70 ± 5
Both drugs	407 ± 8	337 ± 14*	95 ± 5	80 ± 3*	68 ± 5	54 ± 4*

Values are the mean ± s.e.mean, *n* = 12. **P* < 0.05 compared with corresponding value in the control group, one way analysis of variance and modified *t* test (Bonferroni correction).