

# Effects of methyl *p*-hydroxybenzoate (methyl paraben) on $\text{Ca}^{2+}$ concentration and histamine release in rat peritoneal mast cells

<sup>1</sup>Sanae Fukugasako, <sup>\*</sup><sup>1</sup>Shinichi Ito & <sup>1</sup>Yoshimi Ikemoto

<sup>1</sup>Department of Dental Anesthesiology, Faculty of Dental Science, Kyushu University, Fukuoka 812-8582, Japan

**1** Mechanisms of methyl *p*-hydroxybenzoate (methyl paraben) action in allergic reactions were investigated by measuring the intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) and histamine release in rat peritoneal mast cells (RPMCs).

**2** In the presence or absence of extracellular  $\text{Ca}^{2+}$ , methyl paraben (0.1–10 mM) increased  $[\text{Ca}^{2+}]_i$ , in a concentration-dependent manner. Under both the conditions, methyl paraben alone did not evoke histamine release.

**3** In RPMCs pretreated with a protein kinase C (PKC) activator (phorbol 12-myristate 13-acetate (PMA) 3 and 10 nM), methyl paraben (0.3–3 mM) induced histamine release. However, a high concentration (10 mM) of the agent did not increase the histamine release.

**4** U73122 (0.1 and 0.5  $\mu\text{M}$ ), an inhibitor of phospholipase C (PLC), significantly inhibited the methyl paraben-induced histamine release in PMA-pretreated RPMCs. U73343 (0.5  $\mu\text{M}$ ), an inactive analogue of U73122, did not inhibit the histamine release caused by methyl paraben.

**5** In  $\text{Ca}^{2+}$ -free solution, PLC inhibitors (U73122 0.1 and 0.5  $\mu\text{M}$ , D609 1–10  $\mu\text{M}$ ) inhibited the methyl paraben-induced increase in  $[\text{Ca}^{2+}]_i$ , whereas U73343 (0.5  $\mu\text{M}$ ) did not.

**6** Xestospongine C (2–20  $\mu\text{M}$ ) and 2-aminoethoxydiphenyl borate (30 and 100  $\mu\text{M}$ ), blockers of the inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) receptor, inhibited the methyl paraben-induced increase in  $[\text{Ca}^{2+}]_i$  in  $\text{Ca}^{2+}$ -free solution.

**7** In conclusion, methyl paraben causes an increase in  $[\text{Ca}^{2+}]_i$ , which may be due to release of  $\text{Ca}^{2+}$  from storage sites by  $\text{IP}_3$  via activation of PLC in RPMCs. In addition, methyl paraben possibly has some inhibitory effects on histamine release via unknown mechanisms.

*British Journal of Pharmacology* (2003) **139**, 381–387. doi:10.1038/sj.bjp.0705248

**Keywords:** Methyl *p*-hydroxybenzoate; mast cells; intracellular calcium concentration; histamine release; phospholipase C; protein kinase C; inositol 1,4,5-trisphosphate

**Abbreviations:** 2APB, 2-aminoethoxydiphenyl borate;  $[\text{Ca}^{2+}]_i$ , intracellular  $\text{Ca}^{2+}$  concentration; DMSO, dimethyl sulphoxide;  $\text{IP}_3$ , inositol 1,4,5-trisphosphate; PKC, protein kinase C; PLC, phospholipase C; PMA, phorbol 12-myristate 13-acetate; PSS, physiological salt solution; RPMCs, rat peritoneal mast cells

## Introduction

There have been numerous reports on cases of anaphylactic reactions caused by various drugs (Fisher & More, 1981; Mertes & Laxenaire, 2002). Anaphylactic shock is the type I allergy reaction mediated by IgE antibodies and mast cells. The symptom similar to anaphylactic shock is called an anaphylactoid reaction (Fisher & Pennington, 1982), where the mechanism involves activation of mast cells without IgE antibodies (Stellato & Marone, 1995). A preservative, methyl *p*-hydroxybenzoate (methyl paraben), may be responsible for some cases of anaphylactic shock and anaphylactoid reactions caused by various commercially available medicines (Nagel *et al.*, 1977; Wildsmith *et al.*, 1998). Methyl paraben is nonstimulating and nontoxic, and has a broad antibiotic spectrum. The compound is widely used as a preservative for foods, cosmetics and medicines. Those methyl paraben-containing products caused contact dermatitis and drug hypersensitivity (Larson, 1977; Mowad, 2000), but there has

been no fundamental study on allergic reactions related to methyl paraben.

In an immunological mechanism, degranulation of mast cells is triggered off by the aggregation of high-affinity receptor for the Fc region of IgE ( $\text{Fc}\epsilon\text{RI}$ ) caused by crosslinking of IgE by polyvalent antigens. However, specific IgE antibodies for methyl paraben have not been identified (Kokubu *et al.*, 1989). Simple chemicals such as methyl paraben are incapable of producing sensitization and induction of immediate or delayed hypersensitivity without prior conjugation to carrier molecules, usually proteins. The bound methyl paraben is then considered a hapten, whereas its chemical properties are not clear (Soni *et al.*, 2002).

It was reported that methyl paraben activated the ryanodine receptor  $\text{Ca}^{2+}$  release channel in skeletal muscle terminal cisternae (Cavagna *et al.*, 2000). On the other hand, Teraoka *et al.* (1997) reported that caffeine, an activator of the ryanodine receptor, did not increase the intracellular  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_i$ ) in rat peritoneal mast cells, and Soboloff & Berger (2002) described that ryanodine did not

\*Author for correspondence; E-mail: ito@dent.kyushu-u.ac.jp

significantly increase  $[\text{Ca}^{2+}]_i$  in bone marrow-derived mast cells. However, the lack of stimulatory effects of caffeine and ryanodine on  $\text{Ca}^{2+}$  release does not accordingly indicate the absence of ryanodine receptor in many types of nonexcitable cells (Hosoi *et al.*, 2001). The existence of ryanodine receptor is still under controversy, raising the question as to how methyl paraben affects the intracellular events during the allergic reactions.

In the present study, in order to clarify the mechanism of allergic reactions caused by methyl paraben, we investigated the effects of the agent on the changes in  $[\text{Ca}^{2+}]_i$  and histamine release in RPMCs.

## Methods

### *Mast cell isolation and purification*

Male Sprague–Dawley rats weighing 400–800 g were anaesthetized with diethyl ether, and then killed by bleeding from the carotid arteries. Rat peritoneal mast cells were isolated and purified over Percoll density gradient as previously described (Chan *et al.*, 2000). The peritoneal cavity was injected with the physiological salt solution (PSS) containing BSA ( $0.3 \text{ mg ml}^{-1}$ ). After gentle massage of the rat abdominal region, mixed peritoneal cells (30–35 ml of fluid) were obtained by peritoneal lavage. The mixed peritoneal cells were then washed twice by centrifugation (1100 r.p.m., 5 min,  $4^\circ\text{C}$ ) and were resuspended in 1 ml of PSS. The cell suspension was then mixed with 2 ml of 33% Percoll. BSA-supplemented PSS (1 ml) was then carefully layered over the Percoll-cell mixture. Purification was then performed by centrifugation (3000 r.p.m., 20 min,  $4^\circ\text{C}$ ), which allowed cell separation and gradient formation. Harvesting of the mast cells posed no problem since these cells gathered in a layer at the bottom of the tube, whereas other cells formed a rather compact layer on top of the gradient and could easily be removed by aspiration. The cell fraction was then washed twice in PSS by centrifugation and finally resuspended to the desired cell density in PSS.

### *Intracellular calcium measurements*

Intracellular  $\text{Ca}^{2+}$  images were obtained with the confocal laser scanning microscope (IX70, Olympus). Cells were incubated in PSS containing the acetoxymethyl ester of fluo-3 (fluo-3 AM  $5 \mu\text{M}$ ) for 30 min at room temperature ( $22$ – $25^\circ\text{C}$ ). Cells attached to a glass coverslip were mounted on the bottom of a chamber of  $500 \mu\text{l}$  capacity and placed in the microscope for fluorescence measurements. Cells were excited at 488 nm with an argon laser beam through the objective lens (UplanApo40X, Olympus). Fluo-3 fluorescent images (emission 530 nm) were collected with the scan unit (FVX-SU, Olympus) every 0.42 s. To estimate  $[\text{Ca}^{2+}]_i$ , the mean intensity of cell area except nucleus was calculated with the analysing software (FLUOVIEW FV500, Olympus). The data were expressed as ratios of fluorescence intensity changes ( $F$ ) relative to control values before stimulation ( $F_0$ ), namely  $(F-F_0)/F_0$ .

### *Histamine measurements*

Drugs were applied to cell suspensions ( $10^6$  cells  $\text{ml}^{-1}$ ), making a total of 1 ml solution. The histamine-releasing reactions were

terminated by placing the test tube in ice-cold water for 10 min. Cell suspensions were then centrifuged and divided into fractions of supernatant (0.5 ml) and supernatant plus pellets (0.5 ml), both of which were acidified with perchloric acid ( $50 \mu\text{l}$ ) to abolish histamine breakdown. After both fractions were centrifuged at 1100 r.p.m. for 10 min at  $4^\circ\text{C}$  to remove proteins, secreted histamine was determined by the fluorometric method. Histamine release was expressed as a percentage of the total cell contents. All samples were stored at  $-25^\circ\text{C}$  until the histamine level was measured using the high-performance liquid chromatography (HPLC) postlabel system as previously described (Yamatodani *et al.*, 1985; Horinouchi *et al.*, 1993). The system was composed of an intelligent pump (Hitachi, L-6200), a reaction pump (Hitachi, 655-A-13), a fluorescence spectrophotometer (Hitachi, F-1150), an autosampler (Hitachi, AS-4000), a Chromato-integrator (Hitachi, D-2500) and a  $6 \text{ } \varnothing$ , 15 cm column (Catecholepak, Toyosoda, Tokyo, Japan) warmed at  $50^\circ\text{C}$  by a column oven (Hitachi, L-5020). Each  $100 \mu\text{l}$  of supernatant was injected into the HPLC for each sample. The excitation wavelength used was 340 nm and the emission was 450 nm.

### *Chemicals*

The drugs used were BSA, EGTA, methyl paraben, HEPES, phorbol 12-myristate 13-acetate (PMA), compound 48/80, U73343 and U73122 (Sigma Chemicals, St Louis, U.S.A.), dimethyl sulphoxide (DMSO) (Merck, Darmstadt, Germany), A23187, 2APB, xestospongine C and D609 (Calbiochem, San Diego, CA, U.S.A.), Fluo-3 AM (Dojindo Laboratories, Kumamoto, Japan) and Percoll (Pharmacia Fine Chemicals, Uppsala, Sweden).

### *Solutions*

The PSS contained (in mM): NaCl 138.6, KCl 5.0,  $\text{MgCl}_2$  1.5,  $\text{CaCl}_2$  2.0, HEPES 10, glucose 5.6,  $\text{KH}_2\text{PO}_4$  1.0, pH 7.2.  $\text{Ca}^{2+}$ -free solution was made by substituting an equimolar concentration of  $\text{MgCl}_2$  instead of  $\text{CaCl}_2$  and adding 0.5 mM EGTA.


### *Statistics*

Data were expressed as mean  $\pm$  s.e.m. and statistical significance was determined using the paired or unpaired Student's *t*-test. Probabilities less than 5% ( $P < 0.05$ ) were considered significant.

## Results

### *Methyl paraben-induced increase in $[\text{Ca}^{2+}]_i$ in the presence or absence of external $\text{Ca}^{2+}$*

Figure 1a and b shows representative traces of the effects of methyl paraben (3 mM) on  $[\text{Ca}^{2+}]_i$  in rat peritoneal mast cells (RPMCs). Methyl paraben was applied for 75 s. The agent produced a transient increase in  $[\text{Ca}^{2+}]_i$  both in  $\text{Ca}^{2+}$ -containing solution (PSS, 1a) and in  $\text{Ca}^{2+}$ -free solution (1b). Figure 1c shows the average of the peak values of the changes in  $[\text{Ca}^{2+}]_i$  induced by methyl paraben (0.1–10 mM). The  $[\text{Ca}^{2+}]_i$  was increased concentration-dependently in the pre-



**Figure 1** Effects of methyl paraben on  $[\text{Ca}^{2+}]_i$  in RPMCs. The fluo-3 AM-loaded RPMCs were stimulated with methyl paraben for 75 s at room temperature (22–25°C). (a) Methyl paraben (3 mM) increased the  $[\text{Ca}^{2+}]_i$  in an RPMC incubated in PSS containing 2 mM  $\text{Ca}^{2+}$ . (b) Methyl paraben was applied in  $\text{Ca}^{2+}$ -free solution containing 0.5 mM EGTA after a 5 min removal of  $\text{Ca}^{2+}$ . Also in  $\text{Ca}^{2+}$ -free condition, methyl paraben increased the  $[\text{Ca}^{2+}]_i$  in an RPMC. (c) The Concentration – response relation of methyl paraben-induced  $[\text{Ca}^{2+}]_i$  increase in  $\text{Ca}^{2+}$ -containing (open circles) and  $\text{Ca}^{2+}$ -free (filled circles) mediums. The ordinate shows the net maximum  $[\text{Ca}^{2+}]_i$  of the response with baseline subtracted. The data points indicate mean  $\pm$  s.e.m. of 20 experiments. The asterisk shows a significant difference in  $\text{Ca}^{2+}$ -containing and  $\text{Ca}^{2+}$ -free solutions.

sence of extracellular  $\text{Ca}^{2+}$ . A similar result was obtained also in the  $\text{Ca}^{2+}$ -free solution. At a high concentration (10 mM), however, the increase was significantly greater in the  $\text{Ca}^{2+}$ -containing solution.

#### *Methyl paraben did not evoke histamine release from RPMCs*

The amount of histamine release was measured as an index of the degranulation of RPMCs and illustrated in Figure 2. The basal spontaneous release was around 10% of the total contents in RPMCs (control). Compound 48/80 ( $0.3 \mu\text{g ml}^{-1}$ ) potentiated the histamine release both in PSS and in  $\text{Ca}^{2+}$ -free solution, increasing it up to  $63.9 \pm 5.2\%$  ( $n=4$ ) and  $34.6 \pm 1.9\%$  ( $n=4$ ), respectively. A combination of a  $\text{Ca}^{2+}$  ionophore (A23187  $0.1 \mu\text{M}$ ) and the phorbol ester (PMA 10 nM) caused the release of histamine in PSS ( $40.9 \pm 1.2\%$ ,  $n=5$ ). In contrast, methyl paraben did not increase histamine release in PSS ( $4.32 \pm 0.58\%$ ,  $n=4$ , by 3 mM and  $9.84 \pm 1.70\%$ ,  $n=4$ , by 10 mM). It was also the case in  $\text{Ca}^{2+}$ -free solution ( $6.49 \pm 0.91\%$ ,  $n=4$ , by 3 mM and  $8.92 \pm 1.18\%$ ,  $n=4$ , by 10 mM).


#### *Methyl paraben-evoked histamine release from PMA-pretreated RPMCs*

Incapability of methyl paraben to release histamine in the present study may be due to insufficient activation of protein kinase C (PKC), since it has been reported that both increase in  $[\text{Ca}^{2+}]_i$  and activation of PKC are needed for the release (Beaven & Cunha-Melo, 1988). We, therefore, pretreated RPMCs with PMA for 5 min to activate PKC. The sponta-


neous histamine release was normalized as a relative histamine release of 1.0. Application of methyl paraben alone at 0.3–10 mM did not induce histamine release (open circles). In the absence of methyl paraben (control), PMA at 3 nM (closed circles) and 10 nM (closed squares) increased the release to  $1.17 \pm 0.30$  ( $n=4$ ) and  $2.05 \pm 0.54$  ( $n=4$ ) fold, respectively. Pretreatment with PMA for 5 min greatly enhanced the histamine release evoked by methyl paraben (0.3–10 mM). The peak increase was  $2.23 \pm 0.11$  times the control increase in the presence of 3 nM PMA ( $n=4$ , 1 mM methyl paraben) and  $1.53 \pm 0.53$  times the control increase in the presence of 10 nM PMA ( $n=4$ , 3 mM methyl paraben). A high concentration (10 mM) of methyl paraben, however, did not augment or even suppressed the histamine release from the PMA-pretreated RPMCs. The 10 nM PMA-induced increase in histamine release was suppressed to  $0.79 \pm 0.18$  ( $n=4$ ) of the control increase (Figure 3).

#### *Effects of U73122 and U73343 on histamine release induced by combination of PMA and methyl paraben*

To assess the involvement of phospholipase C (PLC) in the histamine release reaction by methyl paraben in the PMA-pretreated RPMCs, effects of a PLC inhibitor U73122 and its inactive analogue U73343 were examined. The spontaneous histamine release in PSS containing 2 mM  $\text{Ca}^{2+}$  was normalized as a relative histamine release of 1.0. A volume of 10 nM of PMA increased the release to  $1.80 \pm 0.17$  fold ( $n=12$ ), and U73122 ( $0.5 \mu\text{M}$ ) did not significantly inhibit the increase ( $n=12$ ). The histamine release caused by methyl paraben (3 mM) applied 5 min after application of PMA (10 nM) was




**Figure 2** Effects of methyl paraben (3 and 10 mM) and various stimuli on histamine release from RPMCs. Cells were preincubated for 5 min at 37°C and subsequently incubated with methyl paraben (3 and 10 mM), compound 48/80 (0.3 µg ml<sup>-1</sup>) or a combination of A23187 (0.1 µM) and PMA (10 nM) for 30 min in the external solution with (open columns) or without (closed columns)  $\text{Ca}^{2+}$ . The released histamine in the 30 min period was calculated as a percentage of the total histamine content of the cells. Results are expressed as mean ± s.e.m. of eight to 10 experiments.



**Figure 3** Effects of combined application of methyl paraben and PMA on histamine release in PSS. Cells were preincubated for 5 min at 37°C in the external solution with  $\text{Ca}^{2+}$  and subsequently incubated without (open circles) or with PMA (3 nM, closed circles or 10 nM, closed squares) for 5 min. The cells were then stimulated with methyl paraben (3 mM) for 30 min in the continuous presence or absence of PMA. The spontaneous histamine release was used for normalization as a relative histamine release of 1.0. The symbols refer to the mean of four to 12 experiments and the error bars represent s.e.m.

$3.86 \pm 0.28$  times the control ( $n = 5$ ). U73122 significantly reduced this augmented histamine release by the combination of PMA and methyl paraben. The normalized inhibitory value by U73122 on the combination of PMA and methyl paraben-induced augmentation was  $0.68 \pm 0.04$  ( $n = 5$ ) and  $0.10 \pm 0.02$  ( $n = 5$ ) at 0.1 and 0.5 µM, respectively. In contrast, U73343 (0.5 µM) did not inhibit the histamine release caused by the combined application of PMA and methyl paraben (Figure 4).




**Figure 4** Effects of U73122 and U73343 on histamine release induced by a combination of PMA and methyl paraben. After preincubation for 5 min at 37°C in PSS, cells were pretreated with PMA (10 nM) or a combination of PMA and U73122 (0.1 and 0.5 µM) or U73343 (0.5 µM) for 5 min. The cells were then stimulated with methyl paraben (closed columns) or not (hatched columns) and incubated for 30 min in PSS with continued treatment of PMA in the continuous presence or absence of U73122 or U73343. The spontaneous histamine release was used for normalization as a relative histamine release of 1.0 (open column). Results are expressed as mean ± s.e.m. of five to 12 experiments. Asterisks indicate a significant difference from the combined application of PMA and methyl paraben in the absence of U73122 or U73343 with  $P < 0.05$ .

#### Effects of U73122, U73343 and D609 on increase in $[\text{Ca}^{2+}]_i$ induced by methyl paraben


We investigated the effects of U73122, U73343 and another PLC inhibitor D609 to confirm that the increase in  $[\text{Ca}^{2+}]_i$  caused by methyl paraben was induced through activation of PLC in RPMCs. The change in  $[\text{Ca}^{2+}]_i$  caused by 3 mM of methyl paraben in the  $\text{Ca}^{2+}$ -free solution containing 0.5 mM EGTA was used as a control (Figure 5a). Pretreatment with U73122 (0.5 µM) for 5 min markedly inhibited the  $[\text{Ca}^{2+}]_i$  increase, while U73343 (0.5 µM) did not (Figure 5b). D609 at 10 µM completely suppressed the increase (Figure 5c). Figure 5d illustrates the effects of U73312 (0.1 and 0.5 µM), U73343 (0.5 µM) and D609 (1–10 µM) on the peak increase in  $[\text{Ca}^{2+}]_i$  caused by 3 mM of methyl paraben. U73122 (0.1 and 0.5 µM) and D609 (1–10 µM) inhibited the increases in  $[\text{Ca}^{2+}]_i$  in a concentration-dependent manner, whereas the inactive analogue (U73343 0.5 µM) did not.

#### Effects of xestospongine C and 2APB on increase in $[\text{Ca}^{2+}]_i$ induced by methyl paraben

PLC increases the cytosolic levels of inositol 1,4,5-trisphosphate ( $\text{IP}_3$ ) and DAG in mast cells (White *et al.*, 1985).  $\text{IP}_3$  stimulates release of  $\text{Ca}^{2+}$  from internal stores (Meyer *et al.*, 1988; Berridge, 1993) and DAG is known to activate PKC (Nishizuka, 1984). We investigated the effects of xestospongine C and 2 aminoethoxydiphenyl borate (2APB), inhibitors of  $\text{IP}_3$ -induced  $\text{Ca}^{2+}$  release, on the methyl paraben-induced increase in  $[\text{Ca}^{2+}]_i$  in RPMCs. Figure 6a depicts a control increase in the  $[\text{Ca}^{2+}]_i$  induced by methyl paraben (3 mM) in the  $\text{Ca}^{2+}$ -free solution containing 0.5 mM EGTA. Pretreat-



**Figure 5** Effects of U73122, U73343 and D609 on the increase in  $[\text{Ca}^{2+}]_i$  induced by methyl paraben in  $\text{Ca}^{2+}$ -free solution containing 0.5 mM EGTA. Methyl paraben (3 mM) was applied for 60–75 s following a 5 min incubation in the  $\text{Ca}^{2+}$ -free solution. (a) Methyl paraben increased the  $[\text{Ca}^{2+}]_i$  in a fluo-3 AM-loaded RPMC. (b) Pretreatment for 5 min with U73122 (0.5  $\mu\text{M}$ ) markedly suppressed the increase, while U73343 (0.5  $\mu\text{M}$ ) had no effects. (c) Pretreatment for 5 min with D609 (10  $\mu\text{M}$ ) completely blocked the effect of methyl paraben. (d) Each column indicates the average of the peak increase above the base line ( $n=8-10$ ). U73122 and D609 suppressed the  $[\text{Ca}^{2+}]_i$  increase dose-dependently. The reduction by U73343 was not significant. Bars indicate s.e.m. Asterisks indicate significant difference from the control with  $P<0.05$ .



**Figure 6** Effects of xestospongine C and 2APB on the increase in  $[\text{Ca}^{2+}]_i$  induced by methyl paraben in  $\text{Ca}^{2+}$ -free solution containing 0.5 mM EGTA. Methyl paraben (3 mM) was applied for 70–75 s in  $\text{Ca}^{2+}$ -free solution after a 5 min removal of  $\text{Ca}^{2+}$ . (a) Methyl paraben increased the  $[\text{Ca}^{2+}]_i$  in a fluo-3 AM-loaded RPMC. (b) Pretreatment for 5 min with xestospongine C (6 or 20  $\mu\text{M}$ ) markedly suppressed the  $[\text{Ca}^{2+}]_i$  increase. (c) Pretreatment for 5 min with 2APB (100  $\mu\text{M}$ ) greatly decreased the  $[\text{Ca}^{2+}]_i$  increase. (d) Each column indicates the average of the peak increase above the base line ( $n=7-10$ ). Xestospongine C and 2APB dose-dependently suppressed the  $[\text{Ca}^{2+}]_i$  increase produced by methyl paraben. Bars indicate s.e.m. Asterisks show significant suppression from the control with  $P<0.05$ .

ment with xestospongine C (6 and 20  $\mu\text{M}$ ; Figure 6b) or 2APB (100  $\mu\text{M}$ ; Figure 6c) markedly inhibited the increase in  $[\text{Ca}^{2+}]_i$ . Effects of the  $\text{IP}_3$  inhibitors on the peak increase in  $[\text{Ca}^{2+}]_i$  are

shown in Figure 6d. Xestospongine C (2–20  $\mu\text{M}$ ) and 2APB (30 and 100  $\mu\text{M}$ ) suppressed the increases in  $[\text{Ca}^{2+}]_i$  in a concentration-dependent manner.

## Discussion

In this study, we investigated the effects of methyl paraben on the changes in  $[\text{Ca}^{2+}]_i$  and histamine release in RPMCs. Methyl paraben at 1–10 mM dose-dependently increased  $[\text{Ca}^{2+}]_i$  in both  $\text{Ca}^{2+}$ -containing and  $\text{Ca}^{2+}$ -free solutions. The peak increase in  $[\text{Ca}^{2+}]_i$  induced by 0.3–3 mM of methyl paraben was not significantly different between the presence and absence of the external  $\text{Ca}^{2+}$ . This finding suggests that the transient increase in  $[\text{Ca}^{2+}]_i$  by the agent was largely due to release of  $\text{Ca}^{2+}$  from intracellular storage sites. At a high concentration (10 mM), however, the peak increase was significantly greater in the  $\text{Ca}^{2+}$ -containing solution. High concentrations of methyl paraben may induce  $\text{Ca}^{2+}$  influx from the extracellular medium (Figure 1).

Many reports have demonstrated that  $\text{Ca}^{2+}$  chelators such as EGTA increase  $[\text{Ca}^{2+}]_i$  in mast cells, although the reason is not known (Teraoka *et al.*, 1997). In the present experiments too, EGTA caused an increase in  $[\text{Ca}^{2+}]_i$  in some of the mast cells. However, this reaction was observed only within 5 min of the exposure to EGTA (data not shown). Hence, measurements were made 5 min after replacement of PSS with  $\text{Ca}^{2+}$ -free solution, and cells showing no change in  $[\text{Ca}^{2+}]_i$  were selected for the following experiments. Actually, changes in  $[\text{Ca}^{2+}]_i$  were negligible probably due to the low concentration of EGTA (0.5 mM) in this study.

Application of the stimulants known to release histamine from mast cells, such as compound 48/80, known as an activator of PLC (Cockcroft & Gomperts, 1979), and the combination of A23187 and PMA (Chew *et al.*, 2002), released histamine in our experimental system, but methyl paraben at 0.3–10 mM did not induce histamine release (Figures 2 and 3). Dvorak *et al.* (1996) demonstrated that the interaction between inositol phospholipids and PKC regulated degranulation of mast cells. In addition, Akita *et al.* (1990) reported that  $\text{Ca}^{2+}$  played an important role in the binding of PKC to the cell membrane. Furthermore, it was suggested that PKC was activated by increased  $[\text{Ca}^{2+}]_i$  and phospholipids, and that the activation was further enhanced by DAG (White *et al.*, 1985). Thus, the increase in  $[\text{Ca}^{2+}]_i$  and the activation of PKC, which may lead to degranulation, are required to release histamine (Heiman & Crews, 1985a, b). Application of methyl paraben alone did not induce histamine release in the present experiments, but after activation of PKC by PMA, the agent evoked histamine release, which might be mediated by the activated PKC and the  $\text{Ca}^{2+}$  mobilized by methyl paraben. The increase in  $[\text{Ca}^{2+}]_i$  induced by methyl paraben was dose-dependent (Figure 1), while the release of histamine by the combined application of PMA and methyl paraben was not dose-dependent but inhibited at the higher concentration (10 mM) (Figure 3). These results suggest that methyl paraben not only mobilizes  $\text{Ca}^{2+}$  but also inhibits the histamine release via some unknown mechanisms.

The PLC inhibitors, U73122 (Bleasdale *et al.*, 1990) and D609 (Ito *et al.*, 2002), inhibited the increase in  $[\text{Ca}^{2+}]_i$  caused by methyl paraben. U73122 also inhibited the histamine release induced by the combined application of PMA and methyl paraben. In contrast, U73343, an inactive analogue of U73122 (Pocock & Bates, 2001), did not show any effects on these reactions. In addition, U73122 did not significantly suppress the histamine release caused by PMA (Figures 4 and

5). Thus the PLC inhibitors suppressed the action of methyl paraben, indicating that methyl paraben causes the increase in  $[\text{Ca}^{2+}]_i$  via activation of PLC. When PLC is activated in mast cells, phosphatidylinositol 4,5-bisphosphate ( $\text{PIP}_2$ ) is hydrolysed to produce  $\text{IP}_3$  and DAG.  $\text{IP}_3$  binds to its receptors on the intracellular  $\text{Ca}^{2+}$  storage site to release  $\text{Ca}^{2+}$  (Meyer *et al.*, 1988; Berridge, 1993), while DAG activates PKC (White & Metzger, 1988).  $\text{IP}_3$  receptor blockers, xestospongin C (Gafni *et al.*, 1997) and 2APB (Maruyama *et al.*, 1997), inhibited the increase in  $[\text{Ca}^{2+}]_i$  induced by methyl paraben (Figure 6), suggesting that methyl paraben produced  $\text{IP}_3$  through PLC activation, which resulted in a release of  $\text{Ca}^{2+}$  from the intracellular storage sites in RPMCs. 4-Chloro-*m*-cresol (0.5 and 1 mM), a potent activator of the ryanodine receptor (Zorzato *et al.*, 1993), induced the transient increase in  $[\text{Ca}^{2+}]_i$ ; however, additional application of any stimulants such as compound 48/80, methyl paraben or A23187 had no effect on  $[\text{Ca}^{2+}]_i$  in RPMCs (data not shown). These results suggest the colocalization of ryanodine receptor and  $\text{IP}_3$  receptor in  $\text{Ca}^{2+}$  store. Although the effect of methyl paraben on the ryanodine receptor in RPMCs is still unclear, PLC inhibitors and  $\text{IP}_3$  receptor blockers almost completely suppressed the action of methyl paraben (Figures 4–6), indicating that  $\text{IP}_3$  is a key messenger on  $\text{Ca}^{2+}$  release and subsequent histamine release induced by methyl paraben in RPMCs. DAG produced by PLC activation is expected to activate PKC and to evoke degranulation by acting together with mobilized  $\text{Ca}^{2+}$ . However, no histamine release was observed by methyl paraben alone in the present study (Figures 2 and 3). In addition, methyl paraben released histamine from the mast cells in which PKC had been activated by PMA (Figure 3). These arguments suggest that although methyl paraben activates PLC, the PKC activity is not enough to release histamine in RPMCs.

Methyl paraben is contained in various pharmaceutical agents at certain concentrations: 0.06–0.25% in injections; 0.015–0.2% in syrups; 0.015–0.05% in instillations; and 0.02–0.3% in external medicines (Anonymous, 1998). In addition, methyl paraben is added to cosmetics at 0.32% at maximum (Rastogi *et al.*, 1995), and to food at 0.03–0.07% in combination with propyl paraben (Krebs-Smith *et al.*, 1997). In this study, methyl paraben showed a significant effect on the  $[\text{Ca}^{2+}]_i$  increase and histamine release at the concentrations of 1–3 mM, which correspond to approximately 0.015–0.046%. Therefore, some agents containing methyl paraben may induce an increase in  $[\text{Ca}^{2+}]_i$  when directly acting on mast cells.

The molecular weight of methyl paraben is as small as 152.14, and hence it has been thought that the agent by itself does not have antigenicity and may become antigenic when bound to a certain kind of proteins (Pressman *et al.*, 1968). In the present experiments, methyl paraben might act on mast cells with no relation to FcεRI on account of the absence of specific IgE and proteins to bind. *In vivo*, however, the situation would be different from *in vitro*, and it is likely that methyl paraben specific IgE may mediate the activation of  $\text{Ca}^{2+}$  release and subsequent histamine release in mast cells. In addition, it is not clear whether methyl paraben bound to proteins has an inhibitory action on histamine release; therefore, the usual view that methyl paraben induces allergic reactions through an immunological mechanism is not excluded. On the other hand, the present results propose the possibility that methyl paraben may cause anaphylactoid

reactions through a nonimmunological mechanism when PKC is simultaneously activated to some extent in the mast cells.

## References

- AKITA, M., SASAKI, S., MATSUYAMA, S. & MIZUSHIMA, S. (1990). SecA interacts with secretory proteins by recognizing the positive charge at the amino terminus of the signal peptide in *Escherichia coli*. *J. Biol. Chem.*, **265**, 8164–8169.
- ANONYMOUS (1998). Preservation, sterilization and stability testing of ophthalmic preparations. *Int. J. Pharm. Compounding*, **2**, 192–195.
- BEAVEN, M.A. & CUNHA-MELO, J.R. (1988). Membrane phosphoinositide-activated signals in mast cells and basophils. *Prog. Allergy*, **42**, 123–184.
- BERRIDGE, M.J. (1993). Inositol trisphosphate and calcium signalling. *Nature*, **361**, 315–325.
- BLEASDALE, J.E., THAKUR, N.R., GREMBAN, R.S., BUNDY, G.L., FITZPATRICK, F.A., SMITH, R.J. & BUNTING, S. (1990). Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J. Pharmacol. Exp. Ther.*, **255**, 756–768.
- CAVAGNA, D., ZORZATO, F., BABINI, E., PRESTIPINO, G. & TREVES, S. (2000). Methyl *p*-hydroxybenzoate (E-218) a preservative for drugs and food is an activator of the ryanodine receptor  $\text{Ca}^{2+}$  release channel. *Br. J. Pharmacol.*, **131**, 335–341.
- CHAN, C.L., JONES, R.L. & LAU, H.Y. (2000). Characterization of prostanoid receptors mediating inhibition of histamine release from anti-IgE-activated rat peritoneal mast cells. *Br. J. Pharmacol.*, **129**, 589–597.
- CHEW, C.S., CHEN, X., PARENTE Jr J.A., TARRER, S., OKAMOTO, C. & QIN, H.Y. (2002). Lasp-1 binds to non-muscle F-actin *in vitro* and is localized within multiple sites of dynamic actin assembly *in vivo*. *J. Cell. Sci.*, **115**, 4787–4799.
- COCKCROFT, S. & GOMPERTS, B.D. (1979). Evidence for a role of phosphatidylinositol turnover in stimulus-secretion coupling. *Biochem. J.*, **178**, 681–687.
- DVORAK, A.M., ACKERMAN, S.J., LETOURNEAU, L., MORGAN, E.S., LICHTENSTEIN, L.M. & MACGLASHAN Jr. D.W. (1996). Vesicular transport of Charcot–Leyden crystal protein in tumor-promoting phorbol diester-stimulated human basophils. *Lab. Invest.*, **74**, 967–974.
- FISHER, M.M. & MORE, D.G. (1981). The epidemiology and clinical features of anaphylactic reactions in anaesthesia. *Anaesth. Intens. Care*, **9**, 226–234.
- FISHER, M.M. & PENNINGTON, J.C. (1982). Allergy to local anaesthesia. *Br. J. Anaesth.*, **54**, 893–894.
- GAFNI, J., MUNSCH, J.A., LAM, T.H., CATLIN, M.C., COSTA, L.G., MOLINSKI, T.F. & PESSACH, I.N. (1997). Xestospongins: potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor. *Neuron*, **19**, 723–733.
- HEIMAN, A.S. & CREWS, F.T. (1985a). Characterization of the effects of phorbol esters on rat mast cell secretion. *J. Immunol.*, **134**, 548–555.
- HEIMAN, A.S. & CREWS, F.T. (1985b). Hydrocortisone inhibits phorbol ester stimulated release of histamine and arachidonic acid from rat mast cells. *Biochem. Biophys. Res. Commun.*, **130**, 640–645.
- HORINOCHI, Y., ABE, K., KUBO, K. & OKA, M. (1993). Mechanisms of vancomycin-induced histamine release from rat peritoneal mast cells. *Agents Actions*, **40**, 28–36.
- HOSOI, E., NISHIZAKI, C., GALLAGHER, K.L., WYRE, H.W., MATSUO, Y. & SEI, Y. (2001). Expression of the ryanodine receptor isoforms in immune cells. *J. Immunol.*, **167**, 4887–4894.
- ITO, J., NAGAYASU, Y., UENO, S. & YOKOYAMA, S. (2002). Apolipoprotein-mediated cellular lipid release requires replenishment of sphingomyelin in a phosphatidylcholine-specific phospholipase C-dependent manner. *J. Biol. Chem.*, **277**, 44709–44714.
- KOKUBU, M., ODA, K. & SHINYA, N. (1989). Detection of serum IgE antibody specific for local anesthetics and methyl paraben. *Anesth. Prog.*, **36**, 186–187.
- KREBS-SMITH, S.M., GUENTHER, P.M., COOK, A., THOMPSON, F.E., CUCINELLI, J. & UDLER, J. (1997). Foods commonly eaten in the United States: quantities consumed per eating occasion and in a day, 1989–91. NTISPB 98-111719.
- LARSON, C.E. (1977). Methyl paraben – an overlooked cause of local anesthetic hypersensitivity. *Anesth. Prog.*, **24**, 72–74.
- MARUYAMA, T., KANAJI, T., NAKADE, S., KANNO, T. & MIKOSHIBA, K. (1997). 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of ins (1,4,5)  $\text{P}_3$ -induced  $\text{Ca}^{2+}$  release. *J. Biochem. (Tokyo)*, **122**, 498–505.
- MERTES, P.M. & LAXENAIRE, M.C. (2002). Allergic reactions occurring during anaesthesia. *Eur. J. Anaesthesiol.*, **19**, 240–262.
- MEYER, T., HOLOWKA, D. & STRYER, L. (1988). Highly cooperative opening of calcium channels by inositol 1,4,5-trisphosphate. *Science*, **240**, 653–656.
- MOWAD, C.M. (2000). Allergic contact dermatitis caused by parabens: 2 case reports and a review. *Am. J. Contact Dermatol.*, **11**, 53–56.
- NAGEL, J.E., FUSCALDO, J.T. & FIREMAN, P. (1977). Paraben allergy. *J. Am. Med. Assoc.*, **237**, 1594–1595.
- NISHIZUKA, Y. (1984). The role of protein kinase C in cell surface signal transduction and tumour promotion. *Nature*, **308**, 693–698.
- POCOCK, T.M. & BATES, D.O. (2001). *In vivo* mechanisms of vascular endothelial growth factor-mediated increased hydraulic conductivity of Rana capillaries. *J. Physiol.*, **534**, 479–488.
- PRESSMAN, D., SIEGEL, M. & HALL, L.A. (1968). The closeness of fit of antibenzoate about haptens and the orientation of the haptens in combination. *J. Am. Chem. Soc.*, **76**, 6336–6342.
- RASTOGI, S.C., SCHOUTEN, A., DE KRUIJFF, N. & WEIJLAND, J.W. (1995). Contents of methyl-, ethyl-, propyl-, butyl- and benzylparaben in cosmetic products. *Contact Dermatitis*, **32**, 28–30.
- SOBOLOFF, J. & BERGER, S.A. (2002). Sustained ER  $\text{Ca}^{2+}$  depletion suppresses protein synthesis and induces activation-enhanced cell death in mast cells. *J. Biol. Chem.*, **277**, 13812–13820.
- SONI, M.G., TAYLOR, S.L., GREENBERG, N.A. & BURDOCK, G.A. (2002). Evaluation of the health aspects of methyl paraben: a review of the published literature. *Food Chem. Toxicol.*, **40**, 1335–1373.
- STELLATO, C. & MARONE, G. (1995). Mast cells and basophils in adverse reactions to drugs used during general anaesthesia. Human basophils and mast cells: clinical aspects. *Chem. Immunol.*, **62**, 108–131.
- TERAOKA, H., AKIBA, H., TAKAKI, R., TANEIKE, T., HIRAGA, T. & OHGA, A. (1997). Inhibitory effects of caffeine on influx and histamine secretion independent of CAMP in rat peritoneal mast cells. *Gen. Pharmac.*, **28**, 237–243.
- WHITE, J.R., PLUZNIK, D.H., ISHIZAKA, K. & ISHIZAKA, T. (1985). Antigen-induced increase in protein kinase C activity in plasma membrane of mast cells. *Proc. Natl. Acad. Sci. U.S.A.*, **82**, 8193–8197.
- WHITE, K.N. & METZGER, H. (1988). Translocation of protein kinase C in rat basophilic leukemic cells induced by phorbol ester or by aggregation of IgE receptors. *J. Immunol.*, **141**, 942–947.
- WILDSMITH, J.A., MASON, A., MCKINNON, R.P. & RAE, S.M. (1998). Alleged allergy to local anaesthetic drugs. *Br. Dent. J.*, **184**, 507–510.
- YAMATODANI, A., FUKUDA, H., WADA, H., IWAEDA, T. & WARANABE, T. (1985). High performance liquid chromatographic determination of plasma and brain histamine without previous purification of biological samples: cation-exchange chromatography coupled with postcolumn derivatization fluorometry. *J. Chromatogr.*, **344**, 115–123.
- ZORZATO, F., SCUTARI, E., TEGAZZIN, V., CLEMENTI, E. & TREVES, S. (1993). Chlorocresol: an activator of ryanodine receptor-mediated  $\text{Ca}^{2+}$  release. *Mol. Pharmacol.*, **44**, 1192–1201.

(Received December 16, 2002

Revised February 10, 2003

Accepted February 19, 2003)