

## RESEARCH PAPER

# Intraplantar PGE<sub>2</sub> causes nociceptive behaviour and mechanical allodynia: the role of prostanoid E receptors and protein kinases

CAL Kassuya<sup>1</sup>, J Ferreira<sup>2</sup>, RF Claudino<sup>1</sup> and JB Calixto<sup>1</sup>

<sup>1</sup>Department of Pharmacology, Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, Brazil and <sup>2</sup>Department of Chemistry, Universidade Federal de Santa Maria, Santa Maria, Rio Grande do Sul, Brazil

**Background and purpose:** Receptor subtypes involved in PGE<sub>2</sub>-induced nociception are still controversial. The present study investigated the prostanoid E receptor (EP) subtypes and the protein kinase (PK) pathways involved in the nociception induced by PGE<sub>2</sub> injection in the mouse paw.

**Experimental approach:** Paw-licking and mechanical allodynia were measured *in vivo* and protein kinase activation *ex vivo* by Western blots of extracts of paw skin.

**Key results:** Intraplantar (i.pl.) injection of PGE<sub>2</sub> into the mouse paw caused nociceptive behaviour of short duration with mean ED<sub>50</sub> of 1.43 nmol. PGE<sub>2</sub> produced a longer-lasting mechanical allodynia, with an ED<sub>50</sub> of 0.05 nmol. Intraplantar injection of antagonists at EP<sub>3</sub> or EP<sub>4</sub>, but not at EP<sub>1</sub> or EP<sub>2</sub> receptors inhibited PGE<sub>2</sub>-induced paw-licking. Paw-licking caused by PGE<sub>2</sub> was blocked by an inhibitor of PKA but only partially decreased by inhibition of the extracellular-regulated kinase (ERK). Selective inhibitors of PKC, c-Jun N-terminal kinase (JNK) or p38, all failed to affect PGE<sub>2</sub>-induced paw-licking. An EP<sub>3</sub> antagonist inhibited PGE<sub>2</sub>-induced mechanical allodynia. However, inhibitors of PKA, PKC or ERK, but not p38 or JNK, also partially inhibited PGE<sub>2</sub>-induced mechanical allodynia. Western blot analyses confirmed that i.pl. injection of PGE<sub>2</sub> activated PKA, PKC $\alpha$ , and mitogen activated kinases (MAPKs) in the paw. Co-treatment with EP<sub>3</sub> or EP<sub>4</sub> receptor antagonists reduced PGE<sub>2</sub>-induced PKA and ERK, but not PKC $\alpha$  activation.

**Conclusions and Implications:** The present results indicate that the nociceptive behaviour and mechanical allodynia caused by i.pl. PGE<sub>2</sub> are mediated through activation of distinct EP receptors and PK-dependent mechanisms.

*British Journal of Pharmacology* (2007) **150**, 727–737. doi:10.1038/sj.bjp.0707149; published online 19 February 2007

**Keywords:** prostaglandin E<sub>2</sub>; prostanoid E receptors; protein kinases; nociception; mitogen-activated protein kinases; mice

**Abbreviations:** DRG, dorsal root ganglion; EP receptor, prostanoid E receptor; ERK, extracellular-regulated kinase; Gs, stimulatory G-protein; i.pl., intraplantar; JNK, c-Jun N-terminal kinase; MAPKs, mitogen-activated protein kinases; MEK, mitogen-activated protein kinase kinase; PBS, phosphate-buffered saline; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; PK, protein kinase; PKA, protein kinase A; PKA RII, protein kinase A regulatory subunit II; PKC, protein kinase C

## Introduction

Symptomatically, pain may be manifested spontaneously (stimulus-independent pain) or following challenge with noxious (hyperalgesia) or innocuous (allodynia) stimulation after damage to, or alterations in, sensory neurons (stimulus-evoked pain) (Woolf and Mannion, 1999). During an inflammatory event, pain generation is a consequence of a complex interaction between a number of inflammatory mediators, including prostaglandins, some of which (notably prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)) are known to exert a critical role in

the generation and maintenance of the nociceptive response (see Samad *et al.* (2002) for review).

A great number of *in vivo* studies have shown that peripherally injected PGE<sub>2</sub> produces hyperalgesia and allodynia both in experimental animals and in humans (Ferreira, 1972; Kuhn and Willis, 1973). This nociceptive effect seems to be related to the ability of PGE<sub>2</sub> to sensitize peripheral terminals of small diameter, high threshold, primary afferent fibers to thermal, chemical and mechanical stimuli (Schaible and Schimdt, 1988; Kumazawa *et al.*, 1993, 1996; Mizumura *et al.*, 1993). Besides sensitization, a direct activation of nociceptors *in vitro* by high concentrations of PGE<sub>2</sub> has been shown (Mense, 1981; Mizumura *et al.*, 1987; Schaible and Schimdt, 1988), as well as the induction of spontaneous nociceptive behaviours following peripheral injection *in vivo* (Hong and Abbott, 1994).

Correspondence: Professor JB Calixto, Departamento de Farmacologia, Universidade Federal de Santa Catarina, Campus Universitário – Trindade, Bloco D-CCB – Cx. Postal: 476 – CEP, 88049-900 – Florianópolis, Santa Catarina, Brazil.

E-mails: calixto@farmaco.ufsc.br or calixto3@terra.com.br

Received 11 May 2006; revised 1 August 2006; accepted 12 December 2006; published online 19 February 2007

The biological actions of PGE<sub>2</sub> are attributed to its ability to interact with G-protein-coupled (prostanoid E receptor) EP receptors that have been classified into four subtypes (EP<sub>1-4</sub>) (see Kobayashi and Narumiya, 2002; Hata and Breyer, 2004 for review). EP receptors can be expressed in various tissues, including sensory neurons (Southall and Vasko, 2001). It has been suggested that EP<sub>2</sub>, EP<sub>3</sub> and EP<sub>4</sub> receptors could mediate the sensitizing effect of PGE<sub>2</sub> in nociceptors and dorsal root ganglion (DRG) neurons (Kumazawa *et al.*, 1993, 1996; Southall and Vasko, 2001). However, the receptor subtype(s) responsible for the stimulus-evoked and nociceptive behaviour *in vivo* produced by peripherally injected PGE<sub>2</sub> still remains unknown.

The stimulation of EP receptors can result in activation of complex signal transduction pathways, depending on the receptor subtype stimulated and the cells being studied. Some studies have demonstrated that the mechanical hyperalgesia caused by peripheral PGE<sub>2</sub> injection in rats is mediated by cAMP-protein kinase A (PKA) pathways (Ferreira and Nakamura, 1979; Taiwo and Levine, 1991; Aley and Levine, 1999). In contrast, thermal hyperalgesia produced by peripheral injection of PGE<sub>2</sub> is only marginally reduced in mice with a targeted mutation of the type I regulatory subunit of PKA, suggesting that other intracellular pathways could also be involved in PGE<sub>2</sub>-induced nociceptive effects (Malmberg *et al.*, 1997). In fact, the activation of EP receptors by PGE<sub>2</sub> can stimulate other protein kinases (PK), including protein kinase C (PKC) and mitogen-activated protein kinases (MAPKs) – both of which exert a critical role in nociceptor excitation and sensitization (Burkey and Regan, 1995; Khasar *et al.*, 1999; Aley *et al.*, 2001; Fiebich *et al.*, 2001).

In the present study we sought to investigate, by the use of pharmacological and Western blot procedures, the receptor subtype and also the signaling pathways involved in mechanical allodynia and nociceptive behaviour produced *in vivo* by intraplantar (i.pl.) injection of PGE<sub>2</sub> in the mouse.

## Methods

### Animals

The experiments were conducted using male Swiss mice (25–35 g) kept in a 12 h light–dark cycle, with controlled humidity (60–80%) and temperature (21 ± 1°C). Food and water were freely available. The animals were acclimatized to the laboratory for at least 2 h before testing and were used only once throughout the experiments. The studies reported in this manuscript were carried out in accordance with current guidelines for the care of laboratory animals and ethical guidelines for the investigation of experimental pain in conscious animals, according to Zimmermann (1983) and approved by the local University Committee (process number 262/CEUA). The number of animals and the intensity of noxious stimuli used here were the minimum necessary to demonstrate consistent effects of the drug treatments.

### PGE<sub>2</sub>-induced paw licking

The procedure used was similar to that described previously (Ferreira *et al.*, 2004). Twenty microliters of solutions of PGE<sub>2</sub> (0.3–10 nmol per paw), 17-phenyl-trinor-PGE<sub>2</sub> (3 nmol per

paw) or a suspension of carrageenan (300 µg per paw) was injected subcutaneously under the plantar surface of the right hindpaw (i.pl. injection). The PGE<sub>2</sub> analogue 17-phenyl-trinor-PGE<sub>2</sub> is an agonist of mouse EP<sub>1</sub> and EP<sub>3</sub> receptors (Kiriyaama *et al.*, 1997). Separate groups of animals received an i.pl. injection of the appropriate vehicle (phosphate-buffered saline (PBS) or PBS plus ethanol 0.5%). The animals were placed individually in chambers (transparent glass cylinders, 20 cm in diameter) and were acclimatized for at least 20 min before algogen or vehicle injection. After challenge, the mice were observed individually for 15 min. The amount of time spent licking the injected paw was measured with a chronometer and was considered as indicative of nociceptive behaviour.

### PGE<sub>2</sub>-induced mechanical allodynia

The mechanical threshold was measured using the up–down paradigm as described previously (Chaplan *et al.*, 1994). Mice were first acclimatized (1–2 h) in individual clear Plexiglass boxes (9 × 7 × 11 cm) on an elevated wire mesh platform to allow the access to the plantar surface of the hind paws. Von Frey filaments of increasing stiffness (0.008–6.0 g) were applied to the same portion of the hind paw before and several times after i.pl. PGE<sub>2</sub> (0.01–0.3 nmol per paw), 17-phenyl-trinor-PGE<sub>2</sub> (0.1 nmol per paw) or carrageenan (300 µg per paw) injection. Mechanical allodynia was considered as a decrease in the threshold when compared with the same paws before injection with agonist. Each round of testing consisted of sets of six stimulations with different increasing filaments. When a filament induced paw withdrawals, a period of 1 min was allowed between stimulations. The mechanical threshold was expressed in milligrams (mg).

### Effect of drug treatments

To assess the involvement of different EP receptor subtypes in the algogen responses induced by PGE<sub>2</sub>, the animals received an i.pl. injection of an EP<sub>1/2</sub> receptor antagonist, AH-6809, (10–100 nmol per paw; Kiriyaama *et al.*, 1997), a selective EP<sub>3</sub> receptor antagonist, L-826266 (0.1–30 nmol per paw; Schlemper *et al.*, 2005) or a selective EP<sub>4</sub> receptor antagonist, L-161982 (1–30 nmol per paw; Machwate *et al.*, 2001) in association with PGE<sub>2</sub>, 17-phenyl-trinor-PGE<sub>2</sub> or carrageenan. To assess the effect of post-treatment with antagonists on PGE<sub>2</sub>-induced nociceptive behaviour, another group of animals was injected with EP receptor antagonists 5 min (for nociceptive behaviour) or 15 min (for mechanical allodynia) after PGE<sub>2</sub> injection.

To investigate some of the signalling pathways involved in PGE<sub>2</sub>-induced nociceptive response, we evaluated the effect of co-administration of selective kinase inhibitors – KT-5720 for PKA (0.1–3 nmol per paw), GF109203X for PKC (10 nmol per paw), SP600125 for c-Jun N-terminal kinase (JNK) (10 or 30 nmol per paw), PD98059 for mitogen-activated protein kinase kinase (MEK) (10 or 30 nmol per paw), or SB203580 for p38 (10 or 30 nmol per paw) – in association with PGE<sub>2</sub>. Another group of animals were injected with kinase inhibitors 5 min (for nociceptive behaviour) or 15 min (for mechanical allodynia) after PGE<sub>2</sub> injection. The choice of

doses for each drug was based on pilot experiments (not shown) or on data reported in the literature (Inoue *et al.*, 2003; Ferreira *et al.*, 2004, 2005; Claudino *et al.*, 2006).

#### Western blot analysis

To confirm the possible activation of PKC, PKA or MAPKs after PGE<sub>2</sub> injection into the mouse paw, western blot analysis was carried out as described previously (André *et al.*, 2004) with minor modifications. The right paws of mice were taken from animals killed at different periods of time (1–60 min) after PGE<sub>2</sub> treatment (3 nmol per paw). Paws were also collected from animals co-treated with L-826266 (10 nmol per paw) or L-161982 (10 nmol per paw) in association with PGE<sub>2</sub> (3 nmol per paw). The skin and connective tissues of the plantar aspect of the hind paws were removed and homogenized in an ice-cold buffer containing protease and phosphatase inhibitors (100 mM Tris-HCl, pH 7.4; 2 mM EDTA; 2 µg/ml aprotinin, 0.1 mM phenylmethanesulphonyl fluoride, 200 mM NaF and 2 mM of sodium orthovanadate). The homogenate was first centrifuged at 1000g for 10 min at 4°C; the pellet was discarded and the supernatant was further centrifuged at 35 000g for 30 min at 4°C. The supernatant was collected as a cytoplasm-rich fraction. The resulting pellet was re-suspended and considered as a membrane-rich fraction. The protein concentration was determined using a protein assay kit (Bio-Rad, Hercules, CA, USA). Equivalent amounts of proteins (10 µg for membrane and 40–70 µg for cytoplasm-rich fractions) were mixed in buffer (Tris 200 mM, glycerol 10%, sodium dodecyl sulphate 2%, β-mercaptoethanol 2.75 mM and bromophenol blue 0.04%) and boiled for 5 min. Proteins were resolved in 10% sodium dodecyl sulphate–polyacrylamide gel by electrophoresis and transferred on to polyvinylidene difluoride membranes. The membranes were blocked by incubation overnight with 10% non-fat dry milk solution and then incubated with anti-PKCα, anti-phosphorylated forms of protein kinase A regulatory subunit II (PKA RII) (phospho-PKA RII), extracellular-regulated kinase (ERK) (phospho-ERK1), JNK (phospho-JNK) or p38 (phospho-p38) antibodies. Following washing, the membranes were incubated with adjusted peroxidase-coupled secondary antibodies. The immunocomplexes were visualized using the enhanced chemiluminescence detection system (Amersham Biosciences, UK). Membranes were then incubated for 10 min in stripping buffer at room temperature and reincubated with anti-actin, which served as a loading control.

#### Statistical analysis

The results are presented as mean ± s.e.m., except for the ED<sub>50</sub> values (i.e. the dose of PGE<sub>2</sub> necessary to produce 50% of the nociceptive response relative to the maximum effect), which are reported as geometric means accompanied by their respective 95% confidence limits. To obtain data that were derived purely by the treatments in algogen-induced nociception, the inhibition values were represented as the difference between the licking times of the vehicle-treated and algogen-treated animals. The statistical significance between groups was assessed by means of one-way analysis

of variance (ANOVA) followed by Dunnett's or Student–Newman–Keuls' test, as appropriate. *P*-values <0.05 were considered as indicative of significance.

#### Drugs and reagents

AH-6809 and 17-phenyl-trinor-PGE<sub>2</sub>, were purchased from Cayman Chemicals (Ann Arbor, MI, USA); PGE<sub>2</sub> was obtained from Sigma-Aldrich (St Louis, MO, USA); PD98059, GF 109203x, SB 203580, SP600125 and KT-5720 were purchased from Calbiochem (Cambridge, MA, USA); polyclonal antibodies anti-PKC-α, anti-phospho-JNK, anti-phospho-p38, anti-phospho-ERK1, anti-actin and anti-rabbit IgG were obtained from Santa Cruz Biotechnology (Santa Cruz, CA, USA); anti-phospho-PKA RII was purchased from Upstate Biotechnology (Lake Placid, NY, USA) and L-826266 and L-161982 were kindly supplied by Merck Frost (Kirkland, Québec, Canada).

Stock solutions of drugs (0.1–1 M) were prepared in absolute ethanol, except for AH-6809, which was prepared in 1% of NaHCO<sub>3</sub> plus 0.9% NaCl and L-161982 that was directly dissolved in PBS. All drugs were stocked in siliconized plastic tubes and maintained at –18°C until use. The final concentration of ethanol did not exceed 0.5%, which alone had no effect on the nociceptive behaviour and on mechanical allodynia, respectively. KT-5720 was protected from light to avoid its photodegradation.


## Results

#### *Nociceptive behaviour and mechanical allodynia produced by i.pl. injection of PGE<sub>2</sub>*


The i.pl. injection of PGE<sub>2</sub> (0.3–10 nmol per paw) into the hindpaw of the mice produced dose-related paw licking when compared with vehicle-injected animals (Figure 1a). Paw licking induced by PGE<sub>2</sub> had a short duration, lasting up to 15 min after its injection (Figure 1a). The calculated mean ED<sub>50</sub> value (and the 95% confidence limits) for this effect was 1.43 (1.20–1.71) nmol per paw. The i.pl. injection of low doses of PGE<sub>2</sub> (0.01–0.3 nmol per paw) produced dose-dependent mechanical allodynia (stimulus-evoked nociception reaction), assessed by the decrease in the mechanical withdrawal thresholds (Figure 1b). In contrast with PGE<sub>2</sub>-induced paw licking, PGE<sub>2</sub>-induced mechanical allodynia had a long duration, peaking within 15 min and lasting for up to 60 min after injection of PGE<sub>2</sub> (Figure 1b). The calculated mean ED<sub>50</sub> value (and 95% confidence limits) for mechanical allodynia was 0.05 (0.03–0.09) nmol per paw. Thus, at the ED<sub>50</sub> level, PGE<sub>2</sub> was about 28-fold more potent in inducing mechanical allodynia in comparison to PGE<sub>2</sub>-induced paw licking. Based on these initial experiments, we chose doses of 3 and 0.1 nmol per paw of PGE<sub>2</sub> for subsequent experiments to evaluate mechanisms involved in paw licking and mechanical allodynia, respectively.

#### *Effects of EP receptor antagonists on PGE<sub>2</sub>-, 17-phenyl-trinor-PGE<sub>2</sub>- or carrageenan-induced nociceptive responses*

PGE<sub>2</sub>-induced paw licking was dose-dependently inhibited by the co-injection of the EP<sub>3</sub> receptor antagonist, L-826266,



**Figure 1** Dose and time dependence for PGE<sub>2</sub>-induced paw licking (0.3–10 nmol per paw) (a) or mechanical allodynia (0.01–0.3 nmol per paw) (b) in mice. The paw licking is expressed as licking time (s) and the mechanical allodynia is expressed as threshold (mg). Each point on the curve represents the mean of 6–10 animals and vertical lines show the s.e.m. Asterisks denote the significance levels in comparison with the vehicle-treated group (one-way ANOVA followed by Dunnett's test). \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ .




**Figure 2** Effect of i.p. treatment with selective receptor antagonists for EP<sub>3</sub> (L-826266, 3–30 nmol per paw (a)) EP<sub>4</sub> (L-161982, 1–30 nmol per paw (b)) EP<sub>1/2</sub> receptors (AH-6809, 10 and 100 nmol per paw (c)) or a combination of EP<sub>3</sub> and EP<sub>4</sub> receptor antagonists (L-826266 plus L-161982, each 10 nmol per paw) (d) on PGE<sub>2</sub>-induced paw licking in mice (PGE<sub>2</sub>, 3 nmol per paw). The drugs were co-injected with PGE<sub>2</sub> and the effects of the drugs are expressed as licking time (s). Each column represents the mean of 7–8 animals and vertical lines show the s.e.m. Asterisks denote significant difference levels, \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , compared with the PGE<sub>2</sub> plus vehicle-injected mice (black bar, one-way ANOVA followed by Dunnett's (a and b) or Student–Newman–Keuls' (c and d) test).


(3–30 nmol per paw) or the EP<sub>4</sub> receptor antagonist, L-161982, (1–30 nmol per paw) (Figure 2a and b, respectively). In contrast, the co-injection of the EP<sub>1/2</sub> receptor antagonist, AH-6809 (10–100 nmol per paw) did not alter the paw licking induced by PGE<sub>2</sub> (Figure 2c). The co-administration of EP<sub>3</sub> receptor antagonist L-826266 (10 nmol per paw) plus the EP<sub>4</sub> receptor antagonist L-161982 (10 nmol per paw) with PGE<sub>2</sub> injection (3 nmol per paw) significantly inhibited PGE<sub>2</sub>-induced paw licking (Figure 2d). On the other hand, L-826266 (an EP<sub>3</sub> receptor antagonist – 0.1–10 nmol per paw) inhibited PGE<sub>2</sub>-induced mechanical allodynia (Figure 3a). Local injection of L-161982 (10 nmol per paw) or AH-6809 (100 nmol per paw) was unable to alter the PGE<sub>2</sub>-induced mechanical allodynia in mice (Figure 3b). To gain further evidence about the role of the EP<sub>3</sub> receptor in PGE<sub>2</sub>-induced mechanical allodynia and paw licking, we carried out some additional experiments using 17-phenyl-trinor-PGE<sub>2</sub>, a se-

lective agonist at mouse EP<sub>1</sub> and EP<sub>3</sub> receptors (Figure 4a and b). Similar to PGE<sub>2</sub>, 17-phenyl-trinor-PGE<sub>2</sub> (3 nmol per paw), produced significant paw licking and mechanical allodynia (0.1 nmol per paw) (Figure 4a and b). Co-injection of the EP<sub>3</sub> receptor antagonist, L-826266 (10 nmol per paw), significantly inhibited both the paw licking and mechanical allodynia induced by 17-phenyl-trinor-PGE<sub>2</sub>. The inhibitions observed were 100 and 54 ± 7%, respectively (Figure 4a and b). Thus, 17-phenyl-trinor-PGE<sub>2</sub>-induced paw licking and mechanical allodynia seem to be mediated, at least partly, through an activation of the EP<sub>3</sub> receptor.

To demonstrate that the EP antagonists are also effective against nociceptive responses induced by stimulating endogenous prostanoid synthesis, we tested the effect of EP<sub>3</sub> and EP<sub>4</sub> receptor antagonists in carrageenan-induced mechanical allodynia (Figure 4d). The i.p. injection of carrageenan (300 µg per paw) caused a significant mechanical allodynia



**Figure 3** Effect of i.pl. treatment with selective receptor antagonists for EP<sub>3</sub> (L-826266, 0.1–10 nmol per paw (a)) EP<sub>1/2</sub> (AH-6809, 100 nmol per paw) or EP<sub>4</sub> receptors (L-161982, 10 nmol per paw (b)) on PGE<sub>2</sub>-induced mechanical allodynia (0.1 nmol per paw) in mice. The drugs were co-injected with PGE<sub>2</sub> and the effects of the drugs are expressed as threshold (mg). Each column represents the mean of 6–12 animals and vertical lines show the s.e.m. <sup>#</sup>Denotes significant difference from basal group ( $P < 0.01$ ) (Student–Newman–Keuls' test). Asterisks denote the significance levels,  $**P < 0.01$ , compared with the PGE<sub>2</sub> plus vehicle-injected mice (black bar, one-way ANOVA followed by Dunnett's (a) or Student–Newman–Keuls' (b) test).




**Figure 4** Effect of i.pl. treatment with the selective EP<sub>3</sub> receptor antagonist (L-826266, 10 nmol per paw) on 17-phenyl-trinor-PGE<sub>2</sub>-induced paw licking (3 nmol per paw (a)) or mechanical allodynia (0.1 nmol per paw (b)). In c and d, the effects of i.pl. treatment with the EP<sub>3</sub> receptor antagonist (L-826266, 10 nmol per paw) or the EP<sub>4</sub> receptor antagonist (L-161982, 10 nmol per paw) on paw-licking or mechanical allodynia induced by carrageenan (300 µg per paw) are shown. The effects of the drugs are expressed as licking time (s) or threshold (mg). Each column represents the mean of six animals and vertical lines show the s.e.m. <sup>#</sup>denotes significant difference from basal group ( $P < 0.01$ ) (Student–Newman–Keuls' test). Asterisks denote the significance levels,  $**P < 0.01$  and  $***P < 0.001$ , compared with the PGE<sub>2</sub> plus vehicle-injected mice (black bar, one-way ANOVA followed by Student–Newman–Keuls' test).

(thresholds of  $630 \pm 50$  and  $11 \pm 2$  mg before and 3 h after injection, respectively) (Figure 4d). The co-administration of the EP<sub>3</sub> receptor antagonist (L-826266, 10 nmol per paw) or the EP<sub>4</sub> receptor antagonist (L-161982, 10 nmol per paw) significantly inhibited the mechanical allodynia induced by carrageenan. The inhibitions observed were  $34 \pm 6$  and  $38 \pm 6\%$ , respectively (Figure 4d). Moreover, the i.pl. injection of carrageenan (300 µg per paw) also induced paw licking that was not significantly altered by the co-administration of the EP<sub>3</sub> receptor antagonist (L-826266, 10 nmol per paw) or the EP<sub>4</sub> receptor antagonist (L-161982, 10 nmol per paw) (Figure 4c).


#### Involvement of PKC, PKA and MAPKs in PGE<sub>2</sub>-induced nociception

PGE<sub>2</sub>-induced paw licking and mechanical allodynia was significantly reduced by co-injection of the PKA inhibitor KT-5720 (0.1–3 nmol per paw) (Figure 5a and c). The inhibitions observed were  $87 \pm 5$  and  $42 \pm 6\%$  for paw licking and mechanical allodynia, respectively. However the post-treatment with KT-5720 (3 nmol per paw) did not alter PGE<sub>2</sub>-induced mechanical allodynia or paw licking (Figure 6a and b).

To further confirm the role played by PKA in PGE<sub>2</sub>-induced paw licking, we carried out western blot analysis on samples of the PGE<sub>2</sub>-injected paws (3 nmol per paw). It has been reported previously that autophosphorylation of the PKA regulatory subunit occurs when PKA is activated by cAMP (Tasken and Aandahl, 2004). Thus, the detection of phosphorylation of the PKA regulatory subunit is useful evidence for PKA activation. Our results showed that phosphorylation




**Figure 5** Effect of i.pl. treatment with an inhibitor of PKA (KT-5720, 0.1–3 nmol per paw) (a), inhibitors of PKC (GF109203X, 10 nmol per paw), ERK (PD98059, 30 nmol per paw) or JNK (SP600125, 30 nmol per paw) on PGE<sub>2</sub>-induced paw licking (3 nmol per paw) (b), inhibitors of PKC (GF109203X, 10 nmol per paw), PKA (KT-5720, 3 nmol per paw), ERK (PD98059, 30 nmol per paw), p38 MAPK (SB203580, 30 nmol per paw) or JNK (SP 600125, 30 nmol per paw) on mechanical allodynia induced by PGE<sub>2</sub> (0.1 nmol per paw) (c). The effects of the drugs are expressed as licking time (s) or threshold (mg). Each column represents the mean of six animals and vertical lines show the s.e.m. #denotes significant difference from basal group,  $P < 0.01$  (Student–Newman–Keuls' test). Asterisks denote the significance levels, \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , compared with the PGE<sub>2</sub> plus vehicle-injected mice (black bar, one-way ANOVA followed by Dunnett's (a) or Student–Newman–Keuls' (b and c) test).



**Figure 6** Effect of post-treatment with GF109203X (10 nmol per paw), KT5720 (3 nmol per paw) or PD98059 (30 nmol per paw) on PGE<sub>2</sub>-induced paw-licking (a) or mechanical allodynia (b) in mice. The effects of the drugs are expressed as licking time (s) or threshold (mg). Each column represents the mean of six animals and vertical lines show the s.e.m. #denotes significant difference from basal group,  $P < 0.01$  (Student–Newman–Keuls' test). Asterisks denote the significance levels, \*\*\* $P < 0.001$ , compared with the PGE<sub>2</sub> plus vehicle-injected mice (black bar, one-way ANOVA followed by Student–Newman–Keuls' test).

of the PKA regulatory subunit II (PKA RII) stimulated by PGE<sub>2</sub> occurred within 5 min after PGE<sub>2</sub> injection, an effect that peaked 15 min after PGE<sub>2</sub> administration and lasted for up to 30 min (Figure 7a). Of note, the observed PKA activation had the same time course as the induction of paw licking following i.pl. injection of PGE<sub>2</sub> (Figures 5a and 7a). The co-administration of the receptor antagonists for the EP<sub>3</sub> receptor, L-826266 (10 nmol per paw), or the EP<sub>4</sub> receptor, L-161982, (10 nmol per paw) significantly reduced PKA activation when assessed 15 min after PGE<sub>2</sub> injection (3 nmol per paw) (Figure 8a).

We next investigated the possible role played by PKC in PGE<sub>2</sub>-induced nociceptive responses. The co-administration of the selective PKC inhibitor GF109203X (10 nmol per paw) significantly inhibited ( $32 \pm 5\%$ ) PGE<sub>2</sub>-induced mechanical allodynia, without altering PGE<sub>2</sub>-induced paw licking (Figure 5b and c). Post-treatment with GF 109203X (10 nmol per paw) also inhibited ( $32 \pm 4\%$ ) PGE<sub>2</sub>-induced mechanical allodynia, but not PGE<sub>2</sub>-induced paw licking (Figure 6a and b). Further, we assessed the time-course response for PKC $\alpha$  isoform activation following i.pl. injection of PGE<sub>2</sub> (3 nmol per paw) as assessed by their translocation from cytosol to membrane fractions collected from skin samples of injected paws. We found that the i.pl. injection of PGE<sub>2</sub> stimulated the translocation of PKC $\alpha$  isoform, an effect that peaked 15 min after injection and lasted for until 30 min after PGE<sub>2</sub> injection (Figure 7c and d). Additionally, co-administration with EP<sub>3</sub> (L-826266, 10 nmol per paw) or the EP<sub>4</sub> (L-161982, 10 nmol per paw) receptor antagonists did not significantly reduce PKC $\alpha$  activation when assessed 15 min after PGE<sub>2</sub> injection (3 nmol per paw) (Figure 8c and d).




**Figure 7** Representative images of western immunoblotting and densitometry analyses showing the time course of phosphorylation of PKA RII (phospho-PKA RII; (a)) ERK (phospho-ERK1, (b)) PKC $\alpha$  translocation from cytosol (c) to membrane (d) in response to i.pl. injection of PGE<sub>2</sub> (3 nmol per paw) into mouse paw. Densities for actin are shown in (e). Mouse paw tissues were obtained from naive (basal, B) or PGE<sub>2</sub>-injected mice at the indicated times. Membrane levels of PKC $\alpha$  and cytosolic levels of phospho-PKA RII, PKC $\alpha$ , phospho-ERK1 and actin were determined using specific antibody and indicated PK activation. Results were normalized by arbitrarily setting the densitometry of the basal group and are expressed as mean  $\pm$  s.e.m. ( $n = 3$ ). \* $P < 0.05$ , as compared with basal values (one-way ANOVA followed by Dunnett's test).

Finally, we evaluated through the use of selective inhibitors and western blot analysis whether or not the different MAPKs are involved in the nociceptive responses produced by i.pl. injection of PGE<sub>2</sub>. The MEK (a kinase upstream to ERK) inhibitor PD98059 (30 nmol per paw) partially, but significantly reduced both the paw licking and the mechanical allodynia produced by i.pl. injection of PGE<sub>2</sub>. The observed inhibitions were  $41 \pm 5$  and  $36 \pm 7\%$ , respectively (Figure 5b and c). Post-treatment with PD98059 (30 nmol per paw) did not significantly inhibit PGE<sub>2</sub>-induced mechanical allodynia or paw licking (Figure 6a and b). Western blot analysis detected ERK activation at 10 and 60 min in PGE<sub>2</sub>-injected tissues (Figure 7b). Co-administration with EP<sub>3</sub> (L-826266, 10 nmol per paw), but not EP<sub>4</sub> (L-161982, 10 nmol per paw), receptor antagonists significantly reduced the ERK activation caused by PGE<sub>2</sub> injection (3 nmol per paw) (Figure 8b). The co-injection of SP600125 (30 nmol

per paw), a selective inhibitor of JNK, did not significantly affect PGE<sub>2</sub>-induced nociceptive responses (Figure 5b and c). The inhibitor of p38 MAK, SB203580 (30 nmol per paw) also failed to significantly affect both paw licking and mechanical allodynia induced by i.pl. injection of PGE<sub>2</sub> (Figure 5b and c).

## Discussion and conclusions

There is now a considerable amount of experimental evidence demonstrating the ability of PGE<sub>2</sub> to induce stimulus-evoked nociception. Furthermore, it has been shown that the peripheral injection of PGE<sub>2</sub> into humans or experimental animals induces both allodynia and hyperalgesia in response to mechanical stimulation (Ferreira, 1972; Taiwo and Levine, 1991). There is also evidence that



**Figure 8** Representative images of western immunoblotting and densitometry analyses showing the effect of EP<sub>3</sub> or EP<sub>4</sub> receptor antagonists in PGE<sub>2</sub>-induced phosphorylation of PKA RII (phospho-PKA RII, (a)) of ERK1 (phospho-ERK1, (b)) or PKC $\alpha$  activation (PKC $\alpha$  membrane (c)) and PKC $\alpha$  cytosol (d)). Mouse paw tissues were obtained from naive (basal, B) or 15 min after PGE<sub>2</sub> injection. Cytosolic levels of phospho-PKA RII were determined using a specific antibody. Results were normalized by arbitrarily setting the densitometry of the basal group and are expressed as mean  $\pm$  s.e.m. ( $n=3$ ). # $P<0.05$ , as compared with basal values (B), \* $P<0.05$ , as compared with vehicle group (one-way ANOVA followed by Student–Newman–Keuls' test).

high concentrations of PGE<sub>2</sub> can directly excite peripheral terminals and also cell bodies of nociceptors *in vitro* (Mense, 1981; Mizumura *et al.*, 1987; Schaible and Schimdt, 1988; Noda *et al.*, 1997). For instance, it was reported that testicular nociceptors were excited by the addition of PGE<sub>2</sub> at 1  $\mu$ M, a concentration about 100 times higher than those necessary to sensitize these nociceptors to heat or to bradykinin (Mizumura *et al.*, 1987, 1993). Confirming the previous *in vitro* experiments, our results show that i.pl. injection of PGE<sub>2</sub> to mice caused dose-related paw licking. However, when compared with PGE<sub>2</sub>-induced mechanical allodynia, this prostanoid was about 28-fold less potent in causing paw licking. In agreement with our data, Hong and Abbott (1994) have shown that i.pl. injection of high doses of PGE<sub>2</sub> into the rat paw produces paw licking.

To date, four distinct EP receptors have been cloned in different animal species and all of them are G-protein-coupled receptors (see Kobayashi and Narumiya, 2002 for review). EP<sub>2</sub> and EP<sub>4</sub> receptors stimulate adenylate cyclase via stimulatory G-protein (G<sub>s</sub>) activation, whereas EP<sub>1</sub> receptor activation normally stimulates phosphoinositide metabolism and mobilizes intracellular calcium through phospholipase C-linked G-protein protein stimulation (see Kobayashi and Narumiya, 2002 for review). On the other hand, EP<sub>3</sub> receptor has several splice variants and its signalling pathways seem to involve a complex interaction with inhibitory G-protein, G<sub>s</sub> and G<sub>13</sub> activation (see Hatae *et al.*, 2002; Hata and Breyer, 2004 for review). Furthermore, the restricted

number of available selective agonists/antagonists for EP receptors has greatly limited the pharmacological strategies to elucidate the EP receptor subtypes involved in the nociceptive responses caused by PGE<sub>2</sub> at a peripheral level.

The nociceptive effect of PGE<sub>2</sub> could be mediated by a direct action on sensory neurons as the mRNA of all EP receptors was found in rat DRG (Southall and Vasko, 2001). Moreover, EP<sub>3</sub> and EP<sub>4</sub> receptors have been found both in DRG and in the peripheral nerve terminals of rats, whereas the EP<sub>4</sub> receptors are colocalized with calcitonin gene-related peptide (Kopp *et al.*, 2004). Studies carried out on mouse DRG have mRNA for the EP<sub>3</sub> receptor, and to a lesser extent, for EP<sub>1</sub> and EP<sub>4</sub> receptors in small-sized neurons (Sugimoto *et al.*, 1994; Oida *et al.*, 1995).

L-826266 and L-161982 are highly potent ( $K_i$  values in nanomolar range) and selective receptor antagonists (affinity about 100–200 times greater on EP<sub>3</sub> or EP<sub>4</sub> receptors when compared with other prostanoid receptors) (Machwate *et al.*, 2001; Michel Gallant, unpublished results). Through the use of these selective EP receptor antagonists, our present results support the participation of both EP<sub>3</sub> and EP<sub>4</sub>, but not EP<sub>1</sub> and EP<sub>2</sub> receptors, in PGE<sub>2</sub>-induced paw licking. These results were not unexpected as PGE<sub>2</sub>, at least in mice, had been found to bind with higher affinity to both EP<sub>3</sub> and EP<sub>4</sub> receptors, when compared with its action at EP<sub>1</sub> and EP<sub>2</sub> receptors (Kiriya *et al.*, 1997). Notably, Southall and Vasko (2001) have shown that the combination of antisense RNAs directed to EP<sub>4</sub> plus EP<sub>3</sub> receptors, but not the individual



antisense RNAs, abolished PGE<sub>2</sub>-induced cAMP production and neuropeptide release in cultured rat DRG. In addition, the EP<sub>3</sub> receptor is also able to augment G<sub>s</sub>-coupled EP<sub>4</sub> receptor-stimulated adenylate cyclase activity (Hatae *et al.*, 2002). Our data extend these previous results by demonstrating that PGE<sub>2</sub>-induced paw licking is consistently reduced by PKA inhibition. The role of PKA in the nociception caused by the EP<sub>1/3</sub> receptor agonist 17-phenyl-trinor-PGE<sub>2</sub> must be elucidated in further studies.

Another new finding of the present study is the demonstration, for the first time, that the activation of PKA, PKC and MAPKs could be detected in PGE<sub>2</sub>-injected paws. In addition, the antagonism of either EP<sub>3</sub> receptors or EP<sub>4</sub> receptors reduced PKA and ERK activation in PGE<sub>2</sub>-injected paws. Notably, the time course of activation of PKA and ERK following i.p.l injection of PGE<sub>2</sub> agreed well with its temporal ability to induce paw licking and mechanical allodynia. Burkey and Regan (1995) have reported that PGE<sub>2</sub> is effective in activating ERK in COS-7 cells heterologously expressing EP<sub>3</sub> receptors. Besides, the stimulation of EP<sub>3</sub> receptors by PGE<sub>2</sub> is known to be able to activate other MAPKs (Fiebich *et al.*, 2001; Mendez and Lapointe, 2005). We have found that the peripheral injection of PGE<sub>2</sub> causes activation of p38 and JNK. However, the activation of these kinases does not seem to be related to the production of nociceptive behaviours since selective inhibitors of p38 and JNK did not significantly alter paw licking or mechanical allodynia even in doses where they significantly reduced PGE<sub>2</sub>-induced paw oedema in mice (Claudino *et al.*, 2006). Overall, our results have demonstrated, for the first time, that PKA stimulation largely accounts for PGE<sub>2</sub>-induced paw licking, whereas ERK has apparently only a minor role in this process.

The current knowledge concerning the EP receptor subtype and the intracellular signalling pathways involved in PGE<sub>2</sub>-induced sensitization of nociceptors and hyperalgesia are still controversial. Electrophysiological studies have suggested that both EP<sub>2</sub> and EP<sub>3</sub>, but not EP<sub>1</sub> receptor subtypes mediate PGE<sub>2</sub>-induced sensitization or activation of nociceptors (Kumazawa *et al.*, 1993, 1996; Smith *et al.*, 1998). Furthermore, the intradermal injection of PGE<sub>2</sub> into rats produces mechanical hyperalgesia, an effect that is greatly reduced by EP<sub>1</sub> receptor antagonism and by inhibitors of PKA, but not of PKC or ERK (Khasar *et al.*, 1995, 1999; Aley and Levine, 1999; Cunha *et al.*, 1999; Aley *et al.*, 2001). Experiments using selective EP agonists have suggested the presence of more than one EP receptor subtype mediating PGE<sub>2</sub>-induced mechanical hyperalgesia in the rat paw (Khasar *et al.*, 1995). The present study has extended these findings to mice by showing that EP<sub>3</sub> receptors mediates the mechanical allodynia caused by PGE<sub>2</sub>, based on the data showing that the EP<sub>3</sub> receptor antagonist L-826266 significantly reduced both PGE<sub>2</sub> and 17-phenyl-trinor-PGE<sub>2</sub>-induced mechanical allodynia. However, a role for EP<sub>1</sub> receptor in allodynia cannot be ruled out as 17-phenyl-trinor-PGE<sub>2</sub>-induced mechanical allodynia was not completely prevented by the EP<sub>3</sub> receptor antagonist. In contrast with our pharmacological data, it has been reported that mechanical allodynia produced by PGE<sub>2</sub> was not altered in EP<sub>3</sub> receptor knockout mice (Reinold *et al.*, 2005). The reason

for such discrepant findings is presently unclear and requires additional study. However, we cannot exclude a role for EP<sub>3</sub> receptor in allodynia, bearing in mind the physiological compensation that is often observed in gene-deleted mice (Mak *et al.*, 2001).

Of note, it has been reported that PGE<sub>2</sub> increases the phospholipase C activity as well as the translocation of PKC $\epsilon$  in cultured DRG (Smith *et al.*, 1998; Vellani *et al.*, 2004). Moreover, evidence also suggests that PKC as well as PKA mediates the increase in sodium currents gated by tetrodotoxin-resistant sodium channels caused by PGE<sub>2</sub>, an effect that is essential to PGE<sub>2</sub>-induced mechanical hyperalgesia in rats (Gold *et al.*, 1998; Khasar *et al.*, 1998). To gain further insight into the signalling mechanisms involved in PGE<sub>2</sub>-induced mechanical allodynia, we have demonstrated by the use of both pharmacological tools and western blot analysis that PKA in conjunction with PKC and ERK largely accounts for the PGE<sub>2</sub>-induced mechanical allodynia in mice. The absence of effect of EP<sub>3</sub> receptor antagonists in PKC $\alpha$  translocation suggests that other PKC subtypes, such as PKC $\epsilon$ , are probably stimulated by the activation of EP<sub>3</sub> receptors and can modulate the nociceptive response.

Finally, we have shown that EP<sub>3</sub> or EP<sub>4</sub> receptor antagonists were able to significantly reduce mechanical allodynia induced by carrageenan. It is now well-known that the levels of PGE<sub>2</sub> are greatly increased in the paw tissue at 3 h but not 15 min after carrageenan injection (Herencia *et al.*, 1998). This temporal profile could explain why EP receptor antagonists were not able to reduce paw licking that was observed at 30 min after carrageenan injection. The effect of EP receptor antagonists in models of pain related with earlier increases in peripheral PGE<sub>2</sub> levels must be carried out to elucidate the role of this endogenous prostanoid in paw licking. In contrast to our data obtained with exogenous PGE<sub>2</sub> injection, we found that the EP<sub>4</sub> receptor antagonist was also capable of reducing mechanical allodynia when PGE<sub>2</sub> was endogenously produced. This result indicates that the receptors and signalling pathways stimulated by PGE<sub>2</sub> can be modulated by the presence of other inflammatory mediators.

On the basis of published data, both EP receptor antagonists and PK inhibitors, in the doses used here, seem to produce selective effects on their targets. It has been previously shown that AH-6809, L-826266 and L-161982 were about 235, 25 and 40 times, respectively, less potent than PGE<sub>2</sub> in binding to EP receptors in cells expressing recombinant receptors (Abramovitz *et al.*, 2000; Machwate *et al.*, 2001; Michel Gallant, unpublished results). On the basis of dose ratios, the doses of EP receptor antagonists we used were maximally 1000 times (for AH-6809) or 100 times (for L-826266 and L-161982) higher than the PGE<sub>2</sub> doses. In our opinion, the antinociceptive action of EP receptor antagonists would not reflect non-specific actions of these compounds as they were not capable of significantly altering the carrageenan-induced paw licking. On the other hand, the doses of AH6809 used in the present study seem to be pharmacologically effective as AH-6809, at 100 nmol per paw, can influence PGE<sub>2</sub>-induced paw oedema in mice (Claudino *et al.*, 2006). The choice of dosage for kinase inhibitors was mainly based on the data reported in the

literature, which previously have shown that these compounds produce pronounced antinociceptive actions not dependent on non-specific effects, when intraplantarly injected (Inoue et al., 2003; Ferreira et al., 2004, 2005; Claudino et al., 2006). Accordingly, KT-5720 was found effective to reduce PGE<sub>2</sub>-induced nociception in doses where it also decreased nocistatin-induced nociceptive responses, but did not alter PGE<sub>2</sub>-induced oedema (Inoue et al., 2003; Claudino et al., 2006). Similarly, PD-98059 inhibited PGE<sub>2</sub>- and phorbol ester-induced nociception in mice, in a dose where it failed to interfere with heat-induced paw withdrawal in rats (Zhuang et al., 2004; Ferreira et al., 2005). On the other hand, p38 and JNK inhibitors did not alter PGE<sub>2</sub>-induced allodynia and paw licking in doses where they markedly inhibit phorbol ester-induced nociceptive behaviour and PGE<sub>2</sub>-induced oedema (Ferreira et al., 2005; Claudino et al., 2006). Thus, based in the above evidence, we believe that the effects observed with the drugs we have used certainly reflect their actions on the selected targets.

In summary, our present results show that the i.pl injection of PGE<sub>2</sub> into mice not only sensitizes nociceptors to produce mechanical allodynia, but also induces robust short-lasting paw licking. Although PGE<sub>2</sub>-induced paw licking is probably mediated through activation of EP<sub>3</sub> and EP<sub>4</sub> receptors, with the involvement of both PKA and ERK signalling pathways, PGE<sub>2</sub>-induced mechanical allodynia is most likely mediated through EP<sub>3</sub> receptor, an action that is associated with activation of the PKA, PKC and ERK signalling mechanisms.

## Acknowledgements

We acknowledge the financial support from CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), PRONEX (Programa de Apoio aos Núcleos de Excelência), FAPESC (Fundação de Apoio a Ciência e Tecnologia do Estado de Santa Catarina) and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior), Brazil. Authors acknowledge the Merck Froost (Kirkland, Québec, Canada) that kindly supplied L-828266 and L-1616982. CALK is a PhD student in Pharmacology holding a grant from CNPq. JF was a holder of post-doctoral fellowship from CNPq and RFC is a MSc student in Pharmacology holding a grant from CAPES.

## Conflict of interest

The authors state no conflict of interest.

## References

- Abramovitz M, Adam M, Boie Y, Carriere M, Denis D, Godbout C et al. (2000). The utilization of recombinant prostanoid receptors to determine the affinities and selectivities of prostaglandins and related analogs. *Biochim Biophys Acta* **1483**: 285–293.
- Aley KO, Levine JD (1999). Role of protein kinase A in the maintenance of inflammatory pain. *J Neurosci* **19**: 2181–2186.
- Aley KO, Martin A, McMahon T, Mok J, Levine JD, Messing RO (2001). Nociceptor sensitization by extracellular signal-regulated kinases. *J Neurosci* **21**: 6933–6939.

- André E, Ferreira J, Malheiros A, Yunes RA, Calixto JB (2004). Evidence for the involvement of vanilloid receptor in the antinociception produced by the dialdehydes unsaturated sesquiterpenes polygodial and drimaniol in rats. *Neuropharmacology* **46**: 590–597.
- Burkey TH, Regan JW (1995). Activation of mitogen-activated protein kinase by the human prostaglandin EP3A receptor. *Biochem Biophys Res Commun* **211**: 152–158.
- Chaplan SR, Bach FW, Pogrel JW, Chung JM, Yaksh TL (1994). Quantitative assessment of tactile allodynia in the rat paw. *J Neurosci Methods* **53**: 55–63.
- Claudino RF, Kassuya CAL, Ferreira J, Calixto JB (2006). Pharmacological and molecular characterization of the mechanisms involved in the prostaglandin E<sub>2</sub> (PGE<sub>2</sub>)-induced edematogenic responses in mice. *J Pharmacol Exp Ther* **318**: 611–618.
- Cunha FQ, Teixeira MM, Ferreira SH (1999). Pharmacological modulation of secondary mediator systems-cyclic AMP and cyclic GMP – on inflammatory hyperalgesia. *Br J Pharmacol* **127**: 671–678.
- Ferreira J, Da Silva GL, Calixto JB (2004). Contribution of vanilloid receptors to the overt nociception induced by B2 kinin receptor activation in mice. *Br J Pharmacol* **141**: 787–794.
- Ferreira J, Triches KM, Medeiros R, Calixto JB (2005). Mechanisms involved in the nociception produced by peripheral protein kinase C activation in mice. *Pain* **117**: 171–181.
- Ferreira SH (1972). Prostaglandin, aspirin-like drugs and analgesia. *Nat New Biol* **240**: 200–203.
- Ferreira SH, Nakamura M (1979). I – Prostaglandin hyperalgesia, a cAMP/Ca<sup>2+</sup> dependent process. *Prostaglandins* **18**: 179–190.
- Fiebig BL, Schleicher S, Spleiss O, Czygan M, Hull M (2001). Mechanisms of prostaglandin E<sub>2</sub>-induced interleukin-6 release in astrocytes: possible involvement of EP<sub>4</sub>-like receptors, p38 mitogenactivated protein kinase and protein kinase C. *J Neurochem* **79**: 950–958.
- Gold MS, Levine JD, Correa AM (1998). Modulation of TTX-R INa by PKC and PKA and their role in PGE<sub>2</sub>-induced sensitization of rat sensory neurons *in vitro*. *J Neurosci* **18**: 10345–10355.
- Hata AN, Breyer RM (2004). Pharmacology and signaling of prostaglandin receptors: multiple roles in inflammation and immune modulation. *Pharmacol Ther* **103**: 147–166.
- Hatae N, Sugimoto Y, Ichikawa A (2002). Prostaglandin receptors: advances in the study of EP<sub>3</sub> receptor signaling. *J Biochem* **131**: 781–784.
- Herencia F, Ubeda A, Ferrandiz ML, Terencio MC, Alcaraz MJ, Garcia-Carrascosa M et al. (1998). Anti-inflammatory activity in mice of extracts from Mediterranean marine invertebrates. *Life Sci* **62**: PL115–PL120.
- Hong Y, Abbott FV (1994). Behavioural effects of intraplantar injection of inflammatory mediators in the rat. *Neuroscience* **63**: 827–836.
- Inoue M, Kawashima T, Allen RG, Ueda H (2003). Nocistatin and prepro-nociceptin/orphanin FQ 160–187 cause nociception through activation of Gi/o in capsaicin-sensitive and of Gs in capsaicin-insensitive nociceptors, respectively. *J Pharmacol Exp Ther* **306**: 141–146.
- Khasar SG, Gold MS, Levine JD (1998). A tetrodotoxin-resistant sodium current mediates inflammatory pain in the rat. *Neurosci Lett* **256**: 17–20.
- Khasar SG, Lin YH, Martin A, Dadgar J, McMahon T, Wang D et al. (1999). A novel nociceptor signaling pathway revealed in protein kinase C epsilon mutant mice. *Neuron* **24**: 253–260.
- Khasar SG, Ouseph AK, Chou B, Ho T, Green PG, Levine JD (1995). Is there more than one prostaglandin E receptor subtype mediating hyperalgesia in the rat hindpaw? *Neuroscience* **64**: 1161–1165.
- Kiriya M, Ushikubi F, Kobayashi T, Hirata M, Sugimoto Y, Narumiya S (1997). Ligand binding specificities of the eight types and subtypes of the mouse prostanoid E receptors expressed in Chinese hamster ovary cells. *Br J Pharmacol* **22**: 217–224.
- Kobayashi T, Narumiya S (2002). Prostanoids in health and disease: lessons from receptor-knockout mice. *Adv Exp Med Biol* **507**: 593–597.
- Kopp UC, Cicha MZ, Nakamura K, Nusing RM, Smith LA, Hokfelt T (2004). Activation of EP<sub>4</sub> receptors contributes to prostaglandin E<sub>2</sub>-mediated stimulation of renal sensory nerves. *Am J Physiol Renal Physiol* **287**: 1269–1282.

- Kuhn DC, Willis AL (1973). Prostaglandin E<sub>2</sub>, inflammation and pain threshold in rat paws. *Br J Pharmacol* **49**: 183–184.
- Kumazawa T, Mizumura K, Koda H (1993). Involvement of EP3 subtype of prostaglandin E receptors in PGE<sub>2</sub>-induced enhancement of the bradykinin response of nociceptors. *Brain Res* **632**: 321–324.
- Kumazawa T, Mizumura K, Koda H, Fukusako H (1996). EP receptor subtypes implicated in the PGE<sub>2</sub>-induced sensitization of polymodal receptors in response to bradykinin and heat. *J Neurophysiol* **75**: 2361–2368.
- Machwate M, Harada S, Leu CT, Seedor G, Labelle M, Gallant M *et al.* (2001). Prostaglandin receptor EP(4) mediates the bone anabolic effects of PGE<sub>2</sub>. *Mol Pharmacol* **60**: 36–41.
- Mak TW, Penninger JM, Ohashi PS (2001). Knockout mice: a paradigm shift in modern immunology. *Nat Rev Immunol* **1**: 11–19.
- Malmberg AB, Brandon EP, Idzerda RL, Liu H, Mcknight GS, Basbaum AI (1997). Diminished inflammation and nociceptive pain with preservation of neuropathic pain in mice with a targeted mutation of the type I regulatory subunit of cAMP-dependent protein kinase. *J Neurosci* **17**: 7462–7470.
- Mendez M, Lapointe MC (2005). PGE<sub>2</sub>-induced hypertrophy of cardiac myocytes involves EP4 receptor-dependent activation of p42/44 MAPK and EGFR transactivation. *Am J Physiol Heart Circ Physiol* **288**: H2111–H2117.
- Mense S (1981). Sensitization of group IV muscle receptors to bradykinin by 5-hydroxytryptamine and prostaglandin E<sub>2</sub>. *Brain Res* **225**: 95–105.
- Mizumura K, Minagawa M, Tsujii Y, Kumazawa T (1993). Prostaglandin E<sub>2</sub>-induced sensitization of the heat response of canine visceral polymodal receptors *in vitro*. *Neurosci Lett* **161**: 117–119.
- Mizumura K, Sato J, Kumazawa T (1987). Effects of prostaglandins and other putative chemical intermediaries on the activity of canine testicular polymodal receptors studied *in vitro*. *Pflug Arch* **408**: 565–572.
- Noda K, Ueda Y, Suzuki K, Yoda K (1997). Excitatory effects of algesic compounds on neuronal processes in murine dorsal root ganglion cell culture. *Brain Res* **751**: 348–351.
- Oida H, Namba T, Sugimoto Y, Ushikubi F, Ohishi H, Ichikawa A *et al.* (1995). *In situ* hybridization studies of prostacyclin receptor mRNA expression in various mouse organs. *Br J Pharmacol* **116**: 2828–2837.
- Reinold H, Ahmadi S, Depner UB, Layh B, Heindl C, Hamza M *et al.* (2005). Spinal inflammatory hyperalgesia is mediated by prostaglandin E receptors of the EP2 subtype. *J Clin Invest* **115**: 673–679.
- Samad TA, Sapirstein A, Woolf CJ (2002). Prostanoids and pain: unraveling mechanism therapeutics targets. *Trends Mol Med* **8**: 390–396.
- Schaible HG, Schimdt RF (1988). Excitation and sensitization of fine articular afferents from cat's knee joint by prostaglandin E<sub>2</sub>. *J Physiol* **403**: 91–104.
- Schlemper V, Medeiros R, Ferreira J, Campos MM, Calixto JB (2005). Mechanisms underlying the relaxation response induced by bradykinin in the epithelium-intact guinea-pig trachea *in vitro*. *Br J Pharmacol* **145**: 740–750.
- Smith JA, Amagasu SM, Eglén RM, Hunter JC, Bley KR (1998). Characterization of prostanoid receptor-evoked responses in rat sensory neurones. *Br J Pharmacol* **124**: 513–523.
- Southall MD, Vasko MR (2001). Prostaglandin receptor subtypes, EP<sub>3C</sub> and EP<sub>4</sub>, mediate the prostaglandin E<sub>2</sub>-induced cAMP production and sensitization of sensory neurons. *J Biol Chem* **276**: 16083–16091.
- Sugimoto Y, Shigemoto R, Namba T, Negishi M, Mizuno N, Narumiya S *et al.* (1994). Distribution of the messenger RNA for the prostaglandin E receptor subtype EP3 in the mouse nervous system. *Neuroscience* **62**: 919–928.
- Taiwo YO, Levine JD (1991). Further confirmation of the role of adenylyl cyclase and of cAMP-dependent protein kinase in primary afferent hyperalgesia. *Neuroscience* **44**: 131–135.
- Tasken K, Aandahl EM (2004). Localized effects of cAMP mediated by distinct routes of protein kinase A. *Physiol Rev* **84**: 137–167.
- Vellani V, Zachrisson O, McNaughton PA (2004). Functional bradykinin B1 receptors are expressed in nociceptive neurones and are upregulated by the neurotrophin GDNF. *J Physiol* **560**: 391–401.
- Woolf CJ, Mannion RJ (1999). Neuropathic pain: aetiology, symptoms, mechanism and management. *Lancet* **353**: 1959–1964.
- Zhuang ZY, Xu H, Clapham DE, Ji RR (2004). Phosphatidylinositol 3-kinase activates ERK in primary sensory neurons and mediates inflammatory heat hyperalgesia through TRPV1 sensitization. *J Neurosci* **24**: 8300–8309.
- Zimmermann M (1983). Ethical guidelines for investigations of experimental pain in conscious animals. *Pain* **16**: 109–110.