

5-HT

Overview: 5-HT receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on 5-HT receptors (Hoyer *et al.*, 1994) and subsequently revised (Hartig *et al.*, 1996)) are, with the exception of the ionotropic 5-HT₃ class, 7TM receptors where the endogenous agonist is 5-HT. The diversity of 5-HT receptors is increased by alternative splicing that produces isoforms of the 5-HT_{2A} (nonfunctional), 5-HT_{2C} (nonfunctional), 5-HT₄ and 5-HT₇ receptors. RNA editing produces 5-HT_{2C} receptor isoforms that differ in function, such as efficiency and specificity of coupling to G_{q/11} (reviewed by Sanders-Bush *et al.*, 2003).

Nomenclature	5-HT_{1A}	5-HT_{1B}	5-HT_{1D}	5-HT_{1E}
Other names	—	5-HT _{1Dβ}	5-HT _{1Dα}	—
Ensembl ID	ENSG00000178394	ENSG00000135321	ENSG00000179546	ENSG00000168830
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Selective agonists	8-OH-DPAT, (R)-UH301, U92016A	Sumatriptan, L694247	PNU10929, sumatriptan, L694247	—
Selective antagonists	(±)WAY100635 (8.7), (S)-UH301, NAD299 (robalzotan, 9.2 Johansson <i>et al.</i> , 1997)	SB236057 (8.9), SB224289 (8.5), GR55562 (7.4)	SB714786 (9.1, Ward <i>et al.</i> , 2005) BRL15572 (7.9)	—
Probes	[³ H]-WAY100635 (0.3 nM, Khawaja <i>et al.</i> , 1997), [³ H]-8-OH-DPAT, [¹¹ C]-WAY100635	[³ H]-sumatriptan, [¹²⁵ I]-GTI, [³ H]-GR125743 (2.6 nM, Xie <i>et al.</i> , 1999), [³ H]-L694247	[³ H]-sumatriptan, [¹²⁵ I]-GTI, [³ H]-GR125743 (2.8 nM, Xie <i>et al.</i> , 1999), [³ H]-L694247	[³ H]-5-HT

Nomenclature	5-HT_{1F}	5-HT_{2A}	5-HT_{2B}	5-HT_{2C}
Other names	5-HT _{1Eβ} , 5-HT ₆	D, 5-HT ₂	5-HT _{2F}	5-HT _{1C}
Ensembl ID	ENSG00000179097	ENSG00000102468	ENSG00000135914	ENSG00000147246
Principal transduction	G _{i/o}	G _{q/11}	G _{q/11}	G _{q/11}
Selective agonists	LY334370, LY334864	DOI	DOI, Ro600175, BW723C86	DOI, Ro600175
Selective antagonists	—	ketanserin (8.5–9.5), M100907 (9.4)	RS127445 (9.5), EGIS-7625 (9.0, Kovacs <i>et al.</i> , 2003)	SB242084 (9.0), RS102221 (8.6)
Probes	[³ H]-LY334370 (0.5 nM, Wainwright <i>et al.</i> , 2005), [¹²⁵ I]-LSD	[³ H]-ketanserin (0.45 nM, Bonhaus <i>et al.</i> , 1995), [³ H]-RP62203 (fananserin, 0.13 nM, Maltouros <i>et al.</i> , 1993), [¹¹ C]-M100907 [¹⁸ F]-altanserin	[³ H]-5-HT	[³ H]-mesulergine (0.67 nM, Bonhaus <i>et al.</i> , 1995), [³ H]-LSD

Nomenclature	5-HT₄	5-HT_{5A}	5-HT_{5B}	5-HT₆
Other names	—	5-HT _{5z}	—	—
Ensembl ID	ENSG00000164270	ENSG00000157219	ENSMUSG00000050534	ENSG00000158748
Principal transduction	G _s	G _{i/o}	None identified	G _s
Selective agonists	BIMU8, ML10302, RS67506	—	—	—
Selective antagonists	GR113808 (9.0–9.5), SB204070 (10.8), RS100235 (11.2)	SB699551 (8.0; Corbett <i>et al.</i> , 2005)	—	SB271046 (8.7), SB357134 (7.6), Ro630563 (7.9)
Probes	[³ H]-GR113808 (0.1 nM, Reynolds <i>et al.</i> , 1995), [¹²⁵ I]-SB207710 (86 pM, Brown <i>et al.</i> , 1993), [³ H]-RS57639	[³ H]-5-CT, [¹²⁵ I]-LSD	[³ H]-5-CT, [¹²⁵ I]-LSD	[¹²⁵ I]-SB258585 (1.0 nM, Hirst <i>et al.</i> , 2000), [³ H]-Ro630563 (5 nM, Boess <i>et al.</i> , 1998), [³ H]-5-CT, [¹²⁵ I]-LSD

Nomenclature	5-HT₇
Other names	5-HT _x , 5-HT ₁ -like
Ensembl ID	ENSG00000148680
Principal transduction	G _s
Selective agonists	—
Selective antagonists	SB656104 (8.5), SB269970 (8.5), SB258719 (7.2)
Probes	[³ H]-SB269970 (1.2 nM, Thomas <i>et al.</i> 2000), [³ H]-5-CT, [¹²⁵ I]-LSD, [³ H]-5-HT

Tabulated K_D values refer to binding to human 5-HT receptors with the exception of SB207710 (piglet) and RP62203 (rat). The nomenclature of 5-HT_{1B}/5-HT_{1D} receptors has been revised (Hartig *et al.*, 1996). Only the non-rodent form of the receptor was previously called 5-HT_{1Dβ}; the human 5-HT_{1B} receptor (tabulated) displays a different pharmacology to the rodent forms of the receptor due to Thr335 of the human sequence being replaced by Asn in rodent receptors. NAS-181 is a selective antagonist of the rodent 5-HT_{1B} receptor. Fananserin binds with high affinity to dopamine D₄, in addition to 5-HT_{2A} receptors. The human 5-HT_{5A} receptor has been claimed to couple to several signal transduction pathways when stably expressed in C6 glioma cells (Noda *et al.*, 2003). The human orthologue of the mouse 5-HT_{5B} receptor is nonfunctional due to interruption of the gene by stop codons. In addition to the receptors listed in the table, an 'orphan' receptor, unofficially termed 5-HT_{1P}, has been described (Gershon, 1999).

Abbreviations: BIMU8, (endo-*N*-8-methyl-8-azabicyclo[3.2.1]oct-3-yl)-2,3-dihydro-3-isopropyl-2-oxo-1*H*-benzimidazol-1-carboxamide hydrochloride; BRL15572, 3-[4-(3-chlorophenyl) piperazin-1-yl]-1,1-diphenyl-2-propanol; BW723C86, 1-[5(2-thienylmethoxy)-1*H*-3-indolyl]propan-2-amine hydrochloride; 5-CT, 5-carbox-

amidotryptamine; **8-OH-DPAT**, 8-hydroxy-2-(di-*n*-propylamino)tetralin; **EGIS-7625**, 1-benzyl-4-[(2-nitro-4-methyl-5-amino)-phenyl]-piperazine; **GR55562**, 3-[3-(dimethylamino)propyl]-4-hydroxy-*N*-[4-(4-pyridinyl)phenyl]benzamide; **GR113808**, [1-2[(methylsulphonyl)amino]ethyl]-4-piperidinylmethyl-1-methyl-1*H*-indole-3-carboxylate; **GR125743**, *n*-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-3-methyl-4-(4-pyridinyl)benzamide; **GTI**, 5-hydroxytryptamine-5-*O*-carboxymethylglycyl-tyrosinamide; **L694247**, 2-[5-[3-(4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1*H*-indol-3-yl] ethanamine; **LY334370**, 5-(4-fluorobenzoyl)amino-3-(1-methylpiperidin-4-yl)-1*H*-indole fumarate; **M100907**, (+/-)-2,3-dimethoxyphenyl-1-[2-(4-piperidine)-methanol]; **NAD299**, (*R*)-3-*N,N*-dicyclobutylamino-8-fluoro-[6-3*H*]-3,4-dihydro-2*H*-1-benzopyran-5-carboxamide; **NAS181**, (*R*)-(+)-2-[[[3-(morpholinomethyl)-2*H*-chromen-8-yl]oxy]methyl] morpholine methane sulphonate; **PNU109291**, (*S*)-3,4-dihydro-1-[2-[4-(4-methoxyphenyl)-1-piperazinyl]ethyl]-*N*-methyl-1-*H*-2-benzopyran-6-carboximide; **Ro600175**, (*S*)-2-(6-chloro-5-fluoroindol-1-yl)-1-methylethylamine; **Ro630563**, 4-amino-*N*-[2,6-bis(methylamino)pyridin-4-yl]benzenesulphonamide; **RP62203**, 2-[3-(4-(4-fluorophenyl)-piperazinyl)propyl]naphtho[1,8-*cd*]isothiazole-1,1-dioxide; **RS127445**, (2-amino-4-(4-fluoronaphthyl-1-yl)-6-isopropylpyrimidine); **RS57639**, 4-amino-5-chloro-2-methoxy benzoic acid 1-(3-[2,3-dihydrobenzo[1,4]dioxin-6-yl]-propyl)-piperidin-4-yl methyl ester; **RS100235**, 1-(8-amino-7-chloro-1,4-benzodioxan-5-yl)-5-((3-(3,4-dimethoxyphenyl)prop-1-yl)piperidin-4-yl)propan-1-one; **RS102221**, 8-[5-(5-amino 2,4-dimethoxyphenyl) 5-oxopentyl]-1,3,8-triazaspiro[4,5]decane-2,4-dione; **SB204070**, 1-butyl-4-piperidinylmethyl-8-amino-7-chloro-1,4-benzodioxan-5-carboxylate; **SB207710**, 1-butyl-4-piperidinylmethyl-8-amino-7-iodo-1,4-benzodioxan-5-carboxylate; **SB224289**, 1'-methyl-5[[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]carbonyl-2,3,6,7-tetrahydrospiro[furo[2,3-*f*]indole-3,4'-piperidine]oxalate; **SB236057**, 1'-ethyl-5-(2'-methyl-4'-(5-methyl-1,3,4-oxadiazol-2-yl)biphenyl-4-carbonyl)-2,3,6,7-tetrahydrospiro[furo[2,3-*f*]indol-3,4'-piperidine; **SB242084**, 6-chloro-5-methyl-1-[2-(2-methylpyridyl-3-oxy)-pyrid-5-yl] carbamoyl] indoline; **SB258585**, 4-iodo-*N*-[4-methoxy-3-(4-methylpiperazin-1-yl)-phenyl]-benzenesulphonamide; **SB258719**, (*R*)-3-*N*-dimethyl-*N*-[1-methyl-3-(4-methylpiperidin-1-yl)propyl]benzene sulphonamide; **SB269970**, (*R*)-3-(2-(2-(4-methylpiperidin-1-yl)ethyl)pyrrolidine-1-sulphonyl)-phenol; **SB271046**, 5-chloro-*N*-(4-methoxy-3-piperazin-1-yl-phenyl)-3-methyl-2-benzothiophenesulphonamide; **SB357134**, *N*-(2,5-dibromo-3-fluorophenyl)-4-methoxy-3-piperazin-1-ylbenzenesulphonamide; **SB656104**, 6-((*R*)-[2-[2-(4-Chloro-phenoxy)-piperidin-1-yl]-ethyl]-pyrrolidine-1-sulphonyl)-1*H*-indole hydrochloride; **SB699551**, 3-cyclopentyl-*N*-[2-(dimethylamino)ethyl]-*N*-[4'-(2-phenylethyl)amino]methyl-4-biphenylmethyl]propanamide dihydrochloride; **SB714786**, 2-methyl-5-((2-[4-(8-quinolinylmethyl)-1-piperazinyl]ethyl)oxy)quinoline; **SR57227**, 4-amino-(6-chloro-2-pyridyl)-1-piperidine hydrochloride; **UH301**, 5-fluoro-8-hydroxy-2-(dipropylamino) tetralin; **U92016A**, (+)-(*R*)-2-cyano-*N,N*-dipropyl-8-amino-6,7,8,9-tetrahydro-3*H*-benz[e]indole; **WAY100635**, *N*-(2-(4-(2-methoxyphenyl)-1-piperazinyl)ethyl)-*N*-(2-pyridyl)-cyclohexanecarboxamide trichloride

Further Reading:

- BARNES, N.M. & SHARP, T. (1999). A review of central 5-HT receptors and their function. *Neuropharmacology*, **38**, 1083–1152.
- BOCKAERT, J., CLAEYSEN, S., COMPAN, V. & DUMUIS, A. (2004). 5-HT₄ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 39–51.
- BONASERA, S.J. & TECOTT, L.H. (2000). Mouse models of serotonin receptor function: toward a genetic dissection of serotonin systems. *Pharmacol. Ther.*, **88**, 133–142.
- GERSHON, M.D. (1999). Roles played by 5-hydroxytryptamine in the physiology of the bowel. *Aliment. Pharmacol. Ther.*, **13** (Suppl. 2), 15–30.
- GLENNON, R.A. (2003). Higher-end serotonin receptors: 5-HT₅, 5-HT₆, and 5-HT₇. *J. Med. Chem.*, **46**, 2795–2812.
- HARTIG, P.R., HOYER, D., HUMPHREY, P.P. & MARTIN, G.R. (1996). Alignment of receptor nomenclature with the human genome: classification of 5-HT_{1B} and 5-HT_{1D} receptor subtypes. *Trends Pharmacol. Sci.*, **17**, 103–105.
- HOYER, D., CLARKE, D.E., FOZARD, J.R., HARTIG, P.R., MARTIN, G.R., MYLECHARANE, E.J., SAXENA, P.R. & HUMPHREY, P.P. (1994). International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (serotonin). *Pharmacol. Rev.*, **46**, 157–203.
- HOYER, D. & MARTIN, G. (1997). 5-HT receptor classification and nomenclature: towards a harmonization with the human genome. *Neuropharmacology*, **36**, 419–428.
- LANFUMEY, L. & HAMON, M. (2004). 5-HT₁ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 1–10.
- LEYSEN, J.E. (2004). 5-HT₂ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 11–26.
- NELSON, D.L. (2004). 5-HT₅ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 53–58.
- PAUWELS, P.J. (2000). Diverse signalling by 5-hydroxytryptamine (5-HT) receptors. *Biochem. Pharmacol.*, **60**, 1743–1750.
- SANDERS-BUSH, E., FENTRESS, H & HAZELWOOD, L. (2003). Serotonin 5-HT₂ receptors: molecular and genomic diversity. *Mol. Interv.*, **3**, 319–330.
- THOMAS, D.R. & HAGAN, J.J. (2004). 5-HT₇ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 81–90.
- WOOLLEY, M.L., MARSDEN, C.A. & FONE, K.C. (2004). 5-HT₆ receptors. *Curr. Drug Targets CNS Neurol. Disord.*, **3**, 59–79.

References:

- BERG, S. et al. (1998). *J. Med. Chem.*, **41**, 1934–1942.
- BOESS, F.G. et al. (1998). *Mol. Pharmacol.*, **54**, 577–583.
- BONHAUS, D.W. et al. (1995). *Br. J. Pharmacol.*, **115**, 622–628.
- BROWN, A.M. et al. (1993). *Br. J. Pharmacol.*, **110**, 10P.
- CORBETT, D.F. et al. (2005). *Bioorg. Med. Chem. Lett.*, **15**, 4014–4018.
- HIRST, W.D. et al. (2000). *Br. J. Pharmacol.*, **130**, 1597–1605.
- JOHANSSON, L. et al. (1997). *J. Pharmacol. Exp. Ther.*, **283**, 216–225.
- KOVACS, A. et al. (2003). *Cardiovasc. Drugs. Ther.*, **17**, 427–434.
- KHAWAJA, X. et al. (1997). *Life Sci.*, **60**, 653–665.
- MALGOURIS, C. et al. (1993). *Eur. J. Pharmacol.*, **233**, 37–45.
- NODA, M. et al. (2003). *J. Neurochem.*, **84**, 222–232.
- REYNOLDS, G.P. et al. (1995). *Br. J. Pharmacol.*, **114**, 993–998.
- THOMAS, D.R. et al. (2000). *Br. J. Pharmacol.*, **130**, 409–417.
- WAINSCOTT, D.B. et al. (2005). *Naunyn Schmiedeberg Arch. Pharmacol.*, **371**, 169–177.
- WARD, S.E. et al. (2005). *J. Med. Chem.*, **48**, 3478–3480.
- XIE, Z. et al. (1999). *FEBS Lett.*, **456**, 63–67.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Acetylcholine (muscarinic)

Overview: Muscarinic acetylcholine receptors (nomenclature as agreed by NC-IUPHAR subcommittee on Muscarinic Acetylcholine Receptors, Caulfield and Birdsall, 1998) are 7TM receptors of the rhodopsin-like family where the endogenous agonist is acetylcholine. In addition to the agents listed in the table, AC-42 and desmethylozapine have recently been described as selective agonists of the M₁ receptor subtype *via* binding to a site distinct to that recognised by nonselective agonists (Spalding *et al.*, 2002; Sur *et al.*, 2003). There are two allosteric sites on muscarinic receptors, one defined by the binding of gallamine, strychnine and brucine and the other binds KT5720, WIN62,577, WIN51,708 and staurosporine (Lazareno *et al.*, 2000, 2002). There are selective enhancers of acetylcholine binding and action; brucine and KT5720 at M₁ receptors, PG135 at M₂ receptors, *N*-chloromethylbrucine and WIN62,577 at M₃ receptors and thiochrome at M₄ receptors (Lazareno *et al.*, 1998, 1999, 2000, 2002, 2004). The allosteric site for gallamine and strychnine on M₂ receptors can be labelled by [³H]-dimethyl-W84 (Tränkle *et al.*, 2003).

Nomenclature	M ₁	M ₂	M ₃
Ensembl ID	ENSG00000168539	ENSG00000181072	ENSG00000133019
Principal transduction	G _{q/11}	G _{i/o}	G _{q/11}
Antagonists	MT7 (9.8), 4-DAMP (8.6–9.2), triptiramine (8.4–8.8), pirenzepine (7.8–8.5), guanylpirenzepine (7.7), darifenacin (7.5–7.8), AFDX384 (7.3–7.5), MT3 (7.1), himbacine (7.0–7.2), PD102807 (5.3)	triptiramine (9.4–9.6), AFDX384 (8.2–9.0), himbacine (8.0–8.3), 4-DAMP (7.8–8.4), darifenacin (7.0–7.4), pirenzepine (6.3–6.7), MT7 (<6), MT3 (<6), PD102807 (5.7), guanylpirenzepine (5.5)	4-DAMP (8.9–9.3), darifenacin (8.4–8.9), AFDX384 (7.2–7.8), triptiramine (7.1–7.4), himbacine (6.9–7.4), pirenzepine (6.7–7.1), guanylpirenzepine (6.5), PD102807 (6.2), MT3 (<6), MT7 (<6)
Probes	[³ H]-NMS (80–150 pM), [³ H]-QNB (15–60 pM), [³ H]-pirenzepine (3–15 nM), [¹¹ C]-xanomeline [¹¹ C]-methylpyridine [¹¹ C]-butylthio-TZTP	[³ H]-NMS (200–400 pM), [³ H]-QNB (20–50 pM), [¹⁸ F]-FP-TZTP	[³ H]-NMS (150–250 pM), [³ H]-QNB (30–90 pM), [³ H]-darifenacin (300 pM)

Nomenclature	M ₄	M ₅
Ensembl ID	ENSG00000180720	ENSG00000184984
Principal transduction	G _{i/o}	G _{q/11}
Antagonists	MT3 (8.7), 4-DAMP (8.4–9.4), himbacine (8.0–8.8), AFDX384 (8.0–8.7), triptiramine (7.8–8.2), darifenacin (7.7–8.0), PD102807 (7.3), pirenzepine (7.1–8.1), guanylpirenzepine (6.5), MT7 (<6)	4-DAMP (8.9–9.0), darifenacin (8.0–8.1), triptiramine (7.3–7.5), guanylpirenzepine (6.8), pirenzepine (6.2–7.1), AFDX384 (6.3), himbacine (6.1–6.3), MT3 (<6), MT7 (<6), PD102807 (5.2)
Probes	[³ H]-NMS (50–100 pM), [³ H]-QNB (20–80 pM)	[³ H]-NMS (500–700 pM), [³ H]-QNB (20–60 pM)

MT3 (m4-toxin) and MT7 (m1-toxin1) are toxins contained with the venom of the Eastern green mamba (*Dendroaspis augusticeps*) (see Bradley, 2000; Potter *et al.*, 2004).

Abbreviations: **AC-42**, 4-*n*-butyl-1-[4-(2-methylphenyl)-4-oxo-1-butyl]-piperidine hydrogen chloride; **AFDX384**, (±)-5,11-dihydro-11-[[2-[2-(dipropylamino)-methyl]-1-piperidinyl]ethyl]amino]carbonyl]-6*H*-pyrido[2,3-*b*](1,4)benzodiazepine-6-one; **Butylthio-TZTP**, butylthio-thiadiazolyltetrahydro-1-methyl-pyridine **Dimethyl-W84**, *N,N'*-bis[3-(1,3-dihydro-1,3-dioxo-4-methyl-2*H*-isoindol-2-yl)propyl]-*N,N,N'*-tetramethyl-1,6-hexanediaminium diiodide; **FP-TZTP**, [3-(3-Fluoropropyl)thio]-1,2,5-thiadiazol-4-yl]-1,2,5,6-tetrahydro-1-methylpyridine; **4-DAMP**, 4-diphenylacetoxy-*N*-methylpiperidine methiodide; **KT5720**, (9*S*,10*S*,12*R*)-2,3,9,10,11,12-hexahydro-10-hydroxy-9-methyl-1-oxo-9,12-epoxy-1*H*-diindolo[1,2,3-*fg*:3',2',1'-*k*]pyrrolo[3,4-*η*][1,6]benzodiazocine-10-carboxylic acid hexyl ester; **NMS**, *N*-methylscopolamine; **PD102807**, 9-methoxy-2-methyl-11,12-dihydro-3*H*,6*αH*,13*H*-6-oxa-3,12*α*-diazabenzocyclopenta(*h*)anthracene-1-carboxylic acid ethyl ester; **PG135**, (3*aS*,12*R*,12*aS*,12*bR*)-2-amino-2,3,3*a*,4,11,12*a*,12*b*-octahydro-10-hydroxyisoquinolo[2,1,8-*lm*]carbazol-5(1*H*)-one hydrochloride; **QNB**, 3-quinuclidinylbenzilate; **WIN51,708**, 17-β-hydroxy-17-α-ethynyl-5-α-androstano[3,2-*b*]pyrimido[1,2-*a*]benzimidazole; **WIN62,577**, 17-β-hydroxy-17-α-ethynyl-Δ⁴-androstano[3,2-*b*]pyrimido[1,2-*a*]benzimidazole

Further Reading:

ABRAMS, P., ANDERSSON, K.E., BUCCAFUSCO, J.J., CHAPPLE, C., DE GROAT, W.C., FRYER, A.D., KAY, G., LATIES, A., NATANSON, N.M., PASRICHA, P.J. and WEIN, A.J. (2006). Muscarinic receptors: their distribution and function in the body systems, and the implications for treating overactive bladder. *Br. J. Pharmacol.*, **148**, 565–578.

BIRDSALL, N.J.M. & LAZARENO, S. (2005). Allosterism at muscarinic receptors: ligands and mechanisms. *Mini Rev. Med. Chem.*, **5**, 523–543.

BRADLEY, K.N. (2000). Muscarinic toxins from the green mamba. *Pharmacol. Ther.*, **85**, 87–109.

CAULFIELD, M.P. & BIRDSALL, N.J.M. (1998). International Union of Pharmacology XVII Classification of muscarinic acetylcholine receptors. *Pharmacol. Rev.*, **50**, 279–290.

EGLIN, R.M. (2005). Muscarinic receptor subtype pharmacology and physiology. *Prog. Med. Chem.*, **43**, 105–136.

EGLIN, R.M. (2006). Muscarinic receptor subtypes in neuronal and non-neuronal cholinergic function. *Auton-Autocoid Pharmacol.*, **26**, 219–233.

ISHII, M. & KURACHI, Y. (2006). Muscarinic acetylcholine receptors. *Curr. Pharm. Des.*, **12**, 3573–3581.

POTTER, L.T., FLYNN, D.D., LIANG, J.S. & MCCOLLUM, M.H. (2004). Studies of muscarinic neurotransmission with antimuscarinic toxins. *Prog. Brain Res.*, **145**, 121–128.

WESS, J. (2004). Muscarinic acetylcholine receptor knockout mice: novel phenotypes and clinical implications. *Annu. Rev. Pharmacol. Toxicol.*, **44**, 423–450.

References:

LAZARENO, S. *et al.* (1998). *Mol. Pharmacol.*, **53**, 573–589.

LAZARENO, S. *et al.* (1999). *Life Sci.*, **64**, 519–526.

LAZARENO, S. *et al.* (2000). *Mol. Pharmacol.*, **58**, 194–207.

LAZARENO, S. *et al.* (2002). *Mol. Pharmacol.*, **62**, 1492–1505.

LAZARENO, S. *et al.* (2004). *Mol. Pharmacol.*, **65**, 257–266.

SPALDING, T.A. *et al.* (2002). *Mol. Pharmacol.*, **61**, 1297–1302.

SUR, C. *et al.* (2003). *Proc. Natl. Acad. Sci. USA*, **100**, 13674–13679.

TRÄNKLE, C. *et al.* (2003). *Mol. Pharmacol.*, **64**, 180–190.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Adenosine

Overview: Adenosine receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Adenosine Receptors; Fredholm *et al.*, 2001) are activated by the endogenous ligand adenosine (potentially inosine also at A₃ receptors). NECA is a nonselective agonist, while XAC and CGS15943 display submicromolar affinity at all four adenosine receptors (Klotz *et al.*, 1998; Ongini *et al.*, 1999).

Nomenclature	A ₁	A _{2A}	A _{2B}	A ₃
Ensembl ID	ENSG00000163485	ENSG00000128271	ENSG00000170425	ENSG00000121933
Principal transduction	G _{i/o}	G _s	G _s	G _{i/o}
Selective agonists	CPA, CCPA, S-ENBA, GR79236	CGS21680, HENECA	—	2-Cl-IB-MECA, IB-MECA
Selective antagonists	DPCPX (8.5)	ZM241385 (9.0), SCH58261 (7.9–9.5)	MRS1754 (8.7), MRS1706 (8.4), PSB1115 (7.7)	MRS1220 (8.8), VUF8504 (7.8, van Muijlwijk-Koezen <i>et al.</i> , 1998), MRS1523 (7.7), MRS1191 (7.0)
Probes	[³ H]-CCPA, [³ H]-DPCPX (0.6–1.2 nM)	[³ H]-CGS21680, [³ H]-ZM241385 (0.8 nM)	[³ H]-MRS1754 (1.1 nM)	[¹²⁵ I]-AB-MECA (0.6 nM)

Adenosine inhibits many intracellular ATP-utilising enzymes, including adenylyl cyclase (P-site). A pseudogene exists for the A_{2B} adenosine receptor (ENSG00000182537) with 79% identity to the A_{2B} adenosine receptor cDNA coding sequence, but which is unable to encode a functional receptor (Jacobson *et al.*, 1995). DPCPX also exhibits antagonism at A_{2B} receptors (pK_i ca. 7, Alexander *et al.*, 1996; Klotz *et al.*, 1998). HENECA also shows activity at A₃ receptors (Varani *et al.*, 1998). Antagonists at A₃ receptors exhibit marked species differences, such that only MRS1523 and MRS1191 are selective at the rat A₃ receptor. In the absence of other adenosine receptors, [³H]-DPCPX and [³H]-ZM241385 can also be used to label A_{2B} receptors (K_D ca. 30 and 60 nM, respectively). [¹²⁵I]-AB-MECA also binds to A₁ receptors (Klotz *et al.*, 1998). [³H]-CGS21680 is relatively selective for A_{2A} receptors, but may also bind to other sites in cerebral cortex (Johansson & Fredholm, 1995; Cunha *et al.*, 1996). [³H]-NECA binds to other nonreceptor elements, which also recognise adenosine (e.g. Lorenzen *et al.*, 1996).

Adenosine has also been reported to act as a partial agonist at the ghrelin receptor (Smith *et al.*, 2000), but this has recently been questioned (Johansson *et al.*, 2005).

Abbreviations: AB-MECA, N⁶-(4-aminobenzyl)-adenosine-5'-N-methyluronamide; CCPA, 2-chloro-N⁶-cyclopentyladenosine; CGS15943, 5-amino-9-chloro-2-(2-furyl)1,2,4-triazolo[1,5-c]quinazoline; CGS21680, 2-(4-[2-carboxyethyl]-phenethylamino)adenosine-5'-N-ethyluronamide; 2Cl-IB-MECA, 2-chloro-N⁶-(3-iodobenzyl)adenosine-5'-N-methyluronamide; CPA, N⁶-cyclopentyladenosine; DPCPX, 8-cyclopentyl-1,3-dipropylxanthine; GR79236, N-[(1s,2s)-2-hydroxycyclopentyl]adenosine; HENECA, 2-(1-(E)-hexenyl)adenosine-5'-N-ethyluronamide; MRS1191, 6-phenyl-1,4-dihydropyridine; MRS1220, 9-chloro-2-(2-furyl)5-phenylacetylaminol[1,2,4]triazolo[1,5-c]quinazoline; MRS1523, 2,3-ethyl-4,5-dipropyl-6-phenylpyridine-3-thiocarboxylate-5-carboxylate; MRS1706, N-(4-acetylphenyl)-2-(4-[2,3,6,7-tetrahydro-2,6-dioxo-1,3-dipropyl-1H-purin-8-yl]phenoxy)acetamide; MRS1754, 8-(4-[(4-cyanophenyl)carbamoylmethyl]oxy)phenyl)-1,3-di(n-propyl)-xanthine; NECA, adenosine-5'-N-ethyluronamide; PSB1115, 4-[2,3,6,7-tetrahydro-2,6-dioxo-1-propyl-1H-purin-8-yl]benzenesulphonic acid; S-ENBA, (2S)-N⁶-(2-endonorbanyl)adenosine; SCH58261, 5-amino-2-(2-furyl)-7-phenylethyl-pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine; VUF8504, 4-methoxy-N-[2-(2-pyridinyl)quinazolin-4-yl]benzamide; XAC, 8-(4-[(2-aminoethyl)amino]carbonylmethyl]oxy)phenyl)-1,3-dipropylxanthine; also known as xanthine amine congener; ZM241385, 4-(2-[7-amino-2-(2-furyl){1,2,4}triazolo{2,3-a}{1,3,5}triazin-5-yl amino]ethyl)phenol

Further Reading:

FREDHOLM, B.B., CHEN, J.F., MASINO, S.A. & VAUGEOIS, J.M. (2005). Actions of adenosine at its receptors in the CNS: insights from knockouts and drugs. *Annu. Rev. Pharmacol. Toxicol.*, **45**, 385–412.

FREDHOLM, B.B., IJZERMAN, A.P., JACOBSON, K.A., KLOTZ, K.N. & LINDEN, J. (2001). International Union of Pharmacology. XXV. Nomenclature and Classification of Adenosine Receptors. *Pharmacol. Rev.*, **53**, 527–552.

JACOBSON, K.A. & GAO, Z.G. (2006). Adenosine receptors as therapeutic targets. *Nat. Rev. Drug Discovery*, **5**, 247–264.

KLINGER, M., FREISSMUTH, M. & NANOFF, C. (2002). Adenosine receptors: G protein-mediated signalling and the role of accessory proteins. *Cell Signal.*, **14**, 99–108.

LATINI, S. & PEDATA, F. (2001). Adenosine in the central nervous system: release mechanisms and extracellular concentrations. *J. Neurochem.*, **79**, 463–484.

YAAR, R., JONES, M.R., CHEN, J.F. & RAVID, K. (2005). Animal models for the study of adenosine receptor function. *J. Cell. Physiol.*, **202**, 9–20.

References:

- ALEXANDER, S.P.H. *et al.* (1996). *Br. J. Pharmacol.*, **119**, 1286–1290.
- CUNHA, R.A. *et al.* (1996). *Naunyn-Schmiedeberg Arch. Pharmacol.*, **353**, 261–271.
- JACOBSON, M.A. *et al.* (1995). *Genomics*, **27**, 374–376.
- JOHANSSON, B. & FREDHOLM, B.B. (1995). *Neuropharmacology*, **34**, 393–403.
- JOHANSSON, S. *et al.* (2005). *Biochem. Pharmacol.*, **70**, 598–605.
- KLOTZ, K.-N. *et al.* (1998). *Naunyn-Schmiedeberg Arch. Pharmacol.*, **357**, 1–9.
- LORENZEN, A. *et al.* (1996). *Biochem. Pharmacol.*, **52**, 1375–1385.
- ONGINI, E. *et al.* (1999). *Naunyn-Schmiedeberg Arch. Pharmacol.*, **359**, 7–10.
- SMITH, R.G. *et al.* (2000). *Biochem. Biophys. Res. Commun.*, **276**, 1306–1313.
- VAN MUIJLWIJK-KOEZEN, J.E. *et al.* (1998). *J. Med. Chem.*, **41**, 3994–4000.
- VARANI, K. *et al.* (1998). *Life Sci.*, **63**, PL81–PL87.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Adiponectin

Overview: Adiponectin receptors (provisional nomenclature, ENSF00000002482) respond to the 30 kDa complement-related protein hormone adiponectin (also known as adipocyte, C1q and collagen domain-containing protein; ACRP30, adipose most abundant gene transcript 1; apM-1; gelatin-binding protein; ENSG00000181092) originally cloned from adipocytes (Maeda *et al.*, 1996). Although sequence data suggest 7TM domains, immunological evidence indicates that, contrary to typical 7TM topology, the carboxyl terminus is extracellular, while the amino terminus is intracellular (Yamauchi *et al.*, 2003). Signalling through these receptors appears to avoid G-proteins and rather activates protein phosphorylation *via* AMP-activated protein kinase and MAP kinase pathways (Yamauchi *et al.*, 2003).

Nomenclature	Adipo1	Adipo2
Other names	AdipoR1, progestin and adipoQ receptor family member I, CGI45	AdipoR2, progestin and adipoQ receptor family member II
Ensembl ID	ENSG00000159346	ENSG00000006831
Rank order of potency	gAdipo > flAdipo	gAdipo = flAdipo

T-Cadherin has also been suggested to be a receptor for (hexameric) adiponectin (Hug *et al.*, 2004).

Abbreviations: flAdipo, full-length adiponectin; gAdipo, globular adiponectin

Further Reading:

DIEZ, J.J. & IGLESIAS, P. (2003). The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur. J. Endocrinol.*, **148**, 293–300.

GIL-CAMPOS, M., CANETE, R.R. & GIL, A. (2004). Adiponectin, the missing link in insulin resistance and obesity. *Clin. Nutr.*, **23**, 963–974.

GOLDSTEIN, B.J. & SCALIA, R. (2004). Adiponectin: a novel adipokine linking adipocytes and vascular function. *J. Clin. Endocrinol. Metab.*, **89**, 2563–2568.

HALUZIK, M., PARIZKOVA, J. & HALUZIK, M.M. (2004). Adiponectin and its role in the obesity-induced insulin resistance and related complications. *Physiol. Res*, **53**, 123–129.

HUG, C. & LODISH, H.F. (2005). The role of the adipocyte hormone adiponectin in cardiovascular disease. *Curr. Opin. Pharmacol.*, **5**, 129–134.

KADOWAKI, T. & YAMAUCHI, T. (2005). Adiponectin and adiponectin receptors. *Endocr. Rev.*, **26**, 439–451.

LAFONTAN, M. (2005). Fat cells: afferent and efferent messages define new approaches to treat obesity. *Annu. Rev. Pharmacol. Toxicol.*, **45**, 119–146.

OUCHI, N., SHIBATA, R., & WALSH, K. (2006). Cardioprotection by adiponectin. *Trends. Cardiovasc. Med.*, **16**, 141–146.

WOLF, G. (2003). Adiponectin: a regulator of energy homeostasis. *Nutr. Rev.*, **61**, 290–292.

References:

HUG, C. *et al.* (2004). *Proc. Natl. Acad. Sci. U.S.A.*, **101**, 10308–10313.

MAEDA, K. *et al.* (1996). *Biochem. Biophys. Res. Commun.*, **221**, 286–289.

YAMAUCHI, T. *et al.* (2003). *Nature*, **423**, 762–769.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Adrenoceptors, α₁

Overview: α₁-Adrenoceptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Adrenoceptors, Bylund *et al.*, 1994) are 7TM receptors, where the endogenous agonists adrenaline and noradrenaline display equal potency. Phenylephrine, methoxamine and cirazoline are examples of agonists selective for α₁-adrenoceptors relative to α₂-adrenoceptors, while prazosin (8.5–10.5) and corynanthine (6.5–7.5) are considered selective for α₁-adrenoceptors relative to α₂-adrenoceptors. [³H]-Prazosin (0.25 nM) and [¹²⁵I]-HEAT (0.1 nM; also known as BE2254) are relatively selective radioligands. Numerous splice variants of the α₁-adrenoceptors exist, some of which may display a different spectrum of signalling properties.

Nomenclature	α _{1A}	α _{1B}	α _{1D}
Other names	α _{1A} , α _{1C}	α _{1b}	α _{1A/D} , α _{1a/d}
Ensembl ID	ENSG00000120907	ENSG00000170214	ENSG00000171873
Principal transduction	G _{q/11}	G _{q/11}	G _{q/11}
Selective agonists	A61603	—	—
Selective antagonists	KMD3213 (10.4), (+)niguldipine (10.0), SNAP5089 (9.7), RS17053 (9.2), SNAP5272 (8.4)	—	BMY7378 (8.4)

The clone originally called the α_{1C}-adrenoceptor corresponds to the pharmacologically defined α_{1A}-adrenoceptor (see Hieble *et al.*, 1995). Some tissues possess α₁-adrenoceptors that display relatively low affinity in functional and binding assays (< 1 nM) for prazosin that might represent different receptor states (termed α_{1L}-adrenoceptors). α_{1A}-adrenoceptor C-terminal splice variants form homo and heterodimers, but fail to generate a functional α_{1A} adrenoceptor (Ramsay *et al.*, 2004). α_{1D}-Adrenoceptors form heterodimers with α_{1B}- or β₂-adrenoceptors that show increased cell-surface expression (Uberty *et al.*, 2005). Heterodimers formed between α_{1D}- and α_{1B}-adrenoceptors have distinct functional properties (Hague *et al.*, 2004). (+)Niguldipine also has high affinity for L-type Ca²⁺ channels.

Abbreviations: **A61603**, *N*-(5-[4,5-dihydro-1*H*-imidazol-2-yl]-2-hydroxy-5,6,7,8-tetrahydronaphthalen-1-yl)methanesulfonamide hydrobromide; **BMY7378**, 8-(2-[4-{2methoxyphenyl}-1-piperazinyl]ethyl)-8-azaspiro[4,5]decane-7,9-dione dihydrochloride; **HEAT**, 2-β-4-hydroxy-3-iodophenylethylaminomethyltetralone; ICI118551: (–)-1-(2,3-[dihydro-7-methyl-1*H*-inden-4-yl]oxy)-3-([1-methylethyl]-amino)-2-butanol; **KMD3213**, (–)-(*R*)-1-(3-hydroxypropyl)-5-(2-[2-(2,2,2-trifluoroethoxy)-phenoxy]ethylamino]propyl)indoline-7-carboxamide; **RS17053**, *N*-[2-(2-cyclopropylmethoxyphenoxy)ethyl]-5-chloro-α,α-dimethyl-1*H*-indole-3-ethanamide; **SNAP5089**, 2,6-dimethyl-4-(4-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate-*N*-[3-(4,4-diphenylpiperidin-1-yl)propyl]amide methyl ester; **SNAP5272**, 5-carboxamide-2,6-diethyl-1,4-dihydro-3-[*N*-(3-[4-hydroxy-4-phenylpiperidinyl]propyl)]carboxamido-4-(4-nitrophenyl)

Further Reading:

BUSCHER, R., HERRMANN, V. & INSEL, P.A. (1999). Human adrenoceptor polymorphisms: evolving recognition of clinical importance. *Trends Pharmacol. Sci.*, **20**, 94–99.

BYLUND, D.B., EIKENBERG, D.C., HIEBLE, J.P., LANGER, S.Z., LEFKOWITZ, R.J., MINNEMAN, K.P., MOLINOFF, P.B., RUFFOLO, R.R. & TRENDLENBURG, U. (1994). International Union of Pharmacology IV. Nomenclature of adrenoreceptors. *Pharmacol. Rev.*, **46**, 121–136.

DOCHERTY, J.R. (1998). Subtypes of functional α₁- and α₂-adrenoceptors. *Eur. J. Pharmacol.*, **361**, 1–15.

GUIMARAES, S. & MOURA, D. (2001). Vascular adrenoceptors: an update. *Pharmacol. Rev.*, **53**, 319–356.

HIEBLE, J.P., BYLUND, D.B., CLARKE, D.E., EIKENBURG, D.C., LANGER, S.Z., LEFKOWITZ, R.J., MINNEMAN, K.P. & RUFFOLO, R.R. (1995). International Union of Pharmacology. X. Recommendation for nomenclature of α₁-adrenoceptors: consensus update. *Pharmacol. Rev.*, **47**, 267–270.

KOCH, W.J., LEFKOWITZ, R.J. & ROCKMAN, H.A. (2000). Functional consequences of altering myocardial adrenergic receptor signaling. *Annu. Rev. Physiol.*, **62**, 237–260.

PIASCIK, M.T. & PEREZ, D.M. (2001). α₁-Adrenergic receptors: new insights and directions. *J. Pharmacol. Exp. Ther.*, **298**, 403–410.

ROHRER, D.K. & KOBILKA, B.K. (1998). Insights from *in vivo* modification of adrenergic receptor gene expression. *Annu. Rev. Pharmacol. Toxicol.*, **38**, 351–373.

TANOUE, A., KOSHIMIZU, T.A., SHIBATA, K., NASA, Y., TAKEO, S. & TSUJIMOTO, G. (2003). Insights into α₁ adrenoceptor function in health and disease from transgenic animal studies. *Trends Endocrinol. Metabol.*, **14**, 107–113.

References:

HAGUE, C. *et al.* (2004). *J. Biol. Chem.*, **279**, 15541–15549.

RAMSAY, D. *et al.* (2004). *Mol. Pharmacol.*, **66**, 228–239.

UBERTI, M.A. *et al.* (2005). *J. Pharmacol., Exp. Ther.*, **313**, 16–23.

Adrenoceptors, α_2

Overview: α_2 -Adrenoceptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Adrenoceptors; Bylund *et al.*, 1994) are 7TM receptors, where the endogenous agonists display a rank order of potency: adrenaline > noradrenaline. UK14304 and BHT920 are examples of agonists selective for α_2 -adrenoceptors relative to α_1 -adrenoceptors. Rauwolscine (9.0) and yohimbine (9.0) are antagonists selective for α_2 -adrenoceptors relative to α_1 -adrenoceptors. [3 H]-rauwolscine (1 nM), [3 H]-UK14304 (5 nM) and [3 H]-RX821002 (0.5 nM and 0.1 nM at α_{2C}) are relatively selective radioligands. There is species variation in the pharmacology of the α_{2A} -adrenoceptor; for example, yohimbine, rauwolscine and oxymetazoline have an ~20-fold lower affinity for rat, mouse and bovine α_{2A} -adrenoceptors. These α_{2A} orthologues are sometimes referred to as α_{2D} -adrenoceptors.

Nomenclature	α_{2A}	α_{2B}	α_{2C}
Other names	α_{2D}	—	—
Ensembl ID	ENSG00000150594	ENSG00000181210	ENSG00000184160
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}
Selective agonists	Oxymetazoline	—	—
Selective antagonists	BRL44408 (8.0)	ARC239 (8.0), prazosin (7.5), imiloxan (7.3)	ARC239 (8.0), prazosin (7.5)

Oxymetazoline is a partial agonist. Binding sites for imidazolines, distinct from α_2 -adrenoceptors, have been identified, but their function is not known; catecholamines have a low affinity for these sites.

Abbreviations: **ARC239**, 2-(2,4-[*O*-methoxyphenyl]-piperazin)-1-yl; **BHT920**, 6-allyl-2-amino-5,6,7,8-tetrahydro-4*H*-thiazolo-[4,5-*d*]-azepine; **BRL44408**, 2-(2*H*-[1-methyl-1,3-dihydroisindol-4-yl]-4,5-dihydroimidazole; **MK912**, (2*S*,12*bS*)1',3'-dimethylspiro(1,3,4,5',6,6',7,12*b*-octahydro-2*H*-benzo[*b*]furo[2,3-*a*]quinolizine)-2,4'-pyrimidin-2'-one; **RX821002**, 2-(2-methoxy-1,4-benzodioxan-2-yl)-2-imidazoline; **UK14304**, 5-bromo-6-[2-imidazolin-2-ylamino]quinoxaline

Further Reading:

- BUSCHER, R., HERRMANN, V. & INSEL, P.A. (1999). Human adrenoceptor polymorphisms: evolving recognition of clinical importance. *Trends Pharmacol. Sci.*, **20**, 94–99.
- BYLUND, D.B., EIKENBERG, D.C., HIEBLE, J.P., LANGER, S.Z., LEFKOWITZ, R.J., MINNEMAN, K.P., MOLINOFF, P.B., RUFFOLO, R.R. & TRENDLENBURG, U. (1994). International Union of Pharmacology IV. Nomenclature of adrenoceptors. *Pharmacol. Rev.*, **46**, 121–136.
- DOCHERTY, J.R. (1998). Subtypes of functional α_1 - and α_2 -adrenoceptors. *Eur. J. Pharmacol.*, **361**, 1–15.
- GUIMARAES, S. & MOURA, D. (2001). Vascular adrenoceptors: an update. *Pharmacol. Rev.*, **53**, 319–356.
- KABLE, J.W., MURRIN, L.C. & BYLUND, D.B. (2000). *In vivo* gene modification elucidates subtype-specific functions of α_2 -adrenergic receptors. *J. Pharmacol. Exp. Ther.*, **293**, 1–7.
- KOCH, W.J., LEFKOWITZ, R.J. & ROCKMAN, H.A. (2000). Functional consequences of altering myocardial adrenergic receptor signaling. *Annu. Rev. Physiol.*, **62**, 237–260.
- PHILIPP, M., BREDE, M. & HEIN, L. (2002). Physiological significance of α_2 -adrenergic receptor subtype diversity: one receptor is not enough. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **283**, R287–R295.
- ROHRER, D.K. & KOBILKA, B.K. (1998). Insights from *in vivo* modification of adrenergic receptor gene expression. *Annu. Rev. Pharmacol. Toxicol.*, **38**, 351–373.

Adrenoceptors, β

Overview: β -Adrenoceptors (nomenclature as agreed by the NC-IUPHAR Subcommittee on Adrenoceptors, Bylund *et al.*, 1994) are 7TM receptors, where the endogenous agonists are adrenaline and noradrenaline. Isoprenaline is an example of an agonist selective for β -adrenoceptors relative to α_1 - and α_2 -adrenoceptors, while propranolol (pKi 8.2–9.2) and cyanopindolol (pKi 10.0–11.0) are relatively selective antagonists. β_3 -Adrenoceptors are relatively resistant to blockade by propranolol (pKi 5.8–7.0), but can be blocked with high concentrations of cyanopindolol (pKi 9.0).

Nomenclature	β_1	β_2	β_3
Other names	—	—	atypical β
Ensembl ID	ENSG00000043591	ENSG00000169252	ENSG00000147477
Principal transduction	G _s	G _s	G _s , G _{i/o}
Rank order of potency	NA > adrenaline	Adrenaline > NA	NA = adrenaline
Selective agonists	Noradrenaline, xamoterol, RO363, denopamine	Procaterol, zinterol, salmeterol, formoterol, terbutaline, fenoterol	BRL37344, CL316243, CGP12177A, carazolol, L742791, SB251023
Selective antagonists	CGP20712A (8.5–9.3), betaxolol (8.5), atenolol (7.6)	ICI118551 (8.3–9.2)	SR59230A (8.8), L748328 (8.5)
Probes	[¹²⁵ I]-ICYP (20–50 pM) + 70 nM ICI118551	[¹²⁵ I]-ICYP (20–50 pM) + 100 nM CGP20712A	[¹²⁵ I]-ICYP (0.5 nM)

Noradrenaline, xamoterol and RO363 show selectivity for β_1 - relative to β_2 -adrenoceptors. Radioligand binding to define β_1 - and β_2 -adrenoceptors can be conducted in the presence of a 'saturating' concentration of the β_1 - or β_2 -adrenoceptor-selective antagonist. [³H]-CGP12177 or [³H]-dihydroalprenolol can be used in place of [¹²⁵I]-ICYP. Many antagonists at β_1 - and β_2 -adrenoceptors are agonists at β_3 -adrenoceptors (CL316243, CGP12177A and carazolol). CGP12177A and carazolol can also show reduced efficacy at β_3 -adrenoceptors. SR59230A has reasonably high affinity at β_3 -adrenoceptors (Manara *et al.*, 1996), but does not discriminate well between the three β -adrenoceptor subtypes (Candelore *et al.*, 1999), has been reported to have lower affinity for the β_3 -adrenoceptor in some circumstances (Kaumann & Molenaar, 1996) and can exhibit agonist properties in some functional assays (Horinouchi & Koike, 2001). Pharmacological differences exist between human and mouse β_3 -adrenoceptors, and the 'rodent selective' agonists BRL37344 and CL316243 have low efficacy at the human β_3 -adrenoceptor (see reviews by Strosberg). The β_3 -adrenoceptor has introns, but splice variants have only been described for the mouse (Evans *et al.*, 1999). The β -adrenoceptor cloned from turkey (termed the β_4 c, t428 SwissProt P43141) has a pharmacology that is intermediate between β_2 - and β_3 -adrenoceptors (Chen *et al.*, 1994).

There is now convincing evidence that the 'putative β_4 -adrenoceptor' is not a novel receptor but is likely to represent an alternative site of interaction of CGP12177A and other nonconventional partial agonists at β_1 -adrenoceptors, since 'putative β_4 -adrenoceptor'-mediated agonist effects of CGP12177A are absent in mice lacking β_1 -adrenoceptors (Konkar *et al.*, 2000; Kaumann *et al.*, 2001).

Numerous polymorphisms exist for the β_1 - and β_2 -adrenoceptors and some of these are associated with alterations in signalling in response to agonists. The polymorphisms may be associated with altered responses to drugs.

Abbreviations: **BRL37344**, sodium 4-(2-[2-hydroxy-3-chlorophenyl]ethylamino)propyl)phenoxyacetate; **CGP12177A**, (–)-4-(3-tert-butylamino-2-hydroxypropoxy)-benzimidazol-2-one; **CGP20712A**, 2-hydroxy-5-(2-[[2-hydroxy-3-(4-[1-methyl-4-trifluoromethyl-2-imidazolyl]phenoxy)propyl]amino]ethoxy)benzamide; **CL316243**, disodium (R,R)-5-(2-[[2-(3-chlorophenyl)-2-hydroxyethyl]-amino]propyl)-1,3-benzodioxole-2,2-dicarboxylate; **ICYP**, iodocyanopindolol; **L742791**, (S)-N-(4-[2-(3-[4-hydroxyphenoxyl]-2-hydroxypropyl)amino]ethyl)phenyl)-4-iodobenzenesulfonamide; **L748328**, (S)-N-(4-[2-(3-[3-(aminosulfonyl)phenoxy]-2-hydroxypropyl)-amino]ethyl)phenyl)benzenesulfonamide; **RO363**, (–)-1-(3,4-dimethoxyphenethylamino)-3-(3,4-dihydroxyphenoxy)-2-propanol)oxalate; **SB251023**, (4-[1-(2-(S)-hydroxy-3-(4-hydroxyphenoxy)-propylamino)cyclopentylmethyl]phenoxy)methyl)phenylphosphonic acid lithium salt; **SR59230A**, 3-(2-ethylphenoxy)-1-[(1S)-1,2,3,4-tetrahydronaphth-1-ylamino]-2S-propanol oxalate

Further Reading:

- ARCH, J.R.S. (2002). β_3 -Adrenoceptor agonists: potential, pitfalls and progress. *Eur. J. Pharmacol.*, **440**, 99–107.
- BUSCHER, R., HERRMANN, V. & INSEL, P.A. (1999). Human adrenoceptor polymorphisms: evolving recognition of clinical importance. *Trends Pharmacol. Sci.*, **20**, 94–99.
- BYLUND, D.B., EIKENBERG, D.C., HIEBLE, J.P., LANGER, S.Z., LEFKOWITZ, R.J., MINNEMAN, K.P., MOLINOFF, P.B., RUFFOLO, R.R. & TRENDLENBURG, U. (1994). International Union of Pharmacology IV. Nomenclature of adrenoceptors. *Pharmacol. Rev.*, **46**, 121–136.
- GUIMARAES, S. & MOURA, D. (2001). Vascular adrenoceptors: an update. *Pharmacol. Rev.*, **53**, 319–356.
- KOCH, W.J., LEFKOWITZ, R.J. & ROCKMAN, H.A. (2000). Functional consequences of altering myocardial adrenergic receptor signaling. *Annu. Rev. Physiol.*, **62**, 237–260.
- LEINWEBER, K., BUSCHER, R., BRUCK, H. & BRODDE, O.E. (2004). β_2 -adrenoceptor polymorphisms. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **369**, 1–22.
- LEINWEBER, K. & BRODDE, O.E. (2004). β_2 -adrenoceptor polymorphisms: relation between *in vitro* and *in vivo* phenotypes. *Life Sci.*, **74**, 2803–2814.
- MICHEL, M.C. & INSEL, P.A. (2003). Receptor gene polymorphisms: lessons on functional relevance from the β_1 -adrenoceptor. *Br. J. Pharmacol.*, **138**, 279–282.
- ROHRER, D.K. & KOBILKA, B.K. (1998). Insights from *in vivo* modification of adrenergic receptor gene expression. *Annu. Rev. Pharmacol. Toxicol.*, **38**, 351–373.
- STROSBERG, A.D. (1997). Association of β_3 -adrenoreceptor polymorphism with obesity and diabetes: current status. *Trends Pharmacol. Sci.*, **18**, 449–454.
- STROSBERG, A.D. (1997). Structure and function of the β_3 -adrenoceptor. *Annu. Rev. Pharmacol. Toxicol.*, **37**, 421–450.

References:

- CANDELORE, M.R. *et al.* (1999). *J. Pharmacol. Exp. Ther.*, **290**, 649–655.
- CHEN, X.H. *et al.* (1994). *J. Biol. Chem.*, **269**, 24810–24819.

- EVANS, B.A. *et al.* (1999). *Br. J. Pharmacol.*, **127**, 1525–1531.
- HORINOUCI, T. & KOIKE, K. (2001). *Eur. J. Pharmacol.*, **416**, 153–163.
- KAUMANN, A.J. & MOLENAAR, P. (1996). *Br. J. Pharmacol.*, **118**, 2085–2098.
- KAUMANN, A.J. *et al.* (2001). *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **363**, 87–93.
- KONKAR, A.A. *et al.* (2000). *Mol. Pharmacol.*, **57**, 252–258.
- MANARA, L. *et al.* (1996). *Br. J. Pharmacol.*, **117**, 435–442.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Anaphylatoxin and chemotactic peptide

Overview: Anaphylatoxin and chemotactic peptide receptors (provisional nomenclature) are activated by the endogenous ~75 aa anaphylatoxin polypeptides C3a (ENSG00000125730) and C5a (ENSG00000106804), generated upon stimulation of the complement cascade. The fMLP receptor responds to exogenous ligands such as the bacterial product formyl-Met-Leu-Phe (fMLP) and endogenous ligands such as annexin I (ENSG00000135046), cathepsin G (ENSG00000100448) and spinorphin, derived from β -haemoglobin (ENSG00000188170).

Nomenclature	C3a	C5a	fMLP
Other names	AZ3B, HNFAG09	CD88	Formyl peptide, FPR
Ensembl ID	ENSG00000171860	ENSG00000134830	ENSG00000171051
Principal transduction	G _{i/o} , G _z	G _{i/o} , G _z , G ₁₆ (Buhl <i>et al.</i> , 1993)	G _{i/o} , G _z
Rank order of potency	C3a > C5a (Ames <i>et al.</i> , 1996)	C5a, C5a des Arg > C3a (Ames <i>et al.</i> , 1996)	fMLP > Cathepsin G > Annexin I (Le <i>et al.</i> , 2002; Sun <i>et al.</i> , 2004)
Selective agonists	Trp-Trp-Gly-Lys-Lys-Tyr-Arg-Ala-Ser-Lys-Leu-Gly-Leu-Ala-Arg (Ames <i>et al.</i> , 1997)	Phe-Lys-Pro-Cha-Cha-Phe-Lys-D-Cha-Cha-D-Arg (Kontekatis <i>et al.</i> , 1994), S19 (Yamamoto, 2000)	fMLP (Le <i>et al.</i> , 1999)
Selective antagonists	SB290157 (pIC ₅₀ 7.5, Ames <i>et al.</i> , 2001)	NMe-Phe-Lys-Pro-D-Cha-Trp-D-Arg (Kontekatis <i>et al.</i> , 1994), AcPhe-Orn-Pro-D-Cha-Trp-Arg (Wong <i>et al.</i> , 1998), W54011 (8.7, Sumichika <i>et al.</i> , 2002), CHIPS (Postma <i>et al.</i> , 2004)	Cyclosporin H (6.3-7.0, Wenzel-Seifert & Seifert, 1993), BOC-PLPLP (6.0-6.5, Wenzel-Seifert and Seifert, 1993), spinorphin (4, Liang <i>et al.</i> , 2001)
Probes	[¹²⁵ I]-C3a	[¹²⁵ I]-C5a	[³ H]-fMLP

SB290157 has also been reported to have agonist properties (Mathieu *et al.*, 2005). A putative chemoattractant receptor termed C5L2 (also known as GPR77, ENSG00000134830) binds [¹²⁵I]-C5a, with no clear signalling function, but a putative role opposing inflammatory responses (Cain & Monk, 2002; Gao *et al.*, 2005; Gavriluk *et al.*, 2005). Binding to this site may be displaced with the rank order C5a des Arg > C5a (Cain & Monk, 2002; Okinaga *et al.*, 2003), while there is controversy over the ability of C3a and C3a des Arg to compete (Kalant *et al.*, 2003; Okinaga *et al.*, 2003; Honczarenko *et al.*, 2005; Kalant *et al.*, 2005).

Abbreviations: BOC-PLPLP, Boc-Phe-Leu-Phe-Leu-Phe; CHIPS, chemotaxis inhibitory protein of *Staphylococcus aureus*; SB290157, N²-([2,2-diphenylethoxy]-acetyl)-L-Arg; W54011, N-([4-dimethylaminophenyl]methyl)-N-(4-isopropylphenyl)-7-methoxy-1,2,3,4-tetrahydronaphthalen-1-carboxamide hydrochloride

Further Reading:

ALLEGRETTI, M., MORICONI, A., BECCARI, A.R., DI BITONDO, R., BIZZARRI, C., BERTINI, R. & COLOTTA, F. (2005). Targeting C5a: recent advances in drug discovery. *Curr. Med. Chem.*, **12**, 217–236.

BARNUM, S.R. (2002). Complement in central nervous system inflammation. *Immunol. Res.*, **26**, 7–13.

CHEN, X., YANG, D., SHEN, W., DONG, H.F., WANG, J.M., OPPENHEIM, J.J. & HOWARD, M.Z. (2000). Characterization of chenodeoxycholic acid as an endogenous antagonist of the G-coupled formyl peptide receptors. *Inflamm. Res.*, **49**, 744–755.

GUO, R.F. & WARD, P.A. (2005). Role of C5a in inflammatory responses. *Annu. Rev. Immunol.*, **23**, 821–852.

GURRATH, M. (2001). Peptide-binding G protein-coupled receptors: new opportunities for drug design. *Curr. Med. Chem.*, **8**, 1605–1648.

KATANAEV, V.L. (2001). Signal transduction in neutrophil chemotaxis. *Biochemistry (Mosc.)*, **66**, 351–368.

PANARO, M.A., ACQUAFREDDA, A., SISTO, M., LISI, S., MAFFIONE, A.B. & MITOLO, V. (2006). Biological role of the N-formal peptide receptors. *Immunopharmacol. Immunotoxicol.*, **28**, 103–127.

PERRETTI, M. (2003). The annexin 1 receptor(s): is the plot unravelling? *Trends Immunol.*, **24**, 574–579.

RICKERT, P., WEINER, O.D., WANG, F., BOURNE, H.R. & SERVANT, G. (2000). Leukocytes navigate by compass: roles of PI3K γ and its lipid products. *Trends Cell Biol.*, **10**, 466–473.

SALLUSTO, F. & MACKAY, C.R. (2004). Chemoattractants and their receptors in homeostasis and inflammation. *Curr. Opin. Immunol.*, **16**, 724–731.

TAYLOR, S.M., SHERMAN, S.A., KIRNARSKY, L. & SANDERSON, S.D. (2001). Development of response-selective agonists of human C5a anaphylatoxin: conformational, biological, and therapeutic considerations. *Curr. Med. Chem.*, **8**, 675–684.

VAN COMPERNOLLE, S.E., CLARK, K.L., RUMMEL, K.A. & TODD, S.C. (2003). Expression and function of formyl peptide receptors on human fibroblast cells. *J. Immunol.*, **171**, 2050–2056.

WARD, P.A. (2004). The dark side of C5a in sepsis. *Nat. Rev. Immunol.*, **4**, 133–142.

WENZEL-SEIFERT, K. & SEIFERT, R. (2003). Functional differences between human formyl peptide receptor isoforms 26, 98, and G6. *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **367**, 509–515.

References:

- AMES, R.S. *et al.* (1996). *FEBS Lett.*, **395**, 157–159.
- AMES, R.S. *et al.* (1997). *Immunopharmacology*, **38**, 87–92.
- AMES, R.S. *et al.* (2001). *J. Immunol.*, **166**, 6341–6348.
- BUHL, A.M. *et al.* (1993). *FEBS Lett.*, **323**, 132–134.
- CAIN, S.A. & MONK, P.N. (2002). *J. Biol. Chem.*, **277**, 7165–7169.
- GAO, H. *et al.* (2005). *FASEB J.*, **19**, 1003–1005.
- GAVRILYUK, V. *et al.* (2005). *J. Neurochem.*, **92**, 1140–1149.
- HONCZARENKO, M. *et al.* (2005). *Leukemia*, **19**, 1682–1683.
- KALANT, D. *et al.* (2003). *J. Biol. Chem.*, **278**, 11123–11129.
- KALANT, D. *et al.* (2005). *J. Biol. Chem.*, **280**, 23936–23944.
- KONTEATIS, Z.D. *et al.* (1994). *J. Immunol.*, **153**, 4200–4205.

- LE, Y. *et al.* (1999). *J. Immunol.*, **163**, 6777–6784.
- LE, Y. *et al.* (2002). *Trends Immunol.*, **23**, 541–548.
- LIANG, T.S. *et al.* (2001). *J. Immunol.*, **167**, 6609–6614.
- MATHIEU, M.C. *et al.* (2005). *Immunol. Lett.*, **100**, 139–145.
- OKINAGA, S. *et al.* (2003). *Biochemistry*, **42**, 9406–9415.
- POSTMA, B. *et al.* (2004). *J. Immunol.*, **172**, 6994–7001.
- SUMICHIKA, H. *et al.* (2002). *J. Biol. Chem.*, **277**, 49403–49407.
- SUN, R. *et al.* (2004). *J. Immunol.*, **173**, 428–436.
- WENZEL-SEIFERT, K. & SEIFERT, R. (1993). *J. Immunol.*, **150**, 4591–4599.
- WONG, A.K. *et al.* (1998). *J. Med. Chem.*, **41**, 3417–3425.
- YAMAMOTO, T. (2000). *Pathol. Int.*, **50**, 863–871.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Angiotensin

Overview: The actions of angiotensin II (Ang II) are mediated by AT₁ and AT₂ receptors (nomenclature agreed by the NC-IUPHAR Subcommittee on Angiotensin Receptors; see de Gasparo *et al.*, 2000), which have around 30% sequence similarity. AT₁ receptors are predominantly coupled to G_{q/11}. Most species express a single AT₁ gene, but two related AT_{1A} and AT_{1B} receptor genes are expressed in rodents. The AT₂ receptor counteracts several of the growth responses initiated by the AT₁ receptors. The AT₂ receptor is much less abundant than the AT₁ receptor in adult tissues and is upregulated in pathological conditions. Endogenous ligands are Ang II and angiotensin III (Ang III), while angiotensin I is weakly active in some systems.

Nomenclature	AT ₁	AT ₂
Ensembl ID	ENSG00000144891	ENSG00000180772
Principal transduction	G _{q/11}	Tyr & Ser/Thr phosphatases
Selective agonists	L162313	[p-NH ₂ -Phe ⁶]-Ang II, CGP42112
Selective antagonists	EXP3174, eprosartan, valsartan, irbesartan, losartan	PD123319, PD123177
Probes	[³ H]-A81988, [³ H]-L158809, [³ H]-eprosartan [³ H]-losartan, [¹²⁵ I]-EXP985	[¹²⁵ I]-CGP42112

There is also evidence for an AT₄ receptor that specifically binds angiotensin IV and is located in the brain and kidney. An additional putative endogenous ligand for the AT₄ receptor has been described (LVV-hemorphin, a globin decapeptide) (Moeller *et al.*, 1997). The AT₁ and bradykinin B2 receptors have been proposed to form a heterodimeric complex (Abdalla *et al.*, 2000). The antagonist activity of CGP42112 has also been reported (Lokuta *et al.*, 1995). Novel AT₁ receptor antagonists bearing substituted 4-phenylquinoline moieties have recently been designed and synthesized. The best of these compounds bind to AT₁ receptors with nanomolar affinity and are slightly more potent than losartan in functional studies (Cappelli *et al.*, 2004).

Abbreviations: **A81988**, 2-(*N*-propyl-*N*-{[2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl]methyl}amino)pyridine-3-carboxylate; **CGP42112A**, nicotinic acid-Tyr-(*N*-benzoyl carbonyl-Arg)-Lys-His-Pro-Ile-OH; **eprosartan**, (E)-α-[(2-butyl-1-[(4-carboxyphenyl)methyl]-1*H*-imidazol-5-yl)methylene]-2-thiophenepropanoate; **EXP3174**, *n*-butyl-4-chloro-1-[(2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxylate; **EXP985**, *N*-(2-[4-hydroxy-3-iodophenyl]ethyl)-4-chloro-2-propyl-1-[(2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole-5-carboxamide; **irbesartan**, 2-butyl-3-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-1,3-diazaspiro[4.4]non-1-en-4-one; **L158809**, 5,7-dimethyl-2-ethyl-3-(2-[1*H*-tetrazol-5-yl]biphenyl-4-yl)imidazo[4,5-*b*]pyridine; **L162313**, 5,7-dimethyl-2-ethyl-3-[[4-[2(*n*-butyloxycarbonylsulfonamido)-5-isobutyl-3-thienyl]phenyl]methyl]imidazo[4,5-*b*]pyridine; **losartan**, 2-*n*-butyl-4-chloro-5-hydroxymethyl-1-[(2'-(1*H*-tetrazol-5-yl)biphenyl-4-yl)methyl]imidazole, also known as Dup 753; **PD123177**, 1-(4-amino-3-methylphenyl)methyl-3-(diphenylacetyl)-4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine-6-carboxylate; **PD123319**, (S)-1-(4-[dimethylamino]-3-methylphenyl)methyl-5-(diphenylacetyl)-4,5,6,7-tetrahydro-1*H*-imidazo[4,5-*c*]pyridine-6-carboxylate; **valsartan**, *N*-(1-oxopentyl)-*N*-[[2'-(1*H*-tetrazol-5-yl)[1,1'-biphenyl]-4-yl]methyl]-L-valine

Further Reading:

- ALDIGIER, J.C. & GHANNAD, E. (2002). Exploring AT₁ and AT₂ angiotensin II receptors in humans. *Drugs*, **62**, 11–19.
- BERNSTEIN, K.E. & MARRERO, M.B. (1996). The importance of tyrosine phosphorylation in angiotensin II signaling. *Trends Cardiovasc. Med.*, **6**, 179–187.
- CAREY, R.M. (2005). Update on the role of the AT₂ receptor. *Curr. Opin. Nephrol. Hypertens.*, **14**, 67–71.
- DE GASPARO, M. (2002). AT₁ and AT₂ angiotensin (II) receptors: key features. *Drugs*, **62**, 1–10.
- DE GASPARO, M., CATT, K.J., INAGAMI, T., WRIGHT, J.W. & UNGER, T. (2000). International Union of Pharmacology. XXIII. The angiotensin II receptors. *Pharmacol. Rev.*, **52**, 415–472.
- FERRARIO, C.M. & CHAPPELL, M.C. (2004). Novel angiotensin peptides. *Cell. Mol. Life Sci.*, **61**, 2720–2727.
- HUNYADY, L., BALLA, T. & CATT, K.J. (1996). The ligand binding site of the angiotensin AT₁ receptor. *Trends Pharmacol. Sci.*, **17**, 135–140.
- KUSSEROW, H. & UNGER, T. (2004). Vasoactive peptides, their receptors and drug development. *Basic Clin. Pharmacol. Toxicol.*, **94**, 5–12.
- NOUET, S. & NAHMIAS, C. (2000). Signal transduction from the angiotensin II AT₂ receptor. *Trends Endocrinol. Metab.*, **11**, 1–6.
- RASHID, A.J., O'DOWD, B.F. & GEORGE, S.R. (2004). Minireview: diversity and complexity of signaling through peptidergic G protein-coupled receptors. *Endocrinology*, **145**, 2645–2652.
- UNGER, T. (1999). The angiotensin type 2 receptor: variations on an enigmatic theme. *J. Hypertens.*, **17**, 1775–1786.
- ZAMAN, M.A., OPARIL, S. & CALHOUN, D.A. (2002). Drugs targeting the renin–angiotensin–aldosterone system. *Nat. Rev. Drug Discov.*, **1**, 621–636.

References:

- ABDALLA, S. *et al.* (2000). *Nature*, **407**, 94–98.
- CAPPELLI, A. *et al.* (2004). *J. Med. Chem.*, **47**, 2574–2586.
- LOKUTA, A.J. *et al.* (1995). *J. Biol. Chem.*, **269**, 4832–4838.
- MOELLER, I. *et al.* (1997). *J. Neurochem.*, **68**, 2530–2537.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Apelin

Overview: The apelin receptor (APJ, provisional nomenclature previously designated as an orphan) responds to apelin, a 36 amino-acid peptide derived initially from bovine stomach. Apelin-36, apelin-13 and (Pyr¹)apelin-13 are the predominant endogenous ligands which are cleaved from a 77 amino-acid precursor peptide (ENSG00000171388) by a so far unidentified enzymatic pathway (Tatemoto *et al.*, 1998).

Nomenclature	APJ
Other names	Apelin receptor, angiotensin receptor-like 1
Ensembl ID	ENSG00000134817
Principal transduction	G _{i/o}
Rank order of potency	[Pyr ¹]apelin-13 > apelin-13 > apelin-36 (Tatemoto <i>et al.</i> , 1998)
Selective agonists	[Pyr ¹]apelin-13, apelin-13, apelin-17, apelin-36
Probes	[¹²⁵ I]-[Pyr ¹]Apelin-13 (0.3 nM, Katugampola <i>et al.</i> , 2001), [³ H]-[Pyr ¹][Met(0 ¹¹)]apelin-13 (Medhurst <i>et al.</i> , 2003), [¹²⁵ I]-[Nle ⁷⁵ ,Tyr ⁷⁷]apelin-36 (Kawatma <i>et al.</i> , 2001)

Potency order determined for heterologously expressed human APJ receptor (pD₂ values range from 9.5 to 8.6). APJ may also act as a co-receptor with CD4 for isolates of human immunodeficiency virus, with apelin blocking this function (Cayabyab *et al.*, 2000).

Further Reading:

KATUGAMPOLA, S. & DAVENPORT, A. (2003). Emerging roles for orphan G-protein-coupled receptors in the cardiovascular system. *Trends Pharmacol. Sci.*, **24**, 30–35.

KLEINZ, M.J. & DAVENPORT, A.P. (2005). Emerging roles for apelin in biology and medicine. *Pharmacol. Ther.*, **107**, 198–211.

LEE, D.K., GEORGE, S.R. & O'DOWD, B.F. (2006). Unravelling the roles of the apelin system: prospective therapeutic applications in heart failure and obesity. *Trends Pharmacol. Sci.*, **27**, 190–194.

MAGUIRE, J.J. (2003). Discovering orphan receptor function using human *in vitro* pharmacology. *Curr. Opin. Pharmacol.*, **3**, 135–139.

MAGUIRE, J.J. & DAVENPORT, A.P. (2005). Regulation of vascular reactivity by established and emerging GPCRs. *Trends Pharmacol. Sci.*, **26**, 448–454.

MASRI, B., KNIBIEHLER, B. & AUDIGIER, Y. (2005). Apelin signalling: a promising pathway from cloning to pharmacology. *Cell Signal.*, 415–426.

References:

CAYABYAB, M. *et al.* (2000). *J. Virol.*, **74**, 11972–11976.

KATUGAMPOLA, S.D. *et al.* (2001). *Br. J. Pharmacol.*, **132**, 1255–1260.

KAWATMA, Y. *et al.* (2001). *Biochim. Biophys. Acta.*, **1538**, 162–171.

MEDHURST, A.D. *et al.* (2003). *J. Neurochem.*, **84**, 1162–1172.

TATEMOTO, K. *et al.* (1998). *Biochem. Biophys. Res. Commun.*, **251**, 471–476.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Bombesin

Overview: Bombesin receptors are activated by the endogenous ligands gastrin-releasing peptide (GRP), neuromedin B (NMB) and GRP-18-27 (previously named neuromedin C). Bombesin is a tetradecapeptide, originally derived from amphibians. These receptors couple, primarily, to the G_{q/11} family of G proteins (but see also Jian *et al.*, 1999). Activation of BB1 and BB2 receptors causes a wide range of physiological actions, including the stimulation of tissue growth, smooth-muscle contraction, secretion and many central nervous system effects (Tokita *et al.*, 2002). A physiological role for the bb3 receptor has yet to be defined.

Nomenclature	BB1	BB2	bb3
Other names	NMB-R	GRP-R	BRS-3
Ensemble ID	ENSG00000135577	ENSG00000126010	ENSG00000102239
Principal transduction	G _{q/11}	G _{q/11}	G _{q/11}
Selective agonists	NMB	GRP	—
Selective antagonists	PD165929, dNal-cyc(Cys-Tyr-dTrp-Orn-Val)-Nal-NH ₂ , dNal-Cys-Tyr-dTrp-Lys-Val-Cys-Nal-NH ₂	1-Naphthoyl-[dAla ²⁴ ,dPro ²⁶ ,ψ ²⁶⁻²⁷]GRP-20-27, kuwanon H, [dPhe ⁶]bombesin-6-13-ethylester, [dPhe ⁶ ,Cpa ¹⁴ ,ψ ¹³⁻¹⁴]bombesin-6-14	—
Probes	[¹²⁵ I]-BH-NMB, [¹²⁵ I]-[Tyr ⁴]bombesin	[¹²⁵ I]-[D-Tyr ⁶]bombesin-6-13-methylester, [¹²⁵ I]-GRP, [¹²⁵ I]-[Tyr ⁴]bombesin	[¹²⁵ I]-[Tyr ⁶ ,βAla ¹¹ ,Phe ¹³ ,Nle ¹⁴] bombesin-6-14

All three subtypes may be activated by [dPhe⁶,βAla¹¹,Phe¹³,Nle¹⁴]bombesin-6-14 (Mantey *et al.*, 1997). One analogue, [D-Tyr⁶, Apa-4Cl, Phe¹³, Nle¹⁴] bombesin-6-14, has more than 200-fold selectivity for bb3 receptors over BB1 and BB2 (Mantey *et al.*, 2004).

Abbreviations: PD165929, 2-[3-(2,6-diisopropylphenyl)-ureido]3-(1*H*-indol-3-yl)-2-methyl-*N*-(1-pyridin-2-yl-cyclohexylmethyl)-propionate

Further Reading:

BATTEY, J. & WADA, E. (1991). Two distinct receptor subtypes for mammalian bombesin receptors. *Trends Neurosci.*, **14**, 524–528.

IWABUCHI, M., UI-TEI, K., YAMADA, K., MATSUDA, Y., SAKAI, Y., TANAKA, K. & OHKI-HAMAZAKI, H. (2003). Molecular cloning and characterisation of avian bombesin-like peptide receptors: new tools for investigating molecular basis of ligand selectivity. *Br. J. Pharmacol.*, **139**, 555–566.

JENSEN, R. & COY, D. (1991). Progress in the development of potent bombesin receptor antagonists. *Trends Pharmacol. Sci.*, **12**, 13–18.

KROOG, G.S., JENSEN, R.T. & BATTEY, J.F. (1995). Bombesin receptors. *Med. Res. Rev.*, **15**, 389–417.

MOODY, T.W. & MERALI, Z. (2004). Bombesin-like peptides and associated receptors within the brain: distribution and behavioural implications. *Peptides*, **25**, 511–520.

MOODY, T.W., MANTEY, S.A., PRADHAN, T.K., SCHUMANN, M., NAKAGAWA, T., MARTINEZ, A., FUSELIER, J., COY, D.H. & JENSEN, R.T. (2004). Development of high affinity camptothecin–bombesin conjugates that have targeted cytotoxicity for bombesin receptor-containing tumor cells. *J. Biol. Chem.*, **279**, 23580–23589.

OHKI-HAMAZAKI, H. (2000). Neuromedin B. *Prog. Neurobiol.*, **62**, 297–312.

OHKI-HAMAZAKI, H., IWABUCHI, M. & MAEKAWA, F. (2005). Development and function of bombesin-like peptides and their receptors. *Int. J. Dev. Biol.*, **49**, 293–300.

TOKITA, K., HOCART, S.J., COY, D.H. & JENSEN, R.T. (2002). Molecular basis of the selectivity of gastrin-releasing peptide receptor for gastrin-releasing peptide. *Mol. Pharmacol.*, **61**, 1435–1443.

WEBER, D. (2003). Design of selective peptidomimetic agonists for the human orphan receptor BRS-3. *J. Med. Chem.*, **46**, 1918–1930.

ZHOU, J., CHEN, J., MOKOTOFF, M. & BALL, E.D. (2004). Targeting gastrin-releasing peptide receptors for cancer treatment. *Anti-cancer Drugs*, **15**, 921–927.

References:

JIAN, X.Y. *et al.* (1999). *J. Biol. Chem.*, **274**, 11573–11581.

MANTEY, S.A. *et al.* (1997). *J. Biol. Chem.*, **272**, 26062–26071.

MANTEY, S.A. *et al.* (2004). *J. Pharmacol. Exp. Ther.*, **310**, 1161–1170.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Bradykinin

Overview: Bradykinin (or kinin) receptors (nomenclature recommended by the NC-IUPHAR Subcommittee on bradykinin (kinin) receptors, Regoli *et al.*, 1998b, Leeb-Lundberg *et al.*, 2005) are activated by the endogenous peptides bradykinin (BK), [des-Arg⁹]BK, Lys-BK (kallidin), Lys-[des-Arg⁹]BK, T-kinin (Ile-Ser-BK), [Hyp³]BK and Lys-[Hyp³]BK. The variation in affinity or inactivity of B₂ receptor antagonists could reflect the existence of species homologues of B₂ receptors.

	B₁	B₂
Nomenclature	B ₁	B ₂
Ensembl ID	ENSG00000100739	ENSG00000168398
Principal transduction	G _{q/11}	G _{q/11}
Rank order of potency	Lys-[des-Arg ⁹]BK > [des-Arg ⁹]BK = Lys-BK > BK	Lys-BK ≥ BK ≫ [des-Arg ⁹]BK, Lys-[des-Arg ⁹]BK
Selective agonists	Lys-[des-Arg ⁹]BK, Sar[D ¹ Phe ⁸][des-Arg ⁹]BK	[Phe ⁸ ,ψ(CH ₂ -NH)Arg ⁹]BK, [Hyp ³ ,Tyr(Me) ⁸]BK
Selective antagonists	B9958 (9.2, Regoli <i>et al.</i> , 1998a), R914 (8.6, Gobeil <i>et al.</i> , 1999), R715 (8.5, Gobeil <i>et al.</i> , 1996a), Lys-[Leu ⁸][des-Arg ⁹]BK (8.0)	Icatibant (8.4, Gobeil <i>et al.</i> , 1996b), FR173657 (8.2, Rizzi <i>et al.</i> , 1997), LF160687 (Puneau <i>et al.</i> , 1999)
Probes	[³ H]-Lys-[des-Arg ⁹]BK (0.4 nM), [³ H]-Lys-[Leu ⁸][des-Arg ⁹]BK, [¹²⁵ I]-Hpp-desArg ⁹ HOE140 (0.1 nM)	[³ H]-BK (0.2 nM), [³ H]-NPC17731 (50–900 pM), [¹²⁵ I]-[Tyr ⁸]BK

Abbreviations: **B9958**, Lys-Lys[Hyp³,Cpg⁵,dTic⁷,Cpg⁸][des-Arg⁹]BK; **FR173657**, (E)-3-(6-acetamido-3-pyridyl)-N-(N-[2,4-dichloro-3{(2-methyl-8-quinolinyl)oxy-methyl} phenyl]-N-methylaminocarbonyl-methyl)acrylamide; **Icatibant**, DArg[Hyp³,Thi⁵,dTic⁷,Oic⁸]BK, also known as HOE140; **LF160687**, 1-([2,4-dichloro-3-([2,4-dimethylquinolin-8-yl]oxy)methyl]phenyl)sulfonyl)-N-(3-[(4-(aminoimethyl)phenyl)carbonylamino]propyl)-2(s)-pyrrolidinecarboxamide; **NPC17731**, DArg[Hyp³,dHypE(transpropyl)⁷,Oic⁸]BK; **R715**, AcLys[D¹Nal⁷,Ile⁸][des-Arg⁹]BK; **R914**, AcLys-Lys-([zMe]Phe⁵,δ-βNal⁷,Ile⁸)desArg⁹BK

Further Reading:

CALIXTO, J.B., MEDEIROS, R., FERNANDES, E.S., FERREIRA, J., CABRINI, D.A. & CAMPOS, M.M. (2004). Kinin B₁ receptors: key G-protein-coupled receptors and their role in inflammatory and painful processes. *Br. J. Pharmacol.*, **143**, 803–818.

COUTURE, R., HARRISSON, M., VIANNA, R.M. & CLOUTIER, F. (2001). Kinin receptors in pain and inflammation. *Eur. J. Pharmacol.*, **429**, 161–176.

DOGGRELL, S.A. (2006). Bradykinin B₂ receptors as a target in diabetic nephropathy. *Curr. Opin. Investig. Drugs*, **7**, 251–255.

FORTIN, J.P. & MARCEAU, F.O. (2006). Advances in the development of bradykinin ligands. *Curr. Top. Med. Chem.*, **6**, 1353–1363.

LEEB-LUNDBERG, L.M.F., MARCEAU, F., MULLER-ESTERL, W., PETTIBONE, D.J. & ZURAW, B.L. (2005). International Union of Pharmacology. XLV. Classification of the kinin receptor family: from molecular mechanisms to pathophysiological consequences. *Pharmacol. Rev.*, **57**, 27–77.

MARCEAU, F., HESS, J.F. & BACHVAROV, D.R. (1998). The B-1 receptors for kinins. *Pharmacol. Rev.*, **50**, 357–386.

MARCEAU, F. & REGOLI, D. (2004). Bradykinin receptor ligands: Therapeutic perspectives. *Nature Rev. Drug Discov.*, **3**, 845–852.

REGOLI, D., ALLOGHO, S.N., RIZZI, A. & GOBEIL, F.J. (1998a). Bradykinin receptors and their antagonists. *Eur. J. Pharmacol.*, **348**, 1–10.

REGOLI, D., GEPPETTI, P., HESS, J.F., MARCEAU, F., MULLER-ESTERL, W. & SCHOELKENS, B.A. (1998b). Bradykinin receptors. In: *The IUPHAR compendium of receptor characterization and classification*, eds. Girdlestone, D. pp. 87–93. London: IUPHAR Media.

RODI, D., COUTURE, R., ONGALI, B. & SIMONATO, M. (2005). Targeting kinin receptors for the treatment of neurological diseases. *Curr. Pharm. Des.*, **11**, 1313–1326.

References:

GOBEIL, F. *et al.* (1996a). *Hypertension*, **28**, 833–839.

GOBEIL, F. *et al.* (1996b). *Br. J. Pharmacol.*, **118**, 289–294.

GOBEIL, F. *et al.* (1999). *Hypertension*, **33**, 823–829.

RIZZI, A. *et al.* (1997). *Hypertension*, **29**, 951–956.

PRUNEAU, D. *et al.* (1999). *Immunopharmacology*, **43**, 187–194.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Calcitonin, amylin, CGRP and adrenomedullin

Overview: Calcitonin (CT), amylin (AMY), calcitonin gene-related peptide (CGRP) and adrenomedullin (AM) receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on CGRPs, AM, AMY, and calcitonin receptors, Poyner *et al.*, 2002) are generated by the genes *CALCR* (which codes for the calcitonin receptor) and *CALCRL* (which codes for the calcitonin receptor-like receptor, CL receptor, previously known as CRLR), whose function and pharmacology are altered in the presence of RAMPs (receptor activity-modifying protein). RAMPs are single TM domain proteins of *ca.* 130 amino acid, identified as a family of three members; RAMP1 (ENSG00000132329), RAMP2 (ENSG00000131477) and RAMP3 (ENSG00000122679). There are splice variants of the CT receptor; these in turn produce variants of the AMY receptor (Poyner *et al.*, 2002). The endogenous agonists are the peptides calcitonin, α CGRP (occasionally termed CGRP-I), β CGRP (occasionally termed CGRP-II), AMY (previously termed islet-amyloid polypeptide, diabetes-associated polypeptide) and AM. There are species differences in peptide sequences, particularly for the calcitonins. AM2/Intermedin (AM2/IMD) is a recently discovered peptide, which can activate CGRP1, AM1, AM2 and AMY1 receptors, albeit less potently than the cognate agonists (Ogoshi *et al.*, 2003; Roh *et al.*, 2004; Hay *et al.*, 2005). Calcitonin receptor-stimulating peptide (CRSP) is another member of the family with selectivity for the calcitonin receptor; it has not been found in humans (Katafuchi *et al.*, 2003). BIBN4096BS is the most selective antagonist available, having a high selectivity for CGRP1 receptors, with a particular preference for those of primate origin. CGRP-(8-37) acts as an antagonist of CGRP (pK_i 6.5–8.0) and inhibits some AM and AMY responses (7.0). It is inactive at calcitonin receptors. Salmon calcitonin-(8-32) is an antagonist at both *amylin* and calcitonin receptors. AC187, a salmon calcitonin analogue, is also an antagonist at *amylin* and calcitonin receptors. Human AM-(22-52) has some selectivity towards AM receptors, but with modest potency, limiting its use.

Nomenclature Composition	Calcitonin <i>CALCR</i>	Amylin <i>AMY1: CALCR + RAMP1</i> <i>AMY2: CALCR + RAMP2</i> <i>AMY3: CALCR + RAMP3</i>	CGRP <i>CGRP1: CALCRL + RAMP1</i>	Adrenomedullin <i>AM1: CALCRL + RAMP2</i> <i>AM2: CALCRL + RAMP3</i>
Ensemble ID	ENSG00000004948	—	ENSG000000064989	—
Principal transduction	G _s /G _q	G _s	G _s /G _q	G _s
Rank order of potency	Salmon CT $\geq$ human CT $\geq$ AMY, CGRP $>$ AM, AM2/IMD	AMY1a: Salmon CT $\geq$ AMY $\geq$ CGRP $>$ AM2/ IMD $>$ human CT $>$ AM AMY3a: Salmon CT $\geq$ AMY $>$ CGRP $>$ AM2/ IMD $>$ human CT $>$ AM	CGRP $>$ AM $\geq$ AM2/ IMD $>$ AMY $\geq$ salmon CT	AM1: AM $\gg$ CGRP, AM2/ IMD $>$ AMY $>$ salmon CT AM2: AM $\geq$ CGRP, AM2/ IMD $>$ AMY $>$ salmon CT
Selective agonists	Human CT	AMY	α CGRP	AM
Selective antagonists	—	—	BIBN4096BS (11, Doods <i>et al.</i> , 2000; Hay <i>et al.</i> , 2003)	AM-(22–52) (7)
Probes	[¹²⁵ I]-CT (salmon, 0.1 nM), [¹²⁵ I]-CT (human, 0.1–1.0 nM)	[¹²⁵ I]-BH-AMY (rat, 0.1–1.0 nM)	[¹²⁵ I]- α CGRP (0.1 nM)	[¹²⁵ I]-AM (rat, 0.1–1.0 nM)

The agonists described represent the best available but their selectivity is limited. AM has appreciable affinity for CGRP receptors and some of its effects can be antagonised by CGRP-(8-37). CGRP can show significant crossreactivity at *amylin* receptors and some AM receptors. Responsiveness to human CT can be affected by splice variation (at the rat C1b receptor it is very weak, Houssami *et al.*, 1994). Particularly for AMY receptors, relative potency can vary with the type and level of RAMP present and can be influenced by other factors such as G-proteins (Tilakaratne *et al.*, 2000).

G_s is a prominent route for effector coupling but other pathways (e.g. Ca²⁺ and nitric oxide) and G-proteins can be activated. The coupling can be affected by splice variants of the CT receptor (e.g. the 490 amino-acid form of the human receptor, CT(b), does not cause an increase in intracellular Ca²⁺ and might have low efficacy in generating cAMP).

There is evidence that CGRP-RCP (a 148 amino-acid hydrophilic protein, ENSG00000126522) is important for the coupling of the CL receptor to adenylyl cyclase (Evans *et al.*, 2000). When coexpressed with RAMP2, the CL receptor produces an AM receptor (AM1). RAMP3 interacts with the CL receptor to give another receptor that is responsive to AM (AM2, Fraser *et al.*, 1999). There is some evidence that these AM receptors are pharmacologically distinct (Hay *et al.*, 2003). Transfection of hCT(a) with any RAMP can give a receptor with a high affinity for both salmon CT and AMY and varying affinity for different antagonists (Christopoulos *et al.*, 1999; Hay *et al.*, 2005). hCT(a)–RAMP1 has a high affinity for CGRP, unlike hCT(a)–RAMP3 (Christopoulos *et al.*, 1999; Hay *et al.*, 2005). However, AMY receptor phenotype is RAMP-type- and cell-line-dependent (Tilakaratne *et al.*, 2000).

[¹²⁵I]-Salmon calcitonin is the most common radioligand for calcitonin receptors but it has high affinity for *amylin* receptors and is also poorly reversible. [¹²⁵I]-Tyr⁰-CGRP is widely used as a radioligand for CGRP receptors.

CGRP₁ and CGRP₂ subtypes have been proposed on the basis of the action of the agonists [Cys(ACM)^{2,7}]CGRP or [Cys(Et)^{2,7}]CGRP (putative CGRP₂-selective agents) and antagonist CGRP-(8-37) (CGRP₁-selective, pK_i 7.0–8.0, Juaneda *et al.*, 2000). CL/RAMP1 represents the CGRP₁ subtype previously described in native tissues and cell lines (Aiyar *et al.*, 1996; McLatchie *et al.*, 1998). There is not yet a clear molecular correlate for the CGRP₂ receptor, although in some cases it may represent CGRP acting via AM2 or *amylin* receptors.

Abbreviations: AC187, acetyl-[Asn³⁰,Tyr³²]salmon CT; BIBN4096BS, 1-piperidinecarboxamide, *N*-(2-[(5-amino-1-[(4-{4-pyridinyl}-1-piperazinyl)carbonyl]pentyl)amino]-1-[(3,5-dibromo-4-hydroxyphenyl)methyl]-2-oxoethyl)-4-(1,4-dihydro-2-oxo-3[2*H*]-quinazolinyl)); [Cys(ACM)^{2,7}]CGRP, [acetamidomethyl-Cys^{2,7}]CGRP; [Cys(Et)^{2,7}]CGRP, [ethylamide-Cys^{2,7}]CGRP

Further Reading:

BRAIN, S.D. & GRANT, A.D. (2004). Vascular actions of calcitonin gene-related peptide and adrenomedullin. *Physiol. Rev.*, **84**, 903–934.

DURHAM, P.L. (2004). CGRP receptor antagonists: a new choice for acute treatment of migraine? *Curr. Opin. Invest. Drugs*, **5**, 731–735.

HAY, D.L., POYNER, D. & SEXTON, P.M. (2006). GPCR modulation by RAMPs. *Pharmacol. Ther.*, **109**, 173–197.

HAY, D.L., POYNER, D. & DICKERSON, I. (2002). CGRP receptor heterogeneity: a role for receptor component protein? *Trends Endocrinol. Metab.*, **14**, 3–4.

LIPTON, R.B. & DODICK, D.W. (2004). CGRP antagonists in the acute treatment of migraine. *Lancet Neurol.*, **3**, 332.

POYNER, D.R., SEXTON, P.M., MARSHALL, I., SMITH, D.M., QUIRION, R., BORN, W., MUFF, R., FISCHER, J.A. & FOORD, S.M. (2002). International Union of Pharmacology. XXXII. The mammalian calcitonin gene-related peptides, adrenomedullin, amylin, and calcitonin receptors. *Pharmacol. Rev.*, **54**, 233–246.

PURDUE, B.W., TILAKARATNE, N. & SEXTON, P.M. (2002). Molecular pharmacology of the calcitonin receptor. *Receptors Channels*, **8**, 243–255.

References:

- AIYAR, N. *et al.* (1996). *J. Biol. Chem.*, **271**, 11325–11329.
- CHRISTOPOULOS, G. *et al.* (1999). *Mol. Pharmacol.*, **56**, 235–242.
- DOODS, H. *et al.* (2000). *Br. J. Pharmacol.*, **129**, 420–423.
- EVANS, B.N. *et al.* (2000). *J. Biol. Chem.*, **275**, 31438–31443.
- FRASER, N.J. *et al.* (1999). *Mol. Pharmacol.*, **55**, 1054–1059.
- HAY, D.L. *et al.* (2003). *Br. J. Pharmacol.*, **140**, 477–486.
- HAY, D.L. *et al.* (2005). *Mol. Pharmacol.*, **67**, 1655–1665.
- HOUSSAMI, S. *et al.* (1994). *Endocrinology*, **135**, 183–190.
- JUANEDA, C. *et al.* (2000). *Trends Pharmacol. Sci.*, **21**, 432–438.
- KATAFUCHI, T. *et al.* (2003). *J. Biol. Chem.*, **278**, 12046–12054.
- MCLATCHIE, L.M. *et al.* (1998). *Nature*, **393**, 333–339.
- OGOSHI, M. *et al.* (2003). *Biochem. Biophys. Res. Commun.*, **311**, 1072–1077.
- POYNER, D.R. *et al.* (2002). *Pharmacol. Rev.*, **54**, 233–246.
- ROH, J. *et al.* (2004). *J. Biol. Chem.*, **279**, 7264–7274.
- TILAKARATNE, N. *et al.* (2000). *J. Pharmacol. Exp. Ther.*, **294**, 61–72.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Calcium-sensing

Overview: The calcium-sensing receptor (CaR, provisional nomenclature) responds to extracellular calcium and magnesium in the millimolar range and to gadolinium and some polycations in the micromolar range (Brown *et al.*, 1993). The sensitivity of CaR to primary agonists can be increased by aromatic L-amino acids (Conigrave *et al.*, 2000).

Nomenclature	CaR
Other names	Parathyroid cell calcium-sensing receptor, CASR
Ensembl ID	ENSG00000036828
Principal transduction	G _{q/11} , G _{i/o} , G _{12/13} (Ward <i>et al.</i> , 2004)
Cation rank order of potency	Gd ³⁺ > Ca ²⁺ > Mg ²⁺ (Brown <i>et al.</i> , 1993)
Polyamine rank order of potency	Spermine > spermidine > putrescine (Quinn <i>et al.</i> , 1997)
Amino-acid rank order of potency	L-Phe, L-Trp, L-His > L-Ala > L-Ser, L-Pro, L-Glu > L-Asp but not L-Lys, L-Arg, L-Leu, and L-Ile (Conigrave <i>et al.</i> , 2000)
Positive allosteric modulators	NPSR568 (Nemeth <i>et al.</i> , 1998), calindol (Petrel <i>et al.</i> , 2004), cinacalcet (Nemeth <i>et al.</i> , 2004)
Negative allosteric modulators	NPS2143, NPS89636 (Nemeth <i>et al.</i> , 2001), Calhex-231 (Petrel <i>et al.</i> , 2004), 2-benzylpyrrolidine derivatives of NPS2143 (Yang <i>et al.</i> , 2005)

Positive allosteric modulators are termed calcimimetics (Hammerland *et al.*, 1998) and can suppress parathyroid hormone secretion, whereas negative allosteric modulators are called calcilytics and can act to increase parathyroid hormone secretion (Nemeth *et al.*, 2001).

The central role of CaR in maintaining whole-body extracellular calcium homeostasis is seen in patients with loss-of-function CaR mutations who develop familial hypocalciuric hypercalcaemia (heterozygous mutation) or neonatal severe hyperparathyroidism (homozygous mutation) and in CaR null mice (Ho *et al.*, 1995), which exhibit similar increases in PTH secretion and blood Ca²⁺ levels. A gain-of-function mutation in the CaR gene is associated with autosomal dominant hypocalcaemia.

Abbreviations: **Calhex-231**, (1S,2S,1'R)-N¹-(4-chlorobenzoyl)-N²-[1-(1-naphthyl)ethyl]-1,2-diaminocyclohexane; **calindol**, (R)-2-[1-(1-naphthyl) ethylaminomethyl]-1H-indole; **NPSR568**, (R)-N-(3-methoxy- α -phenylethyl)-3-(2-chlorophenyl)-1-propylamine hydrochloride; **NPS2143**, N-[(R)-2-hydroxy-3-(2-cyano-3-chlorophenoxy)propyl]-1,1-dimethyl-2-(2-naphthyl)ethylamine

Further Reading:

BROWN, E.M. & MACLEOD, R.J. (2001). Extracellular calcium sensing and extracellular calcium signaling. *Physiol Rev*, **81**, 239–297.

CHEN, R.A. & GOODMAN, W.G. (2004). Role of the calcium-sensing receptor in parathyroid gland physiology. *Am J Physiol Renal Physiol*, **286**, F1005–F1011.

CONIGRAVE, A.D. & BROWN, E.M. (2006). Taste Receptors in the Gastrointestinal Tract II. L-Amino acid sensing by calcium-sensing receptors: implications for GI physiology. *Am. J. Physiol. Gastrointest. Liver Physiol.*, **291**, G753–G761.

HAUACHE, O.M. (2001). Extracellular calcium-sensing receptor: structural and functional features and association with diseases. *Braz. J. Med. Biol. Res.*, **34**, 577–584.

HU, J. & SPIEGEL, A.M. (2003). Naturally occurring mutations of the extracellular Ca²⁺-sensing receptor: implications for its structure and function. *Trends Endocrinol. Metab.*, **14**, 282–288.

RICCARDI, D. (2004). Cell calcium special issue (14 Reviews) on calcium-sensing receptor: physiology, pathology and pharmacological modulation. *Cell Calcium*, **35**, 179–306.

RODRIGUEZ, M., NEMETH, E. & MARTIN, D. (2005). The calcium-sensing receptor: a key factor in the pathogenesis of secondary hyperparathyroidism. *Am. J. Physiol. – Renal Physiol.*, **288**, F253–F264.

STEDDON, S.J. & CUNNINGHAM, J. (2005). Calcimimetics and calcilytics – fooling the calcium receptor. *Lancet*, **365**, 2237–2239.

References:

BROWN, E.M. *et al.* (1993). *Nature*, **366**, 575–580.

CONIGRAVE, A.D. *et al.* (2000). *Proc. Natl. Acad. Sci. USA*, **97**, 4814–4819.

HAMMERLAND, L.G. *et al.* (1998). *Mol. Pharmacol.*, **53**, 1083–1088.

HO, C. *et al.* (1995). *Nat. Genet.*, **11**, 389–394.

NEMETH, E.F. *et al.* (2004). *J. Pharmacol. Exp. Ther.*, **308**, 627–635.

NEMETH, E.F. *et al.* (1998). *Proc. Natl. Acad. Sci. USA*, **95**, 4040–4045.

NEMETH, E.F. *et al.* (2001). *J. Pharmacol. Exp. Ther.*, **299**, 323–331.

PETREL, C. *et al.* (2004). *J. Biol. Chem.*, **279**, 18990–18997.

QUINN, S.J. *et al.* (1997). *Am. J. Physiol. – Cell Physiol.*, **273**, C1315–C1323.

WARD, D.T. (2004). *Cell Calcium*, **35**, 217–228.

YANG, W. *et al.* (2005). *Bioorg. Med. Chem. Lett.*, **15**, 1225–1228.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Cannabinoid

Overview: Cannabinoid receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Cannabinoid Receptors; Howlett *et al.*, 2002) are activated by endogenous ligands that include *N*-arachidonylethanolamine (anandamide), *N*-homo- γ -linolenylethanolamine, *N*-docosatetra-7,10,13,16-enylethanolamine, 2-arachidonoylglycerol and 2-arachidonoylglycerol ether (noladin ether). Potency determinations are complicated by the possibility of differential susceptibility of endogenous ligands to enzymatic conversion. Both CB₁ and CB₂ receptors may be labelled with [³H]-HU243 (45–61 pM; Devane *et al.*, 1992; Bayewitch *et al.*, 1995), [³H]-CP55940 (0.6 nM; Showalter *et al.*, 1996) and [³H]-WIN55212-2 (2–12 nM; Slipetz *et al.*, 1995; Song & Bonner, 1996).

Nomenclature	CB₁	CB₂
Ensembl ID	ENSG00000118432	ENSG00000162562
Principal transduction	G _{i/o}	G _{i/o}
Selective agonists	ACEA (Hillard <i>et al.</i> , 1999), ACPA (Hillard <i>et al.</i> , 1999), methanandamide (Khanolkar <i>et al.</i> , 1996), O1812 (Di Marzo <i>et al.</i> , 2001)	HU308 (Hanus <i>et al.</i> , 1999), JWH133 (Huffman <i>et al.</i> , 1999; Pertwee, 2000), L759633 (Ross <i>et al.</i> , 1999), L759656 (Ross <i>et al.</i> , 1999), AM1241 (Ibrahim <i>et al.</i> , 2003)
Selective antagonists	AM251 (8.1, Lan <i>et al.</i> , 1999a), AM281 (7.9, Lan <i>et al.</i> , 1999b) SR141716A (7.9, Showalter <i>et al.</i> , 1996), LY320135 (6.9, Felder <i>et al.</i> , 1998),	SR144528 (9.2, Rinaldi-Carmona <i>et al.</i> , 1998), AM630 (7.5, Ross <i>et al.</i> , 1999)
Probes	[³ H]-SR141716A (0.6 nM, Rinaldi-Carmona <i>et al.</i> , 1996)	—

Anandamide is also a vanilloid receptor (TRPV1) agonist (Pertwee, 2004; Ross, 2003). A putative 'CB₃' receptor has been suggested following studies of the orphan 7TM receptor GPR55 (ENSG00000135898; Brown *et al.*, 2005; Sjögren *et al.*, 2005). Other pharmacological targets for cannabinoids have also been proposed (Begg *et al.*, 2005; Pertwee, 2004). All of the compounds listed as antagonists behave as inverse agonists in some bioassay systems (Pertwee, 2005). There is evidence for an allosteric site on the CB₁ receptor (Price *et al.*, 2005).

Abbreviations: ACEA, arachidonoyl-2-chloroethylamide; ACPA, arachidonoylcyclopropylamide; **AM251**, *N*-(piperidin-1-yl)-5-(4-iodophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide; **AM281**, 1-(2,4-dichlorophenyl)-5-(4-iodophenyl)-4-methyl-*N*-4-morpholinyl-1*H*-pyrazole-3-carboxamide; **AM630**, 6-iodopravadoline; **AM1241**, (2-iodo-5-nitro-phenyl)-[1-(1-methyl-piperidin-2-ylmethyl)-1*H*-indol-3-yl]-methanone; **CP55940**, (1*R*,3*R*,4*R*)-3-[2-hydroxy-4-(1,1-dimethylheptyl)phenyl]-4-(3-hydroxypropyl)cyclohexan-1-ol; **HU243**, [6*aR*-(6*ax*,9*z*,10*aβ*)]-3-(1,1-dimethylheptyl)-6*a*,7,8,9,10,10*a*-hexahydro-1-hydroxy-6,6-dimethyl-6*H*-dibenzo[*b,d*]pyran-[7,8-3*H*]-9-methanol; **HU308**, {4-[4-(1,1-dimethylheptyl)-2,6-dimethoxy-phenyl]-6,6-dimethyl-bicyclo[3.1.1]hept-2-en-2-yl]-methanol; **JWH133**, (3-(1,1-dimethylbutyl)-6,6,9-trimethyl-6*a*,7,10,10*a*-tetrahydro-6*H*-benzo[*c*]chromene; **L759633**, (6*aR*,10*aR*)-3-(1,1-dimethylheptyl)-1-methoxy-6,6,9-trimethyl-6*a*,7,10,10*a*-tetrahydro-6*H*-benzo[*c*]chromene; **L759656**, (6*aR*,10*aR*)-3-(1,1-dimethylheptyl)-1-methoxy-6,6-dimethyl-9-methylene-6*a*,7,8,9,10,10*a*-hexahydro-6*H*-benzo[*c*]chromene; **LY320135**, (6-methoxy-2-[4-methoxyphenyl]benzo[*b*]thien-3-yl)(4-cyanophenyl)methanone; **methanandamide**, (R)-(+)-arachidonoyl-1'-hydroxy-2'-propylamide; **O1812**, (R)-(20-cyano-16,16-dimethyldocosa-*cis*-5,8,11,14-tetraenyl)-1'-hydroxy-2'-propylamide; **SR141716A**, *N*-(piperidin-1-yl)-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide hydrochloride, also known as rimonabant; **SR144528**, *N*-([1*S*]-endo-1,3,3-trimethylbicyclo[2.2.1]heptan-2-yl)-5-(4-chloro-3-methylphenyl)-1-(4-methylbenzyl)-pyrazole-3-carboxamide; **WIN55212-2**, (R)-(+)-[2,3-dihydro-5-methyl-3-(4-morpholinylmethyl)pyrrolo-[1,2,3-*de*]-1,4-benzoxazin-6-yl]-1-naphthalenylmethanone mesylate

Further Reading:

- BISOGNO, T., LIGRESTI, A. & DI MARZO, V. (2005). The endocannabinoid signalling system: biochemical aspects. *Pharmacol. Biochem. Behav.*, **81**, 224–238.
- BRADSHAW, H.B. & WALKER, J.M. (2005). The expanding field of cannabimimetic and related lipid mediators. *Br. J. Pharmacol.*, **144**, 459–465.
- DAVIES, S.N., PERTWEE, R.G. & RIEDEL, G. (2002). Functions of cannabinoid receptors in the hippocampus. *Neuropharmacology*, **42**, 993–1007.
- DE PETROCELLIS, L., CASCIO, M.G. & DI MARZO, V. (2004). The endocannabinoid system: a general view and latest additions. *Br. J. Pharmacol.*, **141**, 765–774.
- DI MARZO, V. & MATIAS, I. (2005). Endocannabinoid control of food intake and energy balance. *Nat. Neurosci.*, **8**, 585–589.
- DIANA, M.A. & MARTY, A. (2004). Endocannabinoid-mediated short-term synaptic plasticity: depolarization-induced suppression of inhibition (DSI) and depolarization-induced suppression of excitation (DSE). *Br. J. Pharmacol.*, **142**, 9–19.
- FREUND, T.F., KATONA, I. & PIOMELLI, D. (2003). Role of endogenous cannabinoids in synaptic signaling. *Physiol. Rev.*, **83**, 1017–1066.
- HOWLETT, A.C., BARTH, F., BONNER, T.I., CABRAL, G., CASELLAS, P., DEVANE, W.A., FELDER, C.C., HERKENHAM, M., MACKIE, K., MARTIN, B.R., MECOULAM, R. & PERTWEE, R.G. (2002). International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol. Rev.*, **54**, 161–202.
- IDRIS, A.I., VAN 'T HOF, R.J., GREIG, I.R., RIDGE, S.A., BAKER, D., ROSS, R.A. & RALSTON, S.H. (2005). Regulation of bone mass, bone loss and osteoclast activity by cannabinoid receptors. *Nat. Med.*, **11**, 774–779.
- KLEIN, T.W. (2005). Cannabinoid-based drugs as anti-inflammatory therapeutics. *Nat. Rev. Immunol.*, **5**, 400–411.
- LANGE, J.H.M. & KRUSE, C.G. (2005). Medicinal chemistry strategies to CB₁ cannabinoid receptor antagonists. *Drug Discov. Today*, **10**, 693–702.
- LE FOLL, B. & GOLDBERG, S.R. (2005). Cannabinoid CB₁ antagonists as promising new medications for drug dependence. *J. Pharmacol. Exp. Ther.*, **312**, 875–883.
- MCALLISTER, S.D. & GLASS, M. (2002). CB₁ and CB₂ receptor-mediated signalling: a focus on endocannabinoids. *Prostagland. Leukotr. Essent. Fatty Acids*, **66**, 161–171.
- PARK, B., MCPARTLAND, J.M. & GLASS, M. (2004). Cannabis, cannabinoids and reproduction. *Prostagland. Leukotr. Essent. Fatty Acids*, **70**, 189–197.
- PERTWEE, R.G. (2005). Pharmacological actions of cannabinoids. In *Cannabinoids*, ed. Pertwee, R.G. *Handbook Exp. Pharmacol.*. Heidelberg: Springer-Verlag, **168**, 1–51.
- PERTWEE, R.G. (2006). Cannabinoid pharmacology: the first 66 years. *Br. J. Pharmacol.*, **147**, S163–S171.
- PRYCE, G. & BAKER, D. (2005). Emerging properties of cannabinoid medicines in management of multiple sclerosis. *Trends Neurosci.*, **28**, 272–276.
- RODRÍGUEZ DE FONSECA, F., DEL ARCO, I., BERMUDEZ-SILVA, F.J., BILBAO, A., CIPPITELLI, A. & NAVARRO, M. (2005). The endocannabinoid system: physiology and pharmacology. *Alcohol. Alcohol.*, **40**, 2–14.
- WALTER, L. & STELLA, N. (2004). Cannabinoids and neuroinflammation. *Br. J. Pharmacol.*, **141**, 775–785.

References:

- BAYEWITCH, M. *et al.* (1995). *FEBS Lett.*, **375**, 143–147.
- BEGG, M. *et al.* (2005). *Pharmacol. Ther.*, **106**, 133–145.
- BROWN, A.J. *et al.* (2005). *2005 Symposium on the Cannabinoids*. Burlington, Vermont: International Cannabinoid Research Society, p 16.
- DEVANE, W.A. *et al.* (1992). *J. Med. Chem.*, **35**, 2065–2069.
- DI MARZO, V. *et al.* (2001). *Biochem. Biophys. Res. Commun.*, **281**, 444–451.
- FELDER, C.C. *et al.* (1998). *J. Pharmacol. Exp. Ther.*, **284**, 291–297.
- HANUS, L. *et al.* (1999). *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 14228–14233.
- HILLARD, C.J. *et al.* (1999). *J. Pharmacol. Exp. Ther.*, **289**, 1427–1433.
- HOWLETT, A.C. *et al.* (2002). *Pharmacol. Rev.*, **54**, 161–202.
- HUFFMAN, J.W. *et al.* (1999). *Bioorg. Med. Chem.*, **7**, 2905–2914.
- IBRAHIM, M.M. *et al.* (2003). *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 10529–10533.
- KHANOLKAR, A.D. *et al.* (1996). *J. Med. Chem.*, **39**, 4515–4519.
- LAN, R. *et al.* (1999a). *J. Med. Chem.*, **42**, 769–776.
- LAN, R. *et al.* (1999b). *AAPS Pharm. Sci.*, **1**, U13–U24.
- PERTWEE, R.G. (2000). *Exp. Opin. Invest. Drugs*, **9**, 1553–1571.
- PERTWEE, R.G. (2004). *Curr. Neuropharmacol.*, **2**, 9–29.
- PERTWEE, R.G. (2005). *Life Sci.*, **76**, 1307–1324.
- PRICE, M.R. *et al.* (2005). *Mol. Pharmacol.*, **68**, 1484–1495.
- RINALDI-CARMONA, M. *et al.* (1996). *Life Sci.*, **58**, 1239–1247.
- RINALDI-CARMONA, M. *et al.* (1998). *J. Pharmacol. Exp. Ther.*, **284**, 644–650.
- ROSS, R.A. *et al.* (1999). *Br. J. Pharmacol.*, **126**, 665–672.
- ROSS, R.A. (2003). *Br. J. Pharmacol.*, **140**, 790–801.
- SHOWALTER, V.M. *et al.* (1996). *J. Pharmacol. Exp. Ther.*, **278**, 989–999.
- SJÖGREN, S. *et al.* (2005). *2005 Symposium on the Cannabinoids*. Burlington, Vermont: International Cannabinoid Research Society, p 106.
- SLIPETZ, D.M. *et al.* (1995). *Mol. Pharmacol.*, **48**, 352–361.
- SONG, Z.H. & BONNER, T.I. (1996). *Mol. Pharmacol.*, **49**, 891–896.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Chemokine

Overview: Chemokine receptors (nomenclature agreed by NC-IUPHAR Subcommittee on Chemokine Receptors, Murphy *et al.*, 2000; Murphy, 2002) comprise a large subfamily of 7TM receptors activated by one or more of the chemokines, a large family of small cytokines typically possessing chemotactic activity for leukocytes. Chemokines can be divided by structure into four subclasses by the number and arrangement of conserved cysteines. CC (also known as β -chemokines; $n = 28$), CXC (also known as α -chemokines; $n = 16$) and CX₃C ($n = 1$) chemokines all have four conserved cysteines, with zero, one and three amino acids separating the first two cysteines, respectively. C chemokines ($n = 2$) have only the second and fourth cysteines found in other chemokines. Chemokines can also be classified by function into homeostatic and inflammatory subgroups. Most chemokine receptors are able to bind multiple high affinity chemokine ligands, but the ligands for a given receptor are almost always restricted to the same structural subclass. Most chemokines bind to more than one receptor subtype. Receptors for inflammatory chemokines are typically highly promiscuous with regard to ligand specificity, and may lack a selective endogenous ligand. Listed are those human agonists with EC₅₀ values < 50 nM in either Ca²⁺ flux or chemotaxis assays at human recombinant receptors expressed in mammalian cell lines. There can be substantial cross-species differences in the sequences of both chemokines and chemokine receptors, and in the pharmacology and biology of chemokine receptors. Endogenous and HIV-encoded non-chemokine ligands have also been identified for chemokine receptors. Many chemokine receptors function as HIV co-receptors, and at least two, CCR5 and CXCR4, play prominent roles in pathogenesis. The tables include both standard chemokine names (Zlotnik and Yoshie, 2000) and the most commonly used synonyms. Numerical data quoted are typically pKi or pIC₅₀ values from radioligand binding to heterologously expressed receptors.

Nomenclature	CCR1	CCR2	CCR3	CCR4	CCR5
Other names	CCR1, CC CK ₁ , CC CKR1, MIP-1 α R, MIP-1 α /RANTES	CCR2, CC CK ₂ , CC CKR2, MCP-1	CCR3, CC CK ₃ , CC CKR3	CCR4, CC CK ₄ , CC CKR4	CCR5, CC CK ₅ , CC CKR ₅ , CHEMR13
Ensembl ID	ENSG00000163823	ENSG00000121807	ENSG00000183625	ENSG00000183813	ENSG00000160791
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Agonists	CCL3 (MIP-1 α), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (MCP-2), CCL13 (MCP-4), CCL14a (HCC-1), CCL15 (HCC-2), CCL23 (MPIF-1)	CCL2 (MCP-1), CCL7 (MCP-3), CCL8 (MCP-2), CCL13 (MCP-4), CCL16 (HCC-4), HIV-1 Tat	CCL11 (eotaxin), CCL5 (RANTES), CCL7 (MCP-3), CCL8 (MCP-2), CCL13 (MCP-4), CCL15 (HCC-2), CCL24 (eotaxin-2), CCL26 (eotaxin-3), CCL28 (MEC), HIV-1 Tat	CCL22 (MDC), CCL17 (TARC), HHV8 vMIP-III,	CCL3 (MIP-1 α), CCL4 (MIP-1 β), CCL5 (RANTES), CCL8 (MCP-2), CCL11 (eotaxin), CCL14a (HCC-1), CCL16 (HCC-4), R5 HIV-1 gp120
Selective agonists	CCL15 (HCC-2), CCL23 (MPIF-1)	CCL2 (MCP-1)	CCL11 (Eotaxin), CCL24 (eotaxin-2), CCL26 (eotaxin-3),	CCL22 (MDC), CCL17 (TARC)	MIP-1 β , R5-HIV gp120
Selective antagonists	BX471 (8.3-9), 2b-1 (8.7), UCB35625 (8.0), CP-481,715 (8.0), CCL4 (MIP-1 β)	CCL11 (eotaxin), CCL26 (eotaxin-3), GSK Compound 34 (7.6)	Banyu Compound 1b (8.6), SB328437 (8.4), BMS Compound 87b (8.1), CXCL10 (IP10), CXCL9 (Mig), CXCL11 (I-TAC)	—	TAK779 (9.0), CCL7 (MCP-3), SCH C, SCH D, MRK-1, E913 (8.7)
Probes	[¹²⁵ I]-MIP-1 α , [¹²⁵ I]-RANTES, [¹²⁵ I]-MCP-3	[¹²⁵ I]-MCP-1, [¹²⁵ I]-MCP-3	[¹²⁵ I]-RANTES, [¹²⁵ I]-eotaxin, [¹²⁵ I]-MCP-3	[¹²⁵ I]-TARC, [¹²⁵ I]-CTACK/CCL27	[¹²⁵ I]-RANTES, [¹²⁵ I]-MCP-2, [¹²⁵ I]-MIP-1 α , [¹²⁵ I]-MIP-1 β

Nomenclature	CCR6	CCR7	CCR8	CCR9	CCR10
Other names	GPR-CY4, CKR-L3, STRL-22, DRY-6, DCR2, BN-1, GPR29	EBI-1, BLR-2	TER1, CKR-L1, GPR-CY6, ChemR1	GPR 9-6	GPR-2
Ensembl ID	ENSG00000153467	ENSG00000126353	ENSG00000179934	ENSG00000173585	ENSG00000184451
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Agonists	CCL20 (LARC), HBD2	CCL19 (ELC, MIP-3 β), CCL21 (SLC)	CCL1 (I-309), CCL4 (MIP-1 β), CCL16 (HCC-4), CCL17 (TARC), HHV8 vMIP-I	CCL25 (TECK)	CCL27 (Eskine, ALP, CTACK), CCL28 (MEC)
Selective agonists	LARC, HBD2	ELC, SLC	I-309	TECK	Eskine, MEC
Selective antagonists	—	—	MCV MC148R (vMCC-I)	—	—
Probes	[¹²⁵ I]-LARC	[¹²⁵ I]-ELC, [¹²⁵ I]-SLC	[¹²⁵ I]-I309	[¹²⁵ I]-TECK	—

Nomenclature	CXCR1	CXCR2	CXCR3	CXCR4	CXCR5	CXCR6
Other names	IL8R _A , IL-8 receptor type I, IL-8 receptor α	IL8R _B , IL-8 receptor type II, IL-8 receptor β	IP10/Mig R, GPR9	HUMSTR, LESTR, fusin, HM89, LCR1	BLR-1, MDR15	STRL-33, BONZO, TYMSTR
Ensembl ID	ENSG00000163464	ENSG00000180871	SwissProt P49682	ENSG00000121966	ENSG00000160683	ENSG00000172215
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Agonists	CXCL6 (GCP-2), CXCL8 (IL-8), cytokine domain of tyrosyl tRNA synthetase	CXCL1 (GRO α), CXCL2 (GRO β), CXCL3 (GRO γ), CXCL5 (ENA-78), CXCL6 (GCP-2), CXCL7 (NAP-2), CXCL8 (IL-8), HCMV UL146 (vCXCL1)	CXCL9 (Mig), CXCL10 (IP10), CXCL11 (I-TAC)	CXCL12 α & β (SDF-1 α , SDF-1 β)	CXCL13 (BLC, BCA-1)	CXCL16 (SR-PSOX)
Selective agonists	—	GRO α , GRO γ , GRO β , NAP-2, ENA78	IP10, MIG, I-TAC	SDF-1 α , SDF-1 β , X4-HIV gp120	BLC	CXCL16
Selective antagonists	—	SB225002 (7.7)	eotaxin, MCP-3	AMD3100, HIV-1 Tat, T134, ALX41-4C	—	—
Probes	[¹²⁵ I]-IL8	[¹²⁵ I]-IL8, [¹²⁵ I]-GRO α , [¹²⁵ I]-NAP-2, [¹²⁵ I]-ENA78	[¹²⁵ I]-IP10, [¹²⁵ I]-I-TAC/CXCL11	[¹²⁵ I]-SDF-1	—	[¹²⁵ I]-CXCL16

CXCR1 and CXCR2 also couple to phospholipase C when co-transfected with members of the G_{q/11} family of G proteins. Mouse CXCR2 binds iodinated mouse KC and mouse MIP-2 with high affinity (mouse KC and MIP-2 are homologues of human GRO chemokines), but shows low affinity for human IL-8.

Nomenclature	CX₃CR1	XCRI
Other names	CMKBR1, V28	GPR5
Ensembl ID	ENSG00000168329	ENSG00000173578
Principal transduction	G _{i/o}	G _{i/o}
Agonists	CX3CL1 (Fractalkine)	XCL1 α and β (Lymphotoxin α and β)
Selective agonists	Fractalkine	Lymphotoxin
Probes	[¹²⁵ I]-Fractalkine	SEAP-XCL1

Three human 7TM chemokine binding proteins have been identified that lack a known signalling function: D6 (ENSG00000144648), which binds multiple CC chemokines; a molecule previously inappropriately named CCR11 and now known as CCX CKR or the human homologue of the bovine gustatory receptor PPAR1 (ENSG00000118519, ENSG00000129048), which binds ELC, SLC and TECK; and Duffy, a highly promiscuous CC and CXC chemokine binding protein expressed mainly on erythrocytes. Specific chemokine receptors facilitate cell entry by microbes, such as *Plasmodium vivax*, HIV-1 and the poxvirus myxoma virus. Virally encoded chemokine receptors are known (e.g. US28, a homologue of CCR1 from human cytomegalovirus and ECRF3, a homologue of CXCR2 from *Herpesvirus saimiri*), but their role in viral life cycles is not established. Viruses can exploit or subvert the chemokine system by producing chemokine antagonists and scavengers.

Abbreviations: BLC, B-lymphocyte chemokine; ELC, Epstein–Barr virus-induced receptor ligand chemokine; ENA-78, epithelial cell-derived neutrophil-activating factor-78 amino acids; GCP-2, granulocyte chemoattractant protein 2; HBD2, human β defensin 2; HCC, hemofiltrate CC chemokine; IL-8, interleukin 8; IP-10, γ -interferon-inducible protein 10; I-TAC, interferon-inducible T-cell α chemoattractant; LARC, liver and activation-related chemokine (CCL20); MCP, monocyte chemoattractant protein; MDC, macrophage-derived chemokine; MEC, mucosa expressed chemokine; MIG, monokine-induced by γ -interferon; MIP, macrophage inflammatory protein; MPIF-1, myeloid progenitor inhibitory factor 1; NAP-2, neutrophil-activating peptide 2; RANTES, regulated on activation normal T cell expressed and secreted; SDF, stromal cell-derived factor; SLC, secondary lymphoid tissue chemokine; SEAP, secreted alkaline phosphatase; TARC, T-cell and activation-related chemokine; TECK, thymus-expressed chemokine

The CC chemokine family (CCL1–28) includes I309 (CCL1), MCP-1 (CCL2), MIP-1 α (CCL3), MIP-1 β (CCL4), RANTES (CCL5), MCP-3 (CCL7), MCP-2 (CCL8), eotaxin (CCL11), MCP-4 (CCL13), HCC-1 (CCL14), Lkn-1/HCC-2 (CCL15), TARC (CCL17), ELC (CCL19), LARC (CCL20), SLC (CCL21), MDC (CCL22), MPIF-1 (CCL23), eotaxin-2 (CCL24), TECK (CCL25), eotaxin (CCL26), eskine/CTACK (CCL27) and MEC (CCL28). The CXC chemokine family (CXCL1–16) includes GRO α (CXCL1), GRO β (CXCL2), GRO γ (CXCL3), platelet factor 4 (CXCL4), ENA78 (CXCL5), GCP-2 (CXCL6), NAP-2 (CXCL7), IL-8 (CXCL8), MIG (CXCL9), IP10 (CXCL10), I-TAC (CXCL11), SDF-1 (CXCL12), BLC (CXCL13), BRAK (CXCL14), mouse lungkine (CXCL15) and SR-PSOX (CXCL16). The CX₃C chemokine (CX3CL1) is also known as fractalkine (neurotactin in the mouse). Unlike other chemokines, this molecule is multimodular containing a chemokine domain, an elongated mucin-like stalk, a transmembrane domain and a cytoplasmic tail. Both plasma membrane-associated and shed forms have been identified. The C chemokine (XCL1) is also known as lymphotoxin. The non-chemokine family includes the cytokine domain of tyrosyl-tRNA synthetase, HBD2, HIV gp120 and HIV Tat.

Further Reading:

- BOISVERT, W.A. (2004). Modulation of atherogenesis by chemokines. *Trends Cardiovasc. Med.*, **14**, 161–165.
- BOOMKER, J.M., DE LEIJ, L.F., THE, T.H. & HARMSSEN, M.C. (2005). Viral chemokine-modulatory proteins: tools and targets. *Cytokine Growth Factor Rev.*, **16**, 91–103.
- CARTIER, L., HARTLEY, O., DUBOIS-DAUPHIN, M. & KRAUSE, K.H. (2005). Chemokine receptors in the central nervous system: role in brain inflammation and neurodegenerative diseases. *Brain Res. Brain Res. Rev.*, **48**, 16–42.
- CHARO, I.F. & RANSOHOFF, R.M. (2006). The many roles of chemokines and chemokine receptors in inflammation. *N. Engl. J. Med.*, **354**, 610–621.
- COMERFORD, I. & NIBBS, R.J. (2005). Post-translational control of chemokines: a role for decoy receptors? *Immunol. Lett.*, **96**, 163–174.
- CYSTER, J.G. (2005). Chemokines, sphingosine-1-phosphate, and cell migration in secondary lymphoid organs. *Annu. Rev. Immunol.*, **23**, 127–159.
- EBERT, L.M., SCHÄERLI, P. & MOSER, B. (2005). Chemokine-mediated control of T cell traffic in lymphoid and peripheral tissues. *Mol. Immunol.*, **42**, 799–809.

- HANDEL, T.M., JOHNSON, Z., CROWN, S.E., LAU, E.K. & PROUDFOOT, A.E. (2005). Regulation of protein function by glycosaminoglycans – as exemplified by chemokines. *Annu. Rev. Biochem.*, **74**, 385–410.
- MOSER, B., WOLF, M., WALZ, A. & LOETSCHER, P. (2004). Chemokines: multiple levels of leukocyte migration control. *Trends Immunol.*, **25**, 75–84.
- OPPERMANN, M. (2004). Chemokine receptor CCR5: insights into structure, function, and regulation. *Cell Signal.*, **16**, 1201–1210.
- OWENS, T., BABCOCK, A.A., MILLWARD, J.M. & TOFT-HANSEN, H. (2005). Cytokine and chemokine inter-regulation in the inflamed or injured CNS. *Brain Res. Brain Res. Rev.*, **48**, 178–184.
- PUNEET, P., MOOCHHALA, S. & BHATIA, M. (2005). Chemokines in acute respiratory distress syndrome. *Am. J. Physiol. Lung Cell Mol. Physiol.*, **288**, L3–15.
- RIBEIRO, S. & HORUK, R. (2005). The clinical potential of chemokine receptor antagonists. *Pharmacol. Ther.*, **107**, 44–58.
- ROMAGNANI, P., LASAGNI, L., ANNUNZIATO, F., SERIO, M. & ROMAGNANI, S. (2004). CXC chemokines: the regulatory link between inflammation and angiogenesis. *Trends Immunol.*, **25**, 201–209.
- ROSENKILDE, M.M. (2005). Virus-encoded chemokine receptors—putative novel antiviral drug targets. *Neuropharmacology*, **48**, 1–13.
- ROT, A. & VON ANDRIAN, U.H. (2004). Chemokines in innate and adaptive host defense: basic chemokine grammar for immune cells. *Annu. Rev. Immunol.*, **22**, 891–928.
- SCHAERLI, P. & MOSER, B. (2005). Chemokines: control of primary and memory T-cell traffic. *Immunol. Res.*, **31**, 57–74.
- ZLOTNIK, A. (2006). Chemokines and cancer. *Int. J. Cancer*, **119**, 2026–2029.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Cholecystokinin

Overview: Cholecystokinin receptors (nomenclature recommended by the NC-IUPHAR Subcommittee on CCK receptors, Noble *et al.*, 1999) are activated by the endogenous peptides cholecystokinin (CCK)-4, CCK-8, CCK-33 and gastrin. There is evidence for species homologues of CCK₂ receptors distinguished by the relative affinities of the two stereoisomers of devazepide, *R*-L365260 and *S*-L365260, or by the differences in affinity of the agonist BC264 (Durieux *et al.*, 1992).

Nomenclature	CCK₁	CCK₂
Other names	CCK _A	CCK _B , CCK _B /gastrin
Ensembl ID	ENSG00000163394	ENSG00000110148
Principal transduction	G _{q/11} /G _s (Wu <i>et al.</i> , 1997)	G _s
Rank order of potency	CCK-8 ≫ gastrin, des-CCK-8 > CCK-4	CCK-8 ≫ gastrin, des-CCK-8, CCK-4
Selective agonists	A71623, JMV180, GW5823	Desulfated CCK-8, gastrin, CCK-4, BC264, RB400
Selective antagonists	Devazepide (9.8), T0632 (9.6), SR27897 (9.2), IQM95333 (9.2), PD140548 (7.9–8.6), lorglumide (7.2)	YM022 (10.2), L740093 (10.0), GV150013 (9.3), RP73870 (9.3), L365260 (7.5–8.7), LY262691 (7.5)
Probes	[³ H]-Devazepide (0.2 nM)	[³ H]-Propionyl-BC264 (0.15 nM), [³ H]-PD140376 (0.2 nM), [³ H]-L365260 (2 nM), [³ H]- or [¹²⁵ I]-gastrin (1 nM), [¹²⁵ I]-PD142308 (0.25 nM)

A mitogenic gastrin receptor, which can be radiolabelled with [¹²⁵I]-gastrin-(1–17) and which appears to couple to the G_s family of G proteins, has been described in human colon cancer cells (Bold *et al.*, 1994) and other cell lines (e.g. pancreatic AR42J and Swiss 3T3 fibroblasts, Seva *et al.*, 1994; Singh *et al.*, 1995).

Abbreviations: **A71623**, Boc-Trp-Lys(*O*-toluylaminocarbonyl)-Asp-(NMe)Phe-NH₂; **BC264**, Tyr(SO₃H)-gNle-mGly-Trp-(NMe)Nle-Asp-Phe-NH₂; **GV150013**, (+)-*N*-(1-[1-adamantane-1-methyl]-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-3-yl)-*N*'-phenylurea; **GW5823**, 2-[3-(1*H*-indazol-3-ylmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydrobenzo[*b*][1,4]diazepin-1-yl]-*N*-isopropyl-*N*-(methoxyphenyl)acetamide; **IQM95333**, (4*α*,5*R*)-2-benzyl-5[*N*-(*tert*-butoxycarbonyl)-L-Trp]amino-1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine; **JMV180**, Boc-Tyr(SO₃H)Ahx-Gly-Trp-Ahx-Asp²phenylethyl ester; **L365260**, 3*R*(+)-*N*-(2,3-dihydro-1-methyl-2-oxo-5-phenyl-1*H*-1,4-benzodiazepin-3-yl)-*N*'-(3-methylphenyl)urea; **L740093**, *N*-([3*R*]-5-[3-azabicyclo{3.2.2}nonan-3-yl]-2,3-dihydro-1-methyl-2-oxo-1*H*-1,4-benzodiazepin-3-yl)-*N*'-(3-methylphenyl)urea; **LY262691**, *trans*-*N*-(4-bromophenyl)-3-oxo-4,5-diphenyl-1-pyrazolidinecarboxamide(3.3.1.1^{3,7}); **PD140376**, L-3-([4-aminophenyl]methyl)-*N*-(α -methyl-*N*-[tricyclo(3.3.1.1^{0,2}-Trp)- β -Ala]; **PD140548**, *N*-(α -methyl-*N*-[tricyclo(3.3.1.1^{0,2}-Trp)-D-3-(phenylmethyl)- β -Ala]; **PD142308**, iodinated PD140548; **RB400**, HOOC-CH₂-CO-Trp-NMe(Nle)-Asp-Phe-NH₂; **RP73870**, ({[(*R*); **SR27897**, 1-([2-{4-(2-chlorophenyl)thiazole-2-yl}amino-6-methoxy-2-oxo-1*H*-indole]propanoate; **YM022**, (*R*)-1-(2,3-dihydro-1-[2'-methylphenacyl]-2-oxo-5-phenyl-1*H*-1,4-benzodiazepin-3-yl)-3-(3-methylphenyl)urea

Further Reading:

- DE TULLIO, P., DELARGE, J. & PIROTTE, B. (2000). Therapeutic and chemical developments of cholecystokinin receptor ligands. *Expert Opin. Invest. Drugs*, **9**, 129–146.
- HERRANZ, R. (2003). Cholecystokinin antagonists: pharmacological and therapeutic potential. *Med. Res. Rev.*, **23**, 559–605.
- INUI, A. (2003). Neuropeptide gene polymorphisms and human behavioural disorders. *Nat. Rev. Drug Discov.*, **2**, 986–998.
- KOPIN, A.S., MCBRIDE, E.W., SCHAFFER, K. & BEINBORN, M. (2000). CCK receptor polymorphisms: an illustration of emerging themes in pharmacogenomics. *Trends Pharmacol. Sci.*, **21**, 346–353.
- MORAN, T.H. (2000). Cholecystokinin and satiety: current perspectives. *Nutrition*, **16**, 858–865.
- NOBLE, F. & ROQUES, B.P. (1999). CCK_B receptor: chemistry, molecular biology, biochemistry and pharmacology. *Prog. Neurobiol.*, **56**, 1–31.
- NOBLE, F., WANK, S.A., CRAWLEY, J.N., BRADWEJN, J., SEROOGY, K.B., HAMON, M. & ROQUES, B.P. (1999). International Union of Pharmacology. XXI. Structure, distribution, and functions of cholecystokinin receptors. *Pharmacol. Rev.*, **51**, 745–781.
- PETER, S.A., D'AMATO, M. & BEGLINGER, C. (2006). CCKI antagonists: are they ready for clinical use? *Dig. Dis.*, **24**, 70–82.
- ROZENGURT, E. & WALSH, J. (2001). Gastrin, CCK, signaling, and cancer. *Annu. Rev. Physiol.*, **63**, 49–76.

References:

- BOLD, R.J. *et al.* (1994). *Biochem. Biophys. Res. Commun.*, **202**, 1222–1226.
- DURIEUX, C. *et al.* (1992). *Mol. Pharmacol.*, **41**, 1089–1095.
- SEVA, C. *et al.* (1994). *Science*, **265**, 410–412.
- SINGH, P. *et al.* (1995). *J. Biol. Chem.*, **270**, 8429–8438.
- WU, V. *et al.* (1997). *J. Biol. Chem.*, **272**, 9032–9042.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Corticotropin-releasing factor

Overview: Corticotropin-releasing factor (CRF, nomenclature as recommended by the NC-IUPHAR on Corticotropin-releasing Factor Receptors, see Hauger *et al.*, 2003) receptors are activated by the endogenous peptides CRF (also known as corticotropin-releasing hormone [CRH], a 41 aa peptide, ENSG00000147571), urocortin 1 (a 40 aa peptide, ENSG00000163794), urocortin 2 (a 38 aa peptide, ENSG00000145040) and urocortin 3 (a 38 aa peptide, ENSG00000178473). CRF₁ and CRF₂ receptors are activated non-selectively by CRF and urocortin 1. Binding to CRF receptors can be conducted using [¹²⁵I]-Tyr⁰-CRF or [¹²⁵I]-Tyr⁰-sauvagine with *K_d* values of 0.1–0.4 nM. CRF₁ and CRF₂ receptors are non-selectively antagonized by α -helical CRF-(9-41), D-Phe-CRF-(12-41) and astressin.

Nomenclature	CRF ₁	CRF ₂
Other names	CRF-RA, PC-CRF	CRF-RB, HM-CRF
Ensembl ID	ENSG00000120088	ENSG00000106113
Principal transduction	G _s	G _s
Selective agonists	—	Urocortin 2 (Reyes <i>et al.</i> , 2001), urocortin 3 (Lewis <i>et al.</i> , 2001)
Selective antagonists	CP154526 (8.3–9.0, Lundkvist <i>et al.</i> , 1996), NBI27914 (8.3–9.0, Chen <i>et al.</i> , 1996), antalarmin (8.3–9.0, Webster <i>et al.</i> , 1996), CRA1000 (8.3–9.0, Chaki <i>et al.</i> , 1999), DMP696 (8.3–9.0, He <i>et al.</i> , 2000), R121919 (8.3–9.0, Zobel <i>et al.</i> , 2000), SRA125543A (8.7–9.0, Gully <i>et al.</i> , 2002)	K41498 (9.2, Lawrence <i>et al.</i> , 2002), K31440 (8.7–8.8, Ruhmann <i>et al.</i> , 2002), antisauvagine-30 (Ruhmann <i>et al.</i> , 1998)

A CRF binding protein has been identified (CRF-BP, ENSG00000145708) to which both CRF and urocortin 1 bind with high affinities, which has been suggested to bind and inactivate circulating CRF (Perkins *et al.*, 1995).

Abbreviations: antalarmin, *N*-butyl-*N*-ethyl-(2,5,6-trimethyl)-7-[2,4,6-trimethylphenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl-amine; **astressin**, cyc^{30–33}[D-Phe¹²,Nle^{21,38},Glu³⁰,Lys³³]CRF-(12-41); **CP154526**, butyl-ethyl-(2,5-dimethyl-7-[2,4,6-trimethylphenyl]-7*H*-pyrrolo[2,3-*d*]pyrimidin-4-yl)amine; **CRA1000**, 2-(*N*-[2-methylthio-4-isopropylphenyl]-*N*-ethyl-amino-4-[4-(3-fluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl]-6-methylpyrimidine); **DMP696**, 4-(1,3-dimethoxyprop-2-ylamino)-2,7-dimethyl-8-(2,4-dichlorophenyl)pyrazolo[1,5-*a*]-1,3,5-triazine; **D-Phe-CRF-(12-41)**, D-Phe¹²,Nle^{21,38}, α MeLeu³⁷-CRF; **K31440**, Ac-(D-Tyr¹¹,His¹²,Nle¹⁷)sauvagine-(11-40); **K41498**, [D-Phe¹¹,His¹²,Nle¹⁷]sauvagine-(11-40); **NBI27914**, 2-methyl-4-(*N*-propyl-*N*-cyclopropanemethylamino)-5-chloro-6-(2,4,6-trichloroanilino)pyrimidine; **R121919**, 3-[6-(dimethylamino)-4-methyl-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[1,5-*a*]pyrimidin-7-amine; **SRA125543A**, 4-(2-chloro-4-methoxy-5-methylphenyl)-*N*-[(1*S*)-2-cyclopropyl-1-(3-fluoro-4-methylphenyl)ethyl]5-methyl-*N*-(2-propynyl)-1,3-thiazol-2-amine hydrochloride

Further Reading:

- BALE, T.L., GIORDANO, F.J. & VALE, W.W. (2003). A new role for corticotropin-releasing factor receptor-2: suppression of vascularization. *Trends Cardiovasc. Med.*, **13**, 68–71.
- BALE, T.L. & VALE, W.W. (2004). CRF and CRF receptors: role in stress responsivity and other behaviors. *Annu. Rev. Pharmacol. Toxicol.*, **44**, 525–557.
- COSTE, S.C., QUINTOS, R.F. & STENZEL-POORE, M.P. (2002). Corticotropin-releasing hormone-related peptides and receptors: emergent regulators of cardiovascular adaptations to stress. *Trends Cardiovasc. Med.*, **12**, 176–182.
- DAUTZENBERG, F.M. & HAUGER, R.L. (2002). The CRF peptide family and their receptors: yet more partners discovered. *Trends Pharmacol. Sci.*, **23**, 71–77.
- GILLIGAN, P.J. & LI, Y.W. (2004). Corticotropin-releasing factor antagonists: recent advances and exciting prospects for the treatment of human diseases. *Curr. Opin. Drug Discov. Devel.*, **7**, 487–497.
- GRAVANIS, A. & MARGIORIS, A.N. (2005). The corticotropin-releasing factor (CRF) family of neuropeptides in inflammation: potential therapeutic applications. *Curr. Med. Chem.*, **12**, 1503–1512.
- HAUGER, R.L., GRIGORIADIS, D.E., DALLMAN, M.F., PLOTSKY, P.M., VALE, W.W. & DAUTZENBERG, F.M. (2003). International Union of Pharmacology. XXXVI. Current status of the nomenclature for receptors for corticotropin-releasing factor and their ligands. *Pharmacol. Rev.*, **55**, 21–26.
- HEINRICHS, S.C. & KOOB, G.F. (2004). Corticotropin-releasing factor in brain: a role in activation, arousal, and affect regulation. *J. Pharmacol. Exp. Ther.*, **311**, 427–440.
- HILLHOUSE, E.W. & GRAMMATOPOULOS, D.K. (2006). The molecular mechanisms underlying the regulation of the biological activity of corticotropin-releasing hormone receptors: implications for physiology and pathophysiology. *Endocr. Rev.*, **27**, 260–286.
- KEHNE, J. & DE LOMBAERT, S. (2002). Non-peptidic CRF₁ receptor antagonists for the treatment of anxiety, depression and stress disorders. *Curr. Drug Target CNS. Neurol. Disord.*, **1**, 467–493.
- REUL, J.M. & HOLSBOER, F. (2002). Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression. *Curr. Opin. Pharmacol.*, **2**, 23–33.
- SARNYAI, Z., SHAHAM, Y. & HEINRICHS, S.C. (2001). The role of corticotropin-releasing factor in drug addiction. *Pharmacol. Rev.*, **53**, 209–244.
- SAUNDERS, J. & WILLIAMS, J. (2003). Antagonists of the corticotropin releasing factor receptor. *Prog. Med. Chem.*, **41**, 195–247.
- SMAGIN, G.N. & DUNN, A.J. (2000). The role of CRF receptor subtypes in stress-induced behavioural responses. *Eur. J. Pharmacol.*, **405**, 199–206.
- TAKAHASHI, L.K. (2001). Role of CRF₁ and CRF₂ receptors in fear and anxiety. *Neurosci. Biobehav. Rev.*, **25**, 627–636.
- TAO, J. & LI, S. (2005). Urocortin: a cardiac protective peptide? *Biochem. Biophys. Res. Commun.*, **332**, 923–926.
- THEOHARIDES, T.C., DONELAN, J.M., PAPADOPOULOU, N., CAO, J., KEMPURAJ, D. & CONTI, P. (2004). Mast cells as targets of corticotropin-releasing factor and related peptides. *Trends Pharmacol. Sci.*, **25**, 563–568.

References:

- CHAKI, S. *et al.* (1999). *Eur. J. Pharmacol.*, **371**, 205–211.
- CHEN, C. *et al.* (1996). *J. Med. Chem.*, **39**, 4358–4360.
- GULLY, D. *et al.* (2002). *J. Pharmacol. Exp. Ther.*, **301**, 322–332.
- HE, L. *et al.* (2000). *J. Med. Chem.*, **43**, 449–456.

- LAWRENCE, A.J. *et al.* (2002). *Br. J. Pharmacol.*, **136**, 896–904.
- LEWIS, K. *et al.* (2001). *Proc. Natl. Acad. Sci. USA*, **98**, 7570–7575.
- LUNDKVIST, J. *et al.* (1996). *Eur. J. Pharmacol.*, **309**, 195–200.
- PERKINS, A.V. *et al.* (1995). *J. Endocrinol.*, **146**, 395–401.
- REYES, T.M. *et al.* (2001). *Proc. Natl. Acad. Sci. USA*, **98**, 2843–2848.
- RUHMANN, A. *et al.* (1998). *Proc. Natl. Acad. Sci. USA*, **95**, 15264–15269.
- RUHMANN, A. *et al.* (2002). *Peptides*, **23**, 453–460.
- WEBSTER, E.L. *et al.* (1996). *Endocrinology*, **137**, 5747–5750.
- ZOBEL, A.W. *et al.* (2000). *J. Psychiatr. Res.*, **34**, 171–181.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Dopamine

Overview: Dopamine receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Dopamine Receptors, see Schwartz *et al.*, 1998) are commonly divided into D1-like (D1 and D5) and D2-like (D2, D3 and D4) families, where the endogenous agonist is dopamine.

Nomenclature	D1	D2	D3	D4	D5
Other names	D ₁ , D _{1A}	D ₂	D ₃	D ₄	D ₅ , D _{1B}
Ensembl ID	ENSG00000184845	ENSG00000149295	ENSG00000151577	ENSG00000069696	ENSG00000169676
Principal transduction	G _s , G _{olf}	G _{i/o}	G _{i/o}	G _{i/o}	G _s
Selective agonists	R(+)SKF81297, R(+)SKF38393	(+)PHNO	PD128907	PD168077	—
Selective antagonists	SCH23390, SKF83566, SCH39166	Raclopride, domperidone	S33084 (9.6, Millan <i>et al.</i> , 2000), nafadotride (9.5), (+)S14297 (8.7, Millan <i>et al.</i> , 1994), SB277011 (7.5, Reavill <i>et al.</i> , 2000)	L745870 (9.3), U101958 (8.9, Schlachter <i>et al.</i> , 1997), L741742 (8.5)	—
Probes	[³ H]-SCH23390 (0.2 nM), [¹²⁵ I]-SCH23982 (0.7 nM)	[³ H]-Raclopride, [³ H]-spiperone	[³ H]-7-OH-DPAT, [³ H]-PD128907, [³ H]-spiperone	[³ H]-NGD941 (5 nM, Primus <i>et al.</i> , 1997), [¹²⁵ I]-L750667 (1 nM, Patel <i>et al.</i> , 1996), [³ H]-spiperone	[¹²⁵ I]-SCH23982 (0.8 nM) [³ H]-SCH23390 (0.5 nM)

The selectivity of many of these agonists is less than two orders of magnitude. [³H]-Raclopride exhibits similar high affinity for D2 and D3 receptors (low affinity for D4), but has been used to label D2 receptors in the presence of a D3-selective antagonist. [³H]-7-OH-DPAT has similar affinity for D2 and D3 receptors, but labels only D3 receptors in the absence of divalent cations. The pharmacological profile of the D5 receptor is similar to, yet distinct from, that of the D1 receptor. The splice variants of the D2 receptor are commonly termed D2S and D2L (short and long). The *DRD4* gene is highly polymorphic in humans, with allelic variations of the protein from amino acid 387 to 515.

Abbreviations: **L741742**, 5-(4-chlorophenyl)-4-methyl-3-(1-[2-phenethyl]piperidin-4-yl)isoxazole; **L745870**, 3-[(4-(4-chlorophenyl)piperazin-1-yl)methyl]-1*H*-pyrrolo[2,3-*b*]pyridine; **L750667**, iodinated L745870; **NGD941**, 2-phenyl-4(*S*)-(4-[2-pyrimidinyl]-[piperazin-1-yl]-methyl)-imidazole dimaleate; (+)**7-OH-DPAT**, (+)-7-hydroxy-2-aminopropylaminotetralin; **PD128907**, *R*-(+)-*trans*-3,4,4*a*,10*b*-tetrahydro-4-propyl-2*H*,5*H*-[1]benzopyrano[4,3-*b*]-1,4-oxazine-9-ol; **PD168077**, *N*-methyl-4-(2-cyanophenyl)piperazinyl-3-methylbenzamide; (+)**PHNO**, 9-hydroxy-4-propyl-naphthoxazine; (+)**S14297**, (+)-7-(*N,N*-dipropylamino)-5,6,7,8-tetrahydronaphtho(2,3*b*)dihydro-2,3-furane; **S33084**, (3*aR*,9*bS*)-*N*[4-(8-cyano-1,3*a*,4,9*b*-tetrahydro-3*H*-benzopyrano[3,4-*c*]pyrrole-2-yl)-butyl] (4-phenyl)benzamide; **SB277011**, *trans*-*N*-(4-[2-{6-cyano-1,2,3,4-tetrahydroisoquinolin-2-yl}ethyl]cyclohexyl)-4-quinolininecarboxamide; **SCH23390**, 7-chloro-8-hydroxy-3-methyl-5-phenyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; **SCH23982**, 8-iodo-2,3,4,5-tetrahydro-3-methyl-5-phenyl-1*H*-3-benzazepine; **SCH39166**, (−)-*trans*-6,7*a*,8,9,13*b*-hexahydro-3-chloro-2-hydroxy-*N*-ethyl-5*H*-benzo[*d*]naphtho-(2,3*b*)azepine; **R(+)SKF38393**, *R*(+)-7,8-dihydroxy-1-phenyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; **R(+)SKF81297**, *R*(+)-6-chloro-7,8-dihydroxy-1-phenyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine; **SKF83566**, (−)-7-bromo-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-3-benzazepine; **U101958**, 3-isopropoxy-*N*-methyl-*N*-(1-[phenylmethyl]-4-piperidinyl)-2-pyridinylamine

Further Reading:

MISSALE, C., NASH, S.R., ROBINSON, S.W., JABER, M. & CARON, M.G. (1998). Dopamine receptors: from structure to function. *Physiol. Rev.*, **78**, 189–225.

NEVE, K.A. & NEVE, R.L. (eds) (1997). *The Dopamine Receptor*. Humana Press: Totowa, NJ.

SCHWARTZ, J.-C. *et al.* (1998). Dopamine receptors. In: *The IUPHAR Compendium of Receptor Characterization and Classification*, ed. Girdlestone, D. pp. 141–151. London: IUPHAR Media.

References:

MILLAN, M.J. *et al.* (1994). *Eur. J. Pharmacol.*, **260**, R3–R5.

MILLAN, M.J. *et al.* (2000). *J. Pharmacol. Exp. Ther.*, **293**, 1063–1073.

PRIMUS, R.J. *et al.* (1997). *J. Pharmacol. Exp. Ther.*, **282**, 1020–1027.

PATEL, S. *et al.* (1996). *Mol. Pharmacol.*, **50**, 2435–2437.

REAVILL, C. *et al.* (2000). *J. Pharmacol. Exp. Ther.*, **294**, 1154–1165.

SCHLACHTER, S.K. *et al.* (1997). *Eur. J. Pharmacol.*, **322**, 283–286.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Endothelin

Overview: Endothelin receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Endothelin Receptors, Davenport, 2002) are activated by the endogenous 21 amino-acid peptides endothelin-1 (ET-1, ENSG00000078401), ET-2 (ENSG00000127129) and ET-3 (ENSG00000124205). Nonselective peptide (e.g. TAK044, pA₂ 8.4) and nonpeptide (e.g. bosentan, pA₂ 6.0–7.2; SB209670, pA₂ 9.4) antagonists can block both ET_A and ET_B receptors.

Nomenclature	ET_A	ET_B
Ensemble ID	ENSG00000151617	ENSG00000136160
Principal transduction	G _{q/11} , G _s	G _{q/11} , G _{i/o}
Potency order	ET-1, ET-2 > ET-3 (Maguire & Davenport, 1995)	ET-1, ET-2, ET-3
Selective agonists	—	[Ala ^{1,3,11,15}]ET-1 (Molenaar <i>et al.</i> , 1992), sarafotoxin S6c (Russell & Davenport, 1996), IRL1620 (Watakabe <i>et al.</i> , 1992), BQ3020 (Russell & Davenport, 1996)
Selective antagonists	A127722 (9.2–10.5, Opgenorth <i>et al.</i> , 1996), LU135252 (8.9, Riechers <i>et al.</i> , 1996), SB234551 (8.7–9.0, Ohlstein <i>et al.</i> , 1998), PD156707 (8.2–8.5, Maguire <i>et al.</i> , 1997), FR139317 (7.3–7.9, Maguire & Davenport, 1995), BQ123 (6.9–7.4, Maguire & Davenport, 1995)	BQ788 (8.4, Russell & Davenport, 1996), A192621 (8.1, von Geldern <i>et al.</i> , 1999), IRL2500 (7.2, Russell & Davenport, 1996), Ro468443 (7.1, Breu <i>et al.</i> , 1996)
Probes	[³ H]-S0139 (0.6 nM), [³ H]-BQ123 (3.2 nM, Ihara <i>et al.</i> , 1995), [¹²⁵ I]-PD164333 (0.2 nM, Davenport <i>et al.</i> , 1998), [¹²⁵ I]-PD151242 (0.5 nM, Davenport <i>et al.</i> , 1994)	[¹²⁵ I]-IRL1620 (20 pM, Watakabe <i>et al.</i> , 1992), [¹²⁵ I]-BQ3020 (0.1 nM, Molenaar <i>et al.</i> , 1992), [¹²⁵ I]-[Ala ^{1,3,11,15}]ET-1 (0.2 nM, Molenaar <i>et al.</i> , 1992)

Subtypes of the ET_B receptor have been proposed, although gene disruption studies in mice suggest that the heterogeneity results from a single gene product (Mizuguchi *et al.*, 1997).

Abbreviations: **A127722**, *trans-trans*-2-(4-methoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-([N,N-dibutylamino]carbonylmethyl)pyrrolidine-3-carboxylate; **A192621**, (2R,3R,4S)-2-(4-propoxyphenyl)-4-(1,3-benzodioxol-5-yl)-1-(N-[2,6-diethylphenyl]acetamido)pyrrolidine-3-carboxylic acid; **BQ123**, *cyc*(DTrp-DAsp-Pro-D-Val-Leu); **BQ3020**, *N*-acetyl-Leu-Met-Asp-Lys-Glu-Ala-Val-Tyr-Phe-Ala-His-Leu-Asp-Ile-Ile-Trp; **BQ788**, *N*-*cis*-2,6-dimethylpiperidinocarbonyl-L-γ-methylleucyl-D-1-methoxycarbonyl-D-norleucine; **FR139317**, (R)-2-([R-2-((S)-2-([1-{hexahydro-1H-azepinyl}carbonyl]amino)methyl)pentanoyl]amino-3-(3-[methyl-1H-indodolyl]propionylamino-3-(2-pyridyl))propionate; **IRL1620**, Suc[Glu⁹,Ala^{11,15}]ET-1_{10–21}; **IRL2500**, *N*-(3,5-dimethylbenzoyl)-*N*-methyl-(D)-(4-phenylphenyl)-Ala-Trp; **LU135252**, (+)-(-)-S)-2-(4,6-dimethoxypyrimidin-2-yloxy)-3-methoxy-3,3-propionic acid; **PD151242**, (N-[[hexahydro-1-azepinyl]carbonyl]Leu(1-Me)-DTrp-DTyr; **PD156707**, 2-benzo[1,3]dioxol-5-yl-4-(4-methoxyphenyl)-4-oxo-3-(3,4,5-trimethoxybenzyl)-but-2-enoate; **PD164333**, 2-benzo[1,3]dioxol-5-yl-4-(3-[2-(4-hydroxyphenyl)-ethylcarbonyl]-propoxy)-4,5-dimethoxy-phenyl-3-(4-methoxy-benzoyl)-but-2-enoate; **RES7011**, *cyc*(Gly-Asn-Trp-His-Gly-Thr-Ala-Pro-Asp)-Trp-Phe-Phe-Asn-Tyr-Tyr-Trp; **Ro468443**, (R)-4-*tert*-butyl-N-(6-[2,3-dihydroxypropoxy]-5-[2-methoxyphenoxy]-2-[4-methoxyphenyl]-pyrimidin-4-yl)-benzenesulfonamide; **S0139**, 27-O-3-(2-[3-carboxyacryloylamino]-5-hydroxyphenyl)-acryloyloxymyricone, sodium salt; **SB209670**, (+)-1*S*,2*R*,3*S*-(2-carboxymethoxy-4-methoxyphenyl)-1-(3,4-methylenedioxypheyl)-5-prop-1-yloxyindane-2-carboxylate; **SB234551**, (*E*)-α-([1-butyl-5-{2-[(2-carboxyphenyl)methoxy]-4-methoxyphenyl}-1H-pyrazol-4-yl]methylene)-6-methoxy-1,3-benzodioxole-5-propanoic acid; **TAK044**, *cyc*(D-Asp-Asp(Ph)-Asp-D-Thg-Leu-D-Trp)-4-oxobut-2-enoate

Further reading:

- DAVENPORT, A.P. & RUSSELL, F.D. (2001). Endothelin converting enzymes and endothelin receptor localisation in human tissues. *Hdbk. Exp. Pharmacol.*, **152**, 209–237.
- DAVENPORT, A.P. (2002). International Union of Pharmacology. XXIX. Update on endothelin receptor nomenclature. *Pharmacol. Rev.*, **54**, 219–226.
- DAVENPORT, A.P. & MAGUIRE, J.J. (2006). Endothelin. *Handbook Exp. Pharmacol.* Heidelberg: Springer-Verlag, **176**, 295–329.
- D'ORLEANS-JUSTE, P., LABONTE, J., BKAILY, G., CHOUFANI, S., PLANTE, M. & HONORE, J. (2002). Function of the endothelin_B receptor in cardiovascular physiology and pathophysiology. *Pharmacol. Ther.*, **95**, 221–238.
- GALIE, N., MANES, A. & BRANZI, A. (2004). The endothelin system in pulmonary arterial hypertension. *Cardiovasc. Res.*, **61**, 227–237.
- HENRY, P.J. & GOLDIE, R.G. (2001). Endothelin receptors. *Hdbk. Exp. Pharmacol.*, **152**, 69–114.
- IQBAL, J., SANGHIA, R. & DAS, S.K. (2005). Endothelin receptor antagonists: an overview of their synthesis and structure–activity relationship. *Mini Rev. Med. Chem.*, **5**, 381–408.
- KEDZIERSKI, R.M. & YANAGISAWA, M. (2001). Endothelin system: the double-edged sword in health and disease. *Annu. Rev. Pharmacol. Toxicol.*, **41**, 851–876.
- MASAKI, T. (2004). Historical review: endothelin. *Trends Pharmacol. Sci.*, **25**, 219–224.
- NAICKER, S. & BHOOLA, K.D. (2001). Endothelins: vasoactive modulators of renal function in health and disease. *Pharmacol. Ther.*, **90**, 61–88.
- RASHID, A.J., O'DOWD, B.F. & GEORGE, S.R. (2004). Minireview: diversity and complexity of signaling through peptidergic G protein-coupled receptors. *Endocrinology*, **145**, 2645–2652.
- REMUZZI, G., PERICO, N. & BENIGNI, A. (2002). New therapies that antagonize endothelin: promises and frustrations. *Nat. Rev. Drug. Discov.*, **1**, 986–1001.
- SCHINELLI, S. (2006). Pharmacology and physiopathology of the brain endothelin system: an overview. *Curr. Med. Chem.*, **13**, 627–638.

References:

- BREU, V. *et al.* (1996). *FEBS Lett.*, **383**, 37–41.
- DAVENPORT, A.P. *et al.* (1998). *Br. J. Pharmacol.*, **123**, 223–230.
- DAVENPORT, A.P. *et al.* (1994). *Br. J. Pharmacol.*, **111**, 4–6.
- IHARA, M. *et al.* (1995). *Eur. J. Pharmacol.*, **274**, 1–6.
- MAGUIRE, J.J. & DAVENPORT, A.P. (1995). *Br. J. Pharmacol.*, **115**, 191–197.
- MAGUIRE, J.J. *et al.* (1997). *J. Pharmacol. Exp. Ther.*, **280**, 1102–1108.
- MIZUGUCHI, T. *et al.* (1997). *Br. J. Pharmacol.*, **120**, 1427–1430.
- MOLENAAR, P. *et al.* (1992). *Br. J. Pharmacol.*, **107**, 637–639.

- OHLSTEIN, E.H. *et al.* (1998). *J. Pharmacol. Exp. Ther.*, **286**, 650–656.
- OPGENORTH, T.J. *et al.* (1996). *J. Pharmacol. Exp. Ther.*, **276**, 473–481.
- RIECHERS, H. *et al.* (1996). *J. Med. Chem.*, **39**, 2123–2128.
- RUSSELL, F.D. & DAVENPORT, A.P. (1996). *Br. J. Pharmacol.*, **119**, 631–636.
- VON GELDERN, T.W. *et al.* (1999). *J. Med. Chem.*, **42**, 3668–3678.
- WATAKABE, T. *et al.* (1992). *Biochem. Biophys. Res. Commun.*, **185**, 867–873.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Free fatty acid

Overview: Free fatty acid receptors (FFA, provisional nomenclature) are activated by free fatty acids, such that long chain saturated and unsaturated fatty acids (C16:0, C18:0, C18:1, C18:2, C18:3, n-6, C20:4, C20:5, n-3, C22:6, n-3, Briscoe *et al.*, 2003; Itoh *et al.*, 2003; Kotarsky *et al.*, 2003) activate FFA₁ receptors, while short chain fatty acids (C4, C3 & C2) activate FFA₂ (Brown *et al.*, 2003; Le Poul *et al.*, 2003; Nilsson *et al.*, 2003) and FFA₃ (Brown *et al.*, 2003; Le Poul *et al.*, 2003) receptors. The thiazolidinedione PPAR γ antagonist rosiglitazone has also been suggested to activate FFA₁ (Kotarsky *et al.*, 2003).

Nomenclature	FFA₁	FFA₂	FFA₃
Other names	GPR40 (Sawzdargo <i>et al.</i> , 1997)	GPR43 (Sawzdargo <i>et al.</i> , 1997). LSSIG (Senga <i>et al.</i> , 2003)	GPR41 (Sawzdargo <i>et al.</i> , 1997)
Ensembl ID	ENSG00000126266	ENSG00000185897	ENSG00000126262
Principal transduction	G _{q/11} (Briscoe <i>et al.</i> , 2003; Itoh <i>et al.</i> , 2003; Kotarsky <i>et al.</i> , 2003)	G _{q/11} , G _{i/o} (Brown <i>et al.</i> , 2003; Le Poul <i>et al.</i> , 2003; Nilsson <i>et al.</i> , 2003)	G _{q/11} , G _{i/o} (Brown <i>et al.</i> , 2003; Le Poul <i>et al.</i> , 2003)
Selective agonists	Elaidic acid (Briscoe <i>et al.</i> , 2003; Itoh <i>et al.</i> , 2003)	—	—

Gpr120 (ENSG00000186188) has also been suggested to be a target for unsaturated long chain free fatty acids (Hirasawa *et al.*, 2005; Katsuma *et al.*, 2005). GPR42 (ENSG00000126251) is a further member of the family (ENSG00000003273) with as yet undefined characteristics (Brown *et al.*, 2003).

Abbreviations: **C16:0**, palmitic acid; **C18:0**, stearic acid; **C18:1**, oleic acid; **C18:2**, linoleic acid; **C18:3,n-6**, γ -linolenic acid; **C2**, acetic acid; **C20:4**, arachidonic acid; **C20:5,n-3**, eicosapenta(5z,8z,11z,14z,17z)enoic acid, EPA; **C22:6,n-3**, docosahexa(4z,7z,10z,13z,16z,19z)enoic acid, DHA; **C3**, propionic acid; **C4**, butyric acid

Further Reading:

BROWN, A.J., JUPE, S. & BRISCOE, C.P. (2005). A family of fatty acid binding receptors. *DNA Cell Biol.*, **24**, 54–61.

MILLIGAN, G., STODDART, L.J. & BROWN A.J. (2006). G protein-coupled receptors for free fatty acids. *Cell Signal.* **18**, 1360–1365.

References:

BRISCOE, C.P. *et al.* (2003). *J. Biol. Chem.*, **278**, 11303–11311.

BROWN, A.J. *et al.* (2003). *J. Biol. Chem.*, **278**, 11312–11319.

HIRASAWA, A. *et al.* (2005). *Nat. Med.*, **11**, 90–94.

KATSUMA, S. *et al.* (2005). *J. Biol. Chem.*, **280**, 19507–19515.

KOTARSKY, K. *et al.* (2003). *Biochem. Biophys. Res. Commun.*, **301**, 406–410.

LE POUL, E. *et al.* (2003). *J. Biol. Chem.*, **278**, 25481–25489.

NILSSON, N.E. *et al.* (2003). *Biochem. Biophys. Res. Commun.*, **303**, 1047–1052.

SAWZDARGO, M. *et al.* (1997). *Biochem. Biophys. Res. Commun.*, **239**, 543–547.

SENGA, T. *et al.* (2003). *Blood*, **101**, 1185–1187.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

GABA_B

Overview: Functional GABA_B receptors (nomenclature agreed by NC-IUPHAR Subcommittee on GABA_B receptors, Bowery *et al.*, 2002; see also Spedding *et al.*, 2002) are formed from the heterodimerization of two similar 7TM subunits termed GABA_{B1} and GABA_{B2} (Bowery *et al.*, 2002; Pin *et al.*, 2004). The GABA_{B1} subunit, when expressed alone, binds both antagonists and agonists, but the affinity of the latter is generally 10- to 100-fold less than for the native receptor. The GABA_{B1} subunit when expressed alone is not transported to the cell membrane and is nonfunctional. Coexpression of GABA_{B1} and GABA_{B2} subunits allows transport of GABA_{B1} to the cell surface and generates a functional receptor that can couple to signal transduction pathways such as high-voltage-activated Ca²⁺ channels (Ca_v2.1, Ca_v2.2), or inwardly rectifying potassium channels (Kir3) (Bowery & Enna, 2000; Bowery *et al.*, 2002; Bettler *et al.*, 2004). The GABA_{B1} subunit harbours the GABA (orthosteric)-binding site within an extracellular domain (ECD) venus flytrap module (VTM), whereas the GABA_{B2} subunit mediates G-protein coupled signalling (Bowery *et al.*, 2002; Pin *et al.*, 2004). The two subunits interact allosterically in that GABA_{B2} increases the affinity of GABA_{B1} for agonists and reciprocally GABA_{B1} facilitates the coupling of GABA_{B2} to G-proteins (Pin *et al.*, 2004; Kubo & Tateyama, 2005). GABA_{B1} and GABA_{B2} subunits assemble in a 1:1 stoichiometry by means of a coiled-coil interaction between α -helices within their carboxy-termini that masks an endoplasmic reticulum retention motif (RXRR) within the GABA_{B1} subunit but other domains of the proteins also contribute to their heteromerization (Bettler *et al.*, 2004; Pin *et al.*, 2004). Four isoforms of the human GABA_{B1} subunit have been cloned. The predominant GABA_{B1(a)} and GABA_{B1(b)} isoforms, which are most prevalent in neonatal and adult brain tissue, respectively, differ in their ECD sequences as a result of the use of alternative transcription initiation sites. Isoforms generated by alternative splicing are GABA_{B1(c)} that differs in the ECD, and GABA_{B1(e)}, which is a truncated protein that can heterodimerize with the GABA_{B2} subunit but does not constitute a functional receptor. Only the 1a and 1b variants are identified as components of native receptors (Bowery *et al.*, 2002). Additional GABA_{B1} subunit isoforms have been described in rodents (reviewed by Bettler *et al.*, 2004).

Nomenclature	GABA_{B1,2}
Ensembl ID	GABA _{B1} ENSG00000168760; GABA _{B2} ENSG00000136928
Principal transduction	G _{i/o}
Selective agonists	3-APPA (CGP27492, 5 nM), 3-APMPA (CGP35024, 16 nM), (R)(-)-baclofen (32 nM), CGP44532 (45 nM)
Selective antagonists	CGP62349 (2.0 nM), CGP55845 (6 nM), SCH50911 (3 μ M), 2-hydroxy-s(-)-saclofen (11 μ M), CGP35348 (27 μ M)
Positive allosteric modulators	CGP7930, GS39783 (see Bettler <i>et al.</i> , 2004)
Probes	[³ H]-(R)(-)-baclofen, [³ H]-CGP54626 (1.5 nM; Bittiger <i>et al.</i> , 1992), [³ H]-CGP62349 (0.9 nM, Kier <i>et al.</i> , 1999), [¹²⁵ I]-CGP64213 (1 nM, Galvez <i>et al.</i> , 2000), [¹²⁵ I]-CGP71872 (K _i = 0.5 nM, Belley <i>et al.</i> , 1999)

Potencies of agonists and antagonists listed in the table, quantified as IC₅₀ values for the inhibition of [³H]-CGP27492 binding to rat cerebral cortex membranes, are from Froestl & Mickel (1997 and Bowery *et al.* (2002). Radioligand K_D values relate to binding to rat brain membranes. CGP71872 is a photoaffinity ligand for the GABA_{B1} subunit (Belley *et al.*, 1999). In addition to the ligands listed in the table, Ca²⁺ binds to a site on the GABA_{B1} subunit to act as a positive allosteric modulator of GABA (Galvez *et al.*, 2000). In cerebellar Purkinje neurones, the interaction of Ca²⁺ with the GABA_B receptor enhances the activity of mGlu1, probably *via* a direct association between the two receptors (Tabata *et al.*, 2004). Synthetic positive allosteric modulators appear to act on the heptahelical domain of the GABA_{B2} subunit (Pin *et al.*, 2004; Dupuis *et al.*, 2006). Knockout of the GABA_{B1} subunit in C57B mice causes the development of severe tonic-clonic convulsions that prove fatal within a month of birth, whereas GABA_{B1}^{-/-} BALB/c mice, although also displaying spontaneous epileptiform activity, are viable. The phenotype of the latter animals additionally includes hyperalgesia, hyperlocomotion, memory impairment and behaviours indicative of anxiety (Enna & Bowery, 2004). A similar phenotype has been found for GABA_{B2}^{-/-} BALB/c mice (Gassmann *et al.*, 2004).

Abbreviations: 3-APMPA (CGP35024), 3-amino-propyl-(P-methyl)-phosphinic acid; 3-APPA (CGP27492), 3-amino-propyl-phosphinic acid; CGP7930, 2,6-Di-tert-butyl-4-(3-hydroxy-2,2-dimethyl-propyl)-phenol; CGP35348, p-(3-aminopropyl)-P-diethoxymethylphosphinic acid; CGP44532, 3-amino-2-hydroxypropylmethylphosphinic acid; CGP54626, [S-(R,R)]-[3-[[1-(3,4-dichlorophenyl)ethyl]amino]-2-hydroxypropyl](cyclohexylmethyl)phosphinic acid; CGP55845, 3-[1-(S)-(3,4-dichlorophenyl)-ethyl]amino-2(S)-hydroxypropyl-(P-benzyl)-phosphinic acid; CGP62349, [3-[1-R-[[3-(methoxyphenylmethyl)hydroxyphosphinyl]-2(5)-hydroxypropyl]amino]ethyl]-benzoic acid; CGP64213, [3-[1-(R)-[[3-5N-1-[2-[[3-iodo-4-hydroxyphenyl]ethyl]carboxamido]pentyl]hydroxyphosphinyl]-2(S)-hydroxypropyl]amino]ethyl-benzoic acid; CGP71872, 3-(1-(R)-(3-(5-(4-azido-2-hydroxy-5-iodobenzoylamino)pentyl)hydroxyphosphoryl)-2-(S)-hydroxypropylamino)ethyl)-benzoic acid; GS39783, N,N'-dicyclopentyl-2-methylsulfonyl-5-nitro-pyrimidine-4,6-diamine; SCH90511, (+)-(2S)-5,5-dimethyl-2-morpholineacetic acid

Further Reading:

- BETTLER, B., KAUPMANN, K., MOSBACHER, J. & GASSMANN, M. (2004). Molecular structure and physiological functions of GABA_B receptors. *Physiol. Rev.*, **84**, 835–867.
- BETTLER, B. & TIAO, J.Y. (2006). Molecular diversity, trafficking and subcellular localization of GABA_B receptors. *Pharmacol. Ther.*, **110**, 533–543.
- BOWERY, N.G. (2000). Pharmacology of GABA_B receptors. In: *Handbook of Experimental Pharmacology, Pharmacology of GABA and Glycine Neurotransmission*, (ed.) Möhler, H. pp. 150, 311–328. Springer: Berlin.
- BOWERY, N.G. (2006). GABA_B receptor: a site of therapeutic benefit. *Curr. Opin. Pharmacol.*, **6**, 37–43.
- BOWERY, N.G. & ENNA, S.J. (2000). Aminobutyric acid_B receptors: first of the functional metabotropic heterodimers. *J. Pharmacol. Exp. Ther.*, **292**, 2–7.
- BOWERY, N.G., BETTLER, B., FROESTL, W., GALLAGHER, J.P., MARSHALL, F., RAITERI, M., BONNER, T.I. & ENNA, S.J. (2002). International Union of Pharmacology XXXIII. Mammalian γ -aminobutyric acid_B receptors: structure and function. *Pharmacol. Rev.*, **54**, 247–264.
- CRYAN, J.F. & KAUPMANN, K. (2005). Don't worry 'B' happy!: a role for GABA_B receptors in anxiety and depression. *Trends Pharmacol. Sci.*, **26**, 36–43.
- ENNA, S.J. & BOWERY, N.G. (2004). GABA_B receptor alterations as indicators of physiological and pharmacological function. *Biochem. Pharmacol.*, **68**, 1541–1548.
- FROESTL, W. & MICKEL, S.W. (1997). Chemistry of GABA_B modulators. In: *The GABA Receptors*, Enna, S.J. & Bowery, N.G. (eds) pp. 271–296. Humana Press: Totowa.
- HAMMOND, D.L. (2001). GABA_B receptors: new tricks by an old dog. *Curr. Opin. Pharmacol.*, **1**, 26–30.
- KORNAU, H.C. (2006). GABA_B receptors and synaptic modulation. *Cell Tissue Res.*, **326**, 517–533.
- KUBO, Y. & TATEYAMA, M. (2005). Towards a view of functioning dimeric metabotropic receptors. *Curr. Opin. Neurobiol.*, **15**, 289–295.
- MARSHALL, F.H. (2005). Is the GABA B heterodimer a good drug target? *J. Mol. Neurosci.*, **26**, 169–176.
- PIN, J.-P., KNIAZEFF, J., BINET, V., LIU, J., MAUREL, D., GALVEZ, T., DUTHEY, B., HAVLICKOVA, M., BLAHOS, J., PREZEAU, L. & RONDARD, P. (2004). Activation mechanism of the heterodimeric GABA_B receptor. *Biochem. Pharmacol.*, **68**, 1565–1572.
- SPEDDING, M., BONNER, T.I. & WATSON, S.P. (2002). International Union of Pharmacology XXXI. Recommendations for the nomenclature of multimeric G protein-coupled receptors. *Pharmacol. Rev.*, **54**, 231–232.

References:

BELLEY, M. *et al.* (1999). *Bioorg. Med. Chem.*, **7**, 2697–2704.

BITTIGER, H. *et al.* (1992). *Pharmacol. Commun.*, **2**, 23.

DUPUIS, D.S. *et al.* (2006). *Mol. Pharmacol.*, **70**, 2027–2036.

GALVEZ, T. *et al.* (2000). *Mol. Pharmacol.*, **57**, 419–426.

GASSMANN, M. *et al.* (2004). *J. Neurosci.*, **42**, 6086–6097.

KIER, M.J. *et al.* (1999). *Brain Res. Mol. Brain Res.*, **71**, 279–289.

TABATA, T. *et al.* (2004). *Proc. Natl. Acad. Sci. USA*, **101**, 16952–16957.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Galanin

Overview: Galanin receptors (provisional nomenclature) are activated by the endogenous peptides galanin (ENSG00000069482) and galanin-like peptide (GALP, ENSG00000105099). Human galanin is a 30 aa non-amidated peptide (Evans and Shine, 1991); in other species, it is 29 aa long and C-terminally amidated. Amino acids 1–14 of galanin are highly conserved in mammals, birds, reptiles, amphibia and fish. Shorter peptide species (e.g. human galanin-1–19, (Bersani *et al.*, 1991a) and porcine galanin-5–29 (Sillard *et al.*, 1992)) and N-terminally extended forms (e.g. N-terminally seven and nine residue elongated forms of porcine galanin (Bersani *et al.*, 1991b; Sillard *et al.*, 1992)) have been reported.

Nomenclature	GAL1	GAL2	GAL3
Other names	Galanin-1 receptor, GalR1, GALR1	Galanin-2 receptor, GalR2, GALR2	Galanin-3 receptor, GalR3, GALR3
Ensembl ID	ENSG00000166573	ENSG00000182687	ENSG00000128310
Principal transduction	G _{i/o}	G _{i/o} , G _{q/11}	G _{i/o}
Rank order of potency	Galanin > GALP (Ohtaki <i>et al.</i> , 1999)	GALP > galanin (Ohtaki <i>et al.</i> , 1999)	—
Selective agonists	—	Galanin-(2–29) (Fathi <i>et al.</i> , 1997; Wang <i>et al.</i> , 1997), D-Trp ² -galanin-(1–29) (Smith <i>et al.</i> , 1997)	—
Selective antagonists	2,3-Dihydro-dithiin and -dithiepine-1, 1,4,4-tetroxides (Scott <i>et al.</i> , 2000)	—	—

Galanin-(1–11) is a high-affinity agonist at GAL1/GAL2 (pK_i 9) and galanin-(2–11) is selective for GAL2 (pK_i 8.7) compared to GAL1 (pK_i 6.1; Liu *et al.*, 2001). The affinity of GALP, galanin-(1–11) and (2–11) at GAL3 has not been assessed. [¹²⁵I]-[Tyr²⁶]galanin binds to all three subtypes with K_d values ranging from 0.05 to 1 nM (Skofitsch *et al.*, 1986; Smith *et al.*, 1997; 1998; Wang *et al.*, 1997; Fitzgerald *et al.*, 1998). Porcine galanin-(3–29) does not bind to cloned GAL1, GAL2 or GAL3 receptors, but a receptor that is functionally activated by porcine galanin-(3–29) has been reported in pituitary and gastric smooth muscle cells (Wynick *et al.*, 1993; Gu *et al.*, 1995). Additional galanin receptor subtypes are also suggested from studies with chimeric peptides (e.g. M15, M35 and M40), which act as antagonists in functional assays in the cardiovascular system (Ulman *et al.*, 1993), spinal cord (Wiesenfeld-Hallin *et al.*, 1992), locus coeruleus, hippocampus (Bartfai *et al.*, 1991) and hypothalamus (Leibowitz and Kim, 1992; Bartfai *et al.*, 1993), but exhibit agonist activity at some peripheral sites (Bartfai *et al.*, 1993; Gu *et al.*, 1995). The chimeric peptides M15, M32, M35, M40 and C7 are agonists at GAL1 receptors expressed endogenously in Bowes human melanoma cells (Ohtaki *et al.*, 1999), and at heterologously expressed recombinant GAL1, GAL2 and GAL3 receptors (Smith *et al.*, 1997; Fitzgerald *et al.*, 1998; Smith *et al.*, 1998).

Abbreviations: **C7**, galanin-(1–13)-spantide; **M15**, galanin-(1–13)-substance P-5–11 amide, also known as galantide; **M32**, galanin-(1–13)-neuropeptide Y amide-(25–36) amide; **M35**, galanin-(1–13)-bradykinin-(2–9) amide; **M40**, galanin-(1–13)-Pro-Pro-Ala-Leu-Ala-Leu-Ala-Leu-Ala amide

Further Reading:

BRANCHEK, T.A., SMITH, K.E., GERALD, C. & WALKER, M.W. (2000). Galanin receptor subtypes. *Trends Pharmacol. Sci.*, **21**, 109–116.

GOTTSCH, M.L., CLIFTON, D.K. & STEINER, R.A. (2004). Galanin-like peptide as a link in the integration of metabolism and reproduction. *Trends Endocrinol. Metab.*, **15**, 215–221.

GUNDLACH, A.L. (2002). Galanin/GALP and galanin receptors: role in central control of feeding, body weight/obesity and reproduction? *Eur. J. Pharmacol.*, **440**, 255–268.

HOLMES, A., HEILIG, M., RUPNIAK, N.M., STECKLER, T. & GRIEBEL, G. (2003). Neuropeptide systems as novel therapeutic targets for depression and anxiety disorders. *Trends Pharmacol. Sci.*, **24**, 580–588.

IISMAA, T.P. & SHINE, J. (1999). Galanin and galanin receptors. *Results Probl. Cell Differ.*, **26**, 257–291.

JACOBOWITZ, D.M., KRESSE, A. & SKOFITSCH, G. (2004). Galanin in the brain: chemoarchitectonics and brain cartography – a historical review. *Peptides*, **25**, 433–464.

LIU, H.X. & HOKFELT, T. (2002). The participation of galanin in pain processing at the spinal level. *Trends Pharmacol. Sci.*, **23**, 468–474.

LU, X., LUNDSTROM, L., LANGE, U. & BARTFAI, T. (2005). Galanin receptor ligands. *Neuropeptides*, **39**, 143–146.

UBINK, R., CALZA, L. & HOKFELT, T. (2003). 'Neuro'-peptides in glia: focus on NPY and galanin. *Trends Neurosci.*, **26**, 604–609.

WYNICK, D., THOMPSON, S.W. & MCMAHON, S.B. (2001). The role of galanin as a multi-functional neuropeptide in the nervous system. *Curr. Opin. Pharmacol.*, **1**, 73–77.

References:

- BARTFAI, T. *et al.* (1991). *Proc. Natl. Acad. Sci. U.S.A.*, **88**, 10961–10965.
- BARTFAI, T. *et al.* (1993). *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 11287–11291.
- BERSANI, M. *et al.* (1991a). *FEBS Lett.*, **283**, 189–194.
- BERSANI, M. *et al.* (1991b). *Endocrinology*, **129**, 2693–2698.
- EVANS, H.F. & SHINE, J. (1991). *Endocrinology*, **129**, 1682–1684.
- FATHI, Z. *et al.* (1997). *Mol. Brain Res.*, **51**, 49–59.
- FITZGERALD, L.W. *et al.* (1998). *J. Pharmacol. Exp. Ther.*, **287**, 448–456.
- GU, Z.F. *et al.* (1995). *J. Pharmacol. Exp. Ther.*, **272**, 371–378.
- LEIBOWITZ, S.F. & KIM, T. (1992). *Brain Res.*, **599**, 148–152.
- LIU, H.X. *et al.* (2001). *Proc. Natl. Acad. Sci. U.S.A.*, **98**, 9960–9964.
- OHTAKI, T. *et al.* (1999). *J. Biol. Chem.*, **274**, 37041–37045.
- SCOTT, M.K. *et al.* (2000). *Bioorg. Med. Chem.*, **8**, 1383–1391.
- SILLARD, R. *et al.* (1992). *Peptides*, **13**, 1055–1060.
- SKOFITSCH, G. *et al.* (1986). *Peptides*, **7**, 1029–1042.
- SMITH, K.E. *et al.* (1997). *J. Biol. Chem.*, **272**, 24612–24616.
- SMITH, K.E. *et al.* (1998). *J. Biol. Chem.*, **273**, 23321–23326.
- ULMAN, L.G. *et al.* (1993). *J. Physiol.*, **464**, 491–499.
- WANG, S. *et al.* (1997). *Mol. Pharmacol.*, **52**, 337–343.
- WIESENFELD-HALLIN, Z. *et al.* (1992). *Proc. Natl. Acad. Sci. U.S.A.*, **89**, 3334–3337.
- WYNICK, D. *et al.* (1993). *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 4231–4235.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Ghrelin

Overview: Ghrelin receptors (see Davenport *et al.*, 2005) are activated by a 28 amino-acid peptide originally isolated from rat stomach, where it is cleaved from a 117 amino-acid precursor (ENSG00000157017). The human gene encoding the precursor peptide has 83% sequence homology to rat prepro-ghrelin, although the mature peptides from rat and human differ by only two amino acids (Matsumoto *et al.*, 2001). Alternative splicing results in the formation of a second peptide, des-Gln¹⁴-ghrelin with equipotent biological activity (Hosoda *et al.*, 2000). A unique post-translational modification (octanoylation of Ser³) occurs in both peptides, essential for full activity in binding to the ghrelin receptors in the hypothalamus and pituitary; and the release of growth hormone release from the pituitary (Kojima *et al.*, 1999). Structure activity studies showed the first five N-terminal amino acids to be the minimum required for binding (Bednarek *et al.*, 2000).

Nomenclature	Ghrelin
Other names	GHS-R1a (Growth hormone secretagogue receptor type 1), growth hormone-releasing peptide receptor
Ensembl ID	ENSG00000121853
Principal transduction	G _{q/11}
Rank order of potency	Ghrelin = des-Gln-ghrelin (Matsumoto <i>et al.</i> , 2001; Bedendi <i>et al.</i> , 2003)
Probes	[¹²⁵ I-His ⁹]-ghrelin (0.4 nM, Katugampola <i>et al.</i> , 2001), [¹²⁵ I-Tyr ⁴]-ghrelin (0.5 nM, Bedendi <i>et al.</i> , 2003), [¹²⁵ I]-Tyr ⁴ -des-octanoyl (0.7 nM, Bedendi <i>et al.</i> , 2003)

Des-octanoyl ghrelin has recently been shown to bind (as [¹²⁵I]-Tyr⁴-des-octanoyl ghrelin) and have effects in the cardiovascular system (Bedendi *et al.*, 2003), which raises the possible existence of different receptor subtypes in peripheral tissues and the central nervous system. One study has reported constitutive activity of the ghrelin receptor and has identified a potent inverse agonist ([D-Arg¹, D-Phe⁵, D-Trp^{7,9}, Leu¹¹]-substance P, EC₅₀ 5.2 nM; Holst *et al.*, 2003).

Further Reading:

- COWLEY, M.A. & GROVE, K.L. (2004). Ghrelin – satisfying a hunger for the mechanism. *Endocrinology*, **145**, 2604–2606.
- CUMMINGS, D.E. (2006). Ghrelin and the short- and long-term regulation of appetite and body weight. *Physiol. Behav.*, **89**, 71–84.
- DAVENPORT, A.P. (2003). Peptide and trace amine orphan receptors: prospects for new therapeutic targets. *Curr. Opin. Pharmacol.*, **3**, 127–134.
- DAVENPORT, A.P., BONNER, T.L., FOORD, S.M., HARMAR, A.J., NEUBIG, R.R., PIN, J.-P., SPEDDING, M., KOJIMA, M. & KANGAWA, K. (2005). International Union of Basic and Clinical Pharmacology. XLVII. Ghrelin receptor nomenclature, distribution and function. *Pharmacol. Rev.*, **57**, 541–546.
- HOLST, B. & SCHWARTZ, T.W. (2004). Constitutive ghrelin receptor activity as a signaling set-point in appetite regulation. *Trends Pharmacol. Sci.*, **25**, 113–117.
- HORVATH, T.L., CASTANEDA, T., TANG-CHRISTENSEN, M., PAGOTTO, U. & TSCHOP, M.H. (2003). Ghrelin as a potential anti-obesity target. *Curr. Pharm. Des.*, **9**, 1383–1395.
- HORVATH, T.L., DIANO, S., SOTONYI, P., HEIMAN, M. & TSCHOP, M. (2001). Minireview: ghrelin and the regulation of energy balance – a hypothalamic perspective. *Endocrinology*, **142**, 4163–4169.
- HOSODA, M., KOJIMA, M. & KANAGAWA, K. (2006). Biological, physiological and pharmacological aspects of ghrelin. *J. Pharmacol., Sci.*, **100**, 398–410.
- INUI, A. (2001). Ghrelin: an orexigenic and somatotrophic signal from the stomach. *Nat. Rev. Neurosci.*, **2**, 551–560.
- INUI, A., ASAKAWA, A., BOWERS, C.Y., MANTOVANI, G., LAVIANO, A., MEGUID, M.M. & FUJIMIYA, M. (2004). Ghrelin, appetite, and gastric motility: the emerging role of the stomach as an endocrine organ. *FASEB J.*, **18**, 439–456.
- KOJIMA, M., HOSODA, H., MATSUO, H. & KANGAWA, K. (2001). Ghrelin: discovery of the natural endogenous ligand for the growth hormone secretagogue receptor. *Trends Endocrinol. Metab.*, **12**, 118–122.
- KOJIMA, M. & KANAGAWA, K. (2005). Ghrelin: structure and function. *Physiol. Rev.*, **85**, 495–522.
- KORBONITS, M., GOLDSTONE, A.P., GUEORGUIEV, M. & GROSSMAN, A.B. (2004). Ghrelin – a hormone with multiple functions. *Front. Neuroendocrinol.*, **25**, 27–68.
- MAGUIRE, J.J. & DAVENPORT, A.P. (2005). Regulation of vascular reactivity by established and emerging GPCRs. *Trends Pharmacol. Sci.*, **26**, 448–454.
- NAGAYA, N. & KANGAWA, K. (2003). Ghrelin improves left ventricular dysfunction and cardiac cachexia in heart failure. *Curr. Opin. Pharmacol.*, **3**, 146–151.
- VAN DER LELY, A.J., TSCHOP, M., HEIMAN, M.L. & GHIGO, E. (2004). Biological, physiological, pathophysiological, and pharmacological aspects of ghrelin. *Endocr. Rev.*, **25**, 426–457.

References:

- BEDENDI, I. *et al.* (2003). *Eur. J. Pharmacol.*, **476**, 87–95.
- BEDNAREK, M.A. *et al.* (2000). *J. Med. Chem.*, **43**, 4370–4376.
- HOLST, B. *et al.* (2003). *Mol. Endocrinol.*, **17**, 2201–2210.
- HOSODA, H. *et al.* (2000). *J. Biol. Chem.*, **275**, 21995–22000.
- KATUGAMPOLA, S.D. *et al.* (2001). *Br. J. Pharmacol.*, **134**, 143–149.
- KOJIMA, M. *et al.* (1999). *Nature*, **402**, 656–660.
- MATSUMOTO, M. *et al.* (2001). *Biochem. Biophys. Res. Commun.*, **287**, 142–146.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Glucagon, glucagon-like peptide and secretin

Overview: The glucagon family of receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on the Glucagon receptor family, see Mayo *et al.*, 2003) are activated by the endogenous peptide (27–44 aa) hormones glucagon, glucagon-like peptide 1 (GLP-1), glucagon-like peptide 2 (GLP-2), glucose-dependent insulinotropic polypeptide (also known as gastric inhibitory polypeptide or GIP, ENSG00000159224), growth hormone-releasing hormone (GHRH, ENSG00000118702) and secretin (ENSG00000070031). One common precursor (ENSG00000115263) generates glucagon, GLP-1 and GLP-2 peptides (Irwin, 2001).

Nomenclature	Glucagon	GLP-1	GLP-2
Ensembl ID	ENSG00000141558	ENSG00000112164	ENSG00000065325
Principal transduction	G _s	G _s	G _s
Selective agonists	Glucagon	GLP-1-(7-37) (Dillon <i>et al.</i> , 1993); GLP-1-(7-36)amide (Thorens <i>et al.</i> , 1993), exendin-3 (Raufman <i>et al.</i> , 1991), exendin-4 (Thorens <i>et al.</i> , 1993)	GLP-2
Selective antagonists	L168049 (Cascieri <i>et al.</i> , 1999), des-His ¹ -[Glu ⁹]glucagon amide (Post <i>et al.</i> , 1993), BAY27-9955 (Petersen and Sullivan, 2001), xNNC92-1687 (Madsen <i>et al.</i> , 1998)	Exendin-(9-39) (Thorens <i>et al.</i> , 1993); T0632 (Tibaduiza <i>et al.</i> , 2001)	—
Probes	[¹²⁵ I]-glucagon	[¹²⁵ I]-GLP-1-(7-36) amide, [¹²⁵ I]-exendin, [¹²⁵ I]-exendin-(9-39), [¹²⁵ I]-GLP-1-(7-37)	—

Nomenclature	GIP	GHRH	Secretin
Ensembl ID	ENSG00000010310	ENSG00000106128	ENSG00000080293
Principal transduction	G _s	G _s	G _s
Selective agonists	GIP	BIM28011 (Coy <i>et al.</i> , 1996)	Secretin
Selective antagonists	—	JV-1-36 (Schally and Varga, 1999), JV-1-38 (Schally and Varga, 1999)	[(CH ₂ NH) ^{4,5}]secretin (Kim <i>et al.</i> , 1993)
Probes	[¹²⁵ I]-GIP	[¹²⁵ I]-GHRH	[¹²⁵ I]-(Tyr ¹⁰)secretin

The glucagon receptor has been reported to interact with receptor activity modifying proteins (RAMPs), specifically RAMP2, in heterologous expression systems (Christopoulos *et al.*, 2003), although the physiological significance of this has yet to be established.

Abbreviations: **BAY27-9955**, (+)-3,5-diisopropyl-2-(1-hydroxyethyl)-6-propyl-4'-fluoro-1,1'-biphenyl; **BIM28011**, [D-Ala²,Ala^{8,9,15,27},D-Arg²⁹]hGHRH-(1–29)NH₂; **JV-1-36**, [PhAc-Tyr¹,D-Arg²,Phe(4-Cl)⁶,Arg⁹,Abu¹⁵,Nle²⁷,D-Arg²⁸,Har²⁹]hGHRH(1–29)NH₂; **JV-1-38**, [PhAc-Tyr¹,D-Arg²,Phe(4-Cl)⁶,Har⁹,Tyr(Me)¹⁰,Abu¹⁵,Nle²⁷,D-Arg²⁸,Har²⁹]hGHRH(1–29)NH₂; **L168049**, 2-(4-pyridyl)-5-(4-chlorophenyl)-3-(5-bromo-2-propyloxyphenyl)pyrrole; **NNC92-1687**, 2-(benzimidazol-2-ylthio)-1-(3,4-dihydroxyphenyl)-1-ethanone; **T0632**, sodium (S)-3-(1-[2-fluorophenyl]-2,3-dihydro-3-{[3-isoquinolyl]-carbonyl}amino-6-methoxy-2-oxo-1H-indole)propanoate

Further Reading:

- BURCELIN, R. (2005). The incretins: a link between nutrients and well-being. *Br. J. Nutr.*, **93** (Suppl 1), S147–S156.
- D'ALESSIO, D.A. & VAHL, T.P. (2004). Glucagon-like peptide 1: evolution of an incretin into a treatment for diabetes. *Am. J. Physiol. Endocrinol. Metab.*, **286**, E882–E890.
- DOYLE, M.E. & EGAN, J.M. (2001). Glucagon-like peptide-1. *Recent Prog. Hormone Res.*, **56**, 377–399.
- DRUCKER, D.J. (2001a). Glucagon-like peptide 2. *J. Clin. Endocrinol. Metab.*, **86**, 1759–1764.
- DRUCKER, D.J. (2001b). Minireview: the glucagon-like peptides. *Endocrinology*, **142**, 521–527.
- ESTALL, J.L. & DRUCKER, D.J. (2006). Glucagon and glucagon-like peptide receptors as drug targets. *Curr. Pharm. Des.*, **12**, 1731–1750.
- GAULT, V.A., O'HARTE, F.P. & FLATT, P.R. (2003). Glucose-dependent insulinotropic polypeptide (GIP): anti-diabetic and anti-obesity potential? *Neuropeptides*, **37**, 253–263.
- GROMADA, J., BROCK, B., SCHMITZ, O. & RORSMAN, P. (2004). Glucagon-like peptide-1: regulation of insulin secretion and therapeutic potential. *Basic Clin. Pharmacol. Toxicol.*, **95**, 252–262.
- GUTZWILLER, J.P., DEGEN, L., HEUSS, L. & BEGLINGER, C. (2004). Glucagon-like peptide 1 (GLP-1) and eating. *Physiol. Behav.*, **82**, 17–19.
- HOLST, J.J. (2006). Glucagon-like peptide-1: from extract to agent. the Claude Bernard Lecture, 2005. *Diabetologia*, **49**, 253–260.
- KELLER, G., SCHALLY, A.V., GROOT, K., TOLLER, G.L., HAVT, A., KOSTER, F., ARMATIS, P., HALMOS, G., ZARANDI, M., VARGA, J.L. & ENGEL, J.B. (2005). Effective treatment of experimental human non-Hodgkin's lymphomas with antagonists of growth hormone-releasing hormone. *Proc. Natl. Acad. Sci. U.S.A.*, **102**, 10628–10633.
- KNUDSEN, L.B. (2004). Glucagon-like peptide-1: the basis of a new class of treatment for type 2 diabetes. *J. Med. Chem.*, **47**, 4128–4134.
- LETSCH, M., SCHALLY, A.V., BUSTO, R., BAJO, A.M. & VARGA, J.L. (2003). Growth hormone-releasing hormone (GHRH) antagonists inhibit the proliferation of androgen-dependent and -independent prostate cancers. *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 1250–1255.
- LIST, J.F. & HABENER, J.F. (2004). Glucagon-like peptide 1 agonists and the development and growth of pancreatic β -cells. *Am. J. Physiol. Endocrinol. Metab.*, **286**, E875–E881.
- MAYO, K.E., MILLER, L.J., BATAILLE, D., DALLE, S., GOKE, B., THORENS, B. & DRUCKER, D.J. (2003). International Union of Pharmacology. XXXV. The glucagon receptor family. *Pharmacol. Rev.*, **55**, 167–194.
- MEIER, J.J., GALLWITZ, B., SCHMIDT, W.E. & NAUCK, M.A. (2002). Glucagon-like peptide 1 as a regulator of food intake and body weight: therapeutic perspectives. *Eur. J. Pharmacol.*, **440**, 269–279.
- NIELSEN, L.L., YOUNG, A.A. & PARKES, D.G. (2004). Pharmacology of exenatide (synthetic exendin-4): a potential therapeutic for improved glycemic control of type 2 diabetes. *Regul. Peptides*, **117**, 77–88.
- STANLEY, S., WYNNE, K. & BLOOM, S. (2004). Gastrointestinal satiety signals III. Glucagon-like peptide 1, oxyntomodulin, peptide YY, and pancreatic polypeptide. *Am. J. Physiol. Gastrointest. Liver Physiol.*, **286**, G693–G697.

URUSOVA, I.A., FARILLA, L., HUI, H., D'AMICO, E. & PERFETTI, R. (2004). GLP-1 inhibition of pancreatic islet cell apoptosis. *Trends Endocrinol. Metab.*, **15**, 27–33.

References:

CASCIERI, M.A. *et al.* (1999). *J. Biol. Chem.*, **274**, 8694–8697.

CHRISTOPOULOS, A. *et al.* (2003). *J. Biol. Chem.*, **278**, 3293–3297.

COY, D.H. *et al.* (1996). *Ann. NY Acad. Sci.*, **805**, 149–158.

DILLON, J.S. *et al.* (1993). *Endocrinology*, **133**, 1907–1910.

IRWIN, D.M. (2001). *Regul. Peptides*, **98**, 1–12.

KIM, C.D. *et al.* (1993). *Am. J. Physiol. Gastrointest. Liver Physiol.*, **265**, G805–G810.

MADSEN, P. *et al.* (1998). *J. Med. Chem.*, **41**, 5150–5157.

PETERSEN, K.F. & SULLIVAN, J.T. (2001). *Diabetologia*, **44**, 2018–2024.

POST, S.R. *et al.* (1993). *Proc. Natl. Acad. Sci. U.S.A.*, **90**, 1662–1666.

RAUFMAN, J.P. *et al.* (1991). *J. Biol. Chem.*, **266**, 2897–2902.

SCHALLY, A.V. & VARGA, J.L. (1999). *Trends Endocrinol. Metab.*, **10**, 383–391.

THORENS, B. *et al.* (1993). *Diabetes*, **42**, 1678–1682.

TIBADUIZA, E.C. *et al.* (2001). *J. Biol. Chem.*, **276**, 37787–37793.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Glutamate, metabotropic

Overview: Metabotropic glutamate (mGlu) receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Metabotropic Glutamate Receptors, Schoepp *et al.*, 2000) are activated by the endogenous ligands L-glutamate, L-aspartate, L-serine-*O*-phosphate (LSOP), *N*-acetylaspartylglutamate (NAAG) and L-cysteine sulphonic acid. Examples of agonists selective for mGlu receptors compared with ionotropic glutamate receptors are 1*S*,3*R*-ACPD and L-CCG-I, which show limited selectivity for Group II receptors. An example of an antagonist selective for mGlu receptors is LY341495, which blocks mGlu₂ and mGlu₃ at low nanomolar concentrations, mGlu₈ at high nanomolar concentrations, and mGlu₁, mGlu₄, mGlu₅ and mGlu₇ in the micromolar range (Kingston *et al.*, 1998). Currently, three groups of native receptors are distinguishable on the bases of similarities of agonist pharmacology, primary sequence and G-protein effector coupling: Group I (mGlu₁ and mGlu₃); Group II (mGlu₂ and mGlu₃) and Group III (mGlu₄, mGlu₆, mGlu₇ and mGlu₈) (see Further reading). Group I mGlu receptors may be activated by DHPG and 3HPG (Brabet *et al.*, 1995), and antagonized by LY393675 (Baker *et al.*, 1998). Group II mGlu receptors may be activated by LY389795 (Monn *et al.*, 1999), LY379268 (Monn *et al.*, 1999), LY354740 (Schoepp *et al.*, 1997; Wu *et al.*, 1998), DCG-IV and 2*R*,4*R*-APDC (Schoepp *et al.*, 1996), and antagonised by EGLU (4.3, Jane *et al.*, 1996) and LY307452 (Escribano *et al.*, 1998; Wermuth *et al.*, 1996). Group III mGlu receptors may be activated by (*RS*)PPG (Gasparini *et al.*, 1999a).

In addition to orthosteric ligands that directly interact with the glutamate recognition site, allosteric modulators have been described. Negative allosteric modulators are listed separately. The positive allosteric modulators most often act as 'potentiators' of an orthosteric agonist response, without significantly activating the receptor in the absence of agonist. Examples of these unique pharmacological agents have been described for mGlu₁, mGlu₂, mGlu₄ and mGlu₅.

Nomenclature	mGlu ₁	mGlu ₂	mGlu ₃	mGlu ₄
Other names	mGluR ₁	mGluR ₂	mGluR ₃	mGluR ₄
Ensembl ID	ENSG00000152822	ENSG00000164082	ENSG00000105781	ENSG00000124493
Principal transduction	G _{q/11}	G _{i/o}	G _{i/o}	G _{i/o}
Selective agonists	—	—	NAAG (Wroblewska <i>et al.</i> , 1997)	L-AP4, LSOP (Wu <i>et al.</i> , 1998), 1997)
Selective positive allosteric modulators	Ro01-6128, Ro67-4853, Ro67-7476 (Knoflach <i>et al.</i> , 2001)	LY487379 (Johnson <i>et al.</i> , 2003), CBIPES (Johnson <i>et al.</i> , 2005)	—	(-)-PHCCC (Maj <i>et al.</i> , 2003), SIB1893, MPEP (Mathiesen <i>et al.</i> , 2003)
Selective competitive antagonists	3-MATIDA (Moroni <i>et al.</i> , 2002), AIDA (Moroni <i>et al.</i> , 1997), (<i>S</i>)-(+)-CBPG (Mannaioni <i>et al.</i> , 1999), LY367385 (Clark <i>et al.</i> , 1997), (<i>S</i>)-TBPG (Constantino <i>et al.</i> , 2001)	PCCG-4 (Pellicciari <i>et al.</i> , 1996)	—	MAP4
Selective negative allosteric modulators	CPCCOEt (Litschig <i>et al.</i> , 1999), BAY36-7620 (Carroll <i>et al.</i> , 2001), LY456236 (Li <i>et al.</i> , 2002), 3,5-DMPPP (Micheli <i>et al.</i> , 2003), EM-TBPC (Malherbe <i>et al.</i> , 2003), JNJ16259685 (Lavreysen <i>et al.</i> , 2004)	—	—	—

Nomenclature	mGlu ₅	mGlu ₆	mGlu ₇	mGlu ₈
Other names	mGluR ₅	mGluR ₆	mGluR ₇	mGluR ₈
Ensembl ID	ENSG00000168959	ENSG00000113262	ENSG00000168160	ENSG00000179603
Principal transduction	G _{q/11}	G _{i/o}	G _{i/o}	G _{i/o}
Selective agonists	CHPG (Doherty <i>et al.</i> , 1997), (<i>S</i>)-(+)-CBPG (Mannaioni <i>et al.</i> , 1999)	Homo-AMPA (Bräuner-Osborne <i>et al.</i> , 1996), 1-benzyl-APDC (Tuckmantel <i>et al.</i> , 1997)	LSOP (Wu <i>et al.</i> , 1998), L-AP4	LSOP (Wu <i>et al.</i> , 1998), L-AP4, (<i>S</i>)-3,4-DCPG (Thomas <i>et al.</i> , 2001)
Selective positive allosteric modulators	DFB (O'Brien <i>et al.</i> , 2003), CPPHA (O'Brien <i>et al.</i> , 2004), CDPPB (Kinney <i>et al.</i> , 2005)	—	AMN082 (Flor <i>et al.</i> , 2005)	—
Selective competitive antagonists	ACDPP (6.5, Bonnefous <i>et al.</i> , 2005)	MAP4, THPG (Thoreson <i>et al.</i> , 1997)	—	MPPG (Wu <i>et al.</i> , 1998)
Selective negative allosteric modulators	SIB1757 (Varney <i>et al.</i> , 1999), SIB1893 (Varney <i>et al.</i> , 1999), MPEP (Gasparini <i>et al.</i> , 1999b), MTEP (Brodtkin <i>et al.</i> , 2002), fenobam (Porter <i>et al.</i> , 2005), YM298198 (Kohara <i>et al.</i> , 2005)	—	—	—

Radioligand binding using a variety of radioligands has been conducted on recombinant receptors (for example, [³H]-R214127 (Lavreysen *et al.*, 2003) and [³H]-YM298198 (Kohara *et al.*, 2005) at mGlu₁ receptors and [³H]-methoxyMPEP (Gasparini *et al.*, 2002) and [³H]-methoxymethyl-MTEP (Anderson *et al.*, 2002) at mGlu₃ receptors. Although a number of radioligands have been used to examine binding using native tissues, correlation with individual subtypes is limited. Many pharmacological agents have not been fully tested across all known subtypes of mGlu receptors. Potential differences linked to the species (e.g. human *versus* rat or mouse) of the receptors and the receptor splice variants are generally not known. The influence of receptor expression level on pharmacology and selectivity has not been controlled for in most studies, particularly those involving functional assays of receptor coupling.

(*S*)-(+)-CBPG is an antagonist at mGlu₁, but is an agonist (albeit of reduced efficacy) at mGlu₅ receptors. DCG-IV also exhibits agonist activity at NMDA glutamate receptors (Uyama *et al.*, 1997). A potential novel metabotropic glutamate receptor coupled to phosphoinositide turnover has been observed in rat brain; it is activated by 4-methylhomobiotenic acid (ineffective as an agonist at recombinant Group I metabotropic glutamate receptors), but resistant to LY341495 (Chung *et al.*, 1997). There are also reports of a novel metabotropic glutamate receptor coupled to phospholipase D in rat brain, which does not readily fit into the current classification (Klein *et al.*, 1997; Pellegrini-Giampietro *et al.*, 1996).

Abbreviations: **1S,3R-ACPD**, 1-aminocyclopentane-1S,3R-dicarboxylate; **AIDA**, 1-aminoinidan-1,5(*RS*)-dicarboxylic acid; also known as UPF523; **AMN082**, *N*, *N'*-bis(diphenylmethyl)-1,2-ethanediamine dihydrochloride; **L-AP4**, *S*-2-amino-4-phosphonobutyrate; **2R,4R-APDC**, aminopyrrolidine-2*R*,4*R*-dicarboxylate; also known as LY314593; **BAY 36-7620**, (3*aS*,6*aS*)-6*a*-naphtalan-2-ylmethyl-5-methyliden-hexahyrol-cyclopenta[c]furan-1-one; **CBiPES**, *N*-[4'-cyano-biphenyl-3-yl]-*N*-(3-pyridinylmethyl)-ethanesulphonamide hydrochloride; **(S)-(+)-CBPG**, (*s*)-(1)-2-(39-carboxybicyclo[1.1.1]pentyl)glycine; **L-CCG-I**, (2*S*,3*S*,4*S*)- α -(carboxycyclopropyl)glycine; **CDPPB**, 3-cyano-*N*-(1,3-diphenyl-1*H*-[pyrazol-5-yl]benzamide; **CHPG**, 2-chloro-5-hydroxyphenylglycine; **CMPGCPCCOEt**, cyclopropan[*b*]chromen-1*a*-carboxylate; **4CPG**, 4-carboxyphenylglycine; **CPPHA**, *N*-{4-chloro-2-[(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)methyl]phenyl}-2-hydroxybenzamide; **DCG-IV**, (2*S*,1'*R*,2'*R*,3'*R*)-2-(2,3-dicarboxycyclopropyl)glycine; **(S)-3,4-DCPG**, (*S*)-3,4-dicarboxylphenylglycine; **DFB**, 3,3'-difluorobenzaldazine DHPGS-3,5-dihydroxyphenylglycine; **DMPPP**, 3,5-dimethyl pyrrole-2,4-dicarboxylic acid 2-propyl ester 4-(1,2,2-tri-methyl-propyl) ester; **EGLU**, (*s*)- α -ethylglutamate; **fenobam**, *N*-(3-chlorophenyl)-*N'*-(4,5-dihydro-1-methyl-4-oxo-1-*H*-imidazole-2-yl)-urea; **3HPG**, 3-hydroxyphenylglycine; [¹¹C]-**JNJ-16567083**, (3-ethyl-2-[¹¹C]methyl-6-quinolinyl)(*cis*-4-methoxycyclohexyl) methanone; **JNJ16259685**, (3,4-dihydro-2*H*-pyran[2,3-*b*]quinolinyl-7-yl)(*cis*-4-methoxycyclohexyl)methanone; **LY307452**, 2*S*,4*S*-2-amino-4-(4,4-diphenylbut-1-yl)pentan-1,5-dioic acid; **LY341495**, 2*S*-2-amino-2-(1*S*,2*S*-2-carboxycyclopropan-1-yl)-3-(xanth-9-yl)propanoic acid; **LY354740**, (+)-2-aminobicyclo[3.1.0]hexane-2,6-dicarboxylate; **LY367385**, (+)-2-methyl-4-carboxyphenylglycine; **LY379268**, (-)-2-oxa-4-aminobicyclo[3.1.0]hexane-4,6-dicarboxylic acid; **LY389795**, (-)-2-thia-4-aminobicyclo[3.1.0]hexane-4,6-dicarboxylic acid; **LY393675**, α -substituted-cyclobutylglycine; **LY456066**, (2-[4-(indan-2-ylamino)-5,6,7,8-tetrahydro-quinazolin-2-ylsulfanyl]-ethanol)hydrochloride; **LY456236**, [(4-methoxy-phenyl)-(6-methoxy-quinazolin-4-yl)-amine hydrochloride; **LY487379**, 2,2,2-trifluoro-*N*-[4-(2-methoxyphenoxy)phenyl]-*N*-(3-pyridinylmethyl)-ethanesulphonamide; **3-MATIDA**, α -amino-5-carboxy-3-methyl-2-thiopheneacetic acid; **MAP4**, (*S*)-2-methyl-2-amino-4-phosphonobutanoate; **MPEP**, 2-methyl-6-(phenylethynyl)-pyridine; **methoxy-MPEP**, 2-methyl-6-[(3-methoxyphenyl)ethynyl]-pyridine; **methoxy-PEPY**, 3-methoxy-5-(pyridin-2-yl-ethynyl)-pyridine; **MPPG**, (*RS*)- α -methyl-4-phosphonophenylglycine; **MTEP**, 3-[(2-methyl-1,3-thiazol-4-yl)ethynyl]pyridine; **methoxymethyl-MTEP**, 3-(methoxymethyl)-5-[(2-methyl-1,3-thiazol-4-yl)ethynyl]pyridine; **NAAG**, *N*-acetylasparylglutamate, also known as spaglumic acid; **PCCG-4**, (2*S*,1'*S*,2'*S*,3'*R*)-2-(2'-carboxy-3'-phenylcyclopropyl)glycine; **PHCCC**, *N*-phenyl-7-(hydroxylimino)cyclopropan[*b*]chromen-1*a*-carboxamide; **(RS)PPG**, (*RS*)-4-phosphonophenylglycine; **R214127**, 1-(3,4-dihydro-2*H*-pyran[2,3-*b*]quinolin-7-yl)-2-phenyl-1-ethanone; **Ro01-6128**, diphenylacetyl-carbamic acid ethyl ester; **Ro67-4853**, (9*H*-xanthene-9-carbonyl)-carbamic acid butyl ester; **Ro67-7476**, (*S*)-2-(4-fluoro-phenyl)-1-(toluene-4-sulfonyl)-pyrrolidine; **SIB1757**, 6-methyl-2-(phenylazo)-3-pyridinol; **SIB1893**, [(phenylazo)-3-pyridinol]-2-methyl-6-(2-phenylethenyl)pyridine; **S-TBPG**, 2-(3'-(1*H*-tetrazol-5-yl)bicyclo[1.1.1]pent-1-yl)glycine; **THPG**, (*RS*)-3,4,5-trihydroxyphenylglycine; **YM298198**, (6-[(2-methoxyethyl)amino]methyl)-*N*-methyl-*N*-neopentylthiao[3,2-*a*]benzimidazole-2-carboxamide

Further Reading:

- DE BLASI, A., CONN, P.J., PIN, J. & NICOLETTI, F. (2001). Molecular determinants of metabotropic glutamate receptor signaling. *Trends Pharmacol. Sci.*, **22**, 114–120.
- DHAMI, G.K. & FERGUSON, S.S. (2006). Regulation of metabotropic glutamate receptor signaling, desensitization and endocytosis. *Pharmacol. Ther.*, **111**, 260–271.
- FERRAGUTI, F. & SHIGEMOTO, R. (2006). Metabotropic glutamate receptors. *Cell Tissue Res.*, **326**, 483–504.
- GASPARINI, F., KUHN, R. & PIN, J.P. (2002). Allosteric modulators of group I metabotropic glutamate receptors: novel subtype-selective ligands and therapeutic perspectives. *Curr. Opin. Pharmacol.*, **2**, 43–49.
- HERMANS, E. & CHALLISS, R.A.J. (2001). Structural, signalling and regulatory properties of the group I metabotropic glutamate receptors: prototypic family C G-protein-coupled receptors. *Biochem. J.*, **359**, 465–484.
- KUBO, Y. & TATEYAMA, M. (2005). Towards a view of functioning dimeric metabotropic receptors. *Curr. Opin. Neurobiol.*, **15**, 289–295.
- NAKANISHI, S., NAKAJIMA, Y., MASU, M., UEDA, Y., NAKAHARA, K., WATANABE, D., YAMAGUCHI, S., KAWABATA, S. & OKADA, M. (1998). Glutamate receptors: brain function and signal transduction. *Brain Res. Rev.*, **26**, 230–235.
- PELLICCIARI, R. & COSTANTINO, G. (1999). Metabotropic G-protein-coupled glutamate receptors as therapeutic targets. *Curr. Opin. Chem. Biol.*, **3**, 433–440.
- PIN, J.P. & ACHER, F. (2002). The metabotropic glutamate receptors: structure, activation mechanism and pharmacology. *Curr. Drug Target CNS. Neurol. Disord.*, **1**, 297–317.
- SCHOEPP, D.D. (2001). Unveiling the functions of presynaptic metabotropic glutamate receptors in the central nervous system. *J. Pharmacol. Exp. Ther.*, **299**, 12–20.
- SCHOEPP, D.D., ALEXANDER, S.P., BEART, P., CONN, P.J., LODGE, D., NAKANISHI, S., NICOLETTI, A. & PIN, J.-P. (2000). Metabotropic glutamate receptors. In: *IUPHAR Compendium of Receptor Characterization and Classification*, eds. WATSON, S.P. & GIRDLESTONE, D. 2nd edn. London: IUPHAR Press.
- SCHOEPP, D.D., JANE, D.E. & MONN, J.A. (2000). Pharmacological agents acting at subtypes of metabotropic glutamate receptors. *Neuropharmacology*, **38**, 1431–1476.
- SWANSON, C.J., BURES, M., JOHNSON, M.P., LINDEN, A.M., MONN, J.A. & SCHOEPP, D.D. (2005). Metabotropic glutamate receptors as novel targets for anxiety and stress disorders. *Nat. Rev. Drug Discovery*, **4**, 131–144.
- URE, J., BAUDRY, M., & PERASSOLO, M. (2006). Metabotropic glutamate receptors and epilepsy. *J. Neurol. Sci.*, **247**, 1–9.

References:

- ANDERSON, J.J. et al. (2002). *Pharmacol. Exp. Ther.*, **303**, 1044–1051.
- BONNEFOUS, C. et al. (2005). *Bioorg. Med. Chem. Lett.*, **15**, 1197–1200.
- BRABET, L. et al. (1995). *Neuropharmacology*, **34**, 895–905.
- BRÄUNER-OSBORNE, H. et al. (1996). *J. Med. Chem.*, **39**, 3188–3194.
- BRODKIN, J. et al. (2002). *Eur. J. Neurosci.*, **16**, 2241–2244.
- CARROLL, F.Y. et al. (2001). *Mol. Pharmacol.*, **59**, 965–973.
- CHUNG, D.S. et al. (1997). *J. Pharmacol. Exp. Ther.*, **283**, 742–749.
- CLARK, B.P. et al. (1997). *Bioorg. Med. Chem. Lett.*, **7**, 2777–2780.
- CONSTANTINO, G. et al. (2001). *Biorg. Med. Chem.*, **9**, 221–227.
- DOHERTY, A.J. et al. (1997). *Neuropharmacology*, **36**, 265–267.
- ESCRIBANO, A. et al. (1998). *Bioorg. Med. Chem. Lett.*, **8**, 765–770.
- FLOR, P.J. et al. (2005). *Neuropharmacology*, **49S**, 244.
- GASPARINI, F. et al. (1999a). *J. Pharmacol. Exp. Ther.*, **289**, 1678–1687.
- GASPARINI, F. et al. (1999b). *Neuropharmacology*, **38**, 1493–1503.

- GASPARINI, F. *et al.* (2002). *Bioorg. Med. Chem. Lett.*, **12**, 407–409.
- JANE, D.E. *et al.* (1996). *Neuropharmacology*, **35**, 1029–1035.
- JOHNSON, M.P. *et al.* (2003). *J. Med. Chem.*, **46**, 3189–3192.
- JOHNSON, M.P. *et al.* (2005). *Psychopharmacology*, **179**, 271–283.
- KINGSTON, A.E. *et al.* (1998). *Neuropharmacology*, **37**, 1–12.
- KINNEY, G.G. *et al.* (2005). *J. Pharmacol. Exp. Ther.*, **313**, 199–206.
- KLEIN, J. *et al.* (1997). *Neuropharmacology*, **36**, 305–311.
- KNOFLACH, F. *et al.* (2001). *Proc. Natl. Acad. Sci. U.S.A.*, **98**, 13402–13407.
- KOHARA, A. *et al.* (2005). *J. Pharmacol. Exp. Ther.*, **315**, 163–169.
- LAVREYSEN, H. *et al.* (2003). *Mol. Pharmacol.*, **63**, 1082–1093.
- LAVREYSEN, H. *et al.* (2004). *Neuropharmacology*, **47**, 961–972.
- LI, L. *et al.* (2002). *Neuropharmacology*, **43**, 273–318.
- LITSCHIG, S. *et al.* (1999). *Mol. Pharmacol.*, **55**, 453–461.
- MAJ, M. *et al.* (2003). *Neuropharmacology*, **45**, 895–906.
- MALHERBE, P. *et al.* (2003). *J. Biol. Chem.*, **278**, 8340–8347.
- MANNAIONI, G. *et al.* (1999). *Neuropharmacology*, **38**, 917–926.
- MATHIESEN, J.M. *et al.* (2003). *Br. J. Pharmacol.*, **138**, 1026–1030.
- MICHEL, F. *et al.* (2003). *Bioorg. Med. Chem. Letts.*, **13**, 2113–2118.
- MONN, J.A. *et al.* (1999). *J. Med. Chem.*, **42**, 1027–1040.
- MORONI, F. *et al.* (1997). *J. Pharmacol. Exp.*, **281**, 721–729.
- MORONI, F. *et al.* (2002). *Neuropharmacology*, **42**, 741–751.
- O'BRIEN, J.A. *et al.* (2003). *Mol. Pharmacol.*, **64**, 731–740.
- O'BRIEN, J.A. *et al.* (2004). *J. Pharmacol. Exp. Ther.*, **309**, 568–577.
- PELLEGRINI-GIAMPINETRO, D.E. *et al.* (1996). *Br. J. Pharmacol.*, **118**, 1035–1043.
- PELLICCIARI, R. *et al.* (1996). *J. Med. Chem.*, **39**, 2259–2269.
- PORTER, R.H. *et al.* (2005). *J. Pharmacol. Exp. Ther.*, **315**, 711–721.
- SCHOEPP, D.D. *et al.* (1996). *Neuropharmacology*, **35**, 1661–1672.
- SCHOEPP, D.D. *et al.* (1997). *Neuropharmacology*, **36**, 1–11.
- THOMAS, N.K. *et al.* (2001). *Neuropharmacology*, **40**, 311–318.
- THORESON, W.B. *et al.* (1997). *Neuropharmacology*, **36**, 13–20.
- TUCKMANTEL, W. *et al.* (1997). *Bioorg. Med. Chem. Lett.*, **7**, 601–606.
- UYAMA, Y. *et al.* (1997). *Brain Res.*, **752**, 327–330.
- VARNEY, M.A. *et al.* (1999). *J. Pharmacol. Exp. Ther.*, **290**, 170–181.
- WERMUTH, C.G. *et al.* (1996). *J. Med. Chem.*, **39**, 814–816.
- WROBLEWSKA, B. *et al.* (1997). *J. Neurochem.*, **69**, 174–181.
- WU, S. *et al.* (1998). *Mol. Brain Res.*, **53**, 88–97.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Glycoprotein hormone

Overview: Glycoprotein hormone receptors (provisional nomenclature) are activated by a heterodimeric glycoprotein made up of a common α chain (116 aa ENSG00000135346), with a unique β chain that confers the biological specificity to FSH (follicle-stimulating hormone, follitropin, 129 aa, ENSG00000131808), LH (luteinizing hormone, lutropin, 141 aa ENSG00000104826), CG (choriogonadotropin, chorionic gonadotropin, 165 aa, ENSG00000104818/ENSG00000104827) or TSH (thyrotropin, thyroid-stimulating hormone, 138 aa ENSG00000134200). There is binding cross-reactivity across the endogenous agonists for each of the glycoprotein hormone receptors. The deglycosylated hormones appear to exhibit reduced efficacy at these receptors (Sairam, 1989).

Nomenclature	FSH	LH	TSH
Ensembl ID	ENSG00000170820	ENSG00000168546	ENSG00000146013
Principal transduction	G _s	G _s , G _{q/11} and G _i	All four families of G-proteins can be activated by this receptor
Selective agonists	FSH	LH, CG	TSH
Probes	[¹²⁵ I]-FSH	[¹²⁵ I]-LH, [¹²⁵ I]-CG	[¹²⁵ I]-TSH

Animal follitropins are less potent than the human hormone as agonists at the human FSH receptor. Autoimmune antibodies that act as agonists of the TSH receptor are found in patients with Grave's disease (see Rapoport *et al.*, 1998). Gain- and loss-of-function mutations of the FSH receptor are associated with human reproductive disorders (Aittomaki *et al.*, 1995; Beau *et al.*, 1998; Gromoll *et al.*, 1996; Touraine *et al.*, 1999). Loss-of-function mutations of the LH receptor are associated with Leydig cell hypoplasia and gain-of-function mutations are associated with male-limited gonadotropin-independent precocious puberty (e.g. Latronico and Segaloff, 1999; Shenker, 2002) and Leydig cell tumours (Liu *et al.*, 1999). Mutations of the TSH receptor exhibiting constitutive activity underlie hyperfunctioning thyroid adenomas (Parma *et al.*, 1993) and congenital hyperthyroidism (Kopp *et al.*, 1995). TSH receptor loss-of-function mutations are associated with thyrotropin resistance (Sunthornthepvarakul *et al.*, 1995). The rat FSH receptor also stimulates phosphoinositide turnover through an unidentified G protein (Quintana *et al.*, 1994).

Further Reading:

- DAVIES, T., MARIANS, R. & LATIF, R. (2002). The TSH receptor reveals itself. *J. Clin. Invest.*, **110**, 161–164.
- DAVIES, T.F., ANDO, T., LIN, R.Y., TOMER, Y. & LATIF, R. (2005). Thyrotropin receptor-associated diseases: from adenomata to Graves disease. *J. Clin. Invest.*, **115**, 1972–1983.
- FARID, N.R. & SZKUDLINSKI, M.W. (2004). Minireview: structural and functional evolution of the thyrotropin receptor. *Endocrinology*, **145**, 4048–4057.
- KOPP, P. (2001). The TSH receptor and its role in thyroid disease. *Cell. Mol. Life Sci.*, **58**, 1301–1322.
- SCHOTT, M., SCHERBAUM, W.A. & MORGENTHALER, N.G. (2005). Thyrotropin receptor autoantibodies in Graves' disease. *Trends Endocrinol. Metab.*, **16**, 243–248.
- SZKUDLINSKI, M.W., FREMONT, V., RONIN, C. & WEINTRAUB, B.D. (2002). Thyroid-stimulating hormone and thyroid-stimulating hormone receptor structure-function relationships. *Physiol. Rev.*, **82**, 473–502.
- THEMMEN, A.P.N. & HUHTANIEMI, I.T. (2000). Mutations of gonadotropins and gonadotropin receptors: elucidating the physiology and pathophysiology of pituitary-gonadal function. *Endocr. Rev.*, **21**, 551–583.
- VASSART, G., PARDO, L. & COSTAGLIOLA, S. (2004). A molecular dissection of the glycoprotein hormone receptors. *Trends Biochem. Sci.*, **29**, 119–126.
- WONEROW, P., NEUMANN, S., GUDERMANN, T. & PASCHKE, R. (2001). Thyrotropin receptor mutations as a tool to understand thyrotropin receptor action. *J. Mol. Med.*, **79**, 707–721.

References:

- AITTOMAKI, K. *et al.* (1995). *Cell*, **82**, 959–968.
- BEAU, I. *et al.* (1998). *J. Clin. Invest.*, **102**, 1352–1359.
- GROMOLL, J. *et al.* (1996). *J. Clin. Endocrinol. Metab.*, **81**, 1367–1370.
- KOPP, P. *et al.* (1995). *N. Engl. J. Med.*, **332**, 150–154.
- LATRONICO, A.C. & SEGALOFF, D.L. (1999). *Am. J. Hum. Genet.*, **65**, 949–958.
- LIU, G. *et al.* (1999). *N. Engl. J. Med.*, **341**, 1731–1736.
- PARMA, J. *et al.* (1993). *Nature*, **365**, 649–651.
- QUINTANA, J. *et al.* (1994). *J. Biol. Chem.*, **269**, 8772–8779.
- RAPOPORT, B. *et al.* (1998). *Endocr. Rev.*, **19**, 673–716.
- SAIRAM, M.R. (1989). *FASEB J.*, **3**, 1915–1926.
- SHENKER, A. (2002). *Receptors Channels*, **8**, 3–18.
- SUNTHORNTHPEVARAKUL, T. *et al.* (1995). *N. Engl. J. Med.*, **332**, 155–160.
- TOURAIN, P. *et al.* (1999). *Mol. Endocrinol.*, **13**, 1844–1854.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Gonadotrophin-releasing hormone (GnRH)

Overview: Gonadotropin-releasing hormone (GnRH) is a hypothalamic decapeptide (pGlu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pr-Gly-NH₂, also known as luteinising hormone-releasing hormone, gonadoliberin, luliberin, gonadorelin, ENSG00000147437) designated GnRH I, to distinguish it from related peptides such as GnRH II (pGlu-His-Trp-Ser-His-Gly-Trp-Tyr-Plo-Gly-NH₂, also known as chicken GnRH-II, ENSG00000180290) and GnRH III (pGlu-His-Trp-Ser-His-Asp-Trp-Lys-Pro-Gly-NH₂ also known as lamprey GnRH III). Receptors for all three ligands exist in amphibians but only GnRH I and GnRH II (and their cognate receptors) have been found in mammals (Sealfon *et al.*, 1997; Millar, 2002). GnRH receptors have been cloned from numerous species (most of which express two or three types of GnRHR) and grouped phylogenetically (Silver *et al.*, 2005). Type I GnRHRs are expressed primarily by pituitary gonadotrophs in mammals and mediate central control of reproduction. They are selectively activated by GnRH I and all lack the C-terminal tails found in other seven transmembrane region receptors. Type II GnRHRs include all non-mammalian GnRHRs as well as the recently cloned type II primate GnRHRs. They all possess C-terminal tails and (where tested) are selective for GnRH II (over GnRH I). An alternative phylogenetic classification (see Millar *et al.*, 2004) divided these receptors into three classes and includes both GnRH I-selective mammalian type I GnRHRs and GnRH II-selective non-mammalian receptors in type I. Although thousands of peptide analogues of GnRH I have been synthesised and several (agonists and antagonists) are used therapeutically (Kiesel *et al.*, 2002), the potency of most of these peptides at type II GnRHRs is unknown.

Nomenclature	Type I GnRHR	Type II GnRHR
Other names	LHRH receptor, GnRH I receptor	—
Ensembl ID	ENSG00000109163	ENSG00000180290
Principal transduction	G _{q/11}	G _{q/11}
Rank order of potency	GnRH I > GnRH II	GnRH II > GnRH I
Selective agonists	Triptorelin, buserelin, leuporelin, nafarelin, histrelin, goserelin	—
Selective antagonists	Antide (9.0, Neill, 2002), cetrorelix (8.8, Neill, 2002), ganirelix, abarelix	Trptorelix-1 (Maiti <i>et al.</i> , 2003)
Probes	[¹²⁵ I]-GnRH I, [¹²⁵ I]-buserelin	[¹²⁵ I]-GnRH II

Type I (and type II) GnRHRs couple primarily to G_{q/11} (Grosse *et al.*, 2000) but coupling to G_s and G_i is evident in some systems (Krsmanovic *et al.*, 2003). Type II GnRHRs may also mediate (heterotrimeric) G protein-independent signalling to protein kinases (see Caunt *et al.*, 2006). There is increasing evidence for expression of GnRHRs on hormone-dependent cancer cells where they can exert antiproliferative and/or proapoptotic effects and mediate effects of cytotoxins conjugated to GnRH analogues (Limonta *et al.*, 2003; Harrison *et al.*, 2004; Schally and Nagy 2004; Cheng and Leung, 2005). In some human cancer cell models GnRH II is more potent than GnRH I, implying mediation by a type II GnRHR (Grundker *et al.*, 2002). However, type II GnRHRs that are expressed by some primates are probably not expressed in humans because the human type II GnRHR gene contains a frame shift and internal stop codon (Millar, 2002; Morgan *et al.*, 2003). The possibility remains that this gene expresses type II GnRHR-related proteins (other than the full-length receptor) that mediate responses to GnRH II (Neill *et al.*, 2004). Alternatively, there is evidence for multiple active GnRHR conformations (Caunt *et al.*, 2004; Maudsley *et al.*, 2004; Millar *et al.*, 2004) raising the possibility that type I GnRHR-mediated proliferation inhibition in hormone-dependent cancer cells is dependent upon different conformations (with different ligand specificity) than effects on G_{q/11} in pituitary cells (Maudsley *et al.*, 2004). Loss-of-function mutations in the type I GnRHR and deficiency of GnRH I are associated with hypogonadotropic hypogonadism although some 'loss of function' mutations may actually prevent trafficking of 'functional' type I GnRHRs to the cell surface, as evidenced by recovery of function by nonpeptide antagonists (Leanos-Miranda *et al.*, 2003). GnRHR signalling may be dependent upon receptor oligomerisation (Conn *et al.*, 1982; Kroeger *et al.*, 2001).

Further Reading:

- CAUNT, C.J., FINCH, A.R., SEDGLEY, K.R. & MCARDLE, C.A. (2006). GnRH receptor signalling to ERK: kinetics and compartmentalization. *Trends Endocrinol. Metab.*, **17**, 308–313.
- CHENG, C.K. & LEUNG, P.C. (2005). Molecular biology of gonadotropin-releasing hormone (GnRH)-I, GnRH-II, and their receptors in humans. *Endocr. Rev.*, **26**, 283–306.
- HERBST, K.L. (2003). Gonadotropin-releasing hormone antagonists. *Curr. Opin. Pharmacol.*, **3**, 660–666.
- HAPGOOD, J.P., SADIE, H., VAN, B.W. & RONACHER, K. (2005). Regulation of expression of mammalian gonadotrophin-releasing hormone receptor genes. *J. Neuroendocrinol.*, **17**, 619–638.
- KAKAR, S.S., MALIK, M.T. & WINTERS, S.J. (2002). Gonadotropin-releasing hormone receptor: cloning, expression and transcriptional regulation. *Prog. Brain Res.*, **141**, 129–147.
- MCARDLE, C.A., DAVIDSON, J.S. & WILLIAMS, G.B. (1999). The tail of the gonadotrophin-releasing hormone receptor: desensitization at, and distal to, G protein-coupled receptors. *Mol. Cell. Endocrinol.*, **151**, 129–136.
- MCARDLE, C.A., FRANKLIN, J., GREEN, L. & HISLOP, J.N. (2002). Signalling, cycling and desensitisation of gonadotrophin-releasing hormone receptors. *J. Endocrinol.*, **173**, 1–11.
- MILLAR, R.P. (2003). GnRH II and type II GnRH receptors. *Trends Endocrinol. Metab.*, **14**, 35–43.
- MILLAR, R.P., LU, Z.L., PAWSON, A.J., FLANAGAN, C.A., MORGAN, K. & MAUDSLEY, S.R. (2004). Gonadotropin-releasing hormone receptors. *Endocr. Rev.*, **25**, 235–275.
- SHAH, B.H. & CATT, K.J. (2004). Matrix metalloproteinases in reproductive endocrinology. *Trends Endocrinol. Metab.*, **15**, 47–49.

References:

- CAUNT, C.J. *et al.* (2004). *Endocrinology*, **145**, 3594–3602.
- CONN, P.M. *et al.* (1982). *Nature*, **296**, 653–655.
- GROSSE, R. *et al.* (2000). *J. Biol. Chem.*, **275**, 9193–9200.
- GRUNDKER, C. *et al.* (2002). *Am. J. Obstet. Gynecol.*, **187**, 528–537.
- KIESEL, L.A. *et al.* (2002). *Clin. Endocrinol. (Oxford)*, **56**, 677–687.
- KROEGER, K.M. *et al.* (2001). *J. Biol. Chem.*, **276**, 12736–12743.
- KRSMANOVIC, L.Z. *et al.* (2003). *Proc. Natl. Acad. Sci. USA*, **100**, 2969–2974.
- LEANOS-MIRANDA, A. *et al.* (2003). *J. Clin. Endocrinol. Metab.*, **88**, 3360–3367.
- HARRISON, G.S. *et al.* (2004). *Endocr.-Relat. Cancer*, **11**, 725–748.

- LIMONTA, P. (2003). *Frontiers Neuroendocrinol.*, **24**, 279–295.
- MAUDSLEY, S *et al.* (2004). *Cancer Res.*, **64**, 7533–7544.
- MAITI, K. *et al.* (2003). *Mol. Cells.*, **16**, 173–179.
- MILLAR, R.P. (2002). *Trends Endocrinol. Metab.*, **14**, 35–43.
- MORGAN, K. *et al.* (2003). *Endocrinology*, **144**, 423–436.
- NEILL, J.D. (2002). *Endocrinology*, **143**, 737–743.
- NEILL, J.D. *et al.* (2004). *Trends Endocrinol. Metab.*, **15**, 383–392.
- SCHALLY, A.V. & NAGY, A. (2004). *Trends Endocrinol. Metab.*, **15**, 300–310.
- SEALFON, S.C. *et al.* (1997). *Endocr. Rev.*, **18**, 180–205.
- SILVER, M.R. *et al.* (2005). *Endocrinology*, **146**, 3351–3361.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Histamine

Overview: Histamine receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Histamine Receptors, see Hill *et al.*, 1997) are activated by the endogenous ligand histamine. Marked species differences exist between histamine receptor orthologues (see Hill *et al.*, 1997).

Nomenclature	H ₁	H ₂	H ₃	H ₄
Ensembl ID	ENSG00000171088	ENSG00000168546	ENSG00000146013	ENSG00000125861
Principal transduction	G _{q/11}	G _s	G _{i/o}	G _{i/o}
Selective agonists	Histaprodifen, N ^z -methylhistaprodifen	Amthamine	Methimnapip (Kitbunnadaj <i>et al.</i> , 2005),	Clobenpropit, 4-methylhistamine, VUF8430 (Lim <i>et al.</i> , 2006)
Selective antagonists	Triprolidine (9.9), mepyramine (9.1)		Tiotidine (7.8), ranitidine (7.1)	Clobenpropit (9.9), iodophenpropit (9.6), A331440 (8.5, Hancock <i>et al.</i> , 2004) thioperamide (8.4)
Probes	[³ H]-Mepyramine (1 nM), [¹¹ C]-Mepyramine, [¹¹ C]-doxepin	[³ H]-Tiotidine (15 nM), [¹²⁵ I]-iodoaminopotentidine (0.3 nM)	[³ H]-R- α -Methylhistamine (0.5 nM), [³ H]-N ^z -methylhistamine (2 nM), [¹²⁵ I]-iodophenpropit (0.6 nM), [¹²⁵ I]-iodoproxyfan (0.06 nM)	[³ H]-JNJ777120 (3.6 nM)

Histaprodifen and N^z-methylhistaprodifen are reduced efficacy agonists. The H₄ receptor appears to exhibit broadly similar pharmacology to the H₃ receptor for imidazole-containing ligands, although R- α -methylhistamine and N- α -methylhistamine are less potent, while clobenpropit acts as a reduced efficacy agonist (Nakamura *et al.*, 2000; Oda *et al.*, 2000; Liu *et al.*, 2001; Nguyen *et al.*, 2001; Zhu *et al.*, 2001). Moreover, 4-methylhistamine is identified as a high affinity, full agonist for the human H₄ receptor (Lim *et al.*, 2005). [³H]-Histamine has been used to label the H₄ receptor in heterologous expression systems.

Further Reading:

DE ESCH, I.J., THURMOND, R.L., JONGEJAN, A. & LEURS, R. (2005). The histamine H₄ receptor as new therapeutic target for inflammation. *Trends Pharmacol. Sci.*, **26**, 462–469.

HILL, S.J., GANELLIN, C.R., TIMMERMAN, H., SCHWARTZ, J.C., SHANKLEY, N.P., YOUNG, J.M., SCHUNACK, W., LEVI, R. & HAAS, H.L. (1997). International Union of Pharmacology XIII. Classification of histamine receptors. *Pharmacol. Rev.*, **49**, 253–278.

HOUGH, L.B. (2001). Genomics meets histamine receptors: new subtypes, new receptors. *Mol. Pharmacol.*, **59**, 415–419.

LEURS, R., BAKKER, R.A., TIMMERMAN, H. & DE ESCH, I.S. (2005). The histamine H₃ receptor: from gene cloning to H₃ receptor drugs. *Nat. Rev. Drug Discov.*, **4**, 107–120.

References:

- HANCOCK, A.A. *et al.* (2004). *Eur. J. Pharmacol.*, **487**, 183–197.
- JABLONOWSKI, J.A. *et al.* (2003). *J. Med. Chem.*, **46**, 3957–3960.
- KITBUNNADAJ, R. *et al.* (2005). *Bioorg. Med. Chem.*, **13**, 6309–6323.
- LEURS, R. *et al.* (1994). *Br. J. Pharmacol.*, **112**, 847–854.
- LIM, H.D. *et al.* (2005). *J. Pharmacol. Exp. Ther.*, **314**, 1310–1321.
- LIM, H.D. *et al.* (2006). *J. Med. Chem.*, **49**, 6650–6651.
- LIU, C. *et al.* (2001). *Mol. Pharmacol.*, **59**, 420–426.
- NAKAMURA, T. *et al.* (2000). *Biochem. Biophys. Res. Commun.*, **279**, 615–620.
- NGUYEN, T. *et al.* (2001). *Mol. Pharmacol.*, **59**, 427–433.
- ODA, T. *et al.* (2000). *J. Biol. Chem.*, **275**, 36781–36786.
- SMIT, M.J. *et al.* (1996). *J. Neurochem.*, **67**, 1791–1800.
- THURMOND, R.L. *et al.* (2004). *J. Pharmacol. Exp. Ther.*, **309**, 404–413.
- ZHU, Y. *et al.* (2001). *Mol. Pharmacol.*, **59**, 434–441.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Leukotriene

Overview: Leukotriene receptors (nomenclature agreed by NC-IUPHAR on Leukotriene and Lipoxin Receptors, Brink *et al.*, 2003) are activated by the endogenous ligands leukotriene (LT) B₄, LTC₄, LTD₄, LTE₄, 12*R*-HETE and 12*S*-HETE. Leukotrienes bind extensively to enzymes in their metabolic pathways (glutathione-*S*-transferase/LTC₄ synthase, γ -glutamyltranspeptidase and several aminopeptidases) and can also bind to peroxisome proliferator-activated receptor α (PPAR α , Lin *et al.*, 1999) and the ALX lipoxin receptor (Fiore *et al.*, 1994), complicating the interpretation of radioligand binding and functional studies (e.g. LTC₄ is rapidly converted in many systems to LTD₄). Metabolic inhibitors (e.g. serine–borate complex) reduce this problem but can also have nonspecific effects.

Nomenclature	BLT ₁	BLT ₂	CysLT ₁	CysLT ₂
Other names	LTB ₄	—	HG55, HMTMF81, LTD ₄	HPN321, LTC ₄
Ensemble ID	ENSG00000116329	ENSG00000082556	ENSG00000112038	ENSG00000125510
Principal transduction	G _{q/11} , G _{i/o}	G _{q/11} , G _{i/o}	G _{q/11}	G _{q/11}
Rank order of potency	LTB ₄ > 20-hydroxy-LTB ₄ >> 12 <i>R</i> -HETE (Yokomizo <i>et al.</i> , 2001)	LTB ₄ > 12 <i>S</i> -HETE = 12 <i>S</i> -HPETE > 15 <i>S</i> -HETE > 12 <i>R</i> -HETE = 15 <i>S</i> -HETE > 20-hydroxy-LTB ₄ (Yokomizo <i>et al.</i> , 2001)	LTD ₄ > LTC ₄ > LTE ₄ (Sarau <i>et al.</i> , 1999)	LTC ₄ = LTD ₄ >> LTE ₄ (Nothacker <i>et al.</i> , 2000)
Selective agonists	—	12 <i>S</i> -HETE	—	BAYu9773
Selective antagonists	CP105696 (pIC ₅₀ 7.2), U75302 (pIC ₅₀ 6.9)	LY255283 (pIC ₅₀ 6.0)	Zafirlukast (9.5), montelukast (9.3), SR2640 (8.7), pobilukast (8.6), sulukast (8.3)	—
Probes	[³ H]-LTB ₄ (0.2–0.7 nM), [³ H]-CGS23131 (13 nM)	[³ H]-LTB ₄ (0.2–23 nM)	[³ H]-LTD ₄ , [³ H]-ICI198615	[³ H]-LTD ₄

BAYu9773 is an antagonist at CysLT₁ (6.8–7.7) and a reduced efficacy agonist at CysLT₂ receptors. The CysLT₁ and CysLT₂ receptors also respond to uracil nucleotides (Mellor *et al.*, 2001; 2003). GPR17 has recently been described as a 'dualistic' receptor responding to both uracil nucleotides and cysteinyl leukotrienes, responses which may be inhibited by antagonists of either. P2 or CysLT receptors (Ciana *et al.*, 2006).

Abbreviations: 12*R*-HETE, 12(*R*)-hydroxyeicosa-5*Z*,8*Z*,10*E*,14*Z*-tetraenoic acid; BAYu9773, 6(*R*)-(4'-carboxyphenyl-thio)-5(*S*)-hydroxy-7(*E*),11(*Z*),14(*Z*)-eicosatetraenoic acid; CGS23131, (E)-5-(3-carboxybenzoyl)-2-([6-[4-methoxyphenyl]-5-hexenyl]oxy)benzene propanoic acid; also known as LY223982; CP105696, (+)-1-(3*S*,4*R*)-[3-(4-phenylbenzyl)-4-hydroxy-chroman-7-yl]cyclopentane carboxylic acid; ICI198615, (1-[2-methoxy-4-([phenylsulfonylamino]carbonyl)phenyl]-methyl]-1*H*-indazol-6-yl)carbamic acid cyclopentyl ester; LY255283, 1-[5-ethyl-2-hydroxy-4-[[6-methyl-6-(1*H*-tetrazol-5-yl)-heptyl]-oxy]-phenyl]-ethanone; SR2640, 2-[3-[2-quinolylmethoxy]phenylamino]benzoic acid; U75302, 6-(6-(3-hydroxy-1*E*,5*Z*-undecadien-1-yl)-2-pyridinyl)-1,5-hexanediol

Further Reading:

- BACK, M. (2002). Functional characteristics of cysteinyl-leukotriene receptor subtypes. *Life Sci.*, **71**, 611–622.
- BARNES, P.J. (2003). Anti-leukotrienes: here to stay? *Curr. Opin. Pharmacol.*, **3**, 257–263.
- BRINK, C., DAHLEN, S.E., DRAZEN, J., EVANS, J.F., HAY, D.W.P., NICOSIA, S., SERHAN, C.N., SHIMIZU, T. & YOKOMIZO, T. (2003). International Union of Pharmacology. XXXVII. Nomenclature for leukotriene and lipoxin receptors. *Pharmacol. Rev.*, **55**, 195–227.
- EVANS, J.F. (2003). The cysteinyl leukotriene receptors. *Prostaglandins Leukot. Essent. Fatty Acids*, **69**, 117–122.
- HUI, Y. & FUNK, C.D. (2002). Cysteinyl leukotriene receptors. *Biochem. Pharmacol.*, **64**, 1549–1557.
- PETERS-GOLDEN, M., CANETTI, C., MANCUSO, P. & COFFEY, M.J. (2005). Leukotrienes: underappreciated mediators of innate immune responses. *J. Immunol.*, **174**, 589–594.
- TAGER, A.M. & LUSTER, A.D. (2003). BLT₁ and BLT₂: the leukotriene B₄ receptors. *Prostaglandins Leukot. Essent. Fatty Acids*, **69**, 123–134.
- YOKOMIZO, T., IZUMI, T. & SHIMIZU, T. (2001). Leukotriene B₄: metabolism and signal transduction. *Arch. Biochem. Biophys.*, **385**, 231–241.

References:

- CIANA, P. *et al.* (2006). *EMBOJ*, **25**, 4165–4627.
- FIORE, S. *et al.* (1994). *J. Exp. Med.*, **180**, 253–260.
- LIN, Q. *et al.* (1999). *Biochemistry*, **38**, 185–190.
- NOTHACKER, H.P. *et al.* (2000). *Mol. Pharmacol.*, **58**, 1601–1608.
- SARAU, H.M. *et al.* (1999). *Mol. Pharmacol.*, **56**, 657–663.
- YOKOMIZO, T. *et al.* (2001). *J. Biol. Chem.*, **276**, 12454–12459.
- MELLOR, E.A. *et al.* (2001). *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 7964–7969.
- MELLOR, E.A. *et al.* (2003). *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 11589–11593.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Lipoxin

Overview: Lipoxin A₄ receptors (ALX, nomenclature agreed by NC-IUPHAR on Leukotriene and Lipoxin Receptors; Brink *et al.*, 2003) are activated by the endogenous lipid-derived, anti-inflammatory ligands lipoxin A₄ (LXA₄) and 15-epi-LXA₄ (aspirin-triggered lipoxin A₄, ATL). The ALX receptor also interacts with endogenous peptide and protein ligands, such as MHC binding peptide (Chiang *et al.*, 2000) as well as annexin 1 (ANXA1) and its N-terminal peptides (Perretti *et al.*, 2002). In addition, a soluble hydrolytic product of protease action on the urokinase-type plasminogen activator receptor has been reported to activate the ALX receptor (Resnati *et al.*, 2002). Furthermore, ALX has been suggested to act as a receptor mediating proinflammatory actions of the acute-phase reactant, serum amyloid A (Su *et al.*, 1999; Sodin-Semrl *et al.*, 2004).

Nomenclature	ALX
Other names	FPRL1, FPR2, FPRH2, RFP
Ensembl ID	ENSG00000171049
Principal transduction	G _i (Maddox <i>et al.</i> , 1997)
Rank order of potency	LXA ₄ = ATL = ATLa2 > LTC ₄ = LTD ₄ >> 15-deoxy-LXA ₄ >> fMLP (Clish <i>et al.</i> , 1999; Fiore <i>et al.</i> , 1994; Fiore and Serhan, 1995; Gronert <i>et al.</i> , 2001; Takano <i>et al.</i> , 1997)
Selective agonists	LXA ₄ , ATL, ATLa2 (Guilford <i>et al.</i> , 2004)
Probes	[³ H]-LXA ₄ (0.2–1.7 nM; Fiore <i>et al.</i> , 1994; Takano <i>et al.</i> , 1997)

A receptor selective for LXB₄ has been suggested from functional studies (Maddox and Serhan, 1996; Romano *et al.*, 1996; Ariel *et al.*, 2003).

Abbreviations: ANXA1, annexin 1; ATL, aspirin-triggered lipoxin A₄ [15-epi-LXA₄, 5S,6R,15R-trihydroxyl-7,9,13-*trans*-11-*cis*-eicosatetraenoic acid]; ATLa2, ATL analog [15-epi-16-(para-fluoro)-phenoxy-LXA₄]; LTC₄, leukotriene C₄; LTD₄, leukotriene D₄; LXA₄, lipoxin A₄ [5S,6R,15S-trihydroxyl-7,9,13-*trans*-11-*cis*-eicosatetraenoic acid]

Further Reading:

BONNANS, C., CHANEZ, P. & CHAVIS, C. (2004). Lipoxins in asthma: potential therapeutic mediators on bronchial inflammation? *Allergy*, **59**, 1027–1041.

BRINK, C., DAHLEN, S.E., DRAZEN, J., EVANS, J.F., HAY, D.W.P., NICOSIA, S., SERHAN, C.N., SHIMIZU, T. & YOKOMIZO, T. (2003). International Union of Pharmacology. XXXVII. Nomenclature for leukotriene and lipoxin receptors. *Pharmacol. Rev.*, **55**, 195–227.

CHIANG, N., ARITA, M. & SERHAN, C.N. (2005). Anti-inflammatory circuitry: lipoxin, aspirin-triggered lipoxins and their receptor ALX. *Prostaglandins Leukot. Essent. Fatty Acids*, **73**, 163–177.

CHIANG, N., SERHAN, C.N., DAHLEN, S.E., DRAZEN, J.M., HAY, D.W., ROVATI, G.E., SHIMIZU, T., YOKOMIZO, T. & BRINK, C. (2006). The lipoxin receptor ALX: potent ligand-specific and stereoselective actions *in vivo*. *Pharmacol. Rev.*, **58**, 463–487.

SERHAN, C.N. (2005). Lipoxins and aspirin-triggered 15-epi-lipoxins are the first lipid mediators of endogenous anti-inflammation and resolution. *Prostaglandins Leukot. Essent. Fatty Acids*, **73**, 141–162.

SERHAN, C.N. (2005). Novel ω-3-derived local mediators in anti-inflammation and resolution. *Pharmacol. Ther.*, **105**, 7–21.

SERHAN, C.N. (2007). Resolution phases of inflammation: novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu. Rev. Immunol.*, in press.

References:

- ARIEL, A. *et al.* (2003). *J. Immunol.*, **170**, 6266–6272.
- CHIANG, N. *et al.* (2000). *J. Exp. Med.*, **191**, 1197–1208.
- CLISH, C.B. *et al.* (1999). *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 8247–8252.
- FIGLIORE, S. *et al.* (1994). *J. Exp. Med.*, **180**, 253–260.
- FIGLIORE, S. & SERHAN, C.N. (1995). *Biochemistry*, **34**, 16678–16686.
- GRONERT, K. *et al.* (2001). *Am. J. Pathol.*, **158**, 3–9.
- GUILFORD, W.J. *et al.* (2004). *J. Med. Chem.*, **47**, 2157–2165.
- MADDOX, J.F. *et al.* (1997). *J. Biol. Chem.*, **272**, 6972–6978.
- MADDOX, J.F. & SERHAN, C.N. (1996). *J. Exp. Med.*, **183**, 137–146.
- PERRETTI, M. *et al.* (2002). *Nat. Med.*, **8**, 1296–1302.
- RESNATI, M. *et al.* (2002). *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 1359–1364.
- ROMANO, M. *et al.* (1996). *J. Immunol.*, **157**, 2149–2154.
- SODIN-SEMRL, S. *et al.* (2004). *Int. J. Immunopathol. Pharmacol.*, **17**, 145–156.
- SU, S.B. *et al.* (1999). *J. Exp. Med.*, **189**, 395–402.
- TAKANO, T. *et al.* (1997). *J. Exp. Med.*, **185**, 1693–1704.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Lysophosphatidic acid

Overview: Lysophosphatidic acid (LPA) receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Lysophospholipid Receptors; Chun *et al.*, 2002) are activated by the endogenous lipid derivative LPA. The identified receptors can account for most, although not all, LPA-induced phenomena in the literature, indicating that a majority of LPA-dependent phenomena are receptor-mediated. Radioligand binding has been conducted in heterologous expression systems using [³H]-LPA (e.g. Fukushima *et al.*, 1998). In native systems, analysis of binding data is complicated by metabolism and high levels of nonspecific binding, and therefore the relationship between recombinant and endogenously expressed receptors is unclear. Targeted deletion of LPA receptors has clarified signalling pathways and identified physiological and pathophysiological roles. LPA has also been described to be an agonist at PPAR- γ receptors (McIntyre *et al.*, 2003), although the physiological significance of this observation remains unclear (Simon *et al.*, 2005).

Nomenclature	LPA ₁	LPA ₂	LPA ₃	LPA ₄	LPA ₅
Other names	VZG-1, Edg2, <i>lp</i> _{A1}	Edg4, <i>lp</i> _{A2}	Edg7, <i>lp</i> _{A3}	p2y9, gpr23	GPR92
Ensembl ID	ENSG00000119438	ENSG00000064547	ENSG00000171517	ENSG00000147149	ENSG00000184574
Principal transduction	G _{i/o} , G _{q/11} , G _{12/13}	G _{i/o} , G _{q/11} , G _{12/13}	G _{i/o} , G _{q/11} , G _s	G _{q/11} , G _s (Noguchi <i>et al.</i> , 2003)	G _q , G _{12/13} (Kotarsky <i>et al.</i> , 2006; Lee <i>et al.</i> , 2006)
Selective agonists	—	FAP10, FAP12 (Virag <i>et al.</i> , 2003)	OMPT (Hasegawa <i>et al.</i> , 2003)	—	—
Selective antagonists	Ki16425 (Ohta <i>et al.</i> , 2003)	—	DGPP 8:0 (Ohta <i>et al.</i> , 2003)	—	—

FAP12 has antagonist activity at LPA₁ and LPA₃ receptors (Virag *et al.*, 2003). The selectivity of these antagonists is less than two orders of magnitude. None of the currently available chemical tools have validated specificity *in vivo*.

Abbreviations: DGPP 8:0, dioctanoylglycerol pyrophosphate; FAP10, decanol phosphate; FAP12, dodecanol phosphate; Ki16425, 3-(4-[4-((1-[2-chlorophenyl]-ethoxy)carbonylamino)-3-methyl-5-isoxazolyl]benzylsulfanyl)propanoic acid; OMPT, 1-oleoyl-2-O-methyl-rac-glycerophosphothionate

Further Reading:

- ANLIKER, B. & CHUN, J. (2004). Lysophospholipid G protein-coupled receptors. *J. Biol. Chem.*, **279**, 20555–20558.
- CHUN, J., GOETZL, E.J., HLA, T., IGARASHI, Y., LYNCH, K.R., MOOLENAAR, W., PYNE, S. & TIGYI, G. (2002). International Union of Pharmacology. XXXIV. Lysophospholipid Receptor Nomenclature. *Pharmacol. Rev.*, **54**, 265–269.
- CHUN, J. & ROSEN, N. (2006). Lysophospholipid receptors as potential drug targets in tissue transplantation and autoimmune diseases. *Curr. Pharm. Des.*, **12**, 161–171.
- FUKUSHIMA, N. & CHUN, J. (2001). The LPA receptors. *Prostaglandins Other Lipid Mediator*, **64**, 21–32.
- FUKUSHIMA, N., ISHII, I., CONTOS, J.J.A., WEINER, J.A. & CHUN, J. (2001). Lysophospholipid receptors. *Annu. Rev. Pharmacol. Toxicol.*, **41**, 507–534.
- GARDELL, S.E., DUBIN, A.E., & CHUN, J. (2006). Emerging medicinal roles for lysophospholipid signaling. *Trends Mol. Med.*, **12**, 65–75.
- GRALER, M.H. & GOETZL, E.J. (2002). Lysophospholipids and their G protein-coupled receptors in inflammation and immunity. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids*, **1582**, 168–174.
- INOUE, M., RASHID, M.H., FUJITA, R., CONTOS, J.J., CHUN, J. & UEDA, H. (2004). Initiation of neuropathic pain requires lysophosphatidic acid receptor signaling. *Nat. Med.*, **10**, 712–718.
- ISHII, I., FUKUSHIMA, N., YE, X. & CHUN, J. (2004). Lysophospholipid receptors: signaling and biology. *Annu. Rev. Biochem.*, **73**, 321–354.
- KINGSBURY, M.A., REHEN, S.K., CONTOS, J.J., HIGGINS, C.M. & CHUN, J. (2003). Non-proliferative effects of lysophosphatidic acid enhance cortical growth and folding. *Nat. Neurosci.*, **6**, 1292–1299.
- KOSTENIS, E. (2004). Novel clusters of receptors for sphingosine-1-phosphate, sphingosylphosphorylcholine, and (lyso)-phosphatidic acid: new receptors for 'Old' ligands. *J. Cell. Biochem.*, **92**, 923–936.
- LIN, D.A. & BOYCE, J.A. (2006). Lysophospholipids as mediators of immunity. *Adv. Immunol.*, **89**, 141–147.
- OSBORNE, N. & STAINIER, D.Y. (2003). Lipid receptors in cardiovascular development. *Annu. Rev. Physiol.*, **65**, 23–43.
- PAGÈS, C., SIMON, M.F., VALET, P. & SAULNIER-BLACHE, J.S. (2001). Lysophosphatidic acid synthesis and release. *Prostaglandins Other Lipid Mediators*, **64**, 1–10.
- RADEFF-HUANG, J., SEASHOLTZ, T.M., MATTEO, R.G. & BROWN, J.H. (2004). G protein mediated signaling pathways in lysophospholipid induced cell proliferation and survival. *J. Cell. Biochem.*, **92**, 949–966.
- YANG, A.H., ISHII, I. & CHUN, J. (2002). *In vivo* roles of lysophospholipid receptors revealed by gene targeting studies in mice. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids*, **1582**, 197–203.
- YE, X., FUKUSHIMA, N., KINGSBURY, M.A. & CHUN, J. (2002). Lysophosphatidic acid in neural signaling. *Neuroreport*, **13**, 2169–2175.
- YE, X., HAMA, K., CONTOS, J.J., ANLIKER, B., INOUE, A., SKINNER, M.K., SUZUKI, H., AMANO, T., KENNEDY, G., ARAI, H., AOKI, J. & CHUN, J. (2005). LPA3-mediated lysophosphatidic acid signalling in embryo implantation and spacing. *Nature*, **435**, 104–108.
- YE, X., ISHII, I., KINGSBURY, M.A. & CHUN, J. (2002). Lysophosphatidic acid as a novel cell survival/apoptotic factor. *Biochim. Biophys. Acta*, **1585**, 108–113.

References:

- FUKUSHIMA, N. *et al.* (1998). *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 6151–6156.
- HASEGAWA, Y. *et al.* (2003). *J. Biol. Chem.*, **278**, 11962–11969.
- KOTARSKY, K. *et al.* (2006). *J. Pharmacol. Exp. Ther.*, **318**, 619–628.
- LEE, C.N. *et al.* (2006). *J. Biol. Chem.*, **281**, 23589–23597.
- NOGUCHI, K. *et al.* (2003). *J. Biol. Chem.*, **278**, 25600–25606.
- OHATA, H. *et al.* (2003). *Mol. Pharmacol.*, **64**, 994–1005.
- SIMON, M.F. *et al.* (2005). *J. Biol. Chem.*, **280**, 14656–14662.
- VIRAG, T. *et al.* (2003). *Mol. Pharmacol.*, **63**, 1032–1042.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Melanin-concentrating hormone

Overview: Melanin-concentrating hormone (MCH) receptors (provisional nomenclature) are activated by an endogenous nonadecameric cyclic peptide identical in humans and rats (DFDMLRCMLGRVYRCPWQV) generated from a precursor (ENSG00000183395), which also produces neuropeptides EI and GE. The MCH2 receptor appears to be a non-functional pseudogene in rodents (Tan *et al.*, 2002).

Nomenclature	MCH1	MCH2
Other names	SLC-1, GPR24	SLT, GPRv17
Ensembl ID	ENSG00000128285	ENSG00000152034
Principal transduction	G _{q/11} , G _{i/o}	G _{q/11} (Hill <i>et al.</i> , 2001; Mori <i>et al.</i> , 2001; Rodriguez <i>et al.</i> , 2001)
Rank order of potency	Human MCH > salmon MCH	Human MCH = salmon MCH (Hill <i>et al.</i> , 2001)
Selective antagonists	SNAP7941 (9.2, Borowsky <i>et al.</i> , 2002), T226296 (7.5, Takekawa <i>et al.</i> , 2002)	—
Probes	[³ H]-MCH (Burgaud <i>et al.</i> , 1997), [Phe ¹³ , [125I]-Tyr ¹⁹]-MCH (Burgaud <i>et al.</i> , 1997), [¹²⁵ I]-S36057 (0.04 nM, Audinot <i>et al.</i> , 2001)	—

Abbreviations: **S36057**, 3-iodo-tyr-(8-amino-3,6-dioxo-octanoyl)-MCH-(6-17); **SNAP7941**, (+)-methyl(4S)-3-([3-(4-[3-(acetylamino)phenyl]-1-piperidinyl)propyl]amino)carbonyl-4-(3,4-difluorophenyl)-6-(methoxymethyl)-2-oxo-1,2,3,4-tetrahydro-5-pyrimidinecarboxylate hydrochloride; **T226296**, (–)-N-[6-(dimethylamino)-methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide

Further Reading:

- FORRAY, C. (2003). The MCH receptor family: feeding brain disorders? *Curr. Opin. Pharmacol.*, **3**, 85–89.
- GIBSON, W.T., PISSIOS, P., TROMBLY, D.J., LUAN, J., KEOGH, J., WAREHAM, N.J., MARATOS-FLIER, E., O'RAHILLY, S. & FAROOQI, I.S. (2004). Melanin-concentrating hormone receptor mutations and human obesity: functional analysis. *Obes. Res.*, **12**, 743–749.
- HANDLON, A.L. & ZHOU, H. (2006). Melanin-concentrating hormone-1 receptor antagonists for the treatment of obesity. *J. Med. Chem.*, **49**, 4017–4022.
- INUI, A. (1999). Feeding and body-weight regulation by hypothalamic neuropeptides-mediation of the actions of leptin. *Trends Neurosci.*, **22**, 62–67.
- PISSIOS, P. & MARATOS-FLIER, E. (2003). Melanin-concentrating hormone: from fish skin to skinny mammals. *Trends Endocrinol. Metab.*, **14**, 243–248.
- SAITO, Y., NOTHACKER, H.-P. & CIVELLI, O. (2000). Melanin-concentrating hormone receptor: an orphan receptor fits the key. *Trends Endocrinol. Metab.*, **11**, 299–303.

References:

- AUDINOT, V. *et al.* (2001). *Br. J. Pharmacol.*, **133**, 371–378.
- BOROWSKY, B. *et al.* (2002). *Nat. Med.*, **8**, 825–830.
- BURGAUD, J.L. *et al.* (1997). *Biochem. Biophys. Res. Commun.*, **241**, 622–629.
- HILL, J. *et al.* (2001). *J. Biol. Chem.*, **276**, 20125–20129.
- MORI, M. *et al.* (2001). *Biochem. Biophys. Res. Commun.*, **283**, 1013–1018.
- RODRIGUEZ, M. *et al.* (2001). *Mol. Pharmacol.*, **60**, 632–639.
- TAKEKAWA, S. *et al.* (2002). *Eur. J. Pharmacol.*, **438**, 129–135.
- TAN, C.P. *et al.* (2002). *Genomics*, **79**, 785–792.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Melanocortin

Overview: Melanocortin receptors (provisional nomenclature) are activated by members of the melanocortin family (MSH – α , β , and γ forms – δ form is not found in mammals) and adrenocorticotrophin (ACTH). Endogenous antagonists include agouti and agouti-related protein (AGRP).

Nomenclature	MC ₁	MC ₂	MC ₃	MC ₄	MC ₅
Other names	—	ACTH receptor	—	—	—
Ensemble ID	ENSG00000141037	ENSG00000185231	ENSG00000124089	ENSG00000166603	ENSG00000176136
Principal transduction	G _s	G _s	G _s	G _s	G _s
Rank order of potency	α -MSH > β -MSH ≥ ACTH, γ -MSH	ACTH	γ -MSH, β -MSH ≥ ACTH, α -MSH D-Trp ⁸ - γ -MSH (Grieco <i>et al.</i> , 2000)	β -MSH ≥ α -MSH, ACTH > γ -MSH	α -MSH ≥ β -MSH ≥ ACTH > γ -MSH
Selective agonists	—	—	—	THIQ (Van der Ploeg <i>et al.</i> , 2002)	—
Selective antagonists	—	—	—	HS014 (8.5, Schiöth <i>et al.</i> , 1998), MBP10 (Bednarek <i>et al.</i> , 2001)	—
Probes	[¹²⁵ I]-NDP-MSH	[¹²⁵ I]-ACTH-(1–24)	[¹²⁵ I]-NDP-MSH, [¹²⁵ I]-SHU9119	[¹²⁵ I]-NDP-MSH, [¹²⁵ I]-SHU9119	[¹²⁵ I]-NDP-MSH

Polymorphisms of the MC₁ receptor have been linked to variations in skin pigmentation. Defects of the MC₂ receptor underlie familial glucocorticoid deficiency. Polymorphisms of the MC₄ receptor have been linked to obesity (Chagnon *et al.*, 1997).

Abbreviations: **HS014**, *cyc*(S–S)-(Ac-Cys¹¹,D-Nal¹⁴,Cys¹⁸,Asp-NH₂) β -MSH-(11–22); **MBP10**, *cyclo*(6 β →10 ϵ)(succinyl(6)-D-(2')Nal⁷-Arg⁸-Trp⁹-Lys¹⁰)-NH₂; **NDP-MSH**, [Nle⁴,d-Phe⁷] α -MSH; **SHU9119**, Ac-Nle-Asp-His-d-Nal²-Arg-Trp-Lys-NH₂; **THIQ**, *N*-([3R]-1,2,3,4-tetrahydroisoquinolinium-3-ylcarbonyl)-(1R)-1-(4-chlorobenzyl)-2-(4-cyclohexyl-4-[1H-1,2,4-triazol-1-ylmethyl]piperidin-1-yl)-2-oxoethylamine

Further Reading:

ABDEL-MALEK, Z.A. (2001). Melanocortin receptors: their functions and regulation by physiological agonists and antagonists. *Cell. Mol. Life Sci.*, **58**, 434–441.

ADAN, R.A. & GISPEN, W.H. (2000). Melanocortins and the brain: from effects via receptors to drug targets. *Eur. J. Pharmacol.*, **405**, 13–24.

CONE, R.D. (2005). Anatomy and regulation of the central melanocortin system. *Nat. Neurosci.*, **8**, 571–578.

GANTZ, I. & FONG, T.M. (2003). The melanocortin system. *Am. J. Physiol. -Endocrinol. Metabol.*, **284**, E468–E474.

RAMOS, E.J., MEGUID, M.M., CAMPOS, A.C. & COELHO, J.C. (2005). Neuropeptide Y, α -melanocyte-stimulating hormone, and monoamines in food intake regulation. *Nutrition*, **21**, 269–279.

SCHIÖTH, H.B. & WATANOBE, H. (2002). Melanocortins and reproduction. *Brain Res. Rev.*, **38**, 340–350.

SCHIÖTH, H.B. (2001). The physiological role of melanocortin receptors. *Vitam. Horm.*, **63**, 195–232.

WILLIAMS, D.L. & SCHWARTZ, M.W. (2005). The melanocortin system as a central integrator of direct and indirect controls of food intake. *Am. J. Physiol. -Regul. Integr. Comp. Physiol.*, **289**, R2–R3.

References:

BEDNAREK, M.A. *et al.* (2001). *J. Med. Chem.*, **44**, 3665–3672.

CHAGNON, Y.C. *et al.* (1997). *Mol. Med.*, **3**, 663–673.

GRIECO, P. *et al.* (2000). *J. Med. Chem.*, **43**, 4998–5002.

SCHIÖTH, H.B. *et al.* (1998). *Br. J. Pharmacol.*, **124**, 75–82.

VAN DER PLOEG, L.H. *et al.* (2002). *Proc. Natl. Acad. Sci. USA*, **99**, 11381–11386.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Melatonin

Overview: Melatonin receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on melatonin receptors (see Dubocovich *et al.*, 1998, 2000) are activated by the endogenous ligands melatonin and *N*-acetylserotonin.

Nomenclature	MT₁	MT₂	MT₃
Other names	MEL _{1A} , ML _{1A} , Mel _{1a}	MEL _{1B} , ML _{1B} , Mel _{1b}	ML ₂
Ensembl ID	ENSG00000168412	ENSG00000134640	—
Principal transduction	G _{i/o}	G _{i/o}	—
Selective agonists	—	IIK7 (Sugden <i>et al.</i> , 1999), 5-methoxyluzindole (Dubocovich <i>et al.</i> , 1998)	<i>N</i> -acetylserotonin (Eison and Mullins, 1993; Popova and Dubocovich, 1995; Molinari <i>et al.</i> , 1996; Lucchelli <i>et al.</i> , 1997), 5MCA-NAT (Popova and Dubocovich, 1995) Prazosin (Lucchelli <i>et al.</i> , 1997)
Selective antagonists	—	K185 (9.3, Sugden <i>et al.</i> , 1999), 4P-PDOT (8.8, Dubocovich <i>et al.</i> , 1997; Dubocovich <i>et al.</i> , 1998), DH97 (8.0, Teh and Sugden, 1998)	—
Probes	[³ H]-melatonin (Browning <i>et al.</i> , 2000)	[³ H]-melatonin (Browning <i>et al.</i> , 2000)	2-iodo-[¹²⁵ I]-5MCA-NAT (Molinari <i>et al.</i> , 1996)

Melatonin, 2-iodo-melatonin, S20098, GR196429, LY156735 and TAK375 (Kato *et al.*, 2005) are nonselective agonists for MT₁ and MT₂ receptors. 2-Iodo-[¹²⁵I]-melatonin can be used to label all three melatonin receptor subtypes. (–)-AMMTC displays an ~400-fold greater agonist potency than (+)-AMMTC at rat MT₁ receptors (Ting *et al.*, 1999). Luzindole is a non-selective melatonin receptor antagonist with some selectivity for the MT₂ receptor (Dubocovich *et al.*, 1998). The MT₃ binding site of hamster kidney was identified as the hamster homologue of human quinone reductase 2 (ENSG00000124588, Nosjean *et al.*, 2000; Nosjean *et al.*, 2001). Pharmacological investigations of MT₃ binding sites have primarily been conducted in hamster and guinea-pig tissues. A suggested physiological function of the MT₃ receptor is in the control of intraocular pressure in rabbits (Pintor *et al.*, 2003). *Xenopus* melanophores and chick brain express a distinct receptor (x420, P49219; c346, P49288, initially termed Mel_{1C}) coupled to the G_{i/o} family of G proteins, for which a mammalian counterpart has not yet been defined. MT₁/MT₂ heterodimers present different pharmacological profiles from MT₁ and MT₂ receptors (Ayoub *et al.*, 2004).

Abbreviations: 4P-PDOT, 4-phenyl-2-propionamidotetraline; AMMTC, *N*-acetyl-4-aminomethyl-6-methoxy-9-methyl-1,2,3,4-tetrahydrocarbazole; DH97, 2-benzyl-*N*-pentanoyltryptamine; GR196429, *N*-(2-[2,3,7,8-tetrahydro-1*H*-furo(2,3-*g*)indol-1-yl]ethyl)acetamide; IIK7, *N*-butanoyl-2-(2-methoxy-6*H*-isoindolo [2,1-*a*]indol-11-yl) ethanamine; K185, *N*-butanoyl-2-(5,6,7-trihydro-11-methoxybenzo[3,4]cyclohept[2,1-*a*]indol-13-yl)ethanamine; LY156735, β-methyl-6-chloromelatonin; 5MCA-NAT, 5-methoxy-carbonylamino-*N*-acetyltryptamine; S20098, *N*-(2-[7-methoxy-1-naphthalenyl]ethyl)acetamide; TAK375, (S)-*N*-[2(1,6,7,8-tetrahydro-2*H*-indeno [5,4-*b*]furan-8-yl)ethyl]propionamide

Further Reading:

DUBOCOVICH, M.L., CARDINALI, D.P., DELAGRANGE, P.R.M., KRAUSE, D.N., STROSBERG, D., SUGDEN, D. & YOCCA, F.D. (2000). Melatonin Receptors. In: *The IUPHAR Compendium of Receptor Characterization and Classification*. 2nd edn., eds. Girdlestone, D. & Iuphar, pp. 270–277. London: IUPHAR Media.

DUBOCOVICH, M.L., CARDINALI, D.P., GUARDIOLA-LEMAITRE, B., HAGAN, R.M., KRAUSE, D.N., SUGDEN, D., YOCCA, F.D. & VANHOUTTE, P.M. (1998). Melatonin receptors. In: *The IUPHAR compendium of receptor characterization and classification*. eds. Girdlestone, D. & Iuphar, pp. 187–193. London: IUPHAR Media.

DUBOCOVICH, M.L., RIVERA-BERMEDEZ, M.A., GERDIN, M.J. & MASANA, M.I. (2003). Molecular pharmacology, regulation and function of mammalian melatonin receptors. *Front Biosci.*, **8**, d1093–d1108.

LEON, J., ACUNA-CASTROVIEJO, D., SAINZ, R.M., MAYO, J.C., TAN, D.X. & REITER, R.J. (2004). Melatonin and mitochondrial function. *Life Sci.*, **75**, 765–790.

MASANA, M.I. & DUBOCOVICH, M.L. (2001). Melatonin receptor signaling: finding the path through the dark. *Sci. STKE*, **2001**, E39.

NAJI, L., CARRILLO-VICO, A., GUERRERO, J.M. & CALVO, J.R. (2004). Expresión de membrana and nuclear receptors in mouse peripheral organs. *Life Sci.*, **74**, 2227–2236.

SALLINEN, P., SAARELA, S., ILVES, M., VAKKURI, O. & LEPPÄLUOTO, J. (2005). The expression of MT₁ and MT₂ melatonin receptor mRNA in several rat tissues. *Life Sci.*, **76**, 1123–1134.

SUDGEN, D., DAVISON, K., HOUGH, K.A. & THE, M.T. (2004). Melatonin, melatonin receptors and melanophores: a movin story. *Pigment Cell Res.*, **17**, 454–460.

WITT-ENDERBY, P.A., BENNETT, J., JARZYNSKA, M.J., FIRESTINE, S. & MELAN, M.A. (2003). Melatonin receptors and their regulation: biochemical and structural mechanisms. *Life Sci.*, **72**, 2183–2198.

References:

- AYOUB, M. *et al.* (2004). *Mol. Pharmacol.*, **66**, 312–321.
 BROWNING, C. *et al.* (2000). *Br. J. Pharmacol.*, **129**, 877–886.
 DUBOCOVICH, M.L. *et al.* (1997). *Naunyn-Schmiedebergs Arch. Pharmacol.*, **355**, 365–375.
 DUBOCOVICH, M.L. *et al.* (1998). *FASEB J.*, **12**, 1211–1220.
 EISON, A.S. & MULLINS, U.L. (1993). *Life Sci.*, **53**, L393–L398.
 KATO, K. *et al.* (2005). *Neuropharmacol.*, **48**, 301–310.
 KOKKOLA, T. *et al.* (2005). *J. Pine. Res.*, **39**, 1–11.
 LUCCHELLI, A. *et al.* (1997). *Br. J. Pharmacol.*, **121**, 1775–1781.
 MAZNA, P. *et al.* (2004). *J. Neurochem.*, **91**, 836–842.
 MOLINARI, E.J. *et al.* (1996). *Eur. J. Pharmacol.*, **301**, 159–168.
 MULCHAHEY, J.J. *et al.* (2004). *Life Sci.*, **75**, 1843–1856.
 NOSJEAN, O. *et al.* (2000). *J. Biol. Chem.*, **275**, 31311–31317.

- NOSJEAN, O. *et al.* (2001). *Biochem. Pharmacol.*, **61**, 1369–1379.
- PINTOR, J. *et al.* (2003). *Br. J. Pharmacol.*, **138**, 831–836.
- POPOVA, J.S. & DUBOCOVICH, M.L. (1995). *J. Neurochem.*, **64**, 130–138.
- PRADA, C. *et al.* (2005). *J. Neurophysiol.*, **94**, 968–978.
- STEWART, L. & LEUNG, L.S. (2005). *Epilepsia*, **46**, 473–480.
- SUGDEN, D. *et al.* (1999). *Reprod. Nutr. Dev.*, **39**, 335–344.
- SUZUKI, H. *et al.* (1999). *Chem. Pharm. Bull.*, **47**, 1322–1325.
- TEH, M.T. & SUGDEN, D. (1998). *Naunyn-Schmiedeberg's Arch. Pharmacol.*, **358**, 522–528.
- TING, K.N. *et al.* (1999). *Br. J. Pharmacol.*, **127**, 987–995.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Motilin

Overview: Motilin receptors (provisional nomenclature) are activated by a 22 aa peptide derived from a precursor (ENSG00000096395), which also generates motilin-associated peptide. These receptors are suggested to be responsible for the gastrointestinal prokinetic effects of motilides (particular macrolide antibiotics, see Abu-Gharbieh *et al.*, 2004; Inui *et al.*, 2004).

Nomenclature	Motilin
Other names	MTLR1 (Feighner <i>et al.</i> , 1999), GPR38 (Mckee <i>et al.</i> , 1997)
Ensembl ID	ENSG00000102539
Principal transduction	G _{q/11} (Depoortere & Peeters, 1995; Feighner <i>et al.</i> , 1999)
Selective agonists	ABT229 (Lartey <i>et al.</i> , 1995)
Probes	[¹²⁵ I]-motilin (0.1 nM, Feighner <i>et al.</i> , 1999)

Abbreviations: ABT229, 8,9-anhydro-4''-deoxy-3'-N-desmethyl-3'-N-ethylerythromycin B 6,9-hemiacetal

Further Reading:

ABU-GHARBIEH, E., VASINA, V., POLUZZI, E. & DE PONTI, F. (2004). Antibacterial macrolides: a drug class with a complex pharmacological profile. *Pharmacol. Res.*, **50**, 211–222.

INUI, A., ASAKAWA, A., BOWERS, C.Y., MANTOVANI, G., LAVIANO, A., MEGUID, M.M. & FUJIMIYA, M. (2004). Ghrelin, appetite, and gastric motility: the emerging role of the stomach as an endocrine organ. *FASEB J.*, **18**, 439–456.

TAKESHITA, E., MATSUURA, B., DONG, M., MILLER, L.J., MATSUI, H., & ONJI, M. (2006). Molecular characterization and distribution of motilin family receptors in the human gastrointestinal tract. *J. Gastroenterol.*, **41**, 223–230.

References:

DEPOORTERE, I. & PEETERS, T.L. (1995). *Regul. Pept.*, **55**, 227–235.

FEIGHNER, S.D. *et al.* (1999). *Science*, **284**, 2184–2188.

LARTEY, P.A. *et al.* (1995). *J. Med. Chem.*, **38**, 1793–1798.

MCKEE, K.K. *et al.* (1997). *Genomics*, **46**, 426–434.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Neuromedin U

Overview: Neuromedin U receptors (provisional nomenclature) are activated by the endogenous 25 aa peptide neuromedin U (NMU), originally isolated from pig spinal cord (Minamino *et al.*, 1985). In humans, NMU appears to be the sole product of a precursor (ENSG00000109255) showing a broad tissue distribution, but which is expressed at highest levels in the upper gastrointestinal tract, CNS, bone marrow and fetal liver. Shorter versions of NMU are found in some species, being derived at least in some instances from the proteolytic cleavage of the longer NMU. Despite species differences in NMU structure, the C-terminal region (particularly the C-terminal pentapeptide) is highly conserved and contains biological activity. Neuromedin S (NMS) has also been identified as an endogenous agonist (Mori *et al.*, 2005). NMS is a 36 aa product of a precursor protein derived from a single gene (ENSG00000204640) and contains an amidated C-terminal heptapeptide identical to NMU. NMS appears to activate NMU receptors with equivalent potency to NMU.

Nomenclature	NMU ₁	NMU ₂
Other names	GPR66, FM3 (Hedrick <i>et al.</i> , 2000; Kojima <i>et al.</i> , 2000; Szekeres <i>et al.</i> , 2000)	FM4 (Howard <i>et al.</i> , 2000), TGR1 (Hosoya <i>et al.</i> , 2000)
Ensembl ID	ENSG00000171596	ENSG00000132911
Principal transduction	G _{q/11} (Hedrick <i>et al.</i> , 2000)	G _{q/11} (Hosoya <i>et al.</i> , 2000)
Probes	[¹²⁵ I]-NMU-25, Cy3B-tagged NMU-8 (Brighton <i>et al.</i> , 2004a)	—

NMU receptors couple predominantly to G_{q/11}, although there is evidence of coupling to G_{i/o} (see Hosoya *et al.*, 2000; Brighton *et al.*, 2004a).

Abbreviations: NMS, neuromedin S; NMU, neuromedin U

Further Reading:

BRIGHTON, P.J., SZEKERES, P.G. & WILLARS, G.B. (2004b). Neuromedin U and its receptors: structure, function, and physiological roles. *Pharmacol. Rev.*, **56**, 231–248.

References:

BRIGHTON, P.J. *et al.* (2004a). *Mol. Pharmacol.*, **66**, 1544–1556.

FUJII, R. *et al.* (2000). *J. Biol. Chem.*, **275**, 21068–21074.

HEDRICK, J.A. *et al.* (2000). *Mol. Pharmacol.*, **58**, 870–875.

HOSOYA, M. *et al.* (2000). *J. Biol. Chem.*, **275**, 29528–29532.

HOWARD, A.D. *et al.* (2000). *Nature*, **406**, 70–74.

KOJIMA, M. *et al.* (2000). *Biochem. Biophys. Res. Commun.*, **276**, 435–438.

MINAMINO, N. *et al.* (1985). *Biochem. Biophys. Res. Commun.*, **130**, 1078–1085.

MORI, K. *et al.* (2005). *EMBO J.*, **24**, 325–335.

RADDATZ, R. *et al.* (2000). *J. Biol. Chem.*, **275**, 32452–32459.

SHAN, L.X. *et al.* (2000). *J. Biol. Chem.*, **275**, 39482–39486.

SZEKERES, P.G. *et al.* (2000). *J. Biol. Chem.*, **275**, 20247–20250.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Neuropeptide Y

Overview: Neuropeptide Y (NPY) receptors (nomenclature agreed by NC-IUPHAR on Neuropeptide Y Receptors, see Michel *et al.*, 1998) are activated by the endogenous peptides NPY, NPY-(3-36), peptide YY (PYY), PYY-(3-36) and pancreatic polypeptide (PP). The y6 receptor is a functional gene product in mouse, absent in rat, but contains a frame-shift mutation in primates producing a truncated nonfunctional gene (Gregor *et al.*, 1996). Many of the agonists exhibit differing degrees of selectivity dependent on the species examined. For example, the relative potency of PP is greater at the rat Y₄ receptor than at the human receptor (Eriksson *et al.*, 1998). In addition, many agonists lack selectivity for individual subtypes, but can exhibit comparable potency against pairs of NPY receptor subtypes, or have not been examined for activity at all subtypes. [¹²⁵I]-PYY or [¹²⁵I]-NPY can be used to label Y₁, Y₂, Y₅ and Y₆ subtypes non-selectively, while [¹²⁵I]-[cPP(1-7),NPY(19-23),Ala³¹,Aib³²,Glu³⁴]hPP may be used to label Y₅ receptors preferentially.

Nomenclature	Y ₁	Y ₂	Y ₄	Y ₅	Y ₆
Other names	—	—	PP ₁	—	Y ₅ , PP ₂ , Y _{2B}
Ensembl ID	ENSG00000164128	ENSG00000185149	ENSG00000169556	ENSG00000164129	ENSG00000159279
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Rank order of potency	NPY ≥ PYY ≥ PP	NPY ≥ PYY ≥ PP	PP > NPY = PYY	NPY ≥ PYY ≥ PP	NPY = PYY > PP
Selective agonists	[Leu ³¹ ,Pro ³⁴]NPY, [Pro ³⁴]NPY, [Leu ³¹ ,Pro ³⁴]PYY, [Pro ³⁴]PYY	NPY-(3-36), PYY-(3-36)	PP	[Ala ³¹ ,Aib ³²]NPY (Cabrele <i>et al.</i> , 2000)	—
Selective antagonists	BIBO3304 (9.5, Wieland <i>et al.</i> , 1998), BIBP3226 (8.2, Gerald <i>et al.</i> , 1996)	BIIE0246 (8.5, Doods <i>et al.</i> , 1999), JNJ5207787 (Bonaventure <i>et al.</i> , 2004)	—	L152804 (7.6, Kanatani <i>et al.</i> , 2000)	—
Probes	[¹²⁵ I]-[Leu ³¹ ,Pro ³⁴]NPY, [³ H]-BIBP3226 (2.1 nM)	[¹²⁵ I]-PYY-(3-36)	[¹²⁵ I]-PP	[¹²⁵ I]-[cPP(1-7),NY4(19-23),Ala ³¹ ,Aib ³² ,Glu ³⁴]HPP (Dumont <i>et al.</i> , 2004)	—

The Y₁ agonists indicated are selective relative to Y₂ receptors. BIBP3226 is selective relative to Y₂, Y₄ and Y₅ receptors (Gerald *et al.*, 1996). NPY-(13-36) is Y₂ selective relative to Y₁ and Y₅ receptors. PYY-(3-36) is Y₂ selective relative to Y₁ receptors.

Abbreviations: **BIBO3304**, (*R*)-*N*-([4-{aminocarbonylaminoethyl}-phenyl]methyl)-*N*-(diphenylacetyl)-argininamide trifluoroacetate; **BIBP3226**, *R*-*N*-(diphenylacetyl)-*N*-(4-hydroxyphenyl)methyl-argininamide; **BIIE0246**, (*S*)-*N*-(1-[2-(4-[(*r,s*)-5,11-dihydro-6(6*H*)-oxodibenz[*b,e*]azepin-11-yl]-1-piperazinyl)-2-oxoethyl]cyclopentyl]acetyl)-*N*-(2-[1,2-dihydro-3,5(4*H*)-dioxo-1,2-diphenyl-3*H*-1,2,4-triazol-4-yl]ethyl)-argininamide; **JNJ5207787**, *N*-(1-acetyl-2,3-dihydro-1*H*-indol-6-yl)-3-(3-cyano-phenyl)-*N*-[1-(2-cyclopentylethyl)piperidin-4-yl]acrylamide; **L152804**, 2-(3,3-dimethyl-1-oxo-4*H*-1*H*-xanthen-9-yl)-5,5-dimethyl-cyclohexane-1,3-dione

Further Reading:

- LARHAMMAR, D. & SALANECK E. (2004). Molecular evolution of NPY receptor subtypes. *Neuropeptides*, **38**, 141–151.
- MICHEL, M.C. (2004). Neuropeptide Y and related peptides. In: *Handbook of Experimental Pharmacology*, Vol. **162**. Springer Verlag: Heidelberg.
- MICHEL, M.C., BECK-SICKINGER, A., COX, H., DOODS, H.N., HERZOG, H., LARHAMMAR, D., QUIRION, R., SCHWARTZ, T. & WESTFALL, T. (1998). International Union of Pharmacology XVI. Recommendations for the nomenclature of neuropeptide Y, peptide YY, and pancreatic polypeptide receptors. *Pharmacol. Rev.*, **50**, 143–150.
- NYGAARD, R., NIELBO, S., SCHWARTZ, T.W. & POULSEN, F.M. (2006). The PP-fold solution structure of human polypeptide YY and human PYY3-36 as determined by NMR. *Biochemistry*, **45**, 8350–8357.
- PARKER, E., VAN HECK, M. & STAMFORD, A. (2002). Neuropeptide Y receptors as targets for anti-obesity drug development: perspective and current status. *Eur. J. Pharmacol.*, **440**, 173–187.
- PEDRAZZINI, T. (2004). Importance of NPY Y1 receptor-mediated pathways: assessment using NPY Y1 receptor knockouts. *Neuropeptides*, **38**, 267–275.
- RAMOS, E.J., MEGUID, M.M., CAMPOS, A.C. & COELHO, J.C. (2005). Neuropeptide Y, a-melanocyte-stimulating hormone, and monoamines in food intake regulation. *Nutrition*, **21**, 269–279.

References:

- BONAVENTURE, P. *et al.* (2004). *J. Pharmacol. Exp. Ther.* **308**, 1130–1137.
- CABRELE, C. *et al.* (2000). *J. Biol. Chem.*, **275**, 36043–36048.
- DOODS, H. *et al.* (1999). *Eur. J. Pharmacol.*, **384**, R3–R5.
- DUMONT, Y. *et al.* (2004). *Neuropeptides* **38**, 163–174.
- ERIKSSON, H. *et al.* (1998). *Reg. Peptides*, **75–76**, 29–37.
- GERALD, C. *et al.* (1996). *Nature*, **382**, 168–171.
- GREGOR, P. *et al.* (1996). *J. Biol. Chem.*, **271**, 27776–27781.
- KANATANI, A. *et al.* (2000). *Biochem. Biophys. Res. Commun.*, **272**, 169–173.
- WIELAND, H.A. *et al.* (1998). *Br. J. Pharmacol.*, **125**, 549–555.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Neuropeptides B and W

Overview: The neuropeptide BW receptor 1 (NPBW1, provisional nomenclature) is activated by the two 23-amino-acid peptides, neuropeptide W (NPW-23) and neuropeptide B (NPB-23) (Shimomura *et al.*, 2002; Fujii *et al.*, 2002). C-terminally extended forms of the peptides (NPW-30 and NPB-29) also activate NPBW1 (Brezillon *et al.*, 2003). Unique to both forms of NPB is the *N*-terminal bromination of the first tryptophan residue. des-Br-NPB-23 and des-Br-NPB-29 were not found to be major components of bovine hypothalamic tissue extracts. The NPBW2 receptor is activated by the short and C-terminal extended forms of NPB and NPW (Brezillon *et al.*, 2003).

Nomenclature	NPBW1	NPBW2
Other names	GPR7	GPR8
Ensembl ID	ENSG00000183729	ENSG00000125522
Principal transduction	G _{i/o} (Mazzocchi <i>et al.</i> , 2005)	G _{i/o} (Mazzocchi <i>et al.</i> , 2005)
Rank order of potency	NPB-29 > NPB-23 > NPW-23 > NPW-30 (Brezillon <i>et al.</i> , 2003)	NPW-23 > NPW-30 > NPB-29 > NPB-23 (Brezillon <i>et al.</i> , 2003)
Probes	[¹²⁵ I]-NPW-23 (0.44 nM, Singh <i>et al.</i> , 2004)	[¹²⁵ I]-NPW-23

Potency measurements were conducted with heterologously-expressed receptors with a range of 0.14–0.57 nM (NPBW1) and 0.98–21 nM (NPBW2).

Further Reading:

LEE, D.K., GEORGE, S.R. & O'DOWD, B.F. (1999). Two related G protein-coupled receptors: the distribution of GPR7 in rat brain and the absence of GPR8 in rodents. *Mol. Brain Res.*, **23**, 96–103.

LEE, D.K., GEORGE, S.R. & O'DOWD, B.F. (2003). Continued discovery of ligands for G protein-coupled receptors. *Life Sci.*, **74**, 293–297.

SINGH, G. & DAVENPORT, A.P. (2006). Neuropeptide B and W: neurotransmitters in an emerging G-protein-coupled system. *Br. J. Pharmacol.*, **148**, 1033–1041.

References:

BREZILLON, S. *et al.* (2003). *J. Biol. Chem.*, **278**, 776–783.

FUJII, R. *et al.* (2002). *J. Biol. Chem.*, **277**, 34010–34016.

MAZZOCCHI, G. *et al.* (2005). *J. Clin. Endocrinol. Metab.*, **90**, 3466–3471.

SHIMOMURA, Y. *et al.* (2002). *J. Biol. Chem.*, **277**, 35826–35832.

SINGH, G. *et al.* (2004). *Brain Res.*, **1017**, 222–226.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Neurotensin

Overview: Neurotensin receptors (provisional nomenclature) are activated by the endogenous tridecapeptide neurotensin (pGlu-Leu-Tyr-Glu-Asn-Lys-Pro-Arg-Arg-Pro-Tyr-Ile-Leu) derived from a precursor (ENSG00000133636), which also generates neuromedin N, an agonist at the NTS2 receptor. A nonpeptide antagonist, SR142948A, shows high affinity ($pK_i \sim 9$) at both NTS1 and NTS2 receptors (Gully *et al.*, 1997). [^3H]-Neurotensin and [^{125}I]-neurotensin may be used to label NTS1 and NTS2 receptors at 0.1–0.3 and 3–5 nM concentrations, respectively.

Nomenclature	NTS1	NTS2
Other names	High-affinity neurotensin receptor, NTRH, NTR-1, NT ₁	Low-affinity neurotensin receptor, NTRL, NTR-1, NT ₂
Ensembl ID	ENSG00000101188	ENSG00000169006
Principal transduction	G _{q/11}	G _{q/11}
Rank order of potency	Neurotensin > neuromedin N (Hermans <i>et al.</i> , 1997)	Neurotensin = neuromedin N (Mazella <i>et al.</i> , 1996)
Selective agonists	JMV449 (Souaze <i>et al.</i> , 1997)	Levocabastine (Mazella <i>et al.</i> , 1996)
Selective antagonists	SR48692 (7.5–8.2; Gully <i>et al.</i> , 1997)	—
Probes	[^3H]-SR48692 (3.4 nM; Labbe-Jullie <i>et al.</i> , 1995)	—

Neurotensin appears to be a low-efficacy agonist at the NTS2 receptor (Vita *et al.*, 1998). An additional protein, provisionally termed NTS3 (also known as NTR3, gp95 and sortilin; ENSG00000134243), has recently been suggested to bind lipoprotein lipase and mediate its degradation (Nielsen *et al.*, 1999). It has been reported to interact with the NTS1 receptor (Martin *et al.*, 2002) and has been implicated in hormone trafficking and/or neurotensin uptake.

Abbreviations: **JMV449**, *H*-Lys ψ [(CH₂NH)-Lys-Pro-Tyr-Ile-Leu]; **SR142948A**, 2-([5-{2,6-dimethoxyphenyl}-1-{4-(*N*-[3-dimethylaminopropyl]-*N*-methylcarbamoyl)-2-isopropylphenyl}-1*H*-pyrazole-3-carbonyl]amino)adamantane-2-carboxylic acid hydrochloride; **SR48692**, 2-([1-{7-chloro-4-quinolinyl}-5-{2,6-dimethoxyphenyl}-pyrazol-3-yl]carboxylamino)tricyclo(3.3.1.1.1.[3.7])decan-2-carboxylic acid

Further Reading:

- BEROD, A. & ROSTENE, W. (2002). Neurotensin: an endogenous psychostimulant? *Curr. Opin. Pharmacol.*, **2**, 93–98.
- BINDER, E.B., KINKEAD, B., OWENS, M.J. & NEMEROFF, C.B. (2001). Neurotensin and dopamine interactions. *Pharmacol. Rev.*, **53**, 453–486.
- BOULES, M., FREDRICKSON, P. & RICHELSON, E. (2003). Current topics: brain penetrating neurotensin analog. *Life Sci.*, **73**, 2785–2792.
- DOBNER, P.R. (2005). Multitasking with neurotensin in the central nervous system. *Cell Mol. Life Sci.*, **62**, 1946–1963.
- DOBNER, P.R., DEUTCH, A.Y. & FADEL, J. (2003). Neurotensin: dual roles in psychostimulant and antipsychotic drug responses. *Life Sci.*, **73**, 801–811.
- KINKEAD, B. & NEMEROFF, C.B. (2002). Neurotensin: an endogenous antipsychotic? *Curr. Opin. Pharmacol.*, **2**, 99–103.
- KINKEAD, B. & NEMEROFF, C.B. (2004). Neurotensin, schizophrenia, and antipsychotic drug action. *Int. Rev. Neurobiol.*, **59**, 327–349.
- MAZELLA, J. & VINCENT, J.P. (2006). Internalization and recycling properties of neurotensin receptors. *Peptides*, **27**, 2488–2492.
- MCMAHON, B.M., BOULES, M., WARRINGTON, L. & RICHELSON, E. (2002). Neurotensin analogs indications for use as potential antipsychotic compounds. *Life Sci.*, **70**, 1101–1119.
- RICHELSON, E., BOULES, M. & FREDRICKSON, P. (2003). Neurotensin agonists: possible drugs for treatment of psychostimulant abuse. *Life Sci.*, **73**, 679–690.
- TYLER-MCMAHON, B.M., BOULES, M. & RICHELSON, E. (2000). Neurotensin: peptide for the next millennium. *Regul. Pept.*, **93**, 125–136.
- VINCENT, J.P., MAZELLA, J. & KITABGI, P. (1999). Neurotensin and neurotensin receptors. *Trends Pharmacol. Sci.*, **20**, 302–309.

References:

- GULLY, D. *et al.* (1997). *J. Pharmacol. Exp. Ther.*, **280**, 802–812.
- HERMANS, E. *et al.* (1997). *Br. J. Pharmacol.*, **121**, 1817–1823.
- LABBE-JULLIE, C. *et al.* (1995). *Mol. Pharmacol.*, **47**, 1050–1056.
- MARTIN, S. *et al.* (2002). *Gastroenterology*, **123**, 1135–1143.
- MARTIN, S. *et al.* (2003). *J. Neurosci.*, **23**, 1198–1205.
- MAZELLA, J. *et al.* (1996). *J. Neurosci.*, **16**, 5613–5620.
- NIELSEN, M.S. *et al.* (1999). *J. Biol. Chem.*, **274**, 8832–8836.
- PETERSEN, C.M. *et al.* (1997). *J. Biol. Chem.*, **272**, 3599–3605.
- SOUAZE, F. *et al.* (1997). *J. Biol. Chem.*, **272**, 10087–10094.
- VITA, N. *et al.* (1998). *Eur. J. Pharmacol.*, **360**, 265–272.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Opioid and opioid-like

Overview: Opioid and opioid-like receptors are activated by the endogenous peptides [Met]enkephalin (met), [Leu]enkephalin (leu), β -endorphin (β -end), α -neodynorphin, dynorphin A (dynA), dynorphin B (dynB), Big dynorphin (Big dyn), nociceptin/orphanin FQ (N/OFQ), and endomorphin-1 and -2, although several other opioid-like peptides are found in the CNS. The Greek letter names for the opioid receptors, μ , κ , and δ , are well established and, despite digressions into other nomenclatures (Dhawan et al., 1996), IUPHAR appears to now consider these original names most appropriate (Foord et al., 2005). The human N/OFQ receptor is considered 'opioid-related' rather than opioid as it exhibits a high degree of structural homology with the conventional opioid receptors (Mollereau et al., 1994), but displays a distinct pharmacology.

Nomenclature	Delta opioid receptor	Kappa opioid receptor	Mu opioid receptor	N/OFQ receptor
Preferred abbreviation	δ	κ	μ	NOP
Other names	OP ₁ , DOP, DOR	OP ₂ , KOP, KOR	OP ₃ , MOP, MOR	ORL1, OP ₄
Ensembl ID	ENSG00000116329	ENSG00000082556	ENSG00000112038	ENSG00000125510
Principal transduction	G _{i/o}	G _{i/o}	G _{i/o}	G _{i/o}
Rank order of potency	β -End = leu = met > dynA	Big dyn > dynA >> β -end > leu > met	β -End > dynA > met = leu	N/OFQ >> dynA
Selective agonists	DPDPE (Mosberg et al., 1983), DSBULET (Delay-Goyet et al., 1988), [DAla ²]deltorphin I or II (Erspamer et al., 1989), SNC80 (Bilsky et al., 1995)	U69593 (Lahti et al., 1985), CI977 (Hunter et al., 1990), Salvinorin A (Roth et al., 2002)	Endomorphin-1 and -2 (Zadina et al., 1997), morphine (Goldstein & Naidu, 1989), DAMGO (Handa et al., 1981), sufentanil (Yeadon & Kitchen, 1988), PL017 (Costa et al., 1992)	N/OFQ, N/OFQ-(1-13)-NH ₂ (Guerrini et al., 1997), Ro646198 (Jenck et al., 2000), UFP102 (Carra' et al., 2005)
Selective antagonists	Naltrindole (Portoghese et al., 1988), naltriben (Sofuoglu et al., 1991)	Nor-binaltorphimine (Portoghese et al., 1987)	CTAP (Pelton et al., 1986), CTOP (Hawkins et al., 1989)	J113397 (8.3, Kawamoto et al., 1999), SB612111 (8.5, Zaratini et al., 2004), UFP101 (7.2, Calo' et al., 2002)
Probes	[³ H]-DPDPE (Goldstein & Naidu, 1989), [³ H]-naltrindole (Yamamura et al., 1992), [³ H]-Deltorphin II (Gomes et al., 2000), [³ H]-naltriben (Lever & Scheffel, 1998)	[³ H]-U69593 (Lahti et al., 1985), [³ H]-CI977 (Simonin et al., 2001)	[³ H]-DAMGO (Goldstein & Naidu, 1989), [³ H]-PL017 (Hawkins et al., 1987)	[³ H]-N/OFQ (Dooley & Houghten, 1996), [³ H]-Leu-N/OFQ, [¹²⁵ I]-Tyr ¹⁴ -N/OFQ

Subtypes of μ (μ 1, μ 2), δ (δ 1, δ 2) and κ (κ 1, κ 2, κ 3) receptor have been proposed based primarily on binding studies with poorly selective ligands or *in vivo* studies. Only three naloxone-sensitive opioid receptors have been cloned, and most evidence suggests that these receptors undergo minimal alternative splicing, and certainly none that can be shown to give rise to any of the subtypes proposed. There is at present no molecular basis for any of the opioid receptor subtypes, although it remains possible that they are formed by heterodimerization of opioid receptors with each other or with other 7TM receptors (Jordan & Devi, 1999). A distinct met-enkephalin receptor lacking structural resemblance to the opioid receptors listed has been identified (ENSG00000060491) and termed an opioid growth factor receptor (see Zagon et al., 2002).

Abbreviations: CI977, (5*r*)-(5*x*,7*x*,8*β*)-(–)-*N*-methyl-*N*-(7-[1-pyrrolidinyl]-1-oxaspiro[4.5]dec-8-yl)-4-benzofuranacetamide hydrochloride; CTAP, D-Phe-cyc[Cys-Tyr-D-Trp-Arg-Thr-Pen]-Thr-NH₂; CTOP, D-Phe-Cys-Tyr-D-Trp-Oru-Thr-Pen-Thr-NH₂; DAMGO, Tyr-DAla-Gly-[NMePhe]-NH(CH₂)₂; DPDPE, cyc[DPen², DPen⁵]enkephalin; DSBULET, Tyr-DSer(OtBu)-Gly-Phe-Leu-Thr; ICI174864, *N,N*-diallyl-Tyr-Aib-Phe-Leu-OH (Aib is aminoisobutyric acid); J113397, 1-[(3*r*,4*r*)-1-cyclooctylmethyl-3-hydroxymethyl-4-piperidyl]-3-ethyl-1,3-dihydro-2*H*-benzimidazol-2-one; PL017, [N-MePhe³,DPro⁴]morphiceptin; Ro646198, (1*S*,3*aS*)-8-(2,3,3*a*,4,5,6-hexahydro-1*H*-phenalen-1-yl)-1-phenyl-1,3,8-triazaspiro[4.5]decan-4-one; SB612111, (–)-*cis*-1-methyl-7-[[4-(2,6-dichlorophenyl)piperidin-1-yl]methyl]-6,7,8,9-tetrahydro-5*H*-benzocyclohept-5-ol; SNC80, (+)-4-[(α R)- α -(2*S*,5*R*)-4-allyl-2,5-dimethyl-1-piperazinyl]-3-methoxybenzyl]-*N,N*-diethylbenzamide; U69593, 5*x*,7*x*,8*β*-(–)-*N*-methyl-*N*-(7-[1-pyrrolidinyl]-1-oxaspiro[4.5]dec-8-yl)benzene acetamide; UFP101, [Nphe¹,Arg¹⁴,Lys¹⁵]nociceptin-NH₂; UFP-102, [(pF)Phe⁴,Arg¹⁴,Lys¹⁵]N/OFQ-NH₂

Further Reading:

- BAILEY, C.P. & CONNOR, M. (2005). Opioids: cellular mechanisms of tolerance and physical dependence. *Curr. Opin. Pharmacol.*, **5**, 60–68.
- CALO', G., RIZZI, A., BIGONI, R., GUERRINI, R., SALVADORI, S. & REGOLI, D. (2002). Pharmacological profile of nociceptin/orphanin FQ receptors. *Clin. Exp. Pharmacol. Physiol.*, **29**, 223–228.
- FIELDS, H. (2004). State-dependent opioid control of pain. *Nat. Rev. Neurosci.*, **5**, 565–575.
- HEINRICHER, M.M. (2003). Orphanin FQ/nociceptin: from neural circuitry to behavior. *Life Sci.*, **73**, 813–822.
- HOLZER, P. (2004). Opioids and opioid receptors in the enteric nervous system: from a problem in opioid analgesia to a possible new prokinetic therapy in humans. *Neurosci. Lett.*, **361**, 192–195.
- IKEDA, K., IDE, S., HAN, W., HAYASHIDA, M., UHL, G.R. & SORA, I. (2005). How individual sensitivity to opiates can be predicted by gene analyses. *Trends Pharmacol. Sci.*, **26**, 311–317.
- KIEFFER, B.L. & GAVERIAUX-RUFF, C. (2002). Exploring the opioid system by gene knockout. *Prog. Neurobiol.*, **66**, 285–306.
- KREEK, M.J., BART, G., LILLY, C., LAFORGE, K.S. & NIELSEN, D.A. (2005). Pharmacogenetics and human molecular genetics of opiate and cocaine addictions and their treatments. *Pharmacol. Rev.*, **57**, 1–26.
- LEVAC, B.A., O'DOWD, B.F. & GEORGE, S.R. (2002). Oligomerization of opioid receptors: generation of novel signaling units. *Curr. Opin. Pharmacol.*, **2**, 76–81.
- LIU-CHEN, L.Y. (2004). Agonist-induced regulation and trafficking of κ opioid receptors. *Life Sci.*, **75**, 511–536.
- LOTSCH, J. & GEISSLINGER, G. (2005). Are mu-opioid receptor polymorphisms important for clinical opioid therapy? *Trends Mol. Med.*, **11**, 82–89.
- MALINOWSKA, B., GODLEWSKI, G. & SCHLICKER, E. (2002). Function of nociceptin and opioid OP4 receptors in the regulation of the cardiovascular system. *J. Physiol. Pharmacol.*, **53**, 301–324.
- MOGIL, J.S. & PASTERNAK, G.W. (2001). The molecular and behavioral pharmacology of the orphanin FQ/nociceptin peptide and receptor family. *Pharmacol. Rev.*, **53**, 381–415.
- PHILIP, A.E., POUPAERT, J.H. & MCCURDY, C.R. (2005). Opioid receptor-like 1 (ORL1) molecular 'road map' to understanding ligand interaction and selectivity. *Curr. Top. Med. Chem.*, **5**, 325–340.

- SADEE, W., WANG, D. & BILSKY, E.J. (2005). Basal opioid receptor activity, neutral antagonists, and therapeutic opportunities. *Life Sci.*, **76**, 1427–1437.
- SMITH, A.P. & LEE, N.M. (2003). Opioid receptor interactions: local and nonlocal, symmetric and asymmetric, physical and functional. *Life Sci.*, **73**, 1873–1893.
- TEGEDER, I. & GEISLINGER, G. (2004). Opioids as modulators of cell death and survival – unraveling mechanisms and revealing new indications. *Pharmacol. Rev.*, **56**, 351–369.
- VON ZASTROW, M. (2004). A cell biologist's perspective on physiological adaptation to opiate drugs. *Neuropharmacology*, **47** (Suppl 1), 286–292.
- WALDHOER, M., BARTLETT, S.E. & WHISTLER, J.L. (2004). Opioid receptors. *Annu. Rev. Biochem.*, **73**, 953–990.
- WEI, L.N. & LOH, H.H. (2002). Regulation of opioid receptor expression. *Curr. Opin. Pharmacol.*, **2**, 69–75.
- WOLLEMAN, M. & BENYHE, S. (2004). Non-opioid actions of opioid peptides. *Life Sci.*, **75**, 257–270.
- ZAGON, I.S., VERDERAME, M.F. & MCLAUGHLIN, P.J. (2002). The biology of the opioid growth factor receptor (OGFr). *Brain Res. Rev.*, **38**, 351–376.
- ZAVERI, N. (2003). Peptide and nonpeptide ligands for the nociceptin/orphanin FQ receptor ORL1: research tools and potential therapeutic agents. *Life Sci.*, **73**, 663–678.

References:

- BILSKY, E. et al. (1995). *J. Pharmacol. Exp. Ther.*, **273**, 359–366.
- CALO', G. et al. (2002). *Br. J. Pharmacol.*, **136**, 303–311.
- CARRA', G. et al. (2005). *J. Pharmacol. Exp. Ther.*, **312**, 1114–1123.
- COSTA, E.M. et al. (1992). *Life Sci.*, **50**, 73–81.
- COTTON, R. et al. (1984). *Eur. J. Pharmacol.*, **97**, 331–332.
- DELAY-GOYET, P. et al. (1988). *J. Biol. Chem.*, **263**, 4124–4130.
- DHAWAN, B.N. et al. (1996). *Pharmacol. Rev.*, **48**, 567–592.
- DOOLEY, C.T. & HOUGHTEN, R.A. (1996). *Life Sci.*, **59**, L23–L29.
- ERSPAMER, V. et al. (1989). *Proc. Natl. Acad. Sci. U.S.A.*, **86**, 5188–5192.
- FOORD, S.M. et al. (2005). *Pharmacol. Rev.*, **57**, 279–288.
- GOLDSTEIN, A. & NAIDU, A. (1989). *Mol. Pharmacol.*, **36**, 265–272.
- GUERRINI, R. et al. (1997). *J. Med. Chem.*, **40**, 1789–1793.
- HANDA, B.K. et al. (1981). *Eur. J. Pharmacol.*, **70**, 531–540.
- HAWKINS, K.N. et al. (1987). *Eur. J. Pharmacol.*, **133**, 351–352.
- HAWKINS, K.N. et al. (1989). *J. Pharmacol. Exp. Ther.*, **248**, 73–80.
- HUNTER, J. et al. (1990). *Br. J. Pharmacol.*, **101**, 183–189.
- JENCK, F. et al. (2000). *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 4938–4943.
- JORDAN, B.A. & DEVI, L.A. (1999). *Nature*, **399**, 697–700.
- KAWAMOTO, H. et al. (1999). *J. Med. Chem.*, **42**, 5061–5063.
- LAHTI, R.A. et al. (1985). *Eur. J. Pharmacol.*, **109**, 281–284.
- LEVER, J.R. & SCHEFFEL, T.U. (1998). *Eur. J. Pharmacol.*, **350**, 335–344.
- MOLLEREAU, C. et al. (1994). *FEBS Lett.*, **341**, 33–38.
- MOSBER, G. et al. (1983). *Proc. Natl. Acad. Sci. U.S.A.*, **80**, 5871–5874.
- PELTON, J.T. et al. (1986). *J. Med. Chem.*, **29**, 2370–2375.
- PORTOGHESE, P.S. et al. (1988). *Eur. J. Pharmacol.*, **146**, 185–186.
- PORTOGHESE, P.S. et al. (1987). *Life Sci.*, **40**, 1287–1292.
- ROTH, B.L. et al. (2002). *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 11934–11939.
- SIMONIN, F. et al. (2001). *Eur. J. Pharmacol.*, **414**, 189–195.
- SOFUOGLU, M. et al. (1991). *J. Pharmacol. Exp. Ther.*, **257**, 676–680.
- YAMAMURA, H.I. et al. (1992). *Life Sci.*, **50**, PL119–PL124.
- YEADON, M. & KITCHEN, I. (1988). *Neuropharmacology*, **27**, 345–348.
- ZADINA, J.E. et al. (1997). *Nature*, **386**, 499–502.
- ZARATIN, P. et al. (2004). *J. Pharmacol. Exp. Ther.*, **308**, 454–461.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Orexin

Overview: Orexin receptors (provisional nomenclature) are activated by the endogenous polypeptides orexin-A and orexin-B (also known as hypocretin-1 and -2; 33 and 28 aa) derived from a common precursor, orexin (ENSG00000161610), by proteolytic cleavage (Sakurai *et al.*, 1998). Binding to both receptors may be accomplished with [¹²⁵I]-orexin A (Holmqvist *et al.*, 2001).

Nomenclature	OX₁	OX₂
Other names	Hypocretin receptor type 1	Hypocretin receptor type 2
Ensembl ID	ENSG00000121764	ENSG00000137252
Principal transduction	G _{q/11}	G _{q/11}
Rank order of potency	Orexin-A > orexin-B	Orexin-A = orexin-B
Selective agonists	—	[Ala ¹¹ ,D-Leu ¹⁵]orexin-B (Asahi <i>et al.</i> , 2003)
Selective antagonists	SB408124 (7.5, Langmead <i>et al.</i> , 2004), SB334867A (7.2-7.3, Smart <i>et al.</i> , 2001)	—

The *HCRT2* gene encoding the OX₂ receptor has been identified as a possible candidate for inherited narcolepsy (Chemelli *et al.*, 1999; Lin *et al.*, 1999; Siegel, 1999).

Abbreviations: **SB334867A**, 1-(2-methylbenzoxazol-6-yl)-3-[1,5]naphthyridin-4-yl-urea hydrochloride; **SB408124**, 1-(6,8-difluoro-2-methyl-quinolin-4-yl)-3-(4-dimethylamino-phenyl)-urea

Further Reading:

- BERRIDGE, C.W. & ESPANA, R.A. (2005). Hypocretins: waking, arousal, or action? *Neuron*, **46**, 696–698.
- EHMKE, H. & JUST, A. (2003). The orexins: linking circulatory control with behavior. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, **285**, R519–R521.
- HARRIS, G.C. & ASTON-JONES, G. (2006). Arousal and reward: a dichotomy in orexin function. *Trends Neurosci.*, **29**, 571–577.
- KIRCHGESSENER, A.L. (2002). Orexins in the brain-gut axis. *Endocr. Rev.*, **23**, 1–15.
- MIEDA, M. & YANAGISAWA, M. (2002). Sleep, feeding, and neuropeptides: roles of orexins and orexin receptors. *Curr. Opin. Neurobiol.*, **12**, 339–345.
- NISHINO, S. (2003). The hypocretin/orexin system in health and disease. *Biol. Psychiatry*, **54**, 87–95.
- OVEREEM, S., SCAMMELL, T.E. & LAMMERS, G.J. (2002). Hypocretin/orexin and sleep: implications for the pathophysiology and diagnosis of narcolepsy. *Curr. Opin. Neurol.*, **15**, 739–745.
- SIEGEL, J.M. (2004). Hypocretin (orexin): role in normal behavior and neuropathology. *Annu. Rev. Psychol.*, **55**, 125–148.
- SMART, D. & JERMAN, J.C. (2002). The physiology and pharmacology of the orexins. *Pharmacol. Ther.*, **94**, 51–61.
- SPINAZZI, R., ANDREIS, P.G., ROSSI, G.P. & NUSSDORFER, G.G. (2006). Orexins in the regulation of the hypothalamic-pituitary-adrenal axis. *Pharmacol. Rev.*, **58**, 46–57.
- SUTCLIFFE, J.G. & DE LECEA, L. (2002). The hypocretins: setting the arousal threshold. *Nat. Rev. Neurosci.*, **3**, 339–349.
- TAHERI, S., ZEITZER, J.M. & MIGNOT, E. (2002). The role of hypocretins (orexins) in sleep regulation and narcolepsy. *Annu. Rev. Neurosci.*, **25**, 283–313.
- TAYLOR, M.M. & SAMSON, W.K. (2003). The other side of the orexins: endocrine and metabolic actions. *Am. J. Physiol. Endocrinol. Metab.*, **284**, E13–E17.
- VOISIN, T., ROUET-BENZINEB, P., REUTER, N. & LABURTHER, M. (2003). Orexins and their receptors: structural aspects and role in peripheral tissues. *Cell Mol. Life Sci.*, **60**, 72–87.
- ZEITZER, J.M., NISHINO, S. & MIGNOT, E. (2006). The neurobiology of hypocretins (orexins), narcolepsy and related therapeutic interventions. *Trends Pharmacol. Sci.*, **27**, 368–374.

References:

- ASAHI, S. *et al.* (2003). *Bioorg. Med. Chem. Lett.*, **13**, 111–113.
- CHEMELLI, R.M. *et al.* (1999). *Cell*, **98**, 437–451.
- HOLMQVIST, T. *et al.* (2001). *Neurosci. Lett.*, **305**, 177–180.
- LANGMEAD, C.J. *et al.* (2004). *Br. J. Pharmacol.*, **141**, 340–346.
- LIN, L. *et al.* (1999). *Cell*, **98**, 365–376.
- SAKURAI, T. *et al.* (1998). *Cell*, **92**, 573–585.
- SIEGEL, J.M. (1999). *Cell*, **98**, 409–412.
- SMART, D. *et al.* (2001). *Br. J. Pharmacol.*, **132**, 1179–1182.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Oxoeicosanoid

Overview: Oxoeicosanoid receptors (OXE, nomenclature agreed by NC-IUPHAR on Oxoeicosanoid Receptors; Brink *et al.*, 2004) are activated by endogenous chemotactic eicosanoid ligands oxidised at the C-5 position, with 5-oxo-ETE the most potent agonist identified for this receptor.

Nomenclature	OXE
Other names	TG1019 (Hosoi <i>et al.</i> , 2002), R527 (Jones <i>et al.</i> , 2003)
Ensembl ID	ENSG00000162881
Principal transduction	G _{i/o} (O'Flaherty <i>et al.</i> , 2000; Hosoi <i>et al.</i> , 2002; Jones <i>et al.</i> , 2003; Hosoi <i>et al.</i> , 2005)
Rank order of potency	5-Oxo-ETE ≫ 5(S)-HpETE > 5(S)-HETE (Hosoi <i>et al.</i> , 2002; Jones <i>et al.</i> , 2003)
Selective agonists	5-Oxo-ETE
Probes	[³ H]-5-oxo-ETE (3.8 nM; O'Flaherty <i>et al.</i> , 1998)

Initial characterization of the heterologously expressed receptor suggested that polyunsaturated fatty acids, such as DHA and EPA, acted as receptor antagonists (Hosoi *et al.*, 2002).

Abbreviations: **DHA**, 4z,7z,10z,13z,16z,19z-docosahexaenoic acid, **EPA**, 5z,8z,11z,14z,17z-eicosapentaenoic acid, **OXE**, oxoeicosanoid; **5-oxo-ETE**, 5-oxo-6E,8z,11z,14z-eicosatetraenoic acid; **5(S)-HpETE**, 5(s)-hydroperoxy-6E,8z,11z,14z-eicosatetraenoic acid; **5(S)-HETE**, 5(s)-hydroxy-6E,8z,11z,14z-eicosatetraenoic acid

Further Reading:

BRINK, C., DAHLEN, S.E., DRAZEN, J., EVANS, J.F., HAY, D.W., ROVATI, G.E., SERHAN, C.N., SHIMIZU, T. & YOKOMIZO, T. (2004). International Union of Pharmacology XLIV. Nomenclature for the oxoeicosanoid receptor. *Pharmacol. Rev.*, **56**, 149–157.

JONES, C.E. (2005). The OXE receptor: a new therapeutic approach for asthma? *Trends Mol. Med.*, **11**, 266–270.

O'FLAHERTY, J.T., ROGERS, L.C., PAUMI, C.M., HANTGAN, R.R., THOMAS, L.R., CLAY, C.E., HIGH, K., CHEN, V.Q., WILLINGHAM, M.C., SMITHIRMAN, P.K., KUTE, T.E., RAO, A., CRAMER, S.D. & MORROW, C.S. (2005). 5-Oxo-ETE analogs and the proliferation of cancer cells. *Biochem. Biophys. Acta*, **1736**, 228–236.

POWELL, W.S. & ROKACH, J. (2005). Biochemistry, biology and chemistry of the 5-lipoxygenase product 5-oxo-ETE. *Prog. Lipid Res.*, **44**, 154–183.

STURM, G.J., SCHULIGOI, R., STURM, E.M., ROYER, J.F., LANG-LOIDOLT, D., STAMMBERGER, H., AMMAN, R., PESKAR, B.A. & HEINEMANN, A. (2005). 5-Oxo-6,8,11,14-eicosatetraenoic acid is a potent chemoattractant for human basophils. *J. Allergy Clin. Immunol.*, **116**, 1014–1019.

SUNDARAM, S. & GHOSH, J. (2006). Expression of 5-oxo-ETE receptor in prostate cancer cells: critical role in survival. *Biochem. Biophys. Res. Commun.*, **339**, 93–98.

References:

HOSOI, T. *et al.* (2002). *J. Biol. Chem.*, **277**, 31459–31465.

HOSOI, T. *et al.* (2005). *Biochem. Biophys. Res. Commun.*, **334**, 987–995.

JONES, C.E. *et al.* (2003). *Mol. Pharmacol.*, **63**, 471–477.

O'FLAHERTY, J.T. *et al.* (2000). *J. Immunol.*, **164**, 3345–3352.

O'FLAHERTY, J.T. *et al.* (1998). *J. Biol. Chem.*, **273**, 32535–32541.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

P2Y

Overview: P2Y receptors (provisional nomenclature as agreed by NC-IUPHAR Subcommittee on P2Y Receptors, Abbracchio *et al.*, 2003; 2006) are activated by the endogenous ligands ATP, ADP, UTP, UDP and UDP-glucose. The relationship of many of the cloned receptors to endogenously expressed receptors is not yet established and so it might be appropriate to use wording such as 'UTP-preferring (or ATP-, etc.) P2Y receptor' or 'P2Y₁-like', etc., until further, as yet undefined, corroborative criteria can be applied.

Nomenclature	P2Y₁	P2Y₂	P2Y₄	P2Y₆
Ensembl ID	ENSG00000169860	ENSG00000175591	ENSG000000	ENSG00000171361
Principal transduction	G _{q/11}	G _{q/11}	G _{q/11}	G _{q/11}
Rank order of potency	ADP > ATP	UTP = ATP	UTP > ATP (at rat recombinant receptors, UTP = ATP)	UDP ≫ UTP > ATP
Selective agonists	2-MeSADP, ADPβS, MRS2365 (Bourdon <i>et al.</i> , 2006)	UTPγS (Lazarowski <i>et al.</i> , 1996), Ap ₄ A (Castro <i>et al.</i> , 1992)	UTPγS (Lazarowski <i>et al.</i> , 1996)	UDP
Selective antagonists	MRS2279 (8.0, Waldo <i>et al.</i> , 2002), MRS2179 (7.0, Boyer <i>et al.</i> , 1996), PIT (6.8, Gao <i>et al.</i> , 2004), MRS2500 (8.8, Kim <i>et al.</i> , 2003)	—	ATP (6.2, Kennedy <i>et al.</i> , 2000)	MRS2578 (pIC ₅₀ 7.4, Mamedova <i>et al.</i> , 2004)
Probes	[³ H]-MRS2279 (8 nM, Waldo <i>et al.</i> , 2002) [³⁵ S]-ADPβS, [³⁵ S]-ATPαS, [³⁵ S]-dATPαS	—	—	—

Nomenclature	P2Y₁₁	P2Y₁₂	P2Y₁₃	P2Y₁₄
Other names	—	P2Y _{ADP} , P2T	GPR86, GPR94, SP174	KIAAA000001, gpr105
Ensembl ID	ENSG00000176130	ENSG00000169313	ENSG00000181631	ENSG00000174944
Principal transduction	G _s , G _{q/11}	G _{i/o}	G _{i/o}	G _{q/11}
Rank order of potency	ATP > UTP	ADP ≫ ATP	ADP ≫ ATP	UDP-glucose
Selective agonists	ARC67085 (Communi <i>et al.</i> , 1999), NAD ⁺ (Moreschi, <i>et al.</i> , 2006)	ADP, 2-MeSADP	—	UDP-glucose
Selective antagonists	—	ATP, ARL66096 (Humphries <i>et al.</i> , 1995)	MRS2211 (Kim <i>et al.</i> , 2005)	—

The recently described 'P2Y₁₅' receptor (Inbe *et al.*, 2004) appears not to be a genuine nucleotide receptor (see Abbracchio *et al.*, 2006), but rather responds to dicarboxylic acids (He *et al.*, 2004). Further P2Y-like receptors have been cloned from nonmammalian sources; a clone from chick brain, termed a p2y₃ receptor (ch328 aa Q98907), couples to the G_{q/11} family of G proteins and shows the rank order of potency ADP > UTP > ATP = UDP (Webb *et al.*, 1996a). In addition, human sources have yielded a clone with a preliminary identification of p2y5 (h328 P43657) and contradictory evidence of responses to ATP (King & Townsend-Nicholson, 2000; Webb *et al.*, 1996b). The clone p2y7 (h352 Q15722), originally suggested to be a P2Y receptor (Akbar *et al.*, 1996), has been shown to encode a leukotriene receptor (Yokomizo *et al.*, 1997). A P2Y receptor that was initially termed a p2y8 receptor (× 1537 P79928) has been cloned from *Xenopus laevis*; it shows the rank order of potency ADPβS > ATP = UTP = GTP = CTP = TTP = ITP > ATPγS and elicits a Ca²⁺-dependent Cl⁻ current in *Xenopus* oocytes (Bogdanov *et al.*, 1997). The clone termed p2y9 has recently been described as an LPA receptor (Noguchi *et al.*, 2003), while the p2y10 (AF000545) clone lacks functional data. Diadenosine polyphosphates also have effects on as yet uncloned P2Y-like receptors with the rank order of potency of Ap₄A > Ap₅A > Ap₃A, coupling via G_{q/11} (Castro *et al.*, 1992). P2Y-like receptors have recently been described on mitochondria (Belous *et al.*, 2004). CysLT₁ and CysLT₂ leukotriene receptors respond to nanomolar concentrations of UDP, although they are activated principally by leukotrienes C₄ and D₄ (Mellor *et al.*, 2001; 2003); Human and rat GPR17, which are structurally related to CysLT and P2Y receptors, are also activated by leukotrienes as well as UDP and UDP-glucose (Ciana *et al.*, 2006). Activity at the rat GPR17 is inhibited by submicromolar concentrations of MRS2179 and ARL6931 (Ciana *et al.*, 2006).

Abbreviations: **ARC67085**, 2-propylthio-β,γ-dichloromethylene-ATP; **2-MeSADP**, 2-methylthio-adenosine-5'-diphosphate; **2-MeSATP**, 2-methylthio-adenosine-5'-triphosphate; **ARL66096**, 2-propylthio-β,γ-difluoromethylene ATP (previously FPL66096); **ATPγS**, adenosine 5'-(3-thio)triphosphate **MRS2211**, pyridoxal-5'-phosphate-6-azo (2-chloro-5-nitrophenyl)-2,4-disulfonate; **MRS2179**, N⁶-methyl-2'-deoxyadenosine-3',5'-bisphosphate; **MRS2279**, 2-chloro-N⁶-methyl-(N)-methanocarba-2'-deoxyadenosine-3',5'-bisphosphate; **MRS2365**, (N)-methanocarba-2-MeSADP; **MRS2500**, N⁶-methyl-(N)-methanocarba-2'-deoxyadenosine-3',5-bisphosphate; **MRS2578**, N₄N''-1,4-butanediyl bis(N'-[3-isothiocynatophenyl]) thiourea; **PIT**, 2,2'-pyridylisatogentosylate

Further Reading:

ABBRACCHIO, M.P., BOEYNAEMS, J.M., BARNARD, E.A., BOYER, J.L., KENNEDY, C., MIRAS-PORTUGAL, M.T., KING, B.F., GACHET, C., JACOBSON, K.A., WEISMAN, G.A. & BURNSTOCK, G. (2003). Characterization of the UDP-glucose receptor (re-named here the P2Y₁₄ receptor) adds diversity to the P2Y receptor family. *Trends Pharmacol. Sci.*, **24**, 52–55.

ABBRACCHIO, M.P., BURNSTOCK, G., BOEYNAEMS, J.M., BARNARD, E.A., BOYER, J.L., KENNEDY, C., KNIGHT, G.E., FUMAGALLI, M., GACHET, C., JACOBSON, K.A., WEISMAN, G.A. (2006) International Union of Pharmacology LVIII: Update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol. Rev.*, **58**, 281–341.

BURNSTOCK, G. (2002). Potential therapeutic targets in the rapidly expanding field of purinergic signalling. *Clin. Med.*, **2**, 45–53.

BURNSTOCK, G. & KNIGHT, G.E. (2004). Cellular distribution and functions of P2 receptor subtypes in different systems. *Int. Rev. Cytol.*, **240**, 31–304.

JACOBSON, K.A., JARVIS, M.F. & WILLIAMS, M. (2002). Purine and pyrimidine (P2) receptors as drug targets. *J. Med. Chem.*, **45**, 4057–4093.

KOLES, L., FURST, S. & ILLES, P. (2005). P2X and P2Y receptors as possible targets of therapeutic manipulations in CNS illnesses. *Drug News Perspect.*, **18**, 85–101.

VON KÜGELGEN, I. (2006). Pharmacological profiles of cloned mammalian P2Y-receptor subtypes. *Pharmacol. Ther.*, **110**, 415–432.

References:

- ABBRACCHIO, M.P. *et al.* (2003). *Trends Pharmacol. Sci.*, **24**, 52–55.
- AKBAR, G.K.M. *et al.* (1996). *J. Biol. Chem.*, **271**, 18363–18367.
- BELOUS, A. *et al.* (2004). *J. Cell. Biochem.*, **92**, 1062–1073.
- BOGDANOV, Y.D. *et al.* (1997). *J. Biol. Chem.*, **272**, 12583–12590.
- BOURDON, D.M. *et al.* (2006). *J. Thromb. Haemost.*, **4**, 861–868.
- BOYER, J.L. *et al.* (1996). *Mol. Pharmacol.*, **50**, 1323–1329.
- CASTRO, E. *et al.* (1992). *Br. J. Pharmacol.*, **106**, 833–837.
- CIANA, P. *et al.* (2006). *EMBOJ*, **25**, 4615–4627.
- COMMUNI, D. *et al.* (1999). *Br. J. Pharmacol.*, **128**, 1199–1206.
- GAO, Z.G. *et al.* (2004). *Biochem. Pharmacol.*, **68**, 231–237.
- HE, W. *et al.* (2004). *Nature*, **429**, 188–193.
- INBE, H. *et al.* (2004). *J. Biol. Chem.*, **279**, 19790–19799.
- HUMPHRIES, R.G. *et al.* (1995). *Br. J. Pharmacol.*, **115**, 1110–1116.
- KENNEDY, C. *et al.* (2000). *Mol. Pharmacol.*, **57**, 926–931.
- KIM, H.S. *et al.* (2003). *J. Med. Chem.*, **46**, 4974–4987.
- KIM, Y.C. *et al.* (2005). *Biochem. Pharmacol.*, **70**, 266–274.
- KING, B.F. & TOWNSEND-NICHOLSON, A. (2000). *J. Auton. Nerv. System*, **81**, 164–170.
- LAZAROWSKI, E.R. *et al.* (1996). *Br. J. Pharmacol.*, **117**, 203–209.
- MAMEDOVA, L.K. *et al.* (2004). *Biochem. Pharmacol.*, **67**, 1763–1770.
- MELLOR, E.A. *et al.* (2001). *Proc. Natl. Acad. Sci. U.S.A.*, **98**, 7964–7969.
- MELLOR, E.A. *et al.* (2003). *Proc. Natl. Acad. Sci. U.S.A.*, **100**, 11589–11593.
- MORESCHI, I. *et al.* (2006). *J. Biol. Chem.*, **281**, 31419–31429.
- NOGUCHI, K. *et al.* (2003). *J. Biol. Chem.*, **278**, 25600–25606.
- WEBB, T.E. *et al.* (1996a). *Mol. Pharmacol.*, **50**, 258–265.
- WEBB, T.E. *et al.* (1996b). *Biochem. Biophys. Res. Commun.*, **219**, 105–110.
- YOKOMIZO, T. *et al.* (1997). *Nature*, **387**, 620–624.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Platelet-activating factor

Overview: Platelet-activating factor (PAF, 1-*O*-alkyl-2-acetyl-*sn*-glycero-3-phosphocholine) is a biologically active phospholipid mediator. PAF acts by binding to a unique G-protein-coupled seven transmembrane receptor (PAF-R) and activates multiple intracellular signaling pathways by coupling to the G_{q/11} and G_{i/o} families of G-proteins. PAF-R is activated by PAF and its metabolically stable analogue mc-PAF. Other suggested endogenous ligands are oxidized phosphatidylcholine (Marathe *et al.*, 1999) and lysophosphatidylcholine (Ogita *et al.*, 1997). It may also be activated by bacterial lipopolysaccharide (Nakamura *et al.*, 1992).

Nomenclature	PAF-R
Ensembl ID	ENSG00000169403
Principal transduction	G _{q/11} , G _i , G _o
Selective agonists	mc-PAF
Selective antagonists	CV-6209 (9.5), SR27417 (10.0), WEB2086 (8.0), L659989 (8.1), ginkgolide B (6.4)
Radioligand	[³ H]-PAF (1.6 nM, Fukunaga <i>et al.</i> , 2001)

Note that a previously recommended radioligand ([³H]-WEB2086; K_d 44.6 nM) is currently unavailable.

Abbreviations: **CV-6209**, 2-(*N*-acetyl-*N*-[2-methoxy-3-octadecylcarbamoyloxypropoxycarbamoyl]aminomethyl)-1-ethylpyridinium chloride; **L659989**, *trans*-2-(3-methoxy-5-methylsulphonyl-4-propoxyphenyl)-5-(3,4,5-trimethoxyphenyl)tetrahydrofuran; **mc-PAF**, 1-*O*-alkyl-2-*N*-methylcarbamoyl-*sn*-glycero-3-phosphocholine; also known as (methyl)carbam(o)yl-PAF or c-PAF; **SR27417**, *N*-(2-dimethylaminoethyl)-*N*-(3-pyridinylmethyl)(4-[2,4,6-trisopropylphenyl]thiazol-2-yl)amine; **WEB2086**, 3-[4-[2-chlorophenyl]-9-methyl-6*H*-thieno[3,2-*f*][1,2,4]triazolo[4,3-*a*][1,4]diazepine-2-yl]-1-(4-morpholinyl)-1-propanone; also known as apafant

Further Reading:

- HONDA, Z., ISHII, S. & SHIMIZU, T. (2002). Platelet-activating factor receptor. *J. Biochem.*, **1331**, 773–779.
- ISHII, S., NAGASE, T. & SHIMIZU, T. (2002). Platelet-activating factor receptor. *Prostaglandins Other Lipid Mediators*, **68-9**, 599–609.
- ISHII, S. & SHIMIZU, T. (2000). Platelet-activating factor (PAF) receptor and genetically engineered PAF receptor mutant mice. *Prog. Lipid Res.*, **39**, 41–82.
- IZUMI, T. & SHIMIZU, T. (1995). Platelet-activating factor receptor; gene expression and signal transduction. *Biochim. Biophys. Acta*, **1259**, 317–333.
- MONTRUCCHIO, G., ALLOATTI, G. & CAMUSSI, G. (2000). Role of platelet-activating factor in cardiovascular pathophysiology. *Physiol. Rev.*, **80**, 1669–1699.
- MACLENNAN, K.M., SMITH, P.F. & DARLINGTON, C.L. (1996). Platelet-activating factor in the CNS. *Prog. Neurobiol.*, **50**, 585–596.
- PRESCOTT, S.M., ZIMMERMAN, G.A., STAFFORINI, D.M. & MCINTYRE, T.M. (2000). Platelet-activating factor and related lipid mediators. *Annu. Rev. Biochem.*, **69**, 419–445.
- STAFFORINI, D.M., MCINTYRE, T.M., ZIMMERMAN, G.A. & PRESCOTT, S.M. (2003). Platelet-activating factor, a pleiotropic mediator of physiological and pathological processes. *Crit. Rev. Clin. Lab. Sci.*, **40**, 643–672.
- SUMMERS, J.B. & ALBERT, D.H. (1995). Platelet activating factor antagonists. *Adv. Pharmacol.*, **32**, 67–168.

References:

- FUKUNAGA, K. *et al.* (2001). *J. Biol. Chem.*, **276**, 43025–43030.
- MARATHE, G.K. *et al.* (1999). *J. Biol. Chem.*, **274**, 28395–28404.
- NAKAMURA, M. *et al.* (1992). *FEBS Lett.*, **314**, 125–129.
- OGITA, T. *et al.* (1997). *Am. J. Physiol.*, **272**, H17–H24.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Prostanoid

Overview: Prostanoid receptors (nomenclature agreed by the NC-IUPHAR Subcommittee on Prostanoid Receptors, see Coleman *et al.*, 1994) are activated by the endogenous ligands prostaglandin (PG) D₂ (D), PGE₂ (E), PGF_{2α} (F), PGH₂ (H), prostacyclin [PGI₂ (I)] and thromboxane A₂ (T). Measurement of the potency of PGI₂ and TXA₂ is hampered by their instability in physiological salt solution; they are often replaced by cicaprost and U46619, respectively, in receptor characterization studies.

Nomenclature	DP ₁	DP ₂	FP	IP	TP
Other names	—	CRTh2, GPR44	—	—	—
Ensembl ID	ENSG00000168229	ENSG00000183134	ENSG00000122420	ENSG00000160013	ENSG00000006638
Principal transduction	G _s	G _{i/o}	G _{q/11}	G _s	G _{q/11}
Rank order of potency	D >> E > F > I, T	D >> F, E > I, T	F > D > E > I, T	I >> D, E, F > T	T = H >> D, E, F, I
Selective agonists	L644698, BW245C, ZK118182, RS93520, SQ27986	13,14-dihydro-15-oxoPGD ₂ , 15R-methyl PGD ₂ (Hata <i>et al.</i> , 2003; Monneret <i>et al.</i> , 2003)	Fluprostenol, latanoprost	Cicaprost, AFP-07, BMY45778 (Seiler <i>et al.</i> , 1997)	U46619, STA ₂ , I-BOP, AGN192093
Selective antagonists	BWA868C (9.3, Giles, 1989), S5751 (8.8 Arimura <i>et al.</i> , 2001), ZK138357 (7.3, Lydford <i>et al.</i> , 1996)	Ramatroban (Sugimoto <i>et al.</i> , 2003)	—	RO1138452 (8.8, Bley <i>et al.</i> , 2006)	BMS180291 (9.3–10.0), ONO3708 (8.9), GR32191 (8.3–9.4, Lumley <i>et al.</i> , 1989), SQ29548 (8.1–9.1, Swayne <i>et al.</i> , 1988)
Probes	[³ H]-PGD ₂ (13–34 nM)	[³ H]-PGD ₂ (6–11 nM)	[³ H]-PGF _{2α} (2–4 nM), [³ H]-(+)-fluprostenol (34 nM)	[³ H]-Iloprost (1–20 nM)	[³ H]-SQ29548 (5–40 nM), [¹²⁵ I]-SAP (0.2–1.0 nM), [¹²⁵ I]-I-BOP (0.3–5.0 nM)

Ramatroban is also a TP receptor antagonist. Cicaprost exhibits moderate EP₄ receptor agonist potency (Abramovitz *et al.*, 2000). Iloprost also binds to EP₁ receptors. The TP receptor exists in α and β isoforms due to alternative splicing of the cytoplasmic tail (Raychowdhury *et al.*, 1994).

Nomenclature	EP ₁	EP ₂	EP ₃	EP ₄
Ensembl ID	ENSG00000160951	ENSG00000125384	ENSG00000050628	ENSG00000171522
Principal transduction	G _{q/11}	G _s	G _{i/o}	G _s
Rank order of potency	E > F, I > D, T	E > F, I > D, T	E > F, I > D, T	E > F, I > D, T
Selective agonists	17-Ph- ω -trilor-PGE ₂ , ONO-DI-004	Butaprost, AH13205, ONO-AE1–259	Sulprostone, SC46275, ONO-AE-248	ONO-AE1–329
Selective antagonists	ONO8711 (9.2), SC51322 (8.8)	—	L798106 (7.7)	GW627368 (9.2), ONO-AE3–208 (8.5), L161982 (7.6)
Probes	[³ H]-PGE ₂ (1–25 nM)	[³ H]-PGE ₂ (5–22 nM)	[³ H]-PGE ₂ (0.3–7 nM)	[³ H]-PGE ₂ (0.6–24 nM)

17-Ph- ω -trilor-PGE₂ also shows agonist activity at EP₃ receptors. Sulprostone also has affinity for EP₁ receptors. Butaprost and SC46275 may require deesterification within tissues to attain full agonist potency. There is evidence for subtypes of FP (Liljebris *et al.*, 1995), IP (Takechi *et al.*, 1996; Wise *et al.*, 1995) and TP (Krauss *et al.*, 1996) receptors. mRNA for the EP₁ and EP₃ receptors undergo alternative splicing to produce two (Okuda-Ashitaka *et al.*, 1996) and at least six variants, respectively, which can interfere with signalling (Okuda-Ashitaka *et al.*, 1996) or generate complex patterns of G-protein (G_{i/o}, G_{q/11}, G_s and G_{12/13}) coupling (e.g. Kotani *et al.*, 1995; Negishi *et al.*, 1995). The possibility of additional receptors for the isoprostanes has been suggested (Pratico *et al.*, 1996).

Abbreviations: AFP-07, 7,7-difluoro-16S,20-dimethyl-18,19-didehydro-PGI₂; AGN192093, (Z)-7-[(1E,5Z,6Z,7 β)-7-[(1E,3S)-3-hydroxy-1-octenyl]-3-oxo-2,4-dioxobicyclo[3.2.1]oct-6-yl)-5-heptanol; AH13205, trans-2-(4-[1-hydroxyhexyl]phenyl)-5-oxocyclopentane-heptanoic acid; AH23848, (1 α [z],2 β ,5 α)($\pm$)-7-(5-[(1,1'-biphenyl)-4-yl]methoxy)-2-[4-morpholinyl]-3-oxocyclopentyl)-4-heptenoate; BMS180291, (1S-(1Z,2 α ,3 α ,4 α)-2-[(3-{4-[(pentylamino)carbonyl]-2-oxazoly]-7-oxabicyclo[2.2.1]-hept-2-yl]methyl)benzenepropanoic acid; also known as ifetroban; BMY45778, 3-(4-[4,5-diphenyl-2-oxazoly]-5-oxazoly)phenoxyacetic acid; BW245C, 5-(6-carboxyhexyl)-1-(3-cyclohexyl-3-hydroxypropyl)hydantoin; BWA868C, 3-benzyl-5-(6-carboxyhexyl)-1-(2-cyclohexyl-2-hydroxyethylamino)hydantoin; GR32191, [1R-[1(Z),2 β ,3 β ,5]]-(+)-7-[5-[(1,1'-biphenyl)-4-yl]methoxy]-3-hydroxy-2-(1-piperidinyl)cyclopentyl]-4-heptenoic acid; GW627368, N-(2-[4-(4,9-diethoxy-1-oxo-1,3-dihydro-2H-benzof[1]isindol-2-yl)phenyl]acetyl)benzenesulphonamide; I-BOP, (1S-[1 α ,2 β {5 α },3 α {1 ϵ ,3 α },4 α]-7-(3-[hydroxy-4-{4'-iodophenoxy}-1-butenyl]-7-oxabicyclo[2.2.1]hept-2-yl)-5-heptanoate; L161982, 5-butyl-2,4-dihydro-[2'-[N-(5-methyl-2-thiophenecarbonyl)sulphamoyl]biphenyl-4-yl]methyl]-2-[(2-trifluoromethyl)phenyl]-1,2,4-triazol-3-one; L644698, 4-(3-{3-(3-hydroxyoctyl)-4-oxo-2-thiazolidinyl}propyl)benzoate racemate; L798106, 5-bromo-2-methoxy-N-[3-(naphthalen-2-yl-methylphenyl)acryloyl]-benzenesulphonamide; ONO3708, (9,11)-(11,12)-dideoxa-9 α ,11 α -dimethylmethano-11,12-methano-13,14-dihydro-13-aza-14-oxo-15-cyclopentyl-16,17,18,19,20-pentano-15-epi-TXA₂; ONO8711, 6-[(2S,3S)-3-(4-chloro-2-methylphenylsulphonylaminomethyl)-bicyclo[2.2.2]octan-2-yl]-5Z-hexenoic acid; ONO-DI-004, 17S-17,20-dimethyl-2,5-ethano-6-oxo PGE₁; ONO-AE-248, 11,15-O-dimethyl-PGE₂; ONO-AE1-259, 16S-9-deoxy-9 β -chloro-15-deoxy-16-hydroxy-17,17-propano-19,20-didehydro-PGE₂; ONO-AE1-329, 16-(3-methoxymethyl)phenyl- ω -tetranor-3,7-dithia-PGE₁; ONO-AE3-208, 2-(2-(2-methyl-2-naphth-1-ylacetylaminomethyl)-phenylmethyl)-benzoic acid; RO1138452, 4,5-dihydro-1H-imidazol-Z-yl-[4-(4-isopropoxybenzyl)phenyl]amine; RS93520, Z-4-[(C3'S,1R,2R,3S,6R)-2C3'-cyclohexyl-3'-hydroxyprop-1-ynyl]-3-hydroxybicyclo(4.2.0)oct-7-ylidene butyrate; SAP, 7-[(1R,2S,3S,5R)-6,6-dimethyl-3-benzenesulphonamino-bicyclo[3.1.1]hept-2-yl)-5Z-heptenoic acid; SC46275, methyl-7-(2 β -[6-{1-cyclopenten-1-yl}-4R-hydroxy-4-methyl-1 ϵ ,5 ϵ -hexadienyl]-3 α -hydroxy-5-oxo-1 α -cyclopentyl)-4Z-heptenoic acid; SC51322, 8-chlorodibenz[b,f][1,4]oxazepine-10(11H)-carboxylic acid, 2-[3-{furanlylmethyl}-thio]-1-oxopropylhydrazide; SQ27986, (1S-[1 α ,2 β {5 α },3 β ,4 β]-7-(3-{[2-(phenylamino)carbonyl]hydrazino]methyl}-7-oxabicyclo[2.2.1]hept-2-yl)-5-heptenoate; S5751, ((Z)-7-[1R,2R,3S,5S)-2-(5-hydroxybenzo[b]thiophen-3-ylcarbonylamino)-10-norpinan-3-yl]hept-5-enoic acid; SQ27986, [1S-[1B,2B(5Z),3A(1E,3S)4B]]-7-[3-(3-cyclohexyl-3-hydroxy-1-propenyl)-7-oxabicyclo[2.2.1]hept-2-yl]-5Z-heptenoic acid; STA₂, 11 α -carba-9 α ,11 β -thia-TxA₂; U46619, 11 α ,9 α -epoxymethano-PGH₂; ZK 118182, (5Z,13E)-(9R,11R,15S)-9-chloro-15-cyclohexyl-11,15-dihydroxy-3-oxa-16,17,18,19,20-pentano-5,13-prostadienoic acid; ZK138357, (5Z)-7-[(2R,4S,5S)-2-[2-chlorophenyl]-5-{1E}-{3R,S}-3-hydroxy-3-cyclohexyl-1-propenyl]-1,3-dioxolan-4-yl)-5-heptanoic acid

Further Reading:

BOS, C.L., RICHEL, D.J., RITSEMA, T., PEPPELENBOSCH, M.P. & VERSTEEG, H.H. (2004). Prostanoids and prostanoid receptors in signal transduction. *Int. J. Biochem. Cell Biol.*, **36**, 1187–1205.

BREYER, R.M., BAGDASSARIAN, C.K., MYERS, S.A. & BREYER, M.D. (2001). Prostanoid receptors: subtypes and signaling. *Annu. Rev. Pharmacol. Toxicol.*, **41**, 661–690.

COLEMAN, R.A., SMITH, W.L. & NARUMIYA, S. (1994). VIII. International Union of Pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes. *Pharmacol. Rev.*, **46**, 205–229.

HATA, A.N. & BREYER, R.M. (2004). Pharmacology and signaling of prostaglandin receptors: multiple roles in inflammation and immune modulation. *Pharmacol. Ther.*, **103**, 147–166.

KOSTENIS, E. & ULVEN, T. (2006). Emerging roles of DP and CRTH2 in allergic inflammation, *Trends Mol. Med.*, **12**, 148–158.

NARUMIYA, S., SUGIMOTO, Y. & USHIKUBI, F. (1999). Prostanoid receptors: structures, properties, and functions. *Physiol. Rev.*, **79**, 1193–1226.

WISE, H. & JONES, R.L. (2000). *Prostacyclin and its Receptors*. Dordrecht: Kluwer Academic Publishers.

References:

ABRAMOVITZ, M. et al. (2000). *Biochim. Biophys. Acta*, **1483**, 285–293.

ARIMURA, A. et al. (2001). *J. Pharmacol. Exp. Ther.*, **298**, 411–419.

BLEY, K.R. et al. (2006). *Br. J. Pharmacol.*, **147**, 335–345.

GALLANT, M. et al. (2002). *Bioorg. Med. Chem. Lett.*, **12**, 2583–2586.

GILES, H. et al. (1989). *Br. J. Pharmacol.*, **96**, 291–300.

HATA, A.N. et al. (2003). *J. Pharmacol. Exp. Ther.*, **306**, 463–470.

HIRAI, H. et al. (2001). *J. Exp. Med.*, **193**, 255–261.

KOTANI, M. et al. (1995). *Mol. Pharmacol.*, **48**, 869–879.

KRAUSS, A.H. et al. (1996). *Br. J. Pharmacol.*, **117**, 1171–1180.

LILJEBRIS, C. et al. (1995). *J. Med. Chem.*, **38**, 289–304.

LUMLEY, P. et al. (1989). *Br. J. Pharmacol.*, **97**, 783–794.

LYDFORD, S.J. et al. (1996). *Br. J. Pharmacol.*, **117**, 190P.

MONNERET, G. et al. (2001). *Blood*, **98**, 1942–1948.

MONNERET, G. et al. (2003). *J. Pharmacol. Exp. Ther.*, **304**, 349–355.

NEGISHI, M. et al. (1995). *J. Biol. Chem.*, **270**, 16122–16127.

OKUDA-ASHITAKA, E. et al. (1996). *J. Biol. Chem.*, **271**, 31255–31261.

PRATICO, D. et al. (1996). *J. Biol. Chem.*, **271**, 14916–14924.

RAYCHOWDHURY, M.K. et al. (1994). *J. Biol. Chem.*, **269**, 19256–19261.

SEILER, S.M. et al. (1997). *Prostaglandins*, **53**, 21–35.

SUGIMOTO, H. et al. (2003). *J. Pharmacol. Exp. Ther.*, **305**, 347–352.

SWAYNE, G.T. et al. (1988). *Eur. J. Pharmacol.*, **152**, 311–319.

TAKECHI, H. et al. (1996). *J. Biol. Chem.*, **271**, 5901–5906.

WISE, H. et al. (1995). *Eur. J. Pharmacol.*, **278**, 265–269.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Proteinase-activated

Overview: Proteinase-activated receptors (PARs, nomenclature as agreed by NC-IUPHAR Subcommittee on Protease-activated Receptors, see Hollenberg & Coughlan, 2002) are activated by proteolytic cleavage of their amino terminal exodomains. Alternative endogenous proteinases or ligands to thrombin for PAR1, PAR3 or PAR4 have not been demonstrated. Activation of PAR2 by trypsin or tryptase release *in vivo* is yet to be demonstrated. Several proteases, including cathepsin G and chymotrypsin, have an inhibitory effect at the PAR1 receptor such that they cleave the exodomain of the receptor without inducing activation, thereby preventing activation by thrombin but not by agonist peptides. The role of such an action *in vivo* is unclear. Agonist protease-induced hydrolysis is thought to unmask a tethered ligand at the exposed amino terminus, which acts intramolecularly at the binding site in the body of the receptor to effect transmembrane signalling. Tethered ligand sequences at human PAR1–4 are SFLLRN, SLIGKV, TFRGAP and GYPGQV, respectively. With the exception of PAR3, these synthetic peptide sequences (as carboxyl terminal amides) are able to act as agonists at their respective receptors.

Nomenclature	PAR ₁	PAR ₂	PAR ₃	PAR ₄
Other names	Thrombin receptor, PAR-1, PAR ₁	PAR-2, PAR ₂	Thrombin receptor, PAR-3, PAR ₃	Thrombin receptor, PAR-4, PAR ₄
Ensembl ID	ENSG00000181104	ENSG00000164251	ENSG00000164220	ENSG00000127533
Principal transduction	G _{q/11} /G _{i/o} /G _{12/13}	G _{q/11} /G _{i/o}	G _{q/11} /G _{i/o}	G _{q/11} /G _{i/o}
Agonist proteases	Thrombin, trypsin	Trypsin, tryptase	Thrombin, trypsin, factor Xa	Thrombin, trypsin
Selective agonists	TFLLR-NH ₂	SLIGRL, SLIGKV	—	GYPGQV, GYPGKF
Selective antagonists	RWJ56110 (Andrade-Gordon <i>et al.</i> , 1999)	—	—	—
Probes	[³ H]-haTRAP (Ahn <i>et al.</i> , 1997)	<i>Trans</i> -cinnamoyl-LIGRLO[N- [³ H]-propionyl]-NH ₂ (Al Ani <i>et al.</i> , 1999)	—	—

TFLLR-NH₂ is selective relative to the PAR2 receptor (Blackhart *et al.*, 1996; Kawabata *et al.*, 1999). Thrombin is inactive at the PAR₂ receptor.

Abbreviations: [³H]-haTRAP, Ala-*p*-fluoroPhe-Ala-Arg-cyclohexylAla-homoArg-[³H]-Tyr-amide; **RWJ56110**, (αS)-*N*-([1*S*]-3-amino-1-[(phenylmethyl)amino]propyl)-α-[(1-[(2,6-dichlorophenyl)methyl]-3-[1-pyrrolidinylmethyl]-1*H*-indol-6-yl)amino]carbonylamino)-3,4-difluoro-benzenepropanamide

Further Reading:

ANDERLUH, M. & DOLENC, M.S. (2002). Thrombin receptor antagonists; recent advances in PAR-1 antagonist development. *Curr. Med. Chem.*, **9**, 1229–1250.

CHACKALAMANNIL, S. (2003). G-protein coupled receptor antagonists-1: protease activated receptor-1 (PAR-1) antagonists as novel cardiovascular therapeutic agents. *Curr. Top. Med. Chem.*, **3**, 1115–1123.

CHACKALAMANNIL, S. (2006). Thrombin receptor (protease activated receptor-1) antagonists as potent antithrombotic agents with strong antiplatelet effects. *J. Med. Chem.*, **49**, 5389–5403.

COUGHLIN, S.R. (2001). Protease-activated receptors in vascular biology. *Thromb. Haemost.*, **86**, 298–307.

HOLLENBERG, M.D. & COMPTON, S.J. (2002). International Union of Pharmacology. XXVIII. Proteinase-activated receptors. *Pharmacol. Rev.*, **54**, 203–217.

LEGER, A.J., COVIC, L. & KULIOPULOS, A. (2006). Protease-activated receptors in cardiovascular diseases. *Circulation*, **114**, 1070–1077.

SCHMIDLIN, F. & BUNNETT, N.W. (2001). Protease-activated receptors: how proteases signal to cells. *Curr. Opin. Pharmacol.*, **1**, 575–582.

STEINBERG, S.F. (2005). The cardiovascular actions of protease-activated receptors. *Mol. Pharmacol.*, **67**, 2–11.

References:

AHN, H.S. *et al.* (1997). *Mol. Pharmacol.*, **51**, 350–356.

AL ANI, B. *et al.* (1999). *J. Pharmacol. Exp. Ther.*, **290**, 753–760.

ANDRADE-GORDON, P. *et al.* (1999). *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 12257–12262.

BLACKHART, B.D. *et al.* (1996). *J. Biol. Chem.*, **271**, 16466–16471.

KAWABATA, A. *et al.* (1999). *J. Pharmacol. Exp. Ther.*, **288**, 358–370.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Relaxin family peptide

Overview: Relaxin family peptide receptors (RXFP, nomenclature as agreed by NC-IUPHAR Subcommittee on relaxin family peptide receptors, see Bathgate *et al.*, 2006) are activated by heterodimeric peptide hormones analogous to insulin (H1 relaxin [ENSG00000107018], H2 relaxin [ENSG00000107014], H3 relaxin [also known as INSL7, ENSG00000171136] in higher primates, INSL3, Leydig insulin-like peptide [OTTHUMG00000070952*]) and INSL5 [ENSG00000172410]. Species homologs of relaxin have distinct pharmacology – H2 relaxin interacts with RXFP1 and RXFP2, mouse and rat relaxin selectively bind to and activate RXFP1 (Scott *et al.*, 2005a) and porcine relaxin may have a higher efficacy than H2 relaxin (Halls *et al.*, 2005). H3 relaxin has differential affinity for RXFP2 receptors between species; mouse and rat RXFP2 have a higher affinity for H3 relaxin (Scott *et al.*, 2005b). At least two binding sites have been identified on the RXFP1 and RXFP2 receptors: a high-affinity site in the leucine-rich repeat region of the ectodomain and a somewhat lower-affinity site located in the surface loops of the transmembrane (Halls *et al.*, 2005). The unique N-terminal LDLa module of RXFP1 and RXFP2 directs receptor signalling (Scott *et al.*, 2005c).

Nomenclature Other names	RXFP1 Relaxin receptor, LGR7, leucine-Rich repeat- containing G-protein- coupled receptor 7, RX1	RXFP2 INSL3 receptor, LGR8, leucine-rich repeat- containing G-protein- coupled receptor 8, GREAT, RX2 ENSG00000133105	RXFP3 Relaxin 3 receptor, GPCR135, somatostatin and angiotensin-like peptide receptor SALPR, RX3	RXFP4 INSL5 receptor, GPCR142, GPR100, relaxin 3 receptor 2, RX4
Ensembl ID	ENSG00000171509	ENSG00000133105	ENSG00000182631	ENSG00000173080
Principal transduction	G _s , G _{i/o}	G _s , G _{i/o} (Kawamura <i>et al.</i> , 2004)	G _{i/o} (Matsumoto <i>et al.</i> , 2000)	G _{i/o} (Liu <i>et al.</i> , 2003b)
Rank order of potency	H2 relaxin > H3 relaxin >> INSL3 (Sudo <i>et al.</i> , 2003)	INSL3 > H2 relaxin >> H3 relaxin (Kumagai <i>et al.</i> , 2002; Sudo <i>et al.</i> , 2003)	H3 relaxin > H3 relaxin B chain (Liu <i>et al.</i> , 2003a)	INSL5 = H3 relaxin > H3 relaxin B chain (Liu <i>et al.</i> , 2003b, 2005a)
Antagonists	LGR7-truncate (Scott <i>et al.</i> , 2005c)	INSL3 B-chain analog (Del Borgo <i>et al.</i> , 2004) (des 1-8) A-chain INSL3 analog (Bullesbach & Schwabe, 2005)	INSL5 (Liu <i>et al.</i> , 2005a)	—
Probes	[³³ P]-H2 relaxin (0.2 nM; Sudo <i>et al.</i> , 2003)	[³³ P]-H2 relaxin (1.06 nM; Sudo <i>et al.</i> , 2003) [¹²⁵ I]-INSL3 (0.1 nM; Muda <i>et al.</i> , 2005)	[¹²⁵ I]-H3 relaxin (0.3 nM; Liu <i>et al.</i> , 2003a) [¹²⁵ I]-H3-B/INSL5 A chimera (0.5 nM; Liu <i>et al.</i> , 2005b)	[¹²⁵ I]-H3 relaxin (0.2 nM; Liu <i>et al.</i> , 2003b) [¹²⁵ I]-H3-B/INSL5 A chimera (1.2 nM; Liu <i>et al.</i> , 2005b)

Mutations in *INSL3* and *LGR8* (RXFP2) have been reported in populations of patients with cryptorchidism (Ferlin *et al.*, 2003). Numerous splice variants of the human RXFP1 and RXFP2 receptors have been identified, none of which bind relaxin family peptides (Muda *et al.*, 2005). Splice variants of RXFP1 encoding the N-terminal LDLa module act as antagonists of RXFP1 signalling (Scott *et al.*, 2005b).

Abbreviations: **H2 relaxin**, human gene 2 relaxin; **H3 relaxin**, human gene 3 relaxin; **INSL3**, insulin-like peptide 3; **INSL5**, insulin-like peptide 5

Further Reading:

BATHGATE, R.A., SAMUEL, C.S., BURAZIN, T.C., GUNDLACH, A.L. & TREGEAR, G.W. (2003). Relaxin: new peptides, receptors and novel actions. *Trends Endocrinol. Metab.*, **14**, 207–213.

BATHGATE, R.A.D., HSUEH, A.J.W. & SHERWOOD, O.D. (2005). Physiology and molecular biology of the relaxin peptide family. In: *Knobil and Neill's Physiology of Reproduction*, ed. Neill J.D. New York: Academic Press.

BATHGATE, R.A., IVELL, R., SANBORN, B.M., SHERWOOD, O.D. & SYMMERS, R.J. (2006). International Union of Pharmacology LVII: Recommendations for the Nomenclature of Receptors for Relaxin Family Peptides. *Pharmacol. Rev.*, **58**, 7–31.

CONRAD, K.P. & NOVAK, J. (2004). Emerging role of relaxin in renal and cardiovascular function. *Am. J. Physiol. – Regul. Integr. Comp. Physiol.*, **287**, R250–R261.

DSCHIETZIG, T. & STANGL, K. (2003). Relaxin: a pregnancy hormone as central player of body fluid and circulation homeostasis. *Cell. Mol. Life Sci.*, **60**, 688–700.

HSU, S.Y. (2003). New insights into the evolution of the relaxin–LGR signaling system. *Trends Endocrinol. Metab.*, **14**, 303–309.

HSU, S.Y., NAKABAYASHI, K., NISHI, S., KUMAGAI, J., KUDO, M., SHERWOOD, O.D. & HSUEH, A.J. (2002). Activation of orphan receptors by the hormone relaxin. *Science*, **295**, 637–638.

IVELL, R. & BATHGATE, R.A. (2002). Reproductive biology of the relaxin-like factor (RLF/INSL3). *Biol. Reprod.*, **67**, 699–705.

IVELL, R. & EINSPIANIER, A. (2002). Relaxin peptides are new global players. *Trends Endocrinol. Metab.*, **13**, 343–348.

SAMUEL, C.S., PARRY, L.J. & SUMMERS, R.J. (2003). Physiological or pathological – a role for relaxin in the cardiovascular system? *Curr. Opin. Pharmacol.*, **3**, 152–158.

SAMUEL, C.S., DU, X.J., BATHGATE, R.A. & SUMMERS, R.J. (2006). ‘Relaxin’ the stiffened heart and arteries: the therapeutic potential for relaxin in the treatment of cardiovascular disease. *Pharmacol. Ther.*, **112**, 529–552.

SHERWOOD, O.D. (2004). Relaxin's physiological roles and other diverse actions. *Endocr. Rev.*, **25**, 205–234.

References:

- BULLESBACH, E.E. & SCHWABE, C. (2005). *J. Biol. Chem.*, **280**, 14586–14590.
- DEL BORGO, M.P. *et al.* (2004). *18th American Peptide Symposium*, pp 727–729.
- FERLIN, A. *et al.* (2003). *J. Clin. Endocrinol. Metab.*, **88**, 4273–4279.
- HALLS, M.L. *et al.* (2005). *J. Pharm Exp Ther.*, **313**, 677–687.
- KAWAMURA, K. *et al.* (2004). *Proc. Natl. Acad. Sci. U.S.A.*, **101**, 7323–7328.
- KUMAGAI, J. *et al.* (2002). *J. Biol. Chem.*, **277**, 31283–31286.
- LIU, C. *et al.* (2003a). *J. Biol. Chem.*, **278**, 50754–50764.
- LIU, C. *et al.* (2003b). *J. Biol. Chem.*, **278**, 50765–50770.
- LIU, C. *et al.* (2005a). *J. Biol. Chem.*, **280**, 292–300.

- LIU, C. *et al.* (2005b). *Mol. Pharmacol.*, **67**, 231–240.
- MATSUMOTO, M. *et al.* (2000). *Gene*, **248**, 183–189.
- MUDA, M. *et al.* (2005). *Mol. Hum. Reprod.*, **11**, 591–600.
- SCOTT, D.J. *et al.* (2005a). *Ann. N.Y. Acad. Sci.*, **1041**, 8–12.
- SCOTT, D.J. *et al.* (2005b). *Ann. N.Y. Acad. Sci.*, **1041**, 13–16.
- SCOTT, D.J. *et al.* (2005c). *Ann. N.Y. Acad. Sci.*, **1041**, 22–26.
- SUDO, S. *et al.* (2003). *J. Biol. Chem.*, **278**, 7855–7862.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Resolvin

Overview: Resolvin receptors (provisional nomenclature) are activated by the lipid-derived, anti-inflammatory ligand resolvin E1 (RvE1), which is the result of sequential metabolism of EPA by aspirin-modified cyclooxygenase and lipoxygenase (Arita *et al.*, 2005).

Nomenclature	RvE1
Other names	ChemR23, chemokine receptor-like 1, DEZ
Ensembl ID	ENSG00000174600
Principal transduction	Not yet established
Rank order of potency	RvE1 > chemerin C-terminal peptide (Arita <i>et al.</i> , 2005)

Abbreviations: EPA, eicosapentaenoic acid; **RvE1**, resolvin E1 or 5S,12R,18R-trihydroxy-6Z,8E,10E,14Z,16E-EPA

Further Reading:

SERHAN, C.N., HONG, S., GRONERT, K., COLGAN, S.P., DEVCHAND, P.R., MIRICK, G. & MOUSSIGNAC, R.L. (2002). Resolvins: a family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals. *J. Exp. Med.*, **196**, 1025–1037.

SERHAN, C.N. (2005). Novel ω -3-derived local mediators in anti-inflammation and resolution. *Pharmacol. Ther.*, **105**, 7–21.

SERHAN, C.N. (2007). Resolution phases of inflammation: Novel endogenous anti-inflammatory and proresolving lipid mediators and pathways. *Annu. Rev. Immunol.*, in press.

References :

ARITA, M. *et al.* (2005). *J. Exp. Med.*, **201**, 713–722.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Somatostatin

Overview: Somatostatin (somatotropin release inhibiting factor) is an abundant neuropeptide, which acts on five subtypes of somatostatin receptor (sst1 – sst5; nomenclature approved by the NC-IUPHAR Subcommittee on Somatostatin Receptors, see Hoyer *et al.*, 2000). Activation of these receptors produces a wide range of physiological effects throughout the body. The relationship of the cloned receptors to endogenously expressed receptors is not yet well established in some cases. Endogenous ligands for these receptors are somatostatin-14 (SRIF-14) and somatostatin-28 (SRIF-28). Cortistatin (CST-14) has also been suggested to be an endogenous ligand for somatostatin receptors (Delecea *et al.*, 2000).

Nomenclature	sst ₁	sst ₂	sst ₃	sst ₄	sst ₅
Alternative names	SSTR1, SRIF ₂ , SRIF _{2A}	SSTR2, SRIF ₁ , SRIF _{1A}	SSTR3, SRIF ₁ , SRIF _{1C}	SSTR4, SRIF ₂ , SRIF _{2B}	SSTR5, SRIF ₁ , SRIF _{1B}
Ensemble ID	ENSG00000139874	ENSG00000180616	ENSG00000183473	ENSG00000132671	ENSG00000162009
Principal transduction	G _i	G _i	G _i	G _i	G _i
Selective agonists	des-Ala ^{1,2,5} -[DTrp ⁸ ,Iamp ⁹]SRIF, L797591	Octreotide, seglitide, BIM23027, L054522	L796778	NNC269100, L803087	BIM23268, BIM23052, L817818
Selective antagonists	—	Cyanamid 154806 (7.7–8.0)	—	—	BIM23056 (7.4–8.3)
Probes	—	[¹²⁵ I]-[Tyr ³]octreotide (0.13 nM) [¹²⁵ I]-BIM23027	—	—	[¹²⁵ I]-[Tyr ³]octreotide (0.23 nM)

[¹²⁵I]-[Tyr¹¹]SRIF-14, [¹²⁵I]-LTT-SRIF-28, [¹²⁵I]-CGP23996 and [¹²⁵I]-[Tyr¹⁰]CST-14 may be used to label somatostatin receptors nonselectively; BIM23052 is said to be selective in rat but not human receptor (Patel & Srikant, 1994). A number of nonpeptide subtype-selective agonists have been synthesised (see Rohrer *et al.*, 1998).

Abbreviations: **BIM23027**, *cyc*(N-Me-Ala-Tyr-D-Trp-Lys-Abu-Phe); **BIM23052**, D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂; **BIM23056**, D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-d-Nal-NH₂; **BIM23268**, *cyc*(Cys-Phe-Phe-D-Trp-Lys-Phe-Cys)-NH₂; **CGP23996**, *cyc*(Asn-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Tyr-Ser); **Cyanamid 154806**, Ac-(4-NO₂-Phe)-*cyc*(D-Cys-Tyr-D-Trp-Lys-Thr-Cys)-D-Tyr-NH₂; **L797591**, (2R)-N-(6-amino-2,2,4-trimethylhexyl)-3-(1-naphthyl)-2-([(2-phenylethyl)2-pyridin-2-ylethyl]amino)carbonyl]amino]propanamide; **L054522**, tert-butyl (bS)-b-methyl-N-[[4-(2-oxo-2,3-dihydro-1H-benzimidazol-1-yl)piperidin-1-yl]carbonyl]-D-tryptophyl-L-lysinate; **L796778**, methyl (2S)-6-amino-2-(((2R)-2-(((1S)-1-benzyl-2-[(4-nitrophenyl)amino]-2-oxoethyl)amino)carbonyl]amino)hexanoyl)amino]hexanoate; **L803087**, methyl (2S)-5-[[amino(imino)methyl]amino]-2-[[4-(5,7-difluoro-2-phenyl-1H-indol-3-yl)butanoyl]amino]pentanoate; **L817818**, (2R)-2-aminopropyl N2-[[2-(2-naphthyl)-1H-benzo[g]indol-3-yl]acetyl]-L-lysinate; **LTT-SRIF-28**, [Leu⁸,DTrp²²,DTyr²⁵]SRIF-28; **NNC269100**, 1-[3-[N-(5-bromopyridin-2-yl)-N-(3,4-dichlorobenzyl)amino]propyl]-3-[3-(¹H-imidazol-1-yl)propyl]thiourea.

Further Reading:

- CSABA, Z. & DOURNAUD, P. (2001). Cellular biology of somatostatin receptors. *Neuropeptides*, **35**, 1–23.
- DASGUPTA, P. (2004). Somatostatin analogues: multiple roles in cellular proliferation, neoplasia, and angiogenesis. *Pharmacol. Ther.*, **102**, 61–85.
- HANNON, J.P., NUNN, C., STOLZ, B., BRUNS, C., WECKBECKER, G., LEWIS, I., TROXLER, T., HURTH, K. & HOYER, D. (2002). Drug design at peptide receptors – somatostatin receptor ligands. *J. Mol. Neurosci.*, **18**, 15–27.
- HOYER, D., EPELBAUM, J., FENIUK, W., HUMPHREY, P.P.A., MEYERHOF, W., O'CAROLL, A.M., PATEL, Y., REISINE, T., REUBI, J.C., SCHINDLER, M., SCHONBRUNN, A., TAYLOR, J.E. & VEZZANI, A. (2000). Somatostatin receptors. In: *The IUPHAR Compendium of Receptor Characterization and Classification*, 2nd edn. ed. Watson, S. P. & Girdlestone, D., pp. 354–364. London: IUPHAR Media.
- LAHLOU, H., GUILLERMET, J., HORTALA, M., VERNEJOUL, F., PYRONNET, S., BOUSQUET, C. & SUSINI, C. (2004). Molecular signalling of somatostatin receptors. *Ann. NY Acad. Sci.*, **1014**, 121–131.
- MOLLER, L.N., STIDSEN, C.E., HARTMANN, B. & HOLST, J.J. (2003). Somatostatin receptors. *Biochim. Biophys. Acta-biomembr.*, **1616**, 1–84.
- OLIAS, G., VIOLLET, C., KUSSEROW, H., EPELBAUM, J. & MEYERHOF, W. (2004). Regulation and function of somatostatin receptors. *J. Neurochem.*, **89**, 1057–1091.
- PATEL, Y.C. (1999). Somatostatin and its receptor family. *Front. Neuroendocrinology*, **20**, 157–198.
- PATEL, Y.C., GREENWOOD, M.T., PANETTA, R., DEMCHYSHYN, L., NIZNIK, H. & SRIKANT, C.B. (1995). The somatostatin receptor family. *Life Sci.*, **57**, 1249–1265.
- RASHID, A.J., O'DOWD, B.F. & GEORGE, S.R. (2004). Minireview: Diversity and complexity of signaling through peptidergic G protein-coupled receptors. *Endocrinology*, **145**, 2645–2652.
- REISINE, T. & BELL, G. (1995). Molecular properties of somatostatin receptors. *Neuroscience*, **67**, 777–790.
- VAN DER HOEK, J., HOFLAND, L.J. & LAMBERTS, S.W. (2005). Novel subtype specific and universal somatostatin analogues: clinical potential and pitfalls. *Curr. Pharm. Des.*, **11**, 1573–1592.
- WECKBECKER, G., LEWIS, I., ALBERT, R., SCHMID, H.A., HOYER, D. & BRUNS, C. (2003). Opportunities in somatostatin research: Biological, chemical and therapeutic aspects. *Nat. Rev. Drug Discovery*, **2**, 999–1017.

References:

- DELECEA, L. *et al.* (1996). *Nature*, **381**, 242–245.
- PATEL, Y.C. & SRIKANT, C.B. (1994). *Endocrinology*, **135**, 2814–2817.
- ROHRER, S.P. *et al.* (1998). *Science*, **282**, 737–740.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Sphingosine-1-phosphate

Overview: Sphingosine-1-phosphate (S1P) receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on Lysophospholipid receptors; see Chun *et al.*, 2002) are activated by the endogenous lipid derivatives S1P and sphingosylphosphorylcholine (SPC). S1P has also been described to act at intracellular sites (see Hla *et al.*, 1999; Spiegel & Milstein, 2003), although most cellular phenomena ascribed to S1P can be explained by receptor-mediated mechanisms. The relationship between recombinant and endogenously expressed receptors is unclear. Radioligand binding has been conducted in heterologous expression systems using [³²P]-S1P (e.g. Okamoto *et al.*, 1998). In native systems, analysis of binding data is complicated by metabolism and high levels of nonspecific binding. Targeted deletion of several S1P receptors has clarified signalling pathways and physiological roles.

Nomenclature	S1P ₁	S1P ₂	S1P ₃	S1P ₄	S1P ₅
Other names	edg1, <i>lp</i> _{B1}	edg5, <i>lp</i> _{B2} , AGR16, H218	edg3, <i>lp</i> _{B3}	edg6, <i>lp</i> _{C1}	edg8, <i>lp</i> _{B4} , NRG-1
Ensembl ID	ENSG00000170989	ENSG00000175898	ENSG00000186354	ENSG00000125910	ENSG00000180739
Principal transduction	G _{i/o}	G _q , G _{12/13} , G _s	G _q , G _{i/o} , G _s	G _{i/o} , G _{12/13} , G _s	G _{i/o} , G _{12/13}
Rank order of potency	S1P > SPC	S1P > SPC	S1P > SPC	S1P, SPC	S1P, SPC
Selective agonists	SEW2871 (Sanna <i>et al.</i> , 2004)	—	—	—	—
Selective antagonists	—	JTE013 (Osada <i>et al.</i> , 2002)	—	—	—

The novel immunomodulator fingolimod (FTY720) may be phosphorylated *in vivo* (Albert *et al.*, 2005) to generate a relatively potent agonist at S1P₁, S1P₂, S1P₃, S1P₄ and S1P₅ receptors (Brinkmann *et al.*, 2002; Mandala *et al.*, 2002).

Abbreviations: JTE013, pyrazolopyridine analog; SEW2871, 5-(4-phenyl-5-trifluoromethylthiophen-2-yl)-3-(3-trifluoromethylphenyl)-(1,2,4)-oxadiazole

Further Reading:

ALLENDE, M.L. & PROIA, R.L. (2002). Sphingosine-1-phosphate receptors and the development of the vascular system. *Biochim. Biophys. Acta – Mol. Cell. Biol. Lipids*, **1582**, 222–227.

ANLIKER, B. & CHUN, J. (2004). Lysophospholipid G protein-coupled receptors. *J. Biol. Chem.*, **279**, 20555–20558.

CHUN, J., GOETZL, E.J., HLA, T., IGARASHI, Y., LYNCH, K.R., MOOLENAAR, W., PYNE, S. & TIGYI, G. (2002). International Union of Pharmacology. XXXIV. Lysophospholipid receptor nomenclature. *Pharmacol. Rev.*, **54**, 265–269.

CHUN, J. & ROSEN, N. (2006). Lysophospholipid receptors as potential drug targets in tissue transplantation and autoimmune diseases. *Curr. Pharm. Des.*, **12**, 161–171.

FUKUSHIMA, N., ISHII, I., CONTOS, J.J.A., WEINER, J.A. & CHUN, J. (2001). Lysophospholipid receptors. *Annu. Rev. Pharmacol. Toxicol.*, **41**, 507–534.

GOETZL, E.J. & TIGYI, G. (2004). Lysophospholipids and their G protein-coupled receptors in biology and diseases. *J. Cell. Biochem.*, **92**, 867–868.

HLA, T. (2001). Sphingosine 1-phosphate receptors. *Prostaglandins Other Lipid Mediat.*, **64**, 135–142.

ISHII, I., FRIEDMAN, B., YE, X., KAWAMURA, S., MCGIFFERT, C., CONTOS, J.J., KINGSBURY, M.A., ZHANG, G., BROWN, J.H. & CHUN, J. (2001). Selective loss of sphingosine 1-phosphate signaling with no obvious phenotypic abnormality in mice lacking its G protein-coupled receptor, *lpb3/edg-3*. *J. Biol. Chem.*, **276**, 33697–33704.

ISHII, I., YE, X., FRIEDMAN, B., KAWAMURA, S., CONTOS, J.J., KINGSBURY, M.A., YANG, A.H., ZHANG, G., BROWN, J.H. & CHUN, J. (2002). Marked perinatal lethality and cellular signaling deficits in mice null for the two sphingosine 1-phosphate (S1P) receptors, S1P(2)/LP(B2)/EDG-5 and S1P(3)/LP(B3)/EDG-3. *J. Biol. Chem.*, **277**, 25152–25159.

ISHII, I., FUKUSHIMA, N., YE, X.Q. & CHUN, J. (2004). Lysophospholipid receptors: signaling and biology. *Annu. Rev. Biochem.*, **73**, 321–354.

KLUK, M.J. & HLA, T. (2002). Signaling of sphingosine-1-phosphate via the S1P/EDG-family of G-protein-coupled receptors. *Biochim. Biophys. Acta – Mol. Cell. Biol. Lipids*, **1582**, 72–80.

MILSTIEN, S. & SPIEGEL, S. (2006). Targeting sphingosine-1-phosphate: a novel avenue for cancer transplantation and autoimmune diseases. *Cancer Cell*, **9**, 148–159.

OGRETMEN, B. & HANNUN, Y.A. (2004). Biologically active sphingolipids in cancer pathogenesis and treatment. *Nat. Rev. Cancer*, **4**, 604–616.

ROSEN, H. & GOETZL, E.J. (2005). Sphingosine 1-phosphate and its receptors: an autocrine and paracrine network. *Nat. Rev. Immunol.*, **5**, 560–570.

SPIEGEL, S. & MILSTIEN, S. (2003). Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat. Rev. Mol. Cell. Biol.*, **4**, 397–407.

STUNFF, H.L., MILSTIEN, S. & SPIEGEL, S. (2004). Generation and metabolism of bioactive sphingosine-1-phosphate. *J. Cell. Biochem.*, **92**, 882–899.

YANG, A.H., ISHII, I. & CHUN, J. (2002). *In vivo* roles of lysophospholipid receptors revealed by gene targeting studies in mice. *Biochim. Biophys. Acta – Mol. Cell. Biol. Lipids*, **1582**, 197–203.

YATOMI, Y. (2006). Sphingosine-1-phosphate in vascular biology: possible therapeutic strategies to control vascular diseases. *Curr. Pharm. Des.*, **12**, 575–587.

References:

ALBERT, R. *et al.* (2005). *J. Med. Chem.*, **48**, 5373–5377.

BRINKMANN, V. *et al.* (2002). *J. Biol. Chem.*, **277**, 21453–21457.

MANDALA, S. *et al.* (2002). *Science*, **296**, 346–349.

OKAMOTO, H. *et al.* (1998). *J. Biol. Chem.*, **273**, 27104–27110.

OSADA, M. *et al.* (2002). *Biochem. Biophys. Res. Commun.*, **299**, 483–487.

SANNA, M.G. *et al.* (2004). *J. Biol. Chem.*, **279**, 13839–13848.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Tachykinin

Overview: Tachykinin receptors (provisional nomenclature) are activated by the endogenous peptides substance P (SP), neurokinin A (NKA; previously known as substance K, neurokinin α , neuromedin L), neurokinin B (NKB; previously known as neurokinin β , neuromedin K), neuropeptide K and neuropeptide γ (N-terminally extended forms of neurokinin A). The neurokinins (A and B) are mammalian members of the tachykinin family, which includes peptides of mammalian and nonmammalian origin containing the consensus sequence: Phe-x-Gly-Leu-Met. Marked species differences in pharmacology exist for all three receptors, in particular with nonpeptide ligands.

Nomenclature	NK ₁	NK ₂	NK ₃
Other names	Substance P	Substance K	Neurokinin B, neuromedin K
Ensembl ID	ENSG00000115353	ENSG00000075073	ENSG00000169836
Principal transduction	G _{q/11}	G _{q/11}	G _{q/11}
Rank order of potency	SP > NKA > NKB	NKA > NKB > SP	NKB > NKA > SP
Selective agonists	SP methylester, [Sar ⁹ ,Met(O ₂) ¹¹]SP, [Pro ⁹]SP, septide	[β -Ala ⁸]NKA-(4-10), [Lys ⁵ ,Me-Leu ⁹ ,Mle ¹⁰]NKA-(4-10), GR64349	Senktide, [MePhe ⁷]NKB
Selective antagonists	Aprepitant (10.7; Hale <i>et al.</i> , 1998), SR140333 (9.5), LY303870 (9.4), CP99994 (9.3), RP67580 (7.6)	GR94800 (9.6), GR159897 (9.5), MEN10627 (9.2), SR48968 (9.0), MEN11420 (8.6; Catalioto <i>et al.</i> , 1998)	SR142802 (9.2), SB223412 (9.0, Sarau <i>et al.</i> , 1997), PD157672 (7.8)
Probes	[³ H]- or [¹²⁵ I]-SP, [³ H]- or [¹²⁵ I]-BH-[Sar ⁹ ,Met(O ₂) ¹¹]SP, [¹²⁵ I]-L703606 (0.3 nM), [¹⁸ F]-SPA-RQ	[³ H]-SR48968 (0.5 nM), [³ H]-GR100679, [¹²⁵ I]-NKA	[³ H]-Senktide, [¹²⁵ I]-[MePhe ⁷]NKB, [³ H]-SR142801 (0.13 nM)

The NK₁ receptor has also been described to couple to other G proteins (Roush and Kwatra, 1998). The hexapeptide agonist septide appears to bind to an overlapping but nonidentical site to SP on the NK₁ receptor. There are suggestions for additional subtypes of tachykinin receptor; an orphan receptor (SwissProt P30098) with structural similarities to the NK₃ receptor was found to respond to NKB when expressed in *Xenopus* oocytes or Chinese hamster ovary cells (Donaldson *et al.*, 1996; Krause *et al.*, 1997).

Abbreviations: Aprepitant, 5-[[[(2R,3S)-2-[(1R)-1-[3,5-bis(trifluoromethyl) phenyl]ethoxy]-3-(4-fluorophenyl)-4-morpholinyl]methyl]-1,2-dihydro-3H-1,2,4-triazol-3-one (also known as Emend); CP99994, (+)-(2S,3S)-3-(2-methoxybenzylamino)-2-phenylpiperidine; [¹⁸F]-SPA-RQ, ([¹⁸F]-2-fluoromethoxy-5-(5-trifluoromethyl-tetrazol-1-yl)-benzyl)[(2S,3S)-2-phenylpiperidin-3-yl]-amine; GR100679, cyclohexylcarbonyl-Gly-Ala-DTrp-Phe-NMe₂; GR159897, (R)-1-(2-[5-fluoro-1H-indol-3-yl]ethyl)-4-methoxy-4[(phenylsulfenyl)methyl]piperidine; GR64349, Lys-Asp-Ser-Phe-Val-Gly-(r - γ -lactam); GR94800, N- α -benzoyl-Ala-Ala-DTrp-Phe-DPro-Pro-Nle-NH₂; L-703606, *cis*-2-(diphenylmethyl)-N-[(2-iodophenyl)methyl]-1-azabicyclo[2.2.2]octan-3-amine; L-742694, 2(s)-[3,5-bis(trifluoromethyl)benzyl]-oxy-3(S)-phenyl-4-[(3-oxo-1,2,4-triazol-5-yl)methyl]morpholine; LY303870, (r)-1-(N-[2-methoxybenzyl]acetyl)amino-3-(1H-indol-3-yl)-2-(N-[2-(4-(piperidin-1-yl)piperidin-1-yl]acetyl)amino)propane; also known as lanepitant; MEN10627, *cyc*(2 β -5 β)(Met-Asp-Trp-Phe-Dap-Leu); MEN11420, *cyc*(2 β -5 β)[Asn(2-AcNH- β -D-Glc)-Asp-Trp-Phe-Dap-Leu]; also known as nepadutant; PD157672, Boc-(s)Phe-(r) α MePheNH(CH₂)₇NHCONH₂; RP67580, 3 α R,7 α R-(1-imino-2-[2-methoxyphenyl]ethyl)-7,7-diphenyl-4-perhydroisoindolone; SB223412, (s)-(-)-N-(α -ethylbenzyl)-3-hydroxy-2-phenylquinoline-4-carboxamide; SR140333, (s)-1-(2-[3-(3,4-dichlorophenyl)-1-{3-isopropoxyphenylacetyl}piperidin-3-yl]ethyl)-4-phenyl-1-azoniabicyclo(2.2.2)octane chloride; SR142801, (S)-(N)-(1-[3-{1-benzoyl-3-(3,4-dichlorophenyl)piperidin-3-yl}propyl]-4-phenylpiperidin-4-yl)-N-methylacetamide; SR48968, (S)-N-methyl-N-(4-acetyl amino-4-phenylpiperidino)-2-(3,4-dichlorophenyl)butylbenzamide; also known as saredutant

Further Reading:

- CANDENAS, L., LECCI, A., PINTO, F.M., PATAK, E., MAGGI, C.A. & PENNEFATHER, J.N. (2005). Tachykinins and tachykinin receptors: effects in the genitourinary tract. *Life Sci.*, **76**, 835–862.
- GAO, Z. & PEET, N.P. (1999). Recent advances in neurokinin receptor antagonists. *Curr. Med. Chem.*, **6**, 375–388.
- JOOS, G.F. & PAUWELS, R.A. (2001). Tachykinin receptor antagonists: potential in airways diseases. *Curr. Opin. Pharmacol.*, **1**, 235–241.
- PAGE, N.M. (2004). Hemokinins and endokinins. *Cell. Mol. Life Sci.*, **61**, 1652–1663.
- PATACCHINI, R. & MAGGI, C.A. (2001). Peripheral tachykinin receptors as targets for new drugs. *Eur. J. Pharmacol.*, **429**, 13–21.
- PENNEFATHER, J.N., LECCI, A., CANDENAS, M.L., PATAK, E., PINTO, F.M. & MAGGI, C.A. (2004). Tachykinins and tachykinin receptors: a growing family. *Life Sci.*, **74**, 1445–1463.
- SARIA, A. (1999). The tachykinin NK1 receptor in the brain: pharmacology and putative functions. *Eur. J. Pharmacol.*, **375**, 51–60.
- SEVERINI, C., IMPROTA, G., FALCONIERI-ERSPAMER, G., SALVADORI, S. & ERSAMER, V. (2002). The tachykinin peptide family. *Pharmacol. Rev.*, **54**, 285–322.

References:

- BERGSTROM, M. *et al.* (2004). *Biol. Psychiatry*, **55**, 1007–1012.
- CATALIOTO, R.M. *et al.* (1998). *Br. J. Pharmacol.*, **123**, 81–91.
- DONALDSON, L.F. *et al.* (1996). *Biochem. J.*, **320**, 1–5.
- HALE, J.J. *et al.* (1998). *J. Med. Chem.*, **41**, 4607–4614.
- KRAUSE, J.E. *et al.* (1997). *Proc. Natl. Acad. Sci. U.S.A.*, **94**, 310–315.
- ROUSH, E.D. & KWATRA, M.M. (1998). *FEBS Lett.*, **428**, 291–294.
- SARAU, H.M. *et al.* (1997). *J. Pharmacol. Exp. Ther.*, **281**, 1303–1311.
- ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Trace amine-associated

Overview: Trace amine-associated receptors (provisional nomenclature) were initially discovered as a result of a search for novel 5-HT receptors (Borowsky *et al.*, 2001), where 15 mammalian orthologues were identified and divided into two families.

Nomenclature	TA ₁	TA ₂
Other names	TAA1, TaR-1, BO111	TAA2, GPR58
Ensembl ID	ENSG00000146399	ENSG00000146378
Principal transduction	G _s	G _s
Potency order	Tyramine ≥ PEA > octopamine = dopamine (Borowsky <i>et al.</i> , 2001)	PEA > tryptamine (Borowsky <i>et al.</i> , 2001)
Probes	[³ H]-Tyramine (20 nM, Borowsky <i>et al.</i> , 2001)	—

TAA₃ (BO107) and TAA₄ are pseudogenes. The signalling characteristics and pharmacology of TAA₅ (PNR, Putative Neurotransmitter Receptor, ENSG00000135569), TAA₆ (Trace amine receptor 4, TaR-4, ENSG00000146383), TAA₈ (Trace amine receptor 5, GPR102, ENSG00000146385) and TAA₉ (trace amine associated receptor 9, ENSG00000188604) are lacking.

Abbreviations: PEA, 2-phenylethylamine

Further Reading:

BRANCHEK, T.A. & BLACKBURN, T.P. (2003). Trace amine receptors as targets for novel therapeutics: legend, myth and fact. *Curr. Opin. Pharmacol.*, **3**, 90–97.

BURCHETT, S.A. & HICKS, T.P. (2006). The mysterious trace amines: protean neuromodulators of synaptic transmission in mammalian brain. *Prog. Neurobiol.*, **79**, 223–246.

DAVENPORT, A.P. (2003). Peptide and trace amine orphan receptors: prospects for new therapeutic targets. *Curr. Opin. Pharmacol.*, **3**, 127–134.

LIBERLES, S.D. & BUCK, L.B. (2006). A second class of chemosensory receptors in the olfactory epithelium. *Nature*, **442**, 645–650.

LINDEMANN, L. & HOENER, M.C. (2005). A renaissance in trace amines inspired by a novel GPCR family. *Trends Pharmacol. Sci.*, **26**, 274–281.

SOTNIKOVA, T.D., BUDYGIN, E.A., JONES, S.R., DYKSTRA, L.A., CARON, M.G. & GAINETDINOV, R.R. (2004). Dopamine transporter-dependent and -independent actions of trace amine β -phenylethylamine. *J. Neurochem.*, **91**, 362–373.

VANTI, W.B., MUGLIA, P., NGUYEN, T., CHENG, R., KENNEDY, J.L., GEORGE, S.R. & O'DOWD, B.F. (2003). Discovery of a null mutation in a human trace amine receptor gene. *Genomics*, **82**, 531–536.

References:

BOROWSKY, B. *et al.* (2001). *Proc. Natl. Acad. Sci. U.S.A.*, **98**, 8966–8971.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

TRH

Overview: Thyrotropin-releasing hormone (TRH) receptors (provisional nomenclature) are activated by the endogenous tripeptide TRH (pGlu-His-ProNH₂). TRH and TRH analogues fail to distinguish TRH₁ and TRH₂ receptors (see Sun *et al.*, 2003). [³H]-TRH is able to label both TRH₁ and TRH₂ receptors with *K_d* values of 13 and 9 nM, respectively.

	TRH ₁	TRH ₂
Nomenclature	TRH receptor	—
Other names	—	—
Ensembl ID	ENSG00000163485	ENSMUSG00000039079, ENSRNOG00000012789
Principal transduction	G _q	G _q
Selective antagonists	Midazolam (Drummond <i>et al.</i> , 1989), chlordiazepoxide (Straub <i>et al.</i> , 1990), diazepam	—

The human orthologue of the rodent TRH₂ receptor has yet to be identified.

Abbreviations: MeTRH, pGlu-[N^ε-methyl]His-ProNH₂

Further Reading:

COLSON, A.O. & GERSHENGORN, M.C. (2006). Thyrotropin-releasing hormone analogs. *Mini Rev. Med. Chem.*, **6**, 221–226.

GARCIA, S.I & PIROLA, C.J. (2005). Thyrotropin-releasing hormone in cardiovascular pathophysiology. *Regul. Pept.*, **128**, 239–246.

GARY, K.A., SEVARINO, K.A., YARBROUGH, G.G., PRANGE JR, A.J. & WINOKUR, A. (2003). The thyrotropin-releasing hormone (TRH) hypothesis of homeostatic regulation: implications for TRH-based therapeutics. *J. Pharmacol. Exp. Ther.*, **305**, 410–416.

GERSHENGORN, M.C. & OSMAN, R. (2001). Minireview: insights into G protein-coupled receptor function using molecular models. *Endocrinology*, **142**, 2–10.

SUN, Y., LU, X. & GERSHENGORN, M.C. (2003). Thyrotropin-releasing hormone receptors – similarities and differences. *J. Mol. Endocrinol.*, **30**, 87–97.

References:

DRUMMOND, A.H. *et al.* (1989). *Ann. N.Y. Acad. Sci.*, **553**, 197–204.

STRAUB, R.E. *et al.* (1990). *Proc. Natl. Acad. Sci. U.S.A.*, **87**, 9514–9518.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Urotensin-II

Overview: The urotensin-II (U-II) receptor (UT, nomenclature as agreed by NC-IUPHAR, 2005, Douglas and Ohlstein, 2000a) is activated by the endogenous dodecapeptide U-II, originally isolated from the urophysis, the endocrine organ of the caudal neurosecretory system of teleost fish (Bern *et al.*, 1985). Several structural forms of U-II exist in fish and amphibians. The Goby orthologue was used to identify U-II as the cognate ligand for the predicted receptor encoded by the rat gene *gpr14* (Coulouarn *et al.*, 1998; Liu *et al.*, 1999; Mori *et al.*, 1999; Nothacker *et al.*, 1999). Human U-II (derived from ENSG00000049247), an 11-amino-acid peptide (Coulouarn *et al.*, 1998), retains the cyclohexapeptide sequence of goby U-II that is thought to be important in ligand binding (Kinney *et al.*, 2002; Brkovic *et al.*, 2003). This sequence is also conserved in the deduced amino-acid sequence of rat (14 aa) and mouse (14 aa) U-II, although the *N*-terminal is more divergent from the human sequence (Coulouarn *et al.*, 1999).

Nomenclature	UT
Other names	GPR14, SENR, UR-IIR
Ensembl ID	ENSG00000181408
Principal transduction	G _{q/11}
Selective agonists	[Pen ⁵ U-II-(4–11), U-II-(4–11), U-II (Grieco <i>et al.</i> , 2002), AC7954 (Lehmann <i>et al.</i> , 2005)
Selective antagonists	Urantide (8.3, Patacchini <i>et al.</i> , 2003), SB706375 (7.5–8.0, Douglas <i>et al.</i> , 2005), palosuran (pIC ₅₀ 7.1, Clozel <i>et al.</i> , 2004), SB436811 (6.7, Jin <i>et al.</i> , 2005)
Probes	[¹²⁵ I]-hU-II (0.24 nM, Maguire <i>et al.</i> , 2000)

In human vasculature, human urotensin-II elicits both vasoconstrictor (pD₂ 9.3–10.1, Maguire *et al.*, 2000) and vasodilator (pIC₅₀ 10.3–10.4, Stirrat *et al.*, 2001) responses.

Abbreviations: **AC7954**, 3-(4-chlorophenyl)-3-(2-dimethyl-aminoethyl)-isochroman-1-one HCl; **palosuran**, 1-[2-(4-benzyl-4-hydroxy-piperidin-1-yl)-ethyl]-3-(2-methyl-quinolin-4-yl)-urea sulphate, also known as ACT058362; **[Pen⁵]U-II-(4-11]**, [pencillamine, β,β-dimethylcysteine]⁵U-II-(4-11); **SB436811**; **SB706375**, 2-bromo-4,5-dimethoxy-*N*-[3-(*R*)-1-methyl-pyrrolidin-3-yloxy]-4-trifluoromethyl-phenyl]-benzenesulphonamide HCl; **urantide**, [Pen⁵,DTrp⁷,Orn⁸]hU-II(4-11)

Further Reading:

SEE SPECIAL ISSUE OF PEPTIDES (2004). Fish-derived peptides: from fish to human physiology and diseases. In: eds. Takahashi, K. & Kawachi, H. **25**, 1759–1819.

AFFOLTER, J. & WEBB, D.J. (2001). Urotensin II: a new mediator in cardiopulmonary regulation? *Lancet*, **358**, 774–775.

DAVENPORT, A.P. & MAGUIRE, J.J. (2000). Urotensin II: fish neuropeptide catches orphan receptor. *Trends Pharmacol. Sci.*, **21**, 80–82.

DOUGLAS, S.A., DHANAK, D. & JOHNS, D.G. (2004). From 'gills to pills': urotensin-II as a regulator of mammalian cardiorenal function. *Trends Pharmacol. Sci.*, **25**, 76–85.

DOUGLAS, S.A. & OHLSTEIN, E.H. (2000). Urotensin receptors. In: *The IUPHAR Receptor Compendium of Receptor Characterization and Classification*. London: IUPHAR Media Ltd.

GILBERT, R.E., DOUGLAS, S.A. & KRUM, H. (2004). Urotensin-II as a novel therapeutic target in the clinical management of cardiorenal disease. *Curr. Opin. Investig. Drugs*, **5**, 276–282.

KATUGAMPOLA, S. & DAVENPORT, A. (2003). Emerging roles for orphan G-protein-coupled receptors in the cardiovascular system. *Trends Pharmacol. Sci.*, **24**, 30–35.

MAGUIRE, J.J. & DAVENPORT, A.P. (2002). Is urotensin-II the new endothelin? *Br. J. Pharmacol.*, **137**, 579–588.

NOTHACKER, H.P. & CLARK, S. (2005). From heart to mind. The urotensin II system and its evolving neurophysiological role. *FEBS J.*, **272**, 5694–5702.

ONAN, D., HANNAN, R.D. & THOMAS, W.G. (2004). Urotensin II: the old kid in town. *Trends Endocrinol. Metab.*, **15**, 175–182.

RUSSELL, F.D. (2004). Emerging roles of urotensin-II in cardiovascular disease. *Pharmacol. Ther.*, **103**, 223–243.

RUSSELL, F.D. & MOLENAAR, P. (2004). Cardiovascular actions of human urotensin II—considerations for hypertension. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **369**, 271–273.

ZHU, Y.C., ZHU, Y.Z. & MOORE, P.K. (2006). The role of urotensin II in cardiovascular and renal physiology and diseases. *Br. J. Pharmacol.*, **148**, 884–901.

References:

- BERN, H.A. *et al.* (1985). *Recent Prog. Horm. Res.*, **41**, 533–552.
- BRKOVIC, A. *et al.* (2003). *J. Pharmacol. Exp. Therap.*, **306**, 1200–1209.
- CLOZEL, M. *et al.* (2004). *J. Pharmacol. Exp. Therap.*, **311**, 204–212.
- COULOUARN, Y. *et al.* (1998). *Proc. Natl. Acad. Sci. U.S.A.*, **95**, 15803–15808.
- COULOUARN, Y. *et al.* (1999). *FEBS Lett.*, **457**, 28–32.
- DOUGLAS, S.A. *et al.* (2005). *Br. J. Pharmacol.*, **145**, 620–635.
- FOORD, S.M. *et al.* (2005). *Pharmacol. Rev.*, **57**, 279–288.
- GRIECO, P. *et al.* (2002). *J. Med. Chem.*, **45**, 4391–4394.
- JIN, J. *et al.* (2005). *Bioorg. Med. Chem. Lett.*, **15**, 3229–3232.
- KINNEY, W.A. *et al.* (2002). *Angew. Chem. Int. Ed. Engl.*, **41**, 2940–2944.
- LEHMANN, L. *et al.* (2005). *Bioorg. Med. Chem.*, **13**, 3057–3068.
- LIU, Q.Y. *et al.* (1999). *Biochem. Biophys. Res. Commun.*, **266**, 174–178.
- MAGUIRE, J.J. *et al.* (2000). *Br. J. Pharmacol.*, **131**, 441–446.
- MAGUIRE, J.J. *et al.* (2004). *Peptides*, **25**, 1767–1774.
- MORI, M. *et al.* (1999). *Biochem. Biophys. Res. Commun.*, **265**, 123–129.
- NOTHACKER, H.P. *et al.* (1999). *Nat. Cell Biol.*, **1**, 383–385.
- PATACCHINI, R. *et al.* (2003). *Br. J. Pharmacol.*, **140**, 1155–1158.
- STIRRAT, A. *et al.* (2001). *Am. J. Physiol. -Heart Circ. Physiol.*, **280**, H925–H928.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

Vasopressin & Oxytocin

Overview: Vasopressin (AVP) and oxytocin (OT) receptors (nomenclature as agreed by NC-IUPHAR Subcommittee on vasopressin and oxytocin receptors) are activated by the endogenous cyclic nonapeptides AVP and OT. These peptides are derived from precursors (ENSG00000101200 and ENSG00000101405, respectively), which also produce neurophysins.

Nomenclature	V _{1a}	V _{1b}	V ₂	OT
Ensembl ID	ENSG00000166148	P47901	ENSG00000126895	ENSG00000180914
Principal transduction	G _{q/11}	G _{q/11}	G _s	G _{q/11} , G _{i/o}
Rank order of potency	AVP > OT	AVP > OT	AVP > OT	OT ≥ AVP
Selective agonists	F180, [Phe ² , Orn ⁸]VT	d[D-3-Pal ²]VP, d[Cha ⁴]AVP (Derick <i>et al.</i> , 2002)	d[Val ⁴ , DArg ⁸]VP, OPC51803, VNA932	[Thr ⁴ , Gly ⁷]OT (Elands <i>et al.</i> , 1988)
Selective antagonists	d(CH ₂) ₅ [Tyr(Me) ² , Arg ⁸]VP (9.0), SR49059 (8.9), YM087 (8.2)	SSR149415 (8.4; Griebel <i>et al.</i> , 2002; Serradeil-Le Gal <i>et al.</i> , 2002)	VPA985 (8.9, Albright <i>et al.</i> , 1998), d(CH ₂) ₅ [D-Ile ² , Ile ⁴]AVP (8.4), SR121463A (8.4; Serradeil-Le Gal <i>et al.</i> , 1996), OPC31260 (7.6; Yamamura <i>et al.</i> , 1992), YM087 (8.96)	SSR126768A (9.3; Serradeil-Le Gal <i>et al.</i> , 2004) desGlyNH ₂ -d(CH ₂) ₅ [Tyr(Me) ² , Thr ⁴ , Orn ⁸]OT (8.5), L372662 (8.4),
Probes	[³ H]-AVP, [³ H]-SR49059 (1.5 nM), [³ H]-d(CH ₂) ₅ [Tyr(Me) ² , Arg ⁸]AVP (1.1 nM), [¹²⁵ I]-HO-Phaa, D-Tyr(Me)-Phe-Gln-Asn-Arg-Pro-Arg-NH ₂ (50 pM)	[³ H]-AVP	[³ H]-AVP, [³ H]-desGly-NH ₂ [D-Ile ² , Ile ⁴]AVP (2.8 nM), [³ H]-d[D-Arg ⁸]AVP (0.8 nM), [³ H]-SR121463A (4.1 nM)	[³ H]-OT, [³⁵ S]-Non Peptide OT Antagonist (42 pM; Lemaire <i>et al.</i> , 2002), [¹²⁵ I]-d(CH ₂) ₅ [Tyr(Me) ² , Thr ⁴ , Orn ⁸ , Tyr-NH ₂]OVT (90 pM), [¹¹¹ In]-DOTA-dLVT (4.5 nM; Chini <i>et al.</i> , 2003)

The V₂ receptor exhibits marked species differences, such that many ligands (d(CH₂)₅[D-Ile², Ile⁴]VP and [³H]-desGly-NH₂[D-Ile², Ile⁴]VP) exhibit low affinity at human V₂ receptors (Ala *et al.*, 1997). Similarly, [³H]-d[D-Arg⁸]VP is V₂ selective in the rat, not in the human (Saito *et al.*, 1997). The gene encoding the V₂ receptor is polymorphic in man, underlying nephrogenic diabetes insipidus (Bichet, 1998). YM087 display high affinity for both human V_{1a} and V₂ receptors (Tahara *et al.*, 1998).

Abbreviations: **F180**, Hmp-Phe-Ile-Hgn-Asn-Cys-Pro-Dab(Abu)-Gly-NH₂; [¹¹¹In]-DOTA-dLVT, [¹¹¹In]-DOTA-Lys⁸-deamino-vasotocin; **L372662**, 1-(1-{4-[1-(2-methyl-1-oxidopyridin-3-ylmethyl)piperidin-4-yloxy]-2-methoxybenzoyl}piperidin-4-yl)-1,4-dihydrobenz[d[1,3]oxazin-2-one; **OPC31260**, 5-dimethylamino-1-(4-[2-methylbenzoylamino]benzoyl)-2,3,4,5-tetrahydro-1*H*-benzazepine; **OPC51803**, (5*r*)-2-(1-[2-chloro-4-{1-pyrrolidinyl}benzoyl]-2,3,4,5-tetrahydro-1*H*-1-benzazepin-5-yl)-*N*-isopropylacetamide; [³⁵S]-non-peptide OT antagonist, [³⁵S]-(1-(1-(2-(2,2,2-trifluoroethoxy)-4-(1-methylsulfonyl-4-piperidinyl)oxy)phenylacetyl)-4-piperidinyl)-3,4-dihydro-2(1*H*)-quinolinone; **SR121463A**, 1-(4-Boc-2-methoxybenzenesulfonyl)-5-ethoxy-3-spiro-(4-[2-morpholinoethoxycyclohexane]indol-2-one fumarate; equatorial isomer; **SR49059**, (2*s*)-1-([2*r*,3*s*]-[5-chloro-3-{chlorophenyl}-1-{3,4-dimethoxysulfonyl}-3-hydroxy]-2,3-dihydro-1*H*-indole-2-carbonyl)-pyrrolidine-2-carboxamide; **SSR149415**, (2*S*,4*R*)-1-[5-chloro-1-[(2,4-dimethoxyphenyl)sulfonyl]-3-(2-methoxy-phenyl)-2-oxo-2,3-dihydro-1*H*-indol-3-yl]-4-hydroxy-*N,N*-dimethyl-2-pyrrolidine carboxamide; **SSR126768A**, 4-chloro-3-[(3*R*)-(+)5-chloro-1-(2,4-dimethoxybenzyl)-3-methyl-2-oxo-2,3-dihydro-1*H*-indol-3-yl]-*N*-ethyl-*N*-(3-pyridylmethyl)-benzamide, hydrochloride; **VNA932**, (2-chloro-4-[3-methyl-pyrazol-1-yl]-phenyl)-(5*H*,11*H*)-pyrrolo(2,1-*c*)(1,4)benzodiazepin-10-yl-methanone; **VPA985**, 5-fluoro-2-methyl-*N*-(4-[5*H*-pyrrolo[2,1-*c*][1,4]benzodiazepin-10(11*H*)-ylcarbonyl]-3-chlorophenyl)benzamide; **YM087**, (4'-[(2-methyl-1,4,5,6-tetrahydroimidazo[4,5-*d*][1]benzazepin-6-yl) carbonyl]-2-phenylbenzanilide monohydrochloride)

Further Reading:

- BARBERIS, C. & TRIBOLLET, E. (1996). Vasopressin and oxytocin receptors in the central nervous system. *Crit. Rev. Neurobiol.*, **10**, 119–154.
- FREIDINGER, R.M. & PETTIBONE, D.J. (1997). Small molecule ligands for oxytocin and vasopressin receptors. *Med. Res. Rev.*, **17**, 1–16.
- GIMPL, G. & FAHRENHOLZ, F. (2001). The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.*, **81**, 629–683.
- HIBERT, M., HOFACK, J., TRUMPP-KALLMEYER, S., MOUILLAC, B., CHINI, B., MAHE, E., COTTE, N., JARD, S., MANNING, M. & BARBERIS, C. (1999). Functional architecture of vasopressin/oxytocin receptors. *J. Recept. Signal Transduct. Res.*, **19**, 589–596.
- RUSSELL, S.D. & DEWALD, T. (2003). Vasopressin receptor antagonists: therapeutic potential in the management of acute and chronic heart failure. *Am. J. Cardiovasc. Drugs*, **3**, 13–20.
- THIBONNIER, M. (2003). Vasopressin receptor antagonists in heart failure. *Curr. Opin. Pharmacol.*, **3**, 683–687.
- VERBALIS, J.G. (2002). Vasopressin V₂ receptor antagonists. *J. Mol. Endocrinol.*, **29**, 1–9.
- ZINGG, H.H. & LAPORTE, S.A. (2003). The oxytocin receptor. *Trends Endocrinol. Metab.*, **14**, 222–227.

References:

- ALA, Y. *et al.* (1997). *Eur. J. Pharmacol.*, **331**, 285–293.
- ALBRIGHT, J.D. *et al.* (1998). *J. Med. Chem.*, **41**, 2442–2444.
- BICHET, D.G. (1998). *Am. J. Med.*, **105**, 431–442.
- CHINI, B. *et al.* (2003). *Br J Cancer*, **89**, 930–936.
- DERICK, S. *et al.* (2002). *Endocrinology*, **143**, 4655–4664.
- ELANDS, J. *et al.* (1988). *Am J Physiol*, **254**, E31–E38.
- GRIEBEL, G. *et al.* (2002). *Proc. Natl. Acad. Sci. U.S.A.*, **99**, 6370–6375.
- LEMAIRE, W. *et al.* (2002). *Eur. J. Pharmacol.*, **450**, 19–28.
- SAITO, M. *et al.* (1997). *Biochem. Pharmacol.*, **53**, 1711–1717.
- SERRADEIL-LE GAL, C. *et al.* (1996). *J. Clin. Invest.*, **98**, 2729–2738.
- SERRADEIL-LE GAL, C. *et al.* (2002). *J. Pharmacol. Exp. Ther.*, **300**, 1122–1130.
- SERRADEIL-LE GAL, C. *et al.* (2004). *J. Pharmacol. Exp. Ther.*, **309**, 414–424.

TAHARA, A. *et al.* (1998). *Naunyn Schmiedeberg's Arch. Pharmacol.*, **357**, 63–69.

YAMAMURA, Y. *et al.* (1992). *Br. J. Pharmacol.*, **105**, 787–791.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.

VIP & PACAP

Overview: Vasoactive intestinal peptide (VIP) and pituitary adenylate cyclase-activating peptide (PACAP) receptors (nomenclature recommended by the NC-IUPHAR Subcommittee on Vasoactive Intestinal Peptide Receptors, Harmar *et al.*, 1998) are activated by the endogenous peptides VIP, PACAP_{1–38}, PACAP_{1–27}, peptide histidine isoleucineamide (PHI), peptide histidine methionineamide (PHM), peptide histidine valine and growth hormone-releasing factor (GRF). PACAP type II receptors have been defined as those for which PACAP and VIP display comparable affinity. Both VPAC₁ and VPAC₂ meet this definition. [Arg¹⁶]chicken secretin is an agonist at both VPAC₁ and secretin receptors, but can be used as an agonist at VPAC₁ receptors in tissues that do not express secretin receptors (Gourlet *et al.*, 1997a). PACAP_{6–38} also shows significant affinity for VPAC₂ receptors. Helodermin discriminates VPAC₁ and VPAC₂ in a species-dependent manner (Gourlet *et al.*, 1998).

Nomenclature	VPAC ₁	VPAC ₂	PAC ₁
Other names	VIP ₁ /PACAP, VIP, VIP ₁ , PACAP type II, PVR2	VIP ₂ /PACAP, VIP ₂ , PACAP ₃₈ , PVR2	PACAP, PACAP type I, PVR1
Ensembl ID	ENSG00000114812	ENSG00000106018	ENSG00000078549
Principal transduction	G _s	G _s	G _s
Rank order of potency	VIP, PACAP-(1–27) = PACAP-(1–38) > GRF >> PHI >> secretin	VIP, PACAP-(1–38) > PACAP-(1–27) > PHI >> GRF, secretin	PACAP-(1–27), PACAP-(1–38) >> VIP > PHI
Selective agonists	[Arg ¹⁶]chicken secretin, [Lys ¹⁵ , Arg ¹⁶ , Leu ²⁷]VIP-(1–7)-GRF-(8–27)-NH ₂	Ro251553 (Gourlet <i>et al.</i> , 1997a, b), Ro251392 (Xia <i>et al.</i> , 1997)	Maxadilan (Moro and Lerner, 1997)
Selective antagonists	[Ac-His ¹ , D-Phe ² , Lys ¹⁵ , Arg ¹⁶]VIP-(3–7)-GRF-(8–27)-NH ₂ (Gourlet <i>et al.</i> , 1997a)	—	PACAP-(6–38)
Probes	[¹²⁵ I]-VIP, [¹²⁵ I]-PACAP	[¹²⁵ I]-VIP, [¹²⁵ I]-PACAP	[¹²⁵ I]-PACAP

Subtypes of PAC₁ receptors have been proposed based on tissue differences in the potencies of PACAP_{1–27} and PACAP_{1–38}; these might result from differences in G-protein coupling and second messenger mechanisms (Van Ramplebergh *et al.*, 1996), or from alternative splicing of PAC₁ receptor mRNA (Spengler *et al.*, 1993).

Abbreviations: Ro251392, Ac-His¹[Glu⁸, OCH₃-Tyr¹⁰, Lys¹², Nle¹⁷, Ala¹⁹, Asp²⁵, Leu²⁶, Lys^{27,28}]VIP (*cyclo* 21–25); Ro251553, Ac-His¹[Glu⁸, Lys¹², Nle¹⁷, Ala¹⁹, Asp²⁵, Leu²⁶, Lys^{27,28}, Gly^{29,30}, Thr³¹]VIP-NH₂ (*cyclo* 21–25)

Further Reading:

DELGADO, M., POZO, D. & GANEA, D. (2004). The significance of vasoactive intestinal peptide in immunomodulation. *Pharmacol. Rev.*, **56**, 249–290.

GRONEBERG, D.A., RABE, K.F. & FISCHER, A. (2006). Novel concepts of neuropeptide-based drug therapy: vasoactive intestinal polypeptide and its receptors. *Eur. J. Pharmacol.*, **533**, 182–194.

HARMAR, A.J., ARIMURA, A., GOZES, I., JOURNOT, L., LABURTHER, M., PISEGNA, J.R., RAWLINGS, S.R., ROBBERECHT, P., SAID, S.I., SREEDHARAN, S.P., WANK, S.A. & WASCHEK, J.A. (1998). International Union of Pharmacology. XVIII. Nomenclature of receptors for vasoactive intestinal peptide and pituitary adenylate cyclase-activating polypeptide. *Pharmacol. Rev.*, **50**, 265–270.

LABURTHER, M. & COUVINEAU, A. (2002). Molecular pharmacology and structure of VPAC Receptors for VIP and PACAP. *Regul. Pept.*, **108**, 165–173.

LABURTHER, M., COUVINEAU, A. & MARIE, J.C. (2002). VPAC receptors for VIP and PACAP. *Receptors Channels*, **8**, 137–153.

References:

- GOURLET, P. *et al.* (1997a). *Peptides*, **18**, 1539–1545.
 GOURLET, P. *et al.* (1997b). *Peptides*, **18**, 403–408.
 GOURLET, P. *et al.* (1998). *Ann. NY. Acad. Sci.*, **865**, 247–252.
 MORO, O. & LERNER, E.A. (1997). *J. Biol. Chem.*, **272**, 966–970.
 SPENGLER, D. *et al.* (1993). *Nature*, **365**, 170–175.
 VAN RAMPLEBERGH, J. *et al.* (1996). *Mol. Pharmacol.*, **50**, 1596–1604.
 XIA, M. *et al.* (1997). *J. Pharmacol. Exp. Ther.*, **281**, 629–633.

ALEXANDER, S.P.H., MATHIE, A. & PETERS, J.A. (2007). Guide to Receptors and Channels (GRAC), 2nd edition (2007 revision). *Br. J. Pharmacol.*, **150** (Suppl. 1), S1–S168.